
Data

Spring 2016

A Hydrodynamic-Sediment Transport Numerical Model for the Waipaoa Shelf, New Zealand: Model Archive

Julia Moriarty

Virginia Institute of Marine Science, moriarty@vims.edu

Courtney K. Harris

Virginia Institute of Marine Science, ckharris@vims.edu

Mark G. Hadfield

National Institute of Water and Atmospheric Research, mark.hadfield@niwa.co.nz

Follow this and additional works at: <https://scholarworks.wm.edu/data>



Part of the [Geology Commons](#), [Oceanography Commons](#), and the [Sedimentology Commons](#)

Recommended Citation

Moriarty, J.; Harris, C. K.; and Hadfield, M. G., A Hydrodynamic-Sediment Transport Numerical Model for the Waipaoa Shelf, New Zealand: Model Archive. 2016. Virginia Institute of Marine Science, College of William & Mary. <http://doi.org/10.21220/V5159T>

This Data is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Data by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

README

Item Title: A Hydrodynamic-Sediment Transport Numerical Model for the Waipaoa Shelf, New Zealand: Model Archive

Authors: Julia Moriarty¹, Courtney Harris¹, Mark G. Hadfield²

¹Virginia Institute of Marine Science, The College of William & Mary, USA

²National Institute of Water and Atmospheric Research, New Zealand

Author Contact Information: Julia Moriarty (moriarty@vims.edu), Courtney Harris (ckharris@vims.edu)

Date of Issue: 2016.

Keywords: Waipaoa River continental shelf, New Zealand; sediment transport; numerical modeling; oceanography; marine geology; Regional Ocean Modeling System (ROMS).

Description: These files are compressed versions of input files, model code, and output used for two publications:

Moriarty, J. M., Harris, C. K., and Hadfield, M. G. (2014). A hydrodynamic and sediment transport model for the Waipaoa Shelf, New Zealand: Sensitivity of fluxes to spatially-varying erodibility and model nesting. *Journal of Marine Science and Engineering*, 2 (2): 336-369. Doi:10.3390/jmse2020336

Moriarty, J. M., Harris, C. K., and Hadfield, M. G. (2015). Event-to-seasonal sediment dispersal on the Waipaoa River Shelf, New Zealand: A numerical modeling study. *Continental Shelf Research*, 110: 108-123. Doi: 10.1016/j.csr2015.10.005

Compressed files with the .gz file extension can be opened with Gzip GNU software (open source). Compressed files with the .tar file extension can be opened with Gzip Tar software (open source). Many of the input / output files use the NetCDF (Network Common Data Form) file format. These have "nc" as a file extension and can be read using a variety of open source tools: see <http://www.unidata.ucar.edu/software/netcdf/docs/> . For information about the Regional Ocean Modeling System (ROMS), its model code and input / output, see www.myroms.org .

Files include:

File	Description
Input Files	
gridgen_romsformat_redo5.nc.gz	Input File - Model Grid
waipaoa_init_redo7_20100101_new2.nc.gz	Input File – Model Initialization
b_2009_2010.nc.gz.tar	Input File – Data for Open boundary conditions
c_2009_2010.nc.gz.tar	Input File - Data for Model Nudging (i.e. “climatology files”)
waipaoa_swradhr_new.nc.gz waipaoa_windhr_nzlam_new.nc.gz waipaoa_apresshr_new.nc.gz waipaoa_cloud_new.nc.gz waipaoa_rainhr_new.nc.gz waipaoa_relhumhr_new.nc.gz waipaoa_temphr_new.nc.gz	Input File - Atmospheric Forcing
waipaoa_waves3_new.nc	Input File - Wave Forcing
waipaoa_river_newpartitions.nc.gz	Input File - River forcing
pb_tide_PO_new.nc.gz	Input File - Tidal Forcing
sediment_waipaoa.in.gz	Input File - Sediment Information
ocean_waipaoa.in.gz	Input File – Model Run Information
waipaoa.h.gz	Input File – Options for Model Compilation
Model Code	
build.bash.gz	Model Code - Script to Compile Model
trunk4.tar	Model Code - Model Code
Model Output	
*Note that for sensitivity tests, altered input files are also included with the model output	
model_results_standard_part1.gz.tar model_results_standard_part2.gz.tar	Model Output - Standard Model
model_inputs_results_ECM_lowM.gz.tar	Model Output - Low erodibility (i.e. “Low M”) sensitivity test
model_inputs_results_ECM_highM.gz.tar	Model Output - High erodibility (i.e. “High M”) sensitivity test
model_inputs_results_ECM_nesting2.gz.tar	Model Output - “Weakly Nudged” Sensitivity Test
model_inputs_results_ECM_nesting1.gz.tar	Model Output - “Moderately Nudged” Sensitivity Test

Sponsors: Funding was provided by NSF MARGINS grant 0841092 (Moriarty and Harris), a VIMS student fellowship (Moriarty), and NIWA as part of its government-

funded, core research (Hadfield). Many datasets were useful during development and testing of the model, and these were provided by personnel from NIWA (M. Uddstrom, S. Stephens, A. Orpin), VIMS (S. Kuehl, T. Kniskern), the USACoE (J. McNinch), GDC (G. Hall, D. Peacock), East Carolina University (J.P. Walsh, R. Corbett, J. Kiker), and UW (A. Ogston, R. Hale). Thank you to persons who provided technical assistance, including A. Bever (now at Delta Modeling Associates), A. Miller, M.A. Bynum, and D. Weiss (all from VIMS/William & Mary). Computational facilities at VIMS, the SciClone cluster at the College of William & Mary, and the CSDMS computing cluster at the University of Colorado were supported by the NSF, VA Port Authority, Sun Microsystems, and Virginia's Commonwealth Technology Research Fund. High-performance computing facilities at NIWA were supported by the NZ eScience Infrastructure (NESI) and funded by NESI's collaborator institutions and through the Ministry of Business, Innovation & Employment's Research Infrastructure programme.