

3-6-2018

Three doublet lepton-specific model

Marco Merchand

College of William and Mary, mamerchandmedi@email.wm.edu

Marc Sher

College of William and Mary, mtsher@wm.edu

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Merchand, Marco and Sher, Marc, Three doublet lepton-specific model (2018). *PHYSICAL REVIEW D*, 95(5).

10.1103/PhysRevD.95.055004

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



CHORUS

This is the accepted manuscript made available via CHORUS. The article has been published as:

Three doublet lepton-specific model

Marco Merchand and Marc Sher

Phys. Rev. D **95**, 055004 — Published 6 March 2017

DOI: [10.1103/PhysRevD.95.055004](https://doi.org/10.1103/PhysRevD.95.055004)

A Three Doublet Lepton-Specific Model

Marco Merchand* and Marc Sher†

High Energy Theory Group, College of William and Mary, Williamsburg, Virginia 23187, U.S.A.

In the lepton-specific version of two Higgs doublet models, a discrete symmetry is used to couple one Higgs, Φ_2 , to quarks and the other, Φ_1 , to leptons. The symmetry eliminates tree level flavor changing neutral currents (FCNC). Motivated by strong constraints on such currents in the quark sector from meson-antimeson mixing, and by hints of $h \rightarrow \mu\tau$ in the lepton sector, we study a simple three Higgs doublet model in which one doublet couples to quarks and the other two to leptons. Unlike most other studies of three Higgs doublet models, we impose no flavor symmetry and just use a Z_2 symmetry to constrain the Yukawa couplings. We present the model and discuss the various mixing angles. Constraining the parameters to be consistent with observations of the Higgs boson at the LHC, we study the properties of the charged Higgs boson(s) in the model, focusing on the case in which the charged Higgs is above the top threshold. It is found that one can have the branching fraction of the charged Higgs into $\tau\nu_\tau$ comparable to $t\bar{b}$ decay without needing very large values for the ratios of vevs. One can also get a large branching fraction for the much more easily observable $\mu\nu_\tau$ decay.

I. INTRODUCTION

In 2012, the discovery of the Higgs boson [1, 2] was announced. To date, the properties of the Higgs have not significantly deviated from the expectations of the Standard Model. Nonetheless, many expect new physics to be found at the TeV scale. The Standard Model has a metastable vacuum, a hierarchy problem and no dark matter candidate. In most TeV scale extensions of the Standard Model, including supersymmetric models [3], composite Higgs models [4], twin Higgs models [5], and left-right models [6], there are additional Higgs fields. Thus, considerations of extensions of the Higgs sector are well motivated.

The most studied extension of the Higgs sector is the Two Higgs Doublet Model (2HDM); see Ref. [7] for an extensive review. As soon as one extends the Higgs sector, tree level flavor-changing neutral currents (FCNC) become an issue [8, 9]. These can be avoided by imposing a Z_2 discrete symmetry such that all fermions of a given charge couple to the same multiplet. In the 2HDM, there are four models generally discussed. In the type I model, all fermions couple to one Higgs (generally taken to be Φ_2), and in the type II model, the $Q=2/3$ quarks couple to Φ_2 and the $Q=-1/3$ quarks and leptons couple to Φ_1 . In the other two models, the lepton specific and flipped models, the leptons couple to the other Higgs. For the lepton specific model, in particular, the quarks all couple to Φ_2 and the leptons all couple to Φ_1 .

There are hundreds of papers on these four models. However, any discovery of a tree level FCNC would invalidate them. We are intrigued by the recent hints [10] of a nonzero branching ratio of the light Higgs into $\mu\tau$ with 2.4σ significance, with a branching ratio of $0.84 \pm 0.38\%$. More recent results have not seen a signal, but the errors are large, with Ref. [11] giving a branching ratio of $0.76 \pm 0.82\%$ and Ref. [12] giving a branching ratio of $0.53 \pm 0.51\%$. If such a signal is confirmed, then the conventional 2HDMs will be excluded and models with a different structure in the lepton sector will be favored. In the (more likely) event that it is not confirmed, it is still the case that bounds on scalar mediated FCNC in the quark sector are much stronger (due to meson-antimeson mixing) than those in the lepton sector, and thus we wish to study extensions of the Higgs sector with FCNC in the lepton sector but not in the quark sector.

There have been some studies of alternative 2HDMs that can address lepton flavor-violation. For example, Altmanshofer, et al. [13], Ghosh, et al. [14] and Botella et al. [15] consider models in which the Standard Model Higgs couples to the third generation and the other Higgs (which may be composite) to the other two, and they study higher-dimensional operators and the effect on the low energy theory.

Since leptons must transform differently than quarks in this case, the simplest model is to extend the lepton specific model by adding a third doublet, Φ_3 , which behaves under the Z_2 symmetry exactly as Φ_1 . There will then be only one doublet coupling to quarks and the other two to leptons. The three doublet model has two physical pairs of charged Higgs fields, two pseudoscalars and three neutral scalars.

*mamerchandmedi@email.wm.edu

†mtsher@wm.edu

Three Higgs doublet models (3HDMs) have been studied previously. They allow for a very rich structure of discrete symmetry groups in the Higgs and flavor sectors. A detailed analysis of these symmetries has been carried out by Ivanov and collaborators, both for discrete symmetries [16, 17] and Abelian symmetries [18] and Keus, King and Moretti [19] focused on flavor symmetries and analyzed Higgs masses and potentials in several models. A specific model with an S_3 symmetry was considered in Refs. [20, 21] and one with an A_4 symmetry, that also discusses $h \rightarrow \mu\tau$ is in Ref. [22]. Moretti and Yagyu [23] studied perturbative unitarity in 3HDMs in which some of the doublets are inert. More phenomenological analyses were carried out by Aranda, et al. [24, 25] in the context of a 3HDM with a \mathbb{Z}_5 symmetry. A model with a global $U(1)$ flavor symmetry by Crivellin, D'Ambrosio and Heeck [26] not only can account for $h \rightarrow \mu\tau$ but can also resolve anomalies in $B \rightarrow K\mu\mu$. The possibility of a light charged Higgs and constraints from B physics in 3HDMs was considered by Akeroyd, et al. [27]. These authors, and a subsequent article by Yagyu [28] recently considered 3HDMs with no tree level FCNC and studied Higgs boson couplings; these works include a new ‘‘Model Z’’ in which the three doublets couple to the up-type quarks, down-type quarks and leptons, respectively [27]. A nice review can be found in Yagyu’s recent talk [29].

Our model differs from these in that we do have tree level FCNC, albeit only in the lepton sector, the discrete symmetry is a simple Z_2 and we will focus on charged Higgs masses above the top quark threshold. In the lepton specific 2HDM, one can only have $\tau\nu$ charged Higgs decays dominate above the top quark threshold if there is a very large ratio of vacuum expectation values and we wish to see if the requirement that there be such a large ratio still holds in the 3HDM. It will also be noted that the isospin counterpart of $h \rightarrow \mu\tau$ would be $H^+ \rightarrow \mu\nu_\tau$, leading to a substantial and observable enhancement of muonic decays of the charged Higgs.

In the next section, we present the model, scalar mass matrices and mixing angles. In Section 3, we study the charged Higgs bosons, including bounds on the parameters from LHC observations of the light Higgs and focusing on the $\tau\nu$ decay mode which, we will show, can dominate the decays even above the top threshold and in Section 4 we present our conclusions.

II. THE MODEL

The model consists of three Higgs doublets with weak hypercharge $Y = 1/2$. Since FCNC in the quark sector are very small, and we hope to allow tree level FCNC in the lepton sector, quarks and leptons must transform differently under the Z_2 symmetry. In the conventional 2HDMs, one model does treat quarks and leptons differently. This is the lepton specific model (sometimes referred to as the Type X model). In this model, the only fields odd under the Z_2 are Φ_1 and the right-handed leptons, e_R^i . This then forces Φ_1 to couple only to leptons and Φ_2 only to quarks. In this simple extension, we introduce a third doublet, Φ_3 , which is odd under the Z_2 . Thus Φ_2 only couples to quarks, whereas Φ_1 and Φ_3 only couple to leptons. It is the fact that two doublets couple to leptons that will lead to FCNC in the lepton sector.

A. Mass matrices and mixing angles

The scalar potential consistent with the Z_2 symmetry (under which Φ_1, Φ_3 and e_R^i are odd) can be written

$$\begin{aligned}
V = & m_{11}^2 \Phi_1^\dagger \Phi_1 + m_{22}^2 \Phi_2^\dagger \Phi_2 + m_{33}^2 \Phi_3^\dagger \Phi_3 - m_{13}^2 (\Phi_1^\dagger \Phi_3 + \Phi_3^\dagger \Phi_1) + \frac{1}{2} \lambda_{11} (\Phi_1^\dagger \Phi_1)^2 \\
& + \frac{1}{2} \lambda_{22} (\Phi_2^\dagger \Phi_2)^2 + \frac{1}{2} \lambda_{33} (\Phi_3^\dagger \Phi_3)^2 + \lambda_{12} \Phi_1^\dagger \Phi_1 \Phi_2^\dagger \Phi_2 + \lambda_{13} \Phi_1^\dagger \Phi_1 \Phi_3^\dagger \Phi_3 \\
& + \lambda_{23} \Phi_2^\dagger \Phi_2 \Phi_3^\dagger \Phi_3 + \frac{\beta_{12}}{2} \left[(\Phi_1^\dagger \Phi_2)^2 + (\Phi_2^\dagger \Phi_1)^2 \right] + \alpha_{12} \Phi_1^\dagger \Phi_2 \Phi_2^\dagger \Phi_1 \\
& + \frac{\beta_{13}}{2} \left[(\Phi_1^\dagger \Phi_3)^2 + (\Phi_3^\dagger \Phi_1)^2 \right] + \alpha_{13} \Phi_1^\dagger \Phi_3 \Phi_3^\dagger \Phi_1 \\
& + \frac{\beta_{23}}{2} \left[(\Phi_2^\dagger \Phi_3)^2 + (\Phi_3^\dagger \Phi_2)^2 \right] + \alpha_{23} \Phi_2^\dagger \Phi_3 \Phi_3^\dagger \Phi_2.
\end{aligned} \tag{1}$$

In this expression, we have omitted quartic terms odd in either Φ_1 or Φ_3 , such as $\Phi_1^\dagger \Phi_1 \Phi_1^\dagger \Phi_3$. This is entirely done for simplicity, and will avoid needing to solve cubic equations. This simplification will only affect expressions which explicitly have the λ_{ij} , α_{ij} and β_{ij} . The phenomenology only depends on the mixing angles and not on these couplings (which realistically can’t be measured for decades). As a result, nothing more will be learned by adding these extra terms.

We choose parameters such that each doublet acquires a vacuum expectation value (vev). Some models choose parameters so that one or two of the doublets get zero vev – these models are inert models and have a dark matter candidate. Expanding around the minimum we can write

$$\Phi_a = \begin{pmatrix} \phi_a^+ \\ \frac{v_a + \rho_a + i\eta_a}{\sqrt{2}} \end{pmatrix}, \quad a = 1, 2, 3. \quad (2)$$

With this potential and our choice of minimum, one can find the mass matrices. This is done in Appendix A. The charged Higgs, pseudoscalar and scalar mass matrices are all 3×3 . The charged and pseudoscalar mass matrices have a zero eigenvalue, corresponding to the Goldstone bosons.

Before diagonalizing the mass matrices, it is instructive to recall the diagonalization in the more well-known 2HDMs [7]. The minimum of the potential in that model lies along the ray (v_1, v_2) . The mass matrices for the charged scalars and pseudoscalars have a zero eigenvalue, corresponding to the Goldstone bosons. The mass matrix for the scalars does not have a zero eigenvalue, and one eigenvalue is the 125 GeV Higgs boson. If one rotates the basis by a rotation angle given by $\beta \equiv \tan^{-1}(v_2/v_1)$ (which is equivalent to defining a new “x-axis” along the ray), then one has the Higgs basis, in which only one field gets a vev. After this rotation, the zero eigenvalues separate out and decouple. Note that there is only a single rotation angle for both the charged scalar and pseudoscalar matrices, which is not surprising. Note also that in the 2HDM, the charged scalar and pseudoscalar matrices commute, so they are simultaneously diagonalizable.

In the 3HDM, as can be seen from the mass matrices, the charged scalar and pseudoscalar matrices do not commute, and thus they will not be simultaneously diagonalizable. Thus, in general, we will have nine different angles (three for each matrix) that diagonalize the ϕ^+ , η (pseudoscalar) and the ρ (scalar) matrices. It turns out that the charged and the pseudoscalar mass matrices have two common angles which diagonalize the matrices, leaving us with a total of seven different angles. The fact that two of the angles are common is expected. Two angles are needed to take the general vector (v_1, v_2, v_3) to $(v, 0, 0)$, and this will in both cases lead to separating out the Goldstone mode. However, there is still another angle corresponding to an additional rotation about the $(v, 0, 0)$ axis, and there is no reason that this angle should be the same for the charged and pseudoscalar cases.

The first rotation matrix that partially diagonalizes the ϕ^+ and the η matrices is the rotation matrix that takes the ray (v_1, v_2, v_3) to $(v, 0, 0)$. The first transformation is given by (where $v_{ij}^2 \equiv v_i^2 + v_j^2$)

$$R = \begin{pmatrix} \frac{v_1}{v} & \frac{v_2}{v} & \frac{v_3}{v} \\ -\frac{v_2}{v_{12}} & \frac{v_1}{v_{12}} & 0 \\ -\frac{v_1 v_3}{v v_{12}} & -\frac{v_2 v_3}{v v_{12}} & \frac{v_{12}}{v} \end{pmatrix}. \quad (3)$$

This is equivalent to rotating the vector (v_1, v_2, v_3) clockwise around the z-axis by angle ψ and then rotate counter-clockwise around the y-axis by angle $\pi/2 - \theta$. The azimuthal and the polar angles are given in terms of the vev's by

$$\tan \theta = \frac{v_{12}}{v_3}, \quad \tan \psi = \frac{v_2}{v_1}. \quad (4)$$

The third rotation, which is the angle of rotation around the x-axis will be called β_1 and β_2 for the ϕ^+ and the η matrices respectively.

For the scalar mass-squared matrix we denote the Euler angles (in the above convention) that diagonalize it by β_3 , θ_3 and ψ_3 , where we have replaced the standard notation of performing a rotation by angle ϕ first, around the z-axis, by β_3 instead, in order to avoid confusion with the fields.

In summary for the charged and the pseudoscalar matrices we perform the following transformation on the fields

$$T_i = \begin{pmatrix} \cos \psi & -\sin \psi & 0 \\ \sin \psi & \cos \psi & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \cos(\pi/2 - \theta) & 0 & -\sin(\pi/2 - \theta) \\ 0 & 1 & 0 \\ \sin(\pi/2 - \theta) & 0 & \cos(\pi/2 - \theta) \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \beta_i & -\sin \beta_i \\ 0 & \sin \beta_i & \cos \beta_i \end{pmatrix}, \quad (5)$$

where $i = 1, 2$ and the angles θ and ψ are given in (4).

For the scalar we write the transformation as the product of three Euler rotations, namely

$$T_3 = \begin{pmatrix} \cos \psi_3 & -\sin \psi_3 & 0 \\ \sin \psi_3 & \cos \psi_3 & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} \cos \theta_3 & 0 & -\sin \theta_3 \\ 0 & 1 & 0 \\ \sin \theta_3 & 0 & \cos \theta_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \beta_3 & -\sin \beta_3 \\ 0 & \sin \beta_3 & \cos \beta_3 \end{pmatrix}. \quad (6)$$

The field redefinitions are then given by

$$\begin{pmatrix} \phi_1^+ \\ \phi_2^+ \\ \phi_3^+ \end{pmatrix} = T_1 \begin{pmatrix} G^+ \\ H_1^+ \\ H_2^+ \end{pmatrix}, \quad \begin{pmatrix} \eta_1 \\ \eta_2 \\ \eta_3 \end{pmatrix} = T_2 \begin{pmatrix} G^0 \\ A_1 \\ A_2 \end{pmatrix}, \quad \begin{pmatrix} \rho_1 \\ \rho_2 \\ \rho_3 \end{pmatrix} = T_3 \begin{pmatrix} h_1 \\ h_2 \\ h_3 \end{pmatrix}. \quad (7)$$

The spectrum of the theory consists of two charged scalars, two pseudo scalars and three neutral scalars.

B. Gauge boson couplings

The kinetic term of the Higgs doublets has the form

$$(D_\mu \Phi_i)^\dagger D_\mu \Phi_i, \quad (8)$$

where

$$D_\mu = \partial_\mu - igW_\mu^a \tau^a - i\frac{g'}{2}B_\mu \quad (9)$$

is the covariant derivative and W_μ^a and B_μ are the $SU(2)_L$ and $U(1)_Y$ gauge bosons of the electroweak sector respectively. Expanding out in terms of the mass terms given by eq (7) the Lagrangian contains

$$\mathcal{L} \supseteq -\frac{ig}{2} \sec \theta_w (\sin 2\theta_w A^\mu + \cos 2\theta_w Z^\mu) (\partial_\mu H_{a-1}^+ H_{a-1}^- - H_{a-1}^+ \partial_\mu H_{a-1}^-) \quad (10)$$

$$-\frac{ig}{2} (T_1^T \cdot T_3)_{ab} (W_\mu^- (H_{a-1}^+ \partial_\mu h_b - \partial_\mu H_{a-1}^+ h_b) + W_\mu^+ (\partial_\mu H_{a-1}^- h_b - H_{a-1}^- \partial_\mu h_b)) \quad (11)$$

$$+\frac{g^2}{4} (2W_\mu^+ W_\mu^- + \sec^2 \theta_w (Z_\mu \cos 2\theta_w + A_\mu \sin 2\theta_w)^2) H_{a-1}^- H_{a-1}^+ \quad (12)$$

$$+\frac{g^2}{4} (2W_\mu^- W_\mu^+ + Z_\mu^2 \sec^2 \theta_w) (T_3)_{ab} v_a h_b \quad (13)$$

where sum over $a, b = 1, 2, 3$ is implied and $H_0^+ = G^+$ is the Goldstone boson. From this expression we can read off the couplings of the Higgs particles with the gauge bosons. A list of the relevant couplings to gauge bosons are given at the end of the section.

C. The quark sector

As in the lepton-specific model, the RH quarks will couple to Φ_2 and the RH leptons to Φ_1 and Φ_3 . Thus the Yukawa terms in the Lagrangian are

$$-\mathcal{L}_{Yuk} = Y_{ij}^u \bar{Q}^i \tilde{\Phi}_2 u_R^j + Y_{ij}^d \bar{Q}^i \Phi_2 d_R^j + \eta_{ij}^1 \bar{L}^i \Phi_1 e_R^j + \eta_{ij}^2 \bar{L}^i \Phi_3 e_R^j + h.c. \quad (14)$$

where

$$\tilde{\Phi}_2 \equiv i\sigma_2 \Phi_2^*, \quad (15)$$

and

$$Q^i = \begin{pmatrix} u_L^i \\ d_L^i \end{pmatrix}, \quad L_i = \begin{pmatrix} \nu_L^i \\ e_L^i \end{pmatrix}. \quad (16)$$

Here $i = 1, 2, 3$ are the generation indices.

Quarks only couple to the Φ_2 doublet. The Lagrangian of the quark sector is given by

$$\mathcal{L}_{quark} = \bar{Q} \tilde{\Phi}_2 Y^u u_R + \bar{Q} \Phi_2 Y^d d_R. \quad (17)$$

Using equation (2) and expanding out we can write this as

$$\mathcal{L}_{quark} = \frac{\rho_2}{\sqrt{2}} (\bar{u}_L Y^u u_R + \bar{d}_L Y^d d_R) - \frac{i\eta_2}{\sqrt{2}} (\bar{u}_L Y^u u_R - \bar{d}_L Y^d d_R) \quad (18)$$

$$- \phi_2^- \bar{d}_L Y^u u_R + \phi_2^+ \bar{u}_L Y^d d_R + h.c. \quad (19)$$

This is the same as the conventional lepton-specific model, and one can thus immediately write, as in that model,

$$\begin{aligned} \mathcal{L}_{quark} = & \frac{\rho_2}{v_2} (m^u \bar{u}u + m^d \bar{d}d) - \frac{\eta_2}{v_2} (m^u \bar{u}i\gamma_5 u - m^d \bar{d}i\gamma_5 d) \\ & + \left(-\frac{\sqrt{2}}{v_2} \phi_2^- \bar{d}_L V_{CKM}^\dagger m^u u_R + \frac{\sqrt{2}}{v_2} \phi_2^+ \bar{u}_L V_{CKM} m^d d_R + h.c. \right), \end{aligned} \quad (20)$$

m^u and m^d are the entries of the diagonal matrices $\frac{v_2}{\sqrt{2}} M_u$ and $\frac{v_2}{\sqrt{2}} M_d$ respectively.

Expressing this in terms of the physical fields is somewhat more complicated, since there are two charged Higgs, two pseudoscalars and three scalars. In general, one can write

$$\begin{aligned} \mathcal{L}_{quark} = & - \sum_{i=1}^3 \sum_{f=u,d} \frac{m^f}{v} \left(\xi_{h_i}^f \bar{f} f h_i - i \xi_{A_{i-1}}^f \bar{f} \gamma_5 f A_{i-1} \right) \\ & + \left\{ \sum_{i=1}^3 \frac{\sqrt{2} V_{ud}}{v} \bar{u} \left(\xi_{H_{i-1}^+}^u m^u P_L + m^d \xi_{H_{i-1}^+}^d P_R \right) d H_{i-1}^+ + h.c. \right\} \end{aligned} \quad (21)$$

with $A_0 = G^0$, $H_0^+ = G^+$ being the Goldstone bosons. The couplings are given in terms of the matrix elements of the rotations given by equation (7).

D. The Higgs basis and the lepton sector

The lepton sector of the lepton specific 3HDM has the following terms, written in matrix form (with generation indices understood)

$$\mathcal{L}_{Yukawa} \supseteq -\bar{L} (\Phi_1 \eta^1 + \Phi_3 \eta^2) e_R + h.c. \quad (22)$$

where η^1 and η^2 are general complex Yukawa matrices.

We would like to go to the Higgs basis where we make a rotation $(\Phi_1, \Phi_3) \rightarrow (H_1, H_3)$ such that H_1 has zero vev and $\langle H_3 \rangle = v_{13}/\sqrt{2}$. This can be accomplished by performing the field redefinition

$$\begin{pmatrix} \Phi_1 \\ \Phi_3 \end{pmatrix} = \frac{1}{v_{13}} \begin{pmatrix} v_3 & v_1 \\ -v_1 & v_3 \end{pmatrix} \begin{pmatrix} H_1 \\ H_3 \end{pmatrix}. \quad (23)$$

Following the same notation as in [7], we define the matrices

$$N = \frac{1}{\sqrt{2}} (v_3 \eta^1 - v_1 \eta^2), \quad (24)$$

$$M = \frac{1}{\sqrt{2}} (v_1 \eta^1 + v_3 \eta^2). \quad (25)$$

So the Lagrangian of the lepton sector, in the Higgs basis, reads

$$\mathcal{L}_{lepton} = -\frac{\sqrt{2}}{v_{13}} \bar{L} (H_1 N + H_3 M) e_R + h.c. \quad (26)$$

In the Higgs basis, only the Yukawa couplings of the doublet H_3 generate fermion masses, and may be bi-diagonalized so they do not lead to tree-level FCNC.

When passing to the mass basis of the leptons in which the mass matrices are diagonal we need to simultaneously rotate the left-handed and right-handed leptons:

$$e_R \rightarrow U_R e_R, \quad L \rightarrow U_L L. \quad (27)$$

The Lagrangian transforms as

$$-\mathcal{L}_{lepton} = \frac{\sqrt{2}}{v_{13}} \bar{L} \left(U_L^\dagger N U_R H_1 + U_L^\dagger M U_R H_3 \right) e_R + h.c. \quad (28)$$

Naming

$$U_L^\dagger N U_R = N_d, \quad (29)$$

$$U_L^\dagger M U_R = M_d \quad (30)$$

where $M_d = \text{diag}(m_e, m_\mu, m_\tau)$ is diagonal with real and positive diagonal elements.

If after bi-diagonalization, the matrix N_d is not diagonal, then there are scalar tree-level flavour changing neutral interactions in the lepton sector, and the lepton FCNC coupling for those interactions are obtained from the entries of N_d .

Since U_R is completely unknown and N is arbitrary, the N_d coefficients are arbitrary; in order to look at specific processes, some assumptions must be made about their magnitudes.

Motivated by stringent constraints on flavor-changing couplings involving the first two generations, Cheng and Sher [30] showed that if one requires no fine-tuning in the Yukawa matrices, then the flavor-changing couplings should be of the order of the geometric mean of the Yukawa couplings of the two fermions. In other words, the so-called Cheng-Sher ansatz is

$$(N_d)_{ij} = k_{ij} \sqrt{m_i m_j} \quad (31)$$

where k_{ij} are of order one. Since the most severe bounds on FCNC arise from the first two generations and the Yukawa couplings of the first two generations are small, this ansatz can explain these bounds without requiring huge scalar masses.

Using this ansatz we can write the flavor changing matrix to be approximately of the form

$$N_d = \begin{pmatrix} k_{11} m_e & 0 & 0 \\ 0 & k_{22} m_\mu & k_{23} \sqrt{m_\mu m_\tau} \\ 0 & k_{32} \sqrt{m_\mu m_\tau} & k_{33} m_\tau \end{pmatrix}. \quad (32)$$

Note that we have not included the (12), (13), (21) and (31) elements. FCNC involving electrons are tightly constrained and the coefficients cannot be too large, as shown in Ref. [31]. Using the Cheng-Sher ansatz, the current bound from $\mu \rightarrow e\gamma$ requires [31] that $k_{12}, k_{13}, k_{21}, k_{31}$ cannot be much bigger than one. Including these coefficients would make no difference in the physics discussed in the rest of the paper, and thus for simplicity we do not include them here.

The Lagrangian of the lepton sector in the mass basis reads

$$-\mathcal{L}_{lepton} = \frac{\sqrt{2}}{v_{13}} \bar{L} (H_1 N_d + H_3 M_d) e_R + h.c.. \quad (33)$$

We can separate out the lepton sector Lagrangian into two components. One that includes the FCNC couplings and the other is flavor diagonal

$$\mathcal{L}_{lepton} = \mathcal{L}_{FCNC} + \mathcal{L}_{diag}, \quad (34)$$

with

$$\mathcal{L}_{diag} = -\frac{\sqrt{2} m_i}{v_{13}} ((H_1 k_{ii} + H_3) \bar{\nu}_L^i e_R^i + (H_1 k_{ii} + H_3) \bar{e}_L^i e_R^i) + h.c. \quad (35)$$

where the charged component for the first term and the neutral component for the second term are implied for each Higgs doublet.

Expanding out equation (35) in terms of the mass eigenstates we can write in analogous way as eq (21)

$$\mathcal{L}_{diag} = -\sum_{f=e,\mu,\tau} \sum_{b=2}^3 \frac{\sqrt{2}}{v} m_f \left\{ (Z_{H_{b-1}^+} \bar{\nu}_L f_R H_{b-1}^+ + h.c.) + (Z_{h_b} \bar{f} f h_b - Z_{A_{b-1}^i} \bar{f} \gamma_5 f A_{b-1}) \right\}. \quad (36)$$

and the coupling constant Z can be found on the next section.

Similarly for the FCNC component

$$\mathcal{L}_{FCNC} = -\frac{\sqrt{2} k_{23}}{v_{13}} \sqrt{m_\mu m_\tau} (H_1 (\bar{\nu}_{\tau L} \mu_R + \bar{\nu}_{\mu L} \tau_R) + H_1 (\bar{\tau}_L \mu_R + \bar{\mu}_L \tau_R)) + h.c. \quad (37)$$

For now we are only interested in couplings which lead to $h \rightarrow \mu\tau$, thus the relevant terms are

$$\mathcal{L}_{FCNC} \supseteq -\frac{\sqrt{2}}{v_{13}} H_1 \sqrt{m_\mu m_\tau} (k_{23} \bar{\mu}_L \tau_R + k_{32} \bar{\tau}_L \mu_R) + h.c. \quad (38)$$

where the coupling to the neutral component of the doublet is understood. Expanding in terms of the scalars h_i with $i = 1, 2, 3$, (note Eq. 7), we can write this as

$$\mathcal{L}_{FCNC} = -\frac{\sqrt{m_\mu m_\tau}}{v_{13}^2} \{v_3(T_3)_{1i} - v_1(T_3)_{3i}\} h_i \{k_{23} \bar{\mu}_L \tau_R + k_{32} \bar{\tau}_L \mu_R\} + h.c., \quad i = 1, 2, 3. \quad (39)$$

Choosing $k_{ij} = k_{ji}$ and real gives

$$\mathcal{L}_{FCNC} \supseteq \frac{\sqrt{m_\mu m_\tau}}{v_{13}^2} \{v_3(T_3)_{1i} - v_1(T_3)_{3i}\} k_{23} h_i \{\bar{\tau} \mu + \bar{\mu} \tau\} \quad i = 1, 2, 3. \quad (40)$$

which gives the relevant couplings. The flavor changing couplings of the neutral pseudoscalars are found in a similar way, the resulting expression being given by

$$\mathcal{L}_{FCNC} \supseteq \frac{i\sqrt{m_\mu m_\tau}}{v_{13}^2} \{v_3(T_2)_{1i} - v_1(T_2)_{3i}\} k_{23} A_{i-1} \{\bar{\tau} \gamma_5 \mu + \bar{\mu} \gamma_5 \tau\} \quad i = 1, 2, 3. \quad (41)$$

It should be noted that the Cheng-Sher ansatz is completely consistent with the hints from CMS. In the CMS paper[10], they plot their results as a function of the Yukawa couplings $Y_{\mu\tau}$ and $Y_{\tau\mu}$, and also include the line in which the product of the two is $m_\mu m_\tau / v^2$. From this, one can see that the original Cheng-Sher ansatz gives a result slightly below the ‘observed’ value for $k_{\mu\tau} = 1$. Thus, generally the model would predict a branching ratio of the order of a few tenths of a percent, which is consistent with current observations.

One can also discuss the decoupling limit of the model. In the conventional 2HDM, as discussed in detail by Gunion and Haber [32], one notes that $\cos(\alpha - \beta) = 0$ implies that the light Higgs couples to gauge bosons and fermions with the same couplings as in the Standard Model. This is the decoupling limit, and only the light Higgs gets a vev. The other Higgs bosons then have vanishing coupling to gauge bosons, but may still have couplings to fermions. In the decoupling limit, the other Higgs bosons are taken to be sufficiently heavy that they do not affect quark and lepton phenomenology. In this model, one can see that the limit

$$\cos \theta = \sin \theta_3 = \cos \psi_3 = \cos \psi = 0$$

gives the coupling of the light Higgs to WW and to fermions equal to their Standard Model values and gives vanishing couplings of the heavy Higgs to WW and hW. This is certainly sufficient to give the decoupling limit, and means increasing precise bounds on deviations of the light Higgs couplings from the Standard Model values will not eliminate the model, but just restrict the parameter-space somewhat.

The above constraints are certainly sufficient for decoupling, but are not fully necessary. In order for the light Higgs to have SM-equivalent couplings to gauge bosons and for the heavy Higgs to have no couplings to gauge bosons, one needs to have $\psi_3 = \psi$ and $\cos \theta = \sin \theta_3$. This corresponds to the first and third equalities in the above equation. In order for the light Higgs to have SM-equivalent couplings to the fermions, one must add the condition $\cot \theta = \cos \psi$, which corresponds to $v_1 = v_3$. The above equation is consistent with this, of course. Note that the constraints in the above paragraph would lead to quark-phobic charged Higgs bosons, but the more general constraint here does not. Thus, tighter bounds on deviations of the light Higgs couplings from their Standard Model values will not force the charged Higgs to decay primarily leptonically.

E. Summary of the couplings

The gauge, quark and lepton couplings are summarized in the Tables below.

	Gauge boson couplings
$g_{h_1 WW}$	$\frac{g^2}{2} v (\cos \theta \sin \theta_3 + \sin \theta \cos \theta_3 \cos(\psi_3 - \psi))$
$g_{h_2 WW}$	$\frac{g^2}{2} v \cos \theta [\cos \beta_3 \sin(\psi - \psi_3) + \sin \beta_3 (\cot \theta \cos \theta_3 - \sin \theta_3 \cos(\psi_3 - \psi))]$
$g_{h_3 WW}$	$\frac{g^2}{2} v \cos \theta [\sin \beta_3 \sin(\psi_3 - \psi) + \cos \beta_3 (\cot \theta \cos \theta_3 - \sin \theta_3 \cos(\psi_3 - \psi))]$
$g_{H_1 hW}$	$\frac{ig}{2} (\cos \theta \cos \theta_3 \cos(\psi - \psi_3) \sin \beta_1 - \sin \beta_1 \sin \theta \sin \theta_3 + \cos \beta_1 \cos \theta_3 \sin(\psi - \psi_3))$
$g_{H_2 hW}$	$\frac{ig}{2} (\cos \beta_1 (\cos \theta \cos \theta_3 \cos(\psi - \psi_3) - \sin \theta \sin \theta_3) - \cos \theta_3 \sin \beta_1 \sin(\psi - \psi_3))$

Table 1: A list of Higgs couplings to gauge bosons. The last two expressions are multiplied by $(p - p')_\mu$.

$\times \sin\theta \sin\psi$	Quark Couplings
$\xi_{h_1}^{u,d}$	$\cos\theta_3 \sin\psi_3$
$\xi_{h_2}^f$	$\cos\beta_3 \cos\psi_3 - \sin\beta_3 \sin\theta_3 \sin\psi_3$
$\xi_{h_3}^f$	$-\cos\psi_3 \sin\beta_3 - \cos\beta_3 \sin\theta_3 \sin\psi_3$
$\xi_{A_1}^u$	$-\cos\psi \cos\beta_2 + \cos\theta \sin\psi \sin\beta_2$
$\xi_{A_2}^u$	$\cos\psi \sin\beta_2 + \sin\psi \cos\theta \cos\beta_2$
$\xi_{A_1}^d$	$-\xi_{A_1}^u$
$\xi_{A_2}^d$	$-\xi_{A_2}^u$
$\xi_{H_1^+}^u$	$\cos\psi \cos\beta_1 - \cos\theta \sin\psi \sin\beta_1$
$\xi_{H_2^+}^u$	$-\cos\psi \sin\beta_1 - \sin\psi \cos\theta \cos\beta_1$
$\xi_{H_1^+}^d$	$-\xi_{H_1^+}^u$
$\xi_{H_2^+}^d$	$-\xi_{H_2^+}^u$

Table 2: A list of the couplings appearing in equation (21). We do not list the couplings of the Goldstone bosons G^\pm and G^0 .

$\times \frac{v}{k_{23}\sqrt{m_\mu m_\tau}} (1 - \sin^2\theta \sin^2\psi)$	FCNC couplings
$h_1 \bar{\mu} \tau$	$\cos\theta \cos\theta_3 \cos\psi_3 - \sin\theta \cos\psi \sin\theta_3$
$h_2 \bar{\mu} \tau$	$-(\sin\theta \cos\psi \cos\theta_3 \sin\beta_3 + \cos\theta(\sin\beta_3 \cos\psi_3 \sin\theta_3 + \cos\beta_3 \sin\psi_3))$
$h_3 \bar{\mu} \tau$	$-\sin\theta \cos\psi \cos\beta_3 \cos\theta_3 + \cos\theta(\cos\beta_3 \cos\psi_3 \sin\theta_3 - \sin\beta_3 \sin\psi_3)$
$iA_1 \bar{\mu} \gamma_5 \tau$	$-(\cos\psi \cos\beta_2 - \cos\theta \sin\psi \sin\beta_2)$
$iA_2 \bar{\mu} \gamma_5 \tau$	$-(\cos\psi \cos\beta_2 + \cos\theta \sin\psi \cos\beta_2)$
$\sqrt{2}H_1^+ (\bar{\nu}_\tau P_R \mu + \bar{\nu}_\mu P_R \tau)$	$-(\cos\psi \sin\beta_1 + \cos\beta_1 \cos\theta \sin\psi)$
$\sqrt{2}H_2^+ (\bar{\nu}_\tau P_R \mu + \bar{\nu}_\mu P_R \tau)$	$-\cos\beta_1 \cos\psi + \cos\theta \sin\beta_1 \sin\psi$

Table 3: These are the lepton flavor violating couplings.

$\times (1 - \sin^2\theta \sin^2\psi)$	Lepton Couplings
$Z_{H_1^+}$	$-\sin\theta/2 (2 \cos\psi \csc\theta \sin\beta_1 - 2 \cos\theta \sin\beta_1 \sin^2\psi + \cos\beta_1 (2 \cot\theta \sin\psi + \sin 2\psi))$
$Z_{H_2^+}$	$\sin\theta ((\cos\psi + \cot\theta) \sin\beta_1 \sin\psi + \cos\beta_1 (-\cos\psi \csc\theta + \cos\theta \sin^2\psi))$
$\sqrt{2}Z_{h_1}$	$(1 + k_{ff}) \cos\theta \cos\theta_3 \cos\psi_3 + (1 - k_{ff}) \cos\psi \sin\theta \sin\theta_3$
$\sqrt{2}Z_{h_2}$	$(k_{ff} - 1) \cos\theta_3 \cos\psi \sin\beta_3 \sin\theta - (1 + k_{ff}) \cos\theta (\cos\psi_3 \sin\beta_3 \sin\theta_3 + \cos\beta_3 \sin\psi_3)$
$\sqrt{2}Z_{h_3}$	$(k_{ff} - 1) \cos\beta_3 \cos\theta_3 \cos\psi \sin\theta + (k_{ff} + 1) \cos\theta (\sin\beta_3 \sin\psi_3 - \cos\beta_3 \cos\psi_3 \sin\theta_3)$
$\sqrt{2}Z_{A_1}$	$-\left((1 + k_{ff} \cos 2\theta) \cos\psi \sin\beta_2 + (1 + k_{ff}) \cos\beta_2 \cos\theta \sin\psi\right)$
$\sqrt{2}Z_{A_2}$	$-(1 + k_{ff} \cos 2\theta) \cos\beta_2 \cos\psi + (1 + k_{ff}) \cos\theta \sin\beta_2 \sin\psi$

Table 4: These are the couplings to leptons appearing in equation (36).

III. BRANCHING RATIOS AND CONSTRAINTS ON THE CHARGED HIGGS BOSONS

For the analysis of this section it is useful to write down the couplings of the charged Higgs to both, quarks and leptons in one single equation, see (21) and (35),

$$\mathcal{L} = \frac{g}{\sqrt{2}M_W} \sum_{H^+=H_1^+, H_2^+} \{ \xi_{H^+}^u \bar{u}_R V_{ud} m_u d_L - \xi_{H^+}^d \bar{d}_R V_{ud} m_d u_L - Z_{H^+} \bar{\nu}_L m_l l_R \} H^+ + h.c. \quad (42)$$

Note that the charged Higgs interactions depend only on three mixing angles. With the two different charged Higgs masses, this gives a five-dimensional parameter-space, as opposed to the 2HDM in which the interactions depend on the single charged Higgs mass and $\tan\beta \equiv v_2/v_1$.

Since the model has a close similarity to the lepton specific model, one can look at charged Higgs interactions in that model for guidance. Logan and MacLennan [33] studied the constraints on the charged Higgs in this model, looking at direct LEP-II searches and flavor universality in tau decays. Su and Thomas [34] studied constraints from $b \rightarrow s\gamma$, $B^0 - \bar{B}^0$ mixing and several other processes. Aoki, Kanemura, Tsumura and Yagyu [35], in a very comprehensive

study, considered these as well as $B \rightarrow \tau\nu$. These papers and others are discussed in the review article of Ref. [7], where it is noted that the bounds from radiative B decays and R_b are the same as in the type I model, and bounds from flavor universality in tau decays are not significant unless $\tan\beta > 65$. The most intriguing result is the possibility that the $\tau\nu$ decay mode of the charged Higgs can dominate the decay, *even above the $\bar{b}t$ threshold*. For this to occur, one must have $\tan\beta > 10$. Such fairly large values of $\tan\beta$ may have problems with perturbative unitarity [36], although there are allowed regions of parameter-space with larger values of $\tan\beta$. We will see that in the 3HDM, a dominance of the $\tau\nu$ decay mode can occur even if the ratio of vacuum expectations values is not particularly large.

A. Charged Higgs decays

As noted above, the charged Higgs sector depends on three mixing angles and two masses. As a first step, these parameters must be constrained by the requirement that they do not contradict LHC results on production and decay of the SM-Higgs like 125 GeV boson. Given the number of parameters, a full χ -squared analysis is unnecessary, and we will simply require that the couplings of the light Higgs (h_1 in Tables 1 and 2) be within 20% of their Standard Model values for the WW , ZZ and tt couplings and 30% for the bb coupling[37]. The $\gamma\gamma$ coupling will not provide useful information due to unknown contributions of heavy charged Higgs bosons (with arbitrary couplings) in the loop. All of our results below will only consider regions in parameter-space that satisfy those constraints.

We consider the branching ratios of the charged Higgs bosons. In this article we are going to assume that all of the additional neutral scalars are too heavy for the charged Higgs to decay into them. Then the most relevant decay modes to consider are $H^+ \rightarrow hW^+$, $\bar{b}t$, $\bar{\tau}\nu$ and possibly into $\mu\nu$. Focus will be placed mainly on heavy charged scalars $M_{H_i^\pm} \gg m_t$ and therefore the decay mode into a quark-antiquark pair will be dominated by the $\bar{b}t$ decay mode. The leading order expressions for the partial widths are given by

$$\Gamma(H_i^\pm \rightarrow hW^\pm) = \frac{\sqrt{2}G_F}{16\pi} g_{H_i h W}^2 M_{H_i^\pm}^3 \left[1 + \frac{(M_W^2 - m_h^2)^2}{M_{H_i^\pm}^4} - 2 \frac{(M_W^2 + m_h^2)}{M_{H_i^\pm}^2} \right]^{3/2}, \quad (43)$$

$$\Gamma(H_i^+ \rightarrow \bar{b}t) = \frac{3G_F(\xi_{H_i^+}^u)^2}{4\pi\sqrt{2}} M_{H_i^+} m_t^2 \left(1 - \frac{m_t^2}{M_{H_i^+}^2} \right)^2, \quad (44)$$

$$\Gamma(H_i^+ \rightarrow \nu_l \bar{l}) = \frac{G_F}{4\pi\sqrt{2}} M_{H_i^+} \times \begin{cases} m_\tau^2 Z_{H_i^+}^2, & \tau\nu_\tau \\ m_\mu m_\tau C_{H_i^+}^2, & \tau\nu_\mu, \mu\nu_\tau \end{cases} \quad (45)$$

where $h = h_1$ being the SM-like Higgs boson is implied in (43). The b quark and τ lepton masses have been neglected in (44) and (45) respectively. For the leptonic decay (45) we included the flavor changing processes induced by (37) and the respective couplings, $C_{H_i^+}$, are given in the last two rows of table 3.

In the lepton-specific 2HDM the couplings of the charged Higgs boson to quarks are proportional to the SM couplings multiplied by $\cot\beta$. Also in that model, the couplings of the charged Higgs to right handed leptons are proportional to $\tan\beta$. Thus in the limit $\tan\beta \gg 1$, the charged scalar becomes quark-phobic but leptophilic enhancing the possibility of a large decay width into $\bar{\nu}\tau$, even above the $\bar{b}t$ threshold [7].

Here we explore that possibility in the 3HDM. The couplings to the quarks in the 3HDM are given by

$$\xi_{H_1^+}^u = \frac{\cos\beta_1 \cot\psi - \cos\theta \sin\beta_1}{\sin\theta}, \quad (46)$$

$$\xi_{H_2^+}^u = -\frac{\cos\beta_1 \cos\theta + \cot\psi \sin\beta_1}{\sin\theta}. \quad (47)$$

We investigate quark-phobic points in the parameter space of the mixing angles (θ, ψ, β_1) , i.e. points for which either $\xi_{H_1^+}^u$, $\xi_{H_2^+}^u$ or both are very small. Without loss of generality we do this analysis in the region of parameter space $0 < (\theta, \psi, \beta_1) < \pi/2$. The reason is that we want the vevs given in (4) to be real and positive, so that the rotation angles θ, ψ can be restricted to be in the first quadrant. The reason to consider β_1 in that region is by noticing that taking $\beta_1 \rightarrow \beta_1 + \pi/2$ in (46) yields (47).

By considering (46) as a function of θ and ψ and varying β_1 in that region, we find that there is always a surface for which (46) is equal to zero and that (47) never crosses zero in that region. Thus the 3HDM allows for either H_1 or H_2 to be quark-phobic but not both at the same time.

When we consider the decay mode to the SM Higgs and a gauge boson, we see that the decay width (43) is proportional to the square of the couplings $g_{H_i h W}$ given in table 1. These couplings are dependent on the mixing angles θ_3 and ψ_3 , Therefore we also investigate points in the parameter-space that are both quark- and gauge-phobic. We shall consider values of the mixing angles θ_3 , ψ_3 in the same region described above, although they could in principle be larger since they depend on the parameters of the scalar potential. By following the same procedure described above we find that it is always possible for (46) and both gauge couplings to cross zero at the same point. Therefore the 3HDM can always have a quark- and gauge-phobic H_1 decaying mostly into leptons and a gauge-phobic H_2 decaying most of the time to tb .

If one chooses parameters such that the ratio of vevs is large, one must consider unitarity and perturbativity. As remarked in the 2HDM review of Ref. [7], having high ratios of vevs is only allowed for small regions of parameter space. Therefore we consider points for which the ratios in equation (4) are not too big ($\tan \theta, \tan \psi < 6$).

The production cross section for charged Higgs bosons is dominated by the gluon-gluon fusion process $gg \rightarrow \bar{t}bH^+$ and is proportional to the square of the couplings (46) and (47). This mechanism will not produce charged Higgs bosons in the quark-phobic limit. In that case, they can still be produced by vector boson fusion (VBF) which is independent of quark couplings. As shown in Ref. [33], the VBF production cross section is 2 – 3 orders of magnitude smaller than the gluon-gluon fusion cross section for $\tan \beta = 1$, see figure 14 of Ref [53], we thus consider quark-couplings which are 0, 0.03, 0.12 times the $\tan \beta = 1$ coupling.

A novel feature of this model is the appearance of the flavor changing decay modes given in (45). Usually the muonic decay of the charged Higgs is negligible due to the small muon coupling, however, here the tau coupling enters if the ν is a ν_τ . The region of parameter space for which the flavor changing branching ratios given in (45) are bigger than 10% is very small and is concentrated in the upper right corner of the plane (θ, ψ) which corresponds to large ratios of vevs. In the limit $\theta, \psi \rightarrow \pi/2$ these modes will be dominant with H_1 decaying 50% into $\mu\nu_\tau$ and $\tau\nu_\mu$ each.

We find the branching ratios in figure 1. We have chosen three benchmark points which give values for ξ of 0, 0.03, 0.12. These points are $(\theta, \psi, \beta_1, \theta_3, \psi_3) = (1, 1.37, 0.36, 0.92, 1.14)$, $(1.30, 1.36, 0.59, 0.39, 1.36)$, $(1.41, 1.41, 0.25, 0.28, 1.41)$ respectively. Changing the benchmark points, without changing the value of ξ will not substantially alter these results.

For a quark Yukawa coupling of 0.12 times the $\tan \beta = 1$ 2HDM coupling, one sees that the branching ratio into $\tau\nu_\tau$ is substantial, although not dominant. As the quark Yukawa coupling gets smaller, the branching ratio into $\tau\nu_\tau$ becomes dominant for small masses.

The novel flavor changing mode discussed above, into $\mu\nu_\tau$ is small, but would be easier to detect. With the LHC operating with an integrated luminosity of 4000 fb^{-1} and the pair production cross section by vector boson fusion (VBF), which is model independent, is around a few fb as can be seen from figure 8 of Ref. [33], producing approximately 300 events. Note that a discussion of the neutral heavy Higgs decay into $\mu\tau$ can be found in Ref. [38].

B. Constraints on the charged Higgs from B physics

1. $B \rightarrow X_s \gamma$

In this section we investigate bounds on the charged Higgs masses coming from the inclusive radiative decay $B \rightarrow X_s \gamma$ and give the leading order (LO) results at the matching scale M_W at which the full theory is matched into an effective theory with five quark flavours in order to provide insight on the effect of new physics, as is done in Ref. [39]. This is a well suited process to probe new physics. However it suffers from large theoretical uncertainties coming mainly from the choices of the renormalization and matching scales, which are not well defined at the leading order (LO) [41].

The next to leading order calculations (NLO) in the SM [42], [43] and in the 2HDM [41] are well known. The branching fraction of the inclusive radiative decay at NLO is presented as [41]

$$\begin{aligned} \mathcal{B}(B \rightarrow X_s \gamma) = & \mathcal{B}(B \rightarrow X_c e \bar{\nu}_e) \left| \frac{V_{ts}^* V_{tb}}{V_{cb}} \right|^2 \frac{6\alpha_{em}}{\pi f(z)\kappa(z)} \frac{\bar{m}_b^2(\mu_b)}{m_b^2} \\ & \times (|D|^2 + A) \left(1 - \frac{\delta_{SL}^{NP}}{m_b^2} + \frac{\delta_\gamma^{NP}}{m_b^2} + \frac{\delta_c^{NP}}{m_c^2} \right), \end{aligned} \quad (48)$$

where $z = m_c^2/m_b^2$ and f and κ are phase-space suppression factors. The last term in parenthesis includes corrections obtained by the method of the heavy-quark effective theory (HQEFT) which relate the quark decay rate to the




FIG. 1: Branching ratios of H_1 at the benchmark points in the text. ξ is the relative quark coupling to the charged Higgs from Eqs. 46 and 47. $\xi = 0$ corresponds to the pure quark-phobic limit.

hadronic process [45]. The term A is the correction coming from the Bremsstrahlung process $b \rightarrow s\gamma g$. Finally, $|D|^2$ contains the Wilson coefficients at the renormalization scale μ_b relevant to the radiative decay and is given by equation (26) on Ref. [41]. As discussed in section 5 of Ref. [41] the LO 2HDM Wilson coefficients, to be added to SM ones, at the matching scale m_W are given by

$$\delta C_{7,8}^{(0)eff}(m_W) = \frac{A_u^2}{3} F_{7,8}^{(1)}(y) - A_u A_d F_{7,8}^{(2)}(y), \quad (49)$$

with

$$F_7^{(2)}(y) = \frac{y(3-5y)}{12(y-1)^2} + \frac{y(3y-2)}{6(y-1)^3} \ln y, \quad (50)$$

$$F_8^{(2)}(y) = \frac{y(3-y)}{4(y-1)^2} - \frac{y}{2(y-1)^3} \ln y, \quad (51)$$

$$y = \frac{\bar{m}_t^2(m_W)}{M_H^2} \quad (52)$$

and A_u and A_d are the couplings of the charged Higgs to the quarks. In the type I 2HDM these are $A_u = A_d = \cot \beta$.

To obtain the new Wilson coefficients of the 3HDM the only different thing is that we need to add the contribution from each H_1^+ and H_2^+ in the loop and modify the corresponding quark couplings. The result is

$$\delta C_{7,8,NP}^{(0)eff}(m_W) = \sum_{i=1}^2 \left(\xi_{H_i^+}^u \right)^2 \left(\frac{1}{3} F_{7,8}^{(1)}(y_i) - F_{7,8}^{(2)}(y_i) \right), \quad y_i = \frac{\bar{m}_t^2(m_W)}{M_{H_i^+}^2}. \quad (53)$$

Since the above equation is dependent on the three angles (θ, ψ, β_1) and the two charged Higgs masses we see that the branching fraction of the radiative decay will be dependent on five parameters. The Wilson coefficients at the renormalization scale μ_b and at NLO cannot be summarized in a few lines and are extracted from Ref. [42].

Since we are interested in a region of parameter-space in which one of the charged Higgs bosons has suppressed couplings to quarks (we have checked that the suppression is sufficient to make the contribution negligible), the results will only depend on the mass of the other charged Higgs, H_2^+ . For values of $\xi_{H_2^+}^u = 0.2, 0.6, 0.8$ and 1.4 , we find the results in Figure 2. These four points correspond to mixing angles $(\theta, \psi, \beta_1) = (1.41, 1.41, 0.25)$, $(1.00, 1.37, 0.36)$, $(1.20, 1.02, 1.05)$ and $(0.69, 1.21, 0.46)$ respectively. The first two of these correspond to the two values of the points listed in the previous subsection (corresponding to $\xi = 0, 0.12$ (the other point in the last subsection gives results extremely similar to the $\xi = 0.12$ point)). The other two points give results that are somewhat more significant for $B \rightarrow X_s \gamma$.




FIG. 2: Branching fraction for the radiative decay $B \rightarrow X_s \gamma$ as a function of the charged Higgs mass. The horizontal red lines give the 2σ experimentally allowed region and the blue dotted line give the central value of the SM prediction. The points of parameter space, in order of increasing dash length, correspond to values of $\xi_{H_2^+}^u = 0.2, 0.6, 0.8$ and 1.4 .

All black lines asymptote to the SM prediction $BR(B \rightarrow X_s \gamma) = (3.6 \pm 0.36) \times 10^{-4}$ in the limit of high mass values. The experimentally allowed region is $(3.52 \pm 0.23 \pm 0.09) \times 10^{-4}$ [46]. The line corresponding to the first point lies entirely inside the 1σ experimental region and therefore does not yield any bound. Including NLO QCD corrections the lower bounds on the charged Higgs mass corresponding to the points considered in figure 2 at 95% C.L. are given by 295 GeV, 370 GeV and 900 GeV for the values of $\xi_{H_2^+}^u = 0.6, 0.8$ and 1.4 . Thus, the bounds will be fairly weak unless $\xi_{H_2^+}^u$ is unusually large. Note that a more recent analysis by Misiak et al. [44] does the NNLO calculation and finds results that are slightly lower, but well within a single standard deviation from the NLO results.

2. R_b

In this subsection we focus on the observable

$$R_b = \frac{\Gamma(Z \rightarrow b\bar{b})}{\Gamma(Z \rightarrow \text{hadrons})} \quad (54)$$

which is sensitive to radiative corrections. It is shown in Ref. [47] that in non-minimal models containing only doublets, as is our case, the loop corrections due to virtual charged Higgs bosons always worsen agreement with experiment. In that same reference they introduce a parametrization for a general extended Higgs sector and calculate the contribution to $Zb\bar{b}$ from one-loop radiative corrections involving singly charged and neutral Higgs bosons. They obtained general expressions for the corrections to the left- and right-handed $Zb\bar{b}$ couplings, and then use the measurements of R_b and A_b (the coupling asymmetry) to constrain specific models.

We write the interaction as

$$\mathcal{L} \propto Z_\mu \bar{b} \gamma^\mu [\bar{g}_b^L P_L + \bar{g}_b^R P_R] b, \quad (55)$$

the effective couplings are then given by $\bar{g}_b^{L,R} = g_{Zbb}^{L,R} + \delta g^{L,R}$, where g_{Zbb} are the tree-level couplings and δg are the radiative corrections.

The radiative corrections to SM extensions with only doublets and singlets, are given by the second term of equation (4.5) of Ref. [47]

$$\delta g^{L,R} = \pm \frac{1}{32\pi^2} \frac{e}{s_W c_W} \sum_{i \neq G^+} \left(g_{H_i^+}^{L,R} \right)^2 \times \left[\frac{R_i}{(R_i - 1)} - \frac{R_i \log R_i}{(R_i - 1)^2} \right] \quad (56)$$

where $s_W = \sin \theta_W$, $c_W = \cos \theta_W$ are the weak mixing factors, $R_i^2 = m_t^2 / M_{H_i^+}^2$ and $g_{H_i^+}^{L,R}$ are obtained from the Lagrangian by writing the interaction of the charged Higgs to quarks as

$$\bar{t} (g^L P_L + g^R P_R) b H^+ + h.c. \quad (57)$$

The correction due to Goldstone boson exchange is excluded in the sum since is the same as in the SM. In the 3HDM the above coefficients are given by

$$g_{H_i^+}^L = \sqrt{2} \frac{m_t}{v} \xi_{H_i^+}^u, \quad (58)$$

$$g_{H_i^+}^R = -\sqrt{2} \frac{m_b}{v} \xi_{H_i^+}^u. \quad (59)$$

Thus we can write the corrections as

$$\delta g^L = \frac{1}{32\pi^2} \frac{e}{s_W c_W} \left(\frac{\sqrt{2} m_t}{v} \right)^2 \sum_{i=1,2} (\xi_{H_i^+}^u)^2 \left[\frac{R_i}{(R_i - 1)} - \frac{R_i \log R_i}{(R_i - 1)^2} \right] \quad (60)$$

and

$$\delta g^R = -\frac{m_b^2}{m_t^2} \delta g^L. \quad (61)$$

The experimentally allowed range is given by $R_b = 0.21642 \pm 0.00073$ [47]. We find bounds on the charged Higgs mass for the same quark-phobic points listed in figure 2. However the first two points give a prediction that lies well inside the 2σ experimentally allowed region, therefore we only obtain a bound coming from the last point, namely

$$m_{H_2^+} > 395 \text{ GeV} \quad (95\% \text{ C.L.}). \quad (62)$$

Thus we see that the radiative process $B \rightarrow X_s \gamma$ yields stronger constraints than those of R_b .

3. Combining $B \rightarrow X_s \gamma$ and R_b

In the above, we illustrated the bounds with a few benchmark points. But there are only three angles and two masses, and we can scan the parameter-space to determine the allowed regions. In the figures below we have plotted the regions in which the χ^2 tests for both $B \rightarrow X_s \gamma$ and R_b processes are satisfied to 95% significance level. We parametrized the rotation angles θ and ψ in terms of the vevs v_1, v_2 and imposed the constraint $\sum_i v_i^2 = 246 \text{ GeV}$. In Figure 3, we show a contour plot of v_1 and v_2 for a specific value of β_1 chosen to give the quark-phobic point for H_1 .

We then consider different values of β_1 in Figure 4. The patterns are clear. Some regions of parameter-space, such as small v_2 and intermediate v_1 , are excluded for all values of β_1 . This will become relevant when considering the possibility of $B \rightarrow \mu \nu_\tau$. We now turn to the leptonic decays of the B

4. $B^- \rightarrow \tau \bar{\nu}_\tau$

We study the charged current decay $B^- \rightarrow \tau \bar{\nu}$ type of modes which are just tree-level processes mediated by the electroweak gauge bosons W^\pm and the charged Higgs bosons H_i^\pm .




FIG. 3: Allowed region (colored) of the parameter space as a function of the vevs v_1 and v_2 . The green, blue and red regions correspond to the mass values $M_H = 300, 500, 700$ GeV respectively. We take the limit in which H_2 (H_1) effectively decouples from fermions on the left (right). The angle β_1 was chosen at the quark-phobic point of H_1 .




FIG. 4: Allowed region of parameter space for $\beta_1 = 0, \pi/4$. The charged Higgs H_2 (H_1) was taken very heavy on the left (right). The green, blue and red regions correspond to the mass values $M_H = 300, 500, 700$ GeV respectively.

Using the quark and lepton couplings of the charged Higgs mass eigenstates given by (42), the W^\pm and H_i^\pm effectively induce the four-Fermi interaction [49]

$$\mathcal{L} = -2\sqrt{2}G_F V_{ub} [(\bar{u}\gamma^\mu P_L b)(\bar{l}\gamma_\mu P_L \nu_l) - R_{3HDM,l}(\bar{u}P_R b)(\bar{l}P_L \nu_l)], \quad (63)$$

where

$$R_{3HDM,l} = \sum_{i=1}^2 \frac{m_l m_b}{M_{H_i^+}^2} Z_{H_i^+} \xi_{H_i^+}^u, \quad (64)$$

where the first term give the SM contribution, while the second one give that of the charged scalars. In Ref. [48] a study of multi-Higgs doublet models with natural flavor conservation was performed. In that article they assumed that all but the lightest of the charged scalars effectively decouple from fermions and carried out an analysis of phenomenological constraints on the Yukawa couplings.

We do not make those assumptions here and instead modify the 2HDM result by the appropriate couplings and find that the charged Higgs bosons modify the SM expectation by the factor

$$r_H \equiv \frac{BR(B^- \rightarrow l\bar{\nu})}{BR_{SM}(B^- \rightarrow l\bar{\nu})} = \left| 1 - \sum_{i=1}^2 m_B^2 \frac{\xi_{H_i^+}^u Z_{H_i^+}}{M_{H_i^+}^2} \right|^2 \quad (65)$$

which is independent of the lepton mass. We also call that factor r_H , following the notation in Ref. [49]. Notice that in the limit $(\theta, \beta_1) \rightarrow (\pi/2, 0)$, (47) vanishes and H_2 becomes quark-phobic and the sum on the right-hand side collapses to

$$\frac{BR(B \rightarrow l\nu)}{BR_{SM}(B \rightarrow l\nu)} = \left| 1 + \frac{m_B^2}{M_{H^+}^2} \right|^2 \quad (66)$$

which is the expression for the lepton-specific 2HDM. Therefore all the $B^- \rightarrow l\bar{\nu}$ modes are always enhanced in the lepton-specific 2HDM while they could be enhanced or suppressed in the 3HDM by the same factor r_H . The ratio of the measured value to the SM prediction is 1.37 ± 0.39 [50]. We compare this number with the r_H prediction for the 3HDM and we find that in the quark-phobic points of H_1 the 3HDM prediction lies completely inside the 1σ region for all masses of H_2 starting from the threshold. Larger values of the correction factor arise if we take the rotation angles $\theta, \psi \ll 1$, but that would correspond having a hierarchy on the vevs values $v_3 \gg v_1 \gg v_2$. This is no surprise since it is remarked in [49], that for models where d-type quarks and charged leptons derive mass from different doublets there is no interesting effect. This is exactly the case for lepton-specific-type models.

5. Flavor changing processes

This model has another interesting possibility that does not exist in any other versions of the 2HDM. One can study $B \rightarrow \mu\nu\tau$. The Standard Model decay $B \rightarrow \mu\nu\mu$ is very small due to helicity suppression, and the 2HDM charged Higgs contribution is negligible due to the small Yukawa coupling of the muon to the Higgs. But in this model, the flavor-changing couplings are proportional to the geometric mean of the Yukawa couplings, and thus the τ Yukawa coupling can play an important role. Of course, experimenters can not determine the flavor of the neutrino, so this would appear as a contribution to $B \rightarrow \mu\nu$.

From (23) and (37) we can write down the flavor changing interaction of the charged Higgs bosons

$$\mathcal{L}_{FCNC} \supseteq -\frac{\sqrt{2}}{v} \sqrt{m_\mu m_\tau} \sum_{H^+=H_1^+, H_2^+} C_{H^+} [\bar{\nu}_\mu P_R \tau + \bar{\nu}_\tau P_R \mu] H^+ + h.c. \quad (67)$$

where the C_{H^+} are the flavor changing coupling constants given in the last two rows of table 3. The charged Higgs induces the four-Fermi flavor-changing interaction

$$\mathcal{L}_{4F} = \frac{4G_F}{\sqrt{2}} V_{ub} \sum_{H^+} R_{H^+} (\bar{u} P_R b) (\bar{\tau} P_L \nu_\mu + \bar{\mu} P_L \nu_\tau), \quad (68)$$

where the sum is performed over the charged Higgs mass eigenstates and

$$R_{H^+} = \sqrt{m_\mu m_\tau} m_b \frac{\xi_{H^+}^u C_{H^+}}{M_{H^+}^2}, \quad (69)$$

similar to (64). The branching fraction for the flavor changing processes are given by the flavor conserving SM result times a correction factor, i.e.,

$$BR(B \rightarrow \mu\bar{\nu}_\tau, \tau\bar{\nu}_\mu) = BR(B \rightarrow \mu\bar{\nu}_\mu, \tau\bar{\nu}_\tau)_{SM} r_{H,l}, \quad (70)$$

where the l index stands for the charged lepton and the correction factor is

$$r_{H,l} = \left(\frac{\sqrt{m_\mu m_\tau}}{m_l} \sum_{H^+} \frac{m_B^2}{M_{H^+}^2} \xi_{H^+}^u C_{H^+} \right)^2. \quad (71)$$

The SM branching ratio is given by [49], [51], [52]

$$BR_{SM}(B^- \rightarrow l^- \bar{\nu}) = \frac{G_F^2 m_B m_l^2}{8\pi} \left(1 - \frac{m_l^2}{m_B^2}\right)^2 f_B^2 |V_{ub}|^2 \tau_B. \quad (72)$$

Using this formula the SM predictions are [51]

$$BR(B \rightarrow \tau \nu_\tau)_{SM} = (0.84 \pm 0.11) \times 10^{-4}, \quad (73)$$

$$BR(B \rightarrow \mu \nu_\mu)_{SM} = (3.8 \pm 0.5) \times 10^{-7}. \quad (74)$$

The Heavy Flavor Averaging Group (HFAG) found, as of July 2016, the value $BR(B \rightarrow \tau \nu_\tau) = 1.06 \times 10^{-4}$ and the upper limit $BR(B \rightarrow \mu \nu_\mu) < 1.0 \times 10^{-6}$ at 90% C.L.

We investigate the region of the parameter space that allow for an enhancement of the flavor-changing decay $B \rightarrow \mu \nu_\tau$ over the SM flavor-conserving prediction. One special case is the degenerate limit $M_{H_1^+} = M_{H_2^+}$, in which the correction factor vanishes trivially since the coupling constants satisfy the relation

$$\xi_{H_1^+}^u C_{H_1^+} = -\xi_{H_2^+}^u C_{H_2^+} \quad (75)$$

for any rotation angle (θ, ψ, β_1) . Thus one way to maximize the value of the correction factor is to take the decoupling limit in which either charged boson becomes extremely heavy and the other one take its threshold value.

The results are presented in figure 5, where we consider the values of parameters in which the contribution from the charged Higgs is substantial.




Figure 5: Region of the parameter space (θ, ψ) for which the correction factor $r_{H,\mu} > (0.2, 1, 2)$.

We have taken $M_{H_1^+}$ infinitely big and $M_{H_2^+}$ at its threshold value. The chosen value of β_1 maximizes the area.

One natural question that arise is if there exist a region of parameter space that allow a substantial enhancement and is permitted by the χ^2 tests of the previous section. The answer is clearly negative as can be seen from the figure below where the colored region shows the parameter space that is allowed by the radiative processes $B \rightarrow X_s \gamma$ and R_b at 95% C.L. and the same values of M_{H_2} and β_1 were used. That region is concentrated at the upper right corner and there is no overlapping between them.




Figure 6: Allowed region of parameter space by the χ^2 tests of the processes $B \rightarrow X_s \gamma$ and R_b . The decoupling limit of $H_1(H_2)$ has been taken in the left (right).

IV. CONCLUSIONS

Motivated by hints of lepton flavor violation in Higgs decays and the strong constraints on tree level FCNC in the quark sector, we consider a 3HDM in which one doublet couples to quarks and the other two to leptons. This structure can be imposed with a simple Z_2 symmetry. The model has two charged Higgs pairs, two pseudoscalars and three scalars.

The mass matrices and mixing angles are then determined. In the charged Higgs sector, there are two masses and three mixing angles, two of which come from ratios of vevs. We focus on the phenomenology of the charged Higgs bosons, concentrating on the region in which the lightest of the charged Higgs pairs is above the top quark mass. The model is similar to the 2HDM lepton specific model. In that model, the charged Higgs branching ratio into $\tau\nu$ can exceed that of $t\bar{b}$ only if the ratio of vevs ($\tan\beta$ in that model) is quite large, possibly large enough to raise unitarity concerns. In this model, even without such a large ratio of vevs, the decays into $\tau\nu$, $t\bar{b}$ and hW can all be comparable. We also study the constraints from B decay. We find that there is a new decay, $B \rightarrow \mu\nu_\tau$, but the region of parameter-space in which this is substantial is inconsistent with other B decay bounds.

A unique feature of the model is the possibility of charged Higgs decay into $\mu\nu_\tau$. The flavor of the neutrino can of course not be measured, but flavor does feed in through the size of the coupling, expected here to be the geometric mean of the muon and tau Yukawa couplings. The decay branching fraction is typically a few percent, although for extreme regions of parameter-space can be much higher. Although the branching fraction is small, the relative ease of detecting muons makes this decay worthy of careful study.

V. ACKNOWLEDGMENTS

The authors would like to thank Chris Carone, Pedro Ferreira and Keith Thrasher for useful discussion. This work was supported by the NSF under Grant PHY-1519644.

Appendix A

Minimizing the Higgs potential yields:

$$m_{11}^2 = m_{13}^2 \frac{v_3}{v_1} - \lambda_{11} \frac{v_1^2}{2} - \frac{v_2^2}{2} (\lambda_{12} + \beta_{12} + \alpha_{12}) - \frac{v_3^2}{2} (\lambda_{13} + \beta_{13} + \alpha_{13}), \quad (76)$$

$$m_{33}^2 = m_{13}^2 \frac{v_1}{v_3} - \lambda_{33} \frac{v_3^2}{2} - \frac{v_2^2}{2} (\lambda_{23} + \beta_{23} + \alpha_{23}) - \frac{v_1^2}{2} (\lambda_{13} + \beta_{13} + \alpha_{13}), \quad (77)$$

$$m_{22}^2 = -\lambda_{22} \frac{v_2^2}{2} - \frac{v_3^2}{2} (\lambda_{23} + \beta_{23} + \alpha_{23}) - \frac{v_1^2}{2} (\lambda_{12} + \beta_{12} + \alpha_{12}). \quad (78)$$

The mass terms for the charged scalars are given by

$$\mathcal{L} \supseteq (\phi_1^-, \phi_2^-, \phi_3^-) \phi_{matrix} \begin{pmatrix} \phi_1^+ \\ \phi_2^+ \\ \phi_3^+ \end{pmatrix}, \quad (79)$$

where the mass squared matrix for the charged scalars is given by

$$\phi_{matrix} = \begin{pmatrix} m_{13}^2 \frac{v_3}{v_1} - \frac{v_2^2}{2} A_{12} - \frac{v_3^2}{2} A_{13} & \frac{v_1 v_2}{2} A_{12} & -m_{13}^2 + \frac{v_1 v_3}{2} A_{13} \\ \frac{v_1 v_2}{2} A_{12} & -\frac{v_1^2}{2} A_{12} - \frac{v_3^2}{2} A_{23} & \frac{v_2 v_3}{2} A_{23} \\ -m_{13}^2 + \frac{v_1 v_3}{2} A_{13} & \frac{v_2 v_3}{2} A_{23} & m_{13}^2 \frac{v_1}{v_3} - \frac{v_1^2}{2} A_{13} - \frac{v_2^2}{2} A_{23} \end{pmatrix}, \quad (80)$$

and the following definitions where made

$$A_{12} = \alpha_{12} + \beta_{12}, \quad A_{13} = \alpha_{13} + \beta_{13}, \quad A_{23} = \alpha_{23} + \beta_{23}.$$

There is a zero eigenvalue corresponding to the charged Goldstone boson G^\pm which gets eaten by the W^\pm . The mass squared of the charged Higgs particles is given by

$$m_\pm = \frac{1}{4v_1 v_3} \left(2m_{13}^2 v_3^2 - v_1 v_3 (v_1^2 A_{13} + v_2^2 A_{23} + A_{12} v_1^2 + (A_{13} + A_{23}) v_3^2) \right. \\ \left. \pm \sqrt{(-4v_1 v_3 v^2 (A_{23} v_3^2 (-2m_{13}^2 + A_{13} v_1 v_3) + A_{12} v_1 (-2m_{13}^2 v_1 + A_{13} v_1^2 v_3 + A_{23} v_2^2 v_3)) \right.} \\ \left. + (-2m_{13}^2 (v_1^2 + v_3^2) + v_1 v_3 (A_{13} v_1^2 + A_{23} v_2^2 + A_{12} (v_1^2 + v_2^2) + (A_{13} + A_{23}) v_3^2))^2 \right), \quad (81)$$

where we defined $v^2 = v_1^2 + v_2^2 + v_3^2$, $v_{ij}^2 = v_i^2 + v_j^2$, for $i, j = 1, 2, 3$.

The mass terms for the pseudoscalars are given by

$$\mathcal{L} \supseteq (\eta_1, \eta_2, \eta_3) \eta_{matrix} \begin{pmatrix} \eta_1 \\ \eta_2 \\ \eta_3 \end{pmatrix}, \quad (82)$$

with

$$\eta_{matrix} = \begin{pmatrix} m_{13}^2 \frac{v_3}{v_1} - v_2^2 \beta_{12} - v_3^2 \beta_{13} & v_1 v_2 \beta_{12} & -m_{13}^2 + v_1 v_3 \beta_{13} \\ v_1 v_2 \beta_{12} & -v_1^2 \beta_{12} - v_3^2 \beta_{23} & v_2 v_3 \beta_{23} \\ -m_{13}^2 + v_1 v_3 \beta_{13} & v_2 v_3 \beta_{23} & m_{13}^2 \frac{v_1}{v_3} - v_1^2 \beta_{13} - v_2^2 \beta_{23} \end{pmatrix}, \quad (83)$$

there is a zero eigenvalue corresponding to the neutral Goldstone boson G^0 . The mass-squared of the physical pseudoscalar is

$$m_A = -\frac{1}{2v_1 v_3} \left(-m_{13}^2 v_3^2 + v_1 v_3 (v_1^2 (\beta_{12} + \beta_{13}) + v_2^2 (\beta_{12} + \beta_{23}) + v_3^2 (\beta_{13} + \beta_{23})) \right. \\ \left. \pm \sqrt{(-4v_1 v_3 v^2 (-m_{13}^2 (v_1^2 \beta_{12} + v_3^2 \beta_{23}) + v_1 v_3 (v_1^2 \beta_{12} \beta_{13} + v_2^2 \beta_{12} \beta_{23} + v_3^2 \beta_{13} \beta_{23})) \right.} \\ \left. + (m_{13}^2 v_3^2 - v_1 v_3 (v_1^2 (\beta_{12} + \beta_{13}) + v_2^2 (\beta_{12} + \beta_{23}) + v_3^2 (\beta_{13} + \beta_{23})))^2 \right). \quad (84)$$

Finally the mass terms for the scalars read

$$\mathcal{L} \supseteq (\rho_1, \rho_2, \rho_3) \rho_{matrix} \begin{pmatrix} \rho_1 \\ \rho_2 \\ \rho_3 \end{pmatrix}, \quad (85)$$

where the mass matrix is given by

$$\rho_{matrix} = \begin{pmatrix} m_{13}^2 \frac{v_3}{v_1} + v_1^2 \lambda_{11} & B_{12} v_1 v_2 & -m_{13}^2 + B_{13} v_1 v_3 \\ B_{12} v_1 v_2 & v_2^2 \lambda_{22} & B_{23} v_2 v_3 \\ -m_{13}^2 + B_{13} v_1 v_3 & B_{23} v_2 v_3 & m_{13}^2 \frac{v_1}{v_3} + v_3^2 \lambda_{33} \end{pmatrix}, \quad (86)$$

where we defined

$$B_{12} = A_{12} + \lambda_{12}, \quad B_{13} = A_{13} + \lambda_{13} \quad B_{23} = A_{23} + \lambda_{23}.$$

There is no zero eigenvalue in this case and the scalar masses are given by the roots of the cubic characteristic equation of this matrix.

Appendix B: Vacuum stability or bounded from below conditions

As the field value of each of the 12 components of the Higgs doublets go to infinity only the quartic terms of the potential given by (1) become relevant. So we call this term

$$\begin{aligned} V_4 = & \frac{1}{2} \lambda_{11} (\Phi_1^\dagger \Phi_1)^2 + \frac{1}{2} \lambda_{22} (\Phi_2^\dagger \Phi_2)^2 + \frac{1}{2} \lambda_{33} (\Phi_3^\dagger \Phi_3)^2 \\ & + \lambda_{12} \Phi_1^\dagger \Phi_1 \Phi_2^\dagger \Phi_2 + \lambda_{13} \Phi_1^\dagger \Phi_1 \Phi_3^\dagger \Phi_3 + \lambda_{23} \Phi_2^\dagger \Phi_2 \Phi_3^\dagger \Phi_3 \\ & + \frac{\beta_{12}}{2} \left[(\Phi_1^\dagger \Phi_2)^2 + (\Phi_2^\dagger \Phi_1)^2 \right] + \alpha_{12} \Phi_1^\dagger \Phi_2 \Phi_2^\dagger \Phi_1 \\ & + \frac{\beta_{13}}{2} \left[(\Phi_1^\dagger \Phi_3)^2 + (\Phi_3^\dagger \Phi_1)^2 \right] + \alpha_{13} \Phi_1^\dagger \Phi_3 \Phi_3^\dagger \Phi_1 \\ & + \frac{\beta_{23}}{2} \left[(\Phi_2^\dagger \Phi_3)^2 + (\Phi_3^\dagger \Phi_2)^2 \right] + \alpha_{23} \Phi_2^\dagger \Phi_3 \Phi_3^\dagger \Phi_2. \end{aligned} \quad (87)$$

A simple way to obtain necessary conditions on the quartic parameters of the potential is to study its behaviour along specific field directions.

Writing the three Higgs doublets as

$$\Phi_1 = \begin{pmatrix} \phi_1 + i\phi_2 \\ \phi_3 + i\phi_4 \end{pmatrix}, \quad \Phi_2 = \begin{pmatrix} \phi_5 + i\phi_6 \\ \phi_7 + i\phi_8 \end{pmatrix}, \quad \Phi_3 = \begin{pmatrix} \phi_9 + i\phi_{10} \\ \phi_{11} + i\phi_{12} \end{pmatrix}. \quad (88)$$

We consider for example $\phi_3 \rightarrow \infty$ and $\phi_{11} \rightarrow \infty$ and we get

$$V_4 = \phi_{11}^4 \left(\frac{\lambda_{33}}{2} + (\alpha_{13} + \beta_{13} + \lambda_{13}) \left(\frac{\phi_3}{\phi_{11}} \right)^2 + \frac{\lambda_{11}}{2} \left(\frac{\phi_3}{\phi_{11}} \right)^4 \right). \quad (89)$$

By making $\left(\frac{\phi_3}{\phi_{11}} \right)^2 = x$, this can be seen as a simple polynomial of order two, which in order to be positive semi-definite, the following conditions

$$\lambda_{33} \geq 0, \quad \lambda_{11} \geq 0, \quad -\sqrt{\lambda_{11} \lambda_{33}} \leq \alpha_{13} + \beta_{13} + \lambda_{13} < 0, \quad (90)$$

should be satisfied. By doing similarly in all different $\phi_i \phi_j$ planes we get the general necessary stability conditions

$$\lambda_{ii} \geq 0, \quad (91)$$

$$-\sqrt{\lambda_{ii} \lambda_{jj}} \leq B_{ij} < 0, \quad B_{ij} = \alpha_{ij} + \beta_{ij} + \lambda_{ij} \quad (92)$$

$$-\sqrt{\lambda_{ii} \lambda_{jj}} \leq \bar{B}_{ij} < 0, \quad \bar{B}_{ij} = \alpha_{ij} - \beta_{ij} + \lambda_{ij} \quad (93)$$

-
- [1] G. Aad *et al.* [ATLAS Collaboration], Phys. Lett. B **716**, 1 (2012) doi:10.1016/j.physletb.2012.08.020 [arXiv:1207.7214 [hep-ex]].
- [2] S. Chatrchyan *et al.* [CMS Collaboration], Phys. Lett. B **716**, 30 (2012) doi:10.1016/j.physletb.2012.08.021 [arXiv:1207.7235 [hep-ex]].
- [3] S. P. Martin, Adv. Ser. Direct. High Energy Phys. **21**, 1 (2010) [Adv. Ser. Direct. High Energy Phys. **18**, 1 (1998)] [hep-ph/9709356].
- [4] K. Agashe, R. Contino and A. Pomarol, Nucl. Phys. B **719**, 165 (2005) doi:10.1016/j.nuclphysb.2005.04.035 [hep-ph/0412089].
- [5] Z. Chacko, H. S. Goh and R. Harnik, Phys. Rev. Lett. **96**, 231802 (2006) doi:10.1103/PhysRevLett.96.231802 [hep-ph/0506256].
- [6] R. N. Mohapatra and J. C. Pati, Phys. Rev. D **11**, 566 (1975). doi:10.1103/PhysRevD.11.566
- [7] G. C. Branco, P. M. Ferreira, L. Lavoura, M. N. Rebelo, M. Sher and J. P. Silva, Phys. Rept. **516**, 1 (2012) doi:10.1016/j.physrep.2012.02.002 [arXiv:1106.0034 [hep-ph]].
- [8] E. A. Paschos, Phys. Rev. D **15**, 1966 (1977). doi:10.1103/PhysRevD.15.1966
- [9] S. L. Glashow and S. Weinberg, Phys. Rev. D **15**, 1958 (1977). doi:10.1103/PhysRevD.15.1958
- [10] V. Khachatryan *et al.* [CMS Collaboration], Phys. Lett. B **749**, 337 (2015) doi:10.1016/j.physletb.2015.07.053 [arXiv:1502.07400 [hep-ex]].
- [11] CMS Collaboration [CMS Collaboration], CMS-PAS-HIG-16-005.
- [12] G. Aad *et al.* [ATLAS Collaboration], arXiv:1604.07730 [hep-ex].
- [13] W. Altmannshofer, S. Gori, A. L. Kagan, L. Silvestrini and J. Zupan, Phys. Rev. D **93**, no. 3, 031301 (2016) doi:10.1103/PhysRevD.93.031301 [arXiv:1507.07927 [hep-ph]].
- [14] D. Ghosh, R. S. Gupta and G. Perez, Phys. Lett. B **755**, 504 (2016) doi:10.1016/j.physletb.2016.02.059 [arXiv:1508.01501 [hep-ph]].
- [15] F. J. Botella, G. C. Branco, M. N. Rebelo and J. I. Silva-Marcos, Phys. Rev. D **94**, no. 11, 115031 (2016) doi:10.1103/PhysRevD.94.115031 [arXiv:1602.08011 [hep-ph]].
- [16] I. P. Ivanov and E. Vdovin, Phys. Rev. D **86**, 095030 (2012) doi:10.1103/PhysRevD.86.095030 [arXiv:1206.7108 [hep-ph]]. and
- [17] I. P. Ivanov and C. C. Nishi, JHEP **1501**, 021 (2015) doi:10.1007/JHEP01(2015)021 [arXiv:1410.6139 [hep-ph]].
- [18] I. P. Ivanov, V. Keus and E. Vdovin, J. Phys. A **45**, 215201 (2012) doi:10.1088/1751-8113/45/21/215201 [arXiv:1112.1660 [math-ph]].
- [19] V. Keus, S. F. King and S. Moretti, JHEP **1401**, 052 (2014) doi:10.1007/JHEP01(2014)052 [arXiv:1310.8253 [hep-ph]].
- [20] D. Das and U. K. Dey, Phys. Rev. D **89**, no. 9, 095025 (2014) Erratum: [Phys. Rev. D **91**, no. 3, 039905 (2015)] doi:10.1103/PhysRevD.91.039905, 10.1103/PhysRevD.89.095025 [arXiv:1404.2491 [hep-ph]].
- [21] D. Das, U. K. Dey and P. B. Pal, Phys. Lett. B **753**, 315 (2016) doi:10.1016/j.physletb.2015.12.038 [arXiv:1507.06509 [hep-ph]].
- [22] J. Heeck, M. Holthausen, W. Rodejohann and Y. Shimizu, Nucl. Phys. B **896**, 281 (2015) doi:10.1016/j.nuclphysb.2015.04.025 [arXiv:1412.3671 [hep-ph]].
- [23] S. Moretti and K. Yagyu, Phys. Rev. D **91**, 055022 (2015) doi:10.1103/PhysRevD.91.055022 [arXiv:1501.06544 [hep-ph]].
- [24] A. Aranda, C. Bonilla, F. de Anda, A. Delgado and J. Hernandez-Sanchez, Phys. Lett. B **725**, 97 (2013) doi:10.1016/j.physletb.2013.06.047 [arXiv:1302.1060 [hep-ph]].
- [25] A. Aranda, J. E. Barradas-Guevara, A. Cordero-Cid, F. de Anda, A. Delgado, O. Felix-Beltran and J. Hernandez-Sanchez, arXiv:1404.7829 [hep-ph].
- [26] A. Crivellin, G. D'Ambrosio and J. Heeck, Phys. Rev. D **91**, no. 7, 075006 (2015) doi:10.1103/PhysRevD.91.075006 [arXiv:1503.03477 [hep-ph]].
- [27] A. G. Akeroyd, S. Moretti, K. Yagyu and E. Yildirim, arXiv:1605.05881 [hep-ph].
- [28] K. Yagyu, Phys. Lett. B **763**, 102 (2016) doi:10.1016/j.physletb.2016.10.028 [arXiv:1609.04590 [hep-ph]].
- [29] K. Yagyu, talk at the Charged Scalar 2016 meeting, Uppsala, Sweden (2016)
- [30] T. P. Cheng and M. Sher, Phys. Rev. D **35**, 3484 (1987). doi:10.1103/PhysRevD.35.3484
- [31] R. A. Diaz, R. Martinez and J. A. Rodriguez, Phys. Rev. D **67**, 075011 (2003) doi:10.1103/PhysRevD.67.075011 [hep-ph/0208117].
- [32] J. F. Gunion and H. E. Haber, Phys. Rev. D **67**, 075019 (2003) doi:10.1103/PhysRevD.67.075019 [hep-ph/0207010].
- [33] H. E. Logan and D. MacLennan, Phys. Rev. D **79**, 115022 (2009) doi:10.1103/PhysRevD.79.115022 [arXiv:0903.2246 [hep-ph]].
- [34] S. Su and B. Thomas, Phys. Rev. D **79**, 095014 (2009) doi:10.1103/PhysRevD.79.095014 [arXiv:0903.0667 [hep-ph]].
- [35] M. Aoki, S. Kanemura, K. Tsumura and K. Yagyu, Phys. Rev. D **80**, 015017 (2009) doi:10.1103/PhysRevD.80.015017 [arXiv:0902.4665 [hep-ph]].
- [36] A. Arhrib, R. Benbrik, C. H. Chen, R. Guedes and R. Santos, JHEP **0908**, 035 (2009) doi:10.1088/1126-6708/2009/08/035 [arXiv:0906.0387 [hep-ph]].
- [37] A. Djouadi and G. Moreau, Eur. Phys. J. C **73**, no. 9, 2512 (2013) doi:10.1140/epjc/s10052-013-2512-9 [arXiv:1303.6591 [hep-ph]].
- [38] M. Sher and K. Thrasher, Phys. Rev. D **93**, no. 5, 055021 (2016) doi:10.1103/PhysRevD.93.055021 [arXiv:1601.03973

- [hep-ph]].
- [39] C. D. Carone and R. Primulando, Phys. Rev. D **80**, 055020 (2009) doi:10.1103/PhysRevD.80.055020 [arXiv:0908.0342 [hep-ph]].
- [40] B. Grinstein and M. B. Wise, Phys. Lett. B **201**, 274 (1988). doi:10.1016/0370-2693(88)90227-4
- [41] M. Ciuchini, G. Degrossi, P. Gambino and G. F. Giudice, Nucl. Phys. B **527**, 21 (1998) doi:10.1016/S0550-3213(98)00244-2 [hep-ph/9710335].
- [42] K. G. Chetyrkin, M. Misiak and M. Munz, Phys. Lett. B **400**, 206 (1997) Erratum: [Phys. Lett. B **425**, 414 (1998)] doi:10.1016/S0370-2693(97)00324-9 [hep-ph/9612313].
- [43] C. Greub, T. Hurth and D. Wyler, Phys. Rev. D **54**, 3350 (1996) doi:10.1103/PhysRevD.54.3350 [hep-ph/9603404].
- [44] M. Misiak *et al.*, Phys. Rev. Lett. **114**, no. 22, 221801 (2015) doi:10.1103/PhysRevLett.114.221801 [arXiv:1503.01789 [hep-ph]].
- [45] M. Neubert, Int. J. Mod. Phys. A **11**, 4173 (1996) doi:10.1142/S0217751X96001966 [hep-ph/9604412].
- [46] E. Barberio *et al.* [Heavy Flavor Averaging Group Collaboration], arXiv:0808.1297 [hep-ex].
- [47] H. E. Haber and H. E. Logan, Phys. Rev. D **62**, 015011 (2000) doi:10.1103/PhysRevD.62.015011 [hep-ph/9909335].
- [48] Y. Grossman, Nucl. Phys. B **426**, 355 (1994) doi:10.1016/0550-3213(94)90316-6 [hep-ph/9401311].
- [49] W. S. Hou, Phys. Rev. D **48**, 2342 (1993). doi:10.1103/PhysRevD.48.2342
- [50] A. Czarnecki and W. J. Marciano, Phys. Rev. Lett. **81**, 277 (1998) doi:10.1103/PhysRevLett.81.277 [hep-ph/9804252].
- [51] M. Bona *et al.* [UTfit Collaboration], Phys. Lett. B **687**, 61 (2010) doi:10.1016/j.physletb.2010.02.063 [arXiv:0908.3470 [hep-ph]].
- [52] J. Dingfelder and T. Mannel, Rev. Mod. Phys. **88**, no. 3, 035008 (2016). doi:10.1103/RevModPhys.88.035008
- [53] A. Djouadi, J. Ellis, R. Godbole and J. Quevillon, JHEP **1603**, 205 (2016) doi:10.1007/JHEP03(2016)205 [arXiv:1601.03696 [hep-ph]].