

2012

Histological Assessment Of The Lobster (*Homarus Americanus*) In The "100 Lobsters" Project

Jeffrey D. Shields

Virginia Institute of Marine Science

Kersten N. Wheeler

Virginia Institute of Marine Science

Jessica Moss

Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Marine Biology Commons](#)

Recommended Citation

Shields, Jeffrey D.; Wheeler, Kersten N.; and Moss, Jessica, "Histological Assessment Of The Lobster (*Homarus Americanus*) In The "100 Lobsters" Project" (2012). *VIMS Articles*. 340.

<https://scholarworks.wm.edu/vimsarticles/340>

This Article is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

HISTOLOGICAL ASSESSMENT OF THE LOBSTER (*HOMARUS AMERICANUS*) IN THE “100 LOBSTERS” PROJECT

JEFFREY D. SHIELDS, KERSTEN N. WHEELER AND JESSICA A. MOSS

Department of Environmental and Aquatic Animal Health, Virginia Institute of Marine Science,
The College of William & Mary, PO Box 1346, Gloucester Point, VA 23062

ABSTRACT The emergence of epizootic shell disease in the American lobster (*Homarus americanus*) has been devastating to the industry in the coastal waters of southern New England. A comprehensive assessment of the disease syndrome, known as the “100 Lobsters” Project, was initiated to examine health and physiological parameters among laboratories involved in the research on lobster shell disease. A histological study of the 100 lobsters was undertaken as part of that assessment. Tissues from 90 lobsters from Rhode Island and 19 lobsters from Maine were examined as a general health assessment of the 100 lobsters. Approximately half the lobsters from Rhode Island were selected because they had frank epizootic shell disease, whereas none of the lobsters from Maine exhibited the syndrome. In addition to epizootic shell disease, the histological findings revealed 3 other idiopathic syndromes—necrotizing hepatopancreatitis, idiopathic blindness, and nonspecific granulomas—in higher prevalences in lobsters from Rhode Island compared with those from Maine. Necrotizing hepatopancreatitis, a newly described disease syndrome in lobsters, was observed in 15% of the lobsters from Rhode Island. Idiopathic blindness was present in 54% of the lobsters from Rhode Island, and 16% of the animals from Maine. This is the first report of the syndrome in lobsters from Maine. None of the idiopathic syndromes was associated with epizootic shell disease. The detection of multiple disease syndromes such as epizootic shell disease, necrotizing hepatopancreatitis, and idiopathic blindness may be indicative of exposure to environmental stressors in Narragansett Bay, RI.

KEY WORDS: necrotizing hepatopancreatitis, idiopathic blindness, contaminants, epizootic shell disease, histology, American lobster, *Homarus americanus*

INTRODUCTION

Several health issues have recently affected the fisheries and management of the clawed lobster, *Homarus americanus* Milne Edwards, in the United States. Mortality events and high levels of morbidity have occurred in commercially important fishing areas. Some problems appear to be environmentally related, such as increased bottom temperatures during summers and the general effects of eutrophication (Pearce & Balcom 2005). Others are possibly the result of intoxication from anthropogenic substances (Zulkowsky et al. 2005), and yet others may result from newly emerging pathogens. In 1999, the pathogen *Neoparamoeba pemaquidensis* emerged in concert with environmental stressors to decimate the commercially important lobster population in western Long Island Sound (LIS) (Mullen et al. 2004, Pearce & Balcom 2005). In 2002, lobsters suffering from calcinosis, a physiological response to temperature stress, were reported from central LIS (Dove et al. 2004), and, at about the same time, ~50% of the lobsters in portions of western LIS were shown to have some form of blindness (Maniscalco & Shields 2006). Another emergent pathogen, *Vibrio fluvialis*, caused focal mortalities off Maine (Tall et al. 2003). From about 1996 to the present, epizootic shell disease (ESD) has sharply increased in lobsters from eastern LIS (Castro & Angell 2000, Castro et al. 2005, Landers 2005, Powell et al. 2005) and Buzzards Bay (Glenn & Pugh 2005), and the syndrome has apparently spread to lobsters from Cape Cod Bay (Glenn & Pugh 2005). It is extremely rare in the waters off Maine (Wilson 2005).

The newly emergent syndrome, known as ESD (Smolowitz et al. 2005a, Smolowitz et al. 2005b), is different from classic, or endemic, shell disease. Notably, the epizootic disease has not

been shown to be transmitted horizontally to healthy lobsters in laboratory experiments (Chistoserdov et al. 2003, Quinn et al. 2012). It has also been associated with bacterial species other than *Vibrio* spp., such as members of the Flavobacteriaceae, particularly *Aquimarina* sp. (Chistoserdov et al. 2005a, Chistoserdov et al. 2005b, Chistoserdov et al. 2012). Further, the lesions form in a manner quite different from those in classic shell disease, causing deep pits to form early in the infection, followed by friable areas of shell (pillars of lattice crystal) that eventually coalesce into large, broad ulcers (Smolowitz et al. 2005a, Smolowitz et al. 2005b, Kunkel et al. 2012). In addition, the erosion of the shell appears to be the result of bacterial degradation of the interstitial matrix of the epicuticle (Smolowitz et al. 2005a, Smolowitz et al. 2005b). Lobsters with the disease syndrome have increased levels of proteases and lipases, presumably from bacterial degradation (Bell et al. 2012), as well as a dysbiotic microbial flora with imbalances in certain community members (Meres et al. 2012).

ESD appears to be reducing both the quantity and quality of commercial landings of lobster (Cobb & Castro 2006, Wahle et al. 2009). It has spread rapidly in the lobster population of eastern LIS and Cape Cod Bay (Castro & Angell 2000, Cobb & Castro 2006). In addition, the incidence of ESD has increased markedly, with focal outbreaks affecting 25–35% of lobster populations in eastern LIS (Castro et al. 2005, Glenn & Pugh 2005, Howell et al. 2005, Landers 2005). From 1997 to 2004, ESD has shown a north–south cline, wherein the disease is least prevalent in lobsters in the Gulf of Maine and most prevalent in lobsters off Rhode Island (20–30%) (Cobb & Castro 2006). Interestingly, there is little occurrence of ESD in western LIS or off the New York Bight. Tagging studies have shown that lobsters with shell disease may rapidly contract the disease again shortly after molting out of it (Castro et al. 2005). It is not known whether these animals carried the agent with them or

whether they were more susceptible to the development of the disease.

The “100 Lobsters” Project was developed as a means to share insights, tissues, and samples among researchers from several institutions for shared research objectives (Shields et al. 2012). As part of the development of the demonstration project, we took various tissue samples from the 100 lobsters for histological assessment. Histological analysis of these tissues represented an opportunity to match histological data with data from a variety of other analyses. Thus, the objectives of this study were to analyze the histology of the animals in the 100 Lobsters Project to give an overview of the general state of each animal and to document any unusual conditions that might be associated with the newly emerging phenomenon of ESD from eastern LIS. A new disease syndrome, necrotizing hepatopancreatitis, was observed in animals from Rhode Island; therefore, a secondary objective was to describe the pathology of this syndrome and analyze its prevalence in relation to ESD and other conditions. An assessment of idiopathic blindness was also undertaken, and this idiopathic syndrome was highly prevalent in lobsters from Rhode Island waters.

MATERIALS AND METHODS

Collection of Animals

Lobsters were collected from Narragansett Bay, RI, in June, July, and October 2008. Animals from Narragansett Bay were collected by commercial lobstermen and the Rhode Island Department of Environmental Management (RIDEM) using unvented, baited traps. Lobsters from Maine were collected from off Mount Desert Rock, December 2008, by personnel from the Maine Department of Marine Resources using commercial gear as just noted. Shortly after capture, lobsters were shipped in Styrofoam coolers packed with blue-ice bricks to the Virginia Institute of Marine Science (VIMS) via overnight express delivery. On arrival, lobsters were sorted into 3 groups: those few that had died and were not fit for further processing, those that were moribund and required immediate dissection and processing, and those to be housed in a refrigerated aquarium (220 gal.) for short-term holding (<3 wk) at 10°C. Animals to be processed were examined for sex, carapace length (CL), shell condition, and injuries. Lobsters were rated as having no signs of shell disease, or intensities that were light, moderate, or heavy, as in Landers (2005) (Fig. 1). Prior to processing and dissection, lobsters were photographed with a digital camera (Olympus 3000) (Fig. 1).

For dissection and processing, hemolymph was drawn with a 27-G. syringe from the juncture of the basis and ischium of the 5th walking leg. The area was swabbed with 70% ethanol prior to bleeding. A few drops of hemolymph were dotted onto TCBS plates (thiosulfate citrate bile salts sucrose agar, specific for cultivation of *Vibrio* spp.) and incubated at room temperature for 24–48 h. Lobsters were then killed by injection of a 1-mL aliquot of ice-cold saturated KCl into the region of the ventral nerve ganglia, as in Battison et al. (2000). The ligaments around the carapace were then severed with a knife, the carapace was removed, and various tissues were dissected for later processing and analyses. For histology, samples (~1 cm² pieces) of the hepatopancreas, heart, epidermis underlying the dorsal carapace, gonad, gill, cuticle, and the entire left eye were excised,




Figure 1. Lobsters from Narragansett Bay, RI. (A, B) Moderate epizootic shell disease (A; lobster RI023), and heavy epizootic shell disease (B; lobster RI027). Bar = 30 mm. (C–F) Histological sections from lobsters without epizootic shell disease. (C) Low magnification view of normal healthy hepatopancreas showing cross-sections of the hepatopancreatic tubules. In the region to the right, B cells are highly vacuolated for digestion. In the region to the left, B cells are quiescent (lobster RI092). Bar = 300 μ m. (D) Healthy hepatopancreatic tubules from lobster RI004. Note the lack of vacuolated B cells. Fixed phagocytes (arrow) in a “rosette” pattern surround an arteriole in the spongy connective tissues between tubules. Bar = 100 μ m. (E) Fixed phagocytes surround an arteriole in the spongy connective tissues in lobster RI009. The reserve inclusion cells (arrows) are basophilic and sparse. Bar = 50 μ m. (F) A region of the hepatopancreas with highly vacuolated B cells (*) in lobster RI007. Bar = 100 μ m. b, basement layer; h, hepatopancreatic tubule.

placed in cassettes, and fixed in Bouin’s solution (standard formulation; Fisher Scientific, Pittsburgh, PA). Other tissues were sampled occasionally as well, including the thoracic muscle, hematopoietic tissue, and antennal gland. After 48 h, the tissues were washed for at least 3 h in running tap water, then stored in 70% ethanol until further processing. Pieces of the gills were decalcified for 2–3 h using the formic acid–sodium citrate method (Luna 1968). Pieces of the shell and eye were decalcified overnight using the same method. After decalcification, each eye was bisected sagittally with a razor. All tissues were processed using paraffin histological techniques and stained with Mayer’s hematoxylin and eosin (Luna 1968). Tissues from a subset of animals were processed through the Leavander modification of Brown and Brenn’s stain for bacteria as well as Wheatley’s modification to Mallory’s trichrome stain (Humason 1979). Prepared sections were examined with an Olympus BX51 compound microscope, and photographs were taken using a Nikon DXM1200 digital camera with aid of the ACT-1 computer program (Nikon).

RESULTS

General Assessments

Lobsters from Rhode Island were collected for the study based on their sex and status of ESD (none, light, moderate, heavy). There were 57 female and 33 male lobsters, totaling 90 lobsters from Rhode Island. Of these, 23 females and 20 males had no signs of ESD ($n = 43$), and 34 females and 13 males had various intensities of shell disease ($n = 47$). Of the animals with ESD, 12 had light infections, 14 had moderate infections, and 21 had heavy infections. Lobsters from Maine were collected to match the relative sizes and sexes of animals from Rhode Island. There were 9 female and 10 male lobsters, totaling 19 animals from Maine; of these, none had signs of ESD.

From the TCBS plates, *Vibrio*-like bacteria were commonly isolated from the hemolymph of animals from Rhode Island and Maine (Table 1). However, lobsters from Maine had a significantly lower prevalence (6%) of *Vibrio*-like bacteria than those from Rhode Island (37%; chi-square = 7.110, $df = 1$, $P = 0.008$). Excluding the animals from Maine, lobsters with ESD (51%) had a higher prevalence of *Vibrio*-like bacteria in their hemolymph than those without shell disease (23%; chi-square = 7.387, $df = 1$, $P = 0.007$; Table 2). However, this pattern was dependent on whether the animals had sustained other injuries, such as limb loss (autotomy) or other types of lesions associated with fishing. That is, animals without injuries showed no differences in the prevalence of *Vibrio*-like bacteria whether they had ESD (36%) or not (25%; chi-square = 0.705, $df = 1$, $P = 0.401$, $n = 19$, 24), whereas animals with injuries and ESD (60%) had a significantly higher prevalence of *Vibrio*-like bacteria than their injured counterparts without shell disease (21%; chi-square = 7.204, $df = 1$, $P = 0.007$, $n = 19$, 28, respectively).

Histological Assessment and Idiopathic Conditions

Several idiopathic conditions were observed in the histological assessments of lobsters from Rhode Island (Tables 1 and 2). Idiopathic conditions are those with no specific or known causality. The most noteworthy finding was the occurrence of necrosis in the hepatopancreas, which was observed in 15% of the animals from Rhode Island. Focal necrosis of individual

TABLE 1.
Presence of idiopathic conditions in lobsters from Rhode Island and Maine.

Condition	Rhode Island	Maine
No. of lobsters	90	19
Vibriosis	38%*	6%
Granulomas	44%*	5%
Possible early calcinosis	3%	0%
Hepatopancreatitis (any form)	15%*	0%
Focal necrosis	9%	0%
Coalescent necrosis (severe)	6%	0%
Filament necrosis in gill	8%	5%
Idiopathic lesions in eyes	54%*	16%
Severity of eye lesions	21.5% ± 26.5%	1.5% ± 4.7%
Acanthocephalan cystacanth	7%	0%

* Chi-square, $P < 0.05$.

TABLE 2.

Presence of idiopathic conditions in lobsters from Rhode Island with epizootic shell disease compared with those without the syndrome.

Condition	No Shell Disease	Epizootic Shell Disease
No. of lobsters	43	47
Vibriosis in hemolymph	23%	51%*
Granulomas	40%	49%
Possible early calcinosis	5%	9%
Hepatopancreatitis (any form)	16%	15%
Focal necrosis	9%	9%
Coalescent necrosis (severe)	7%	6%
Idiopathic lesions in eyes	55%	53%
Severity of eye lesions	17.0% ± 20.4%	25.6% ± 30.6%

Affected lobsters had light, moderate or heavy infestations of epizootic shell disease. * Chi-square, $P < 0.05$.

hepatopancreatic tubules occurred in 9% of the animals, and a severe, coalescing form, termed necrotizing hepatopancreatitis, was found in 6% of the animals (see description in next paragraph). The most common lesions observed in histology were idiopathic blindness arising from the loss of ommatidia in the eyes, followed by the presence of nonspecific, or idiopathic, granulomas in a variety of tissues. Some of the granulomas were present in large quantities or occurred in a number of different organs, a condition tentatively identified as early calcinosis. Localized, or focal, necrosis of gill filaments was observed in a few animals from both Rhode Island and Maine, but the condition was extensive on 1 gill podobranch in only 2 lobsters from Rhode Island (Table 1). For animals from Rhode Island, there were no differences in idiopathic conditions in lobsters with or without ESD, with the exception of *Vibrio*-like bacterial infections in the hemolymph (Table 2). There were no differences in the histology of the gonads (ovaries or testes) between lobsters with and without shell disease. Normal gonadal development was observed in all male and female lobsters (not shown).

Necrotizing Hepatopancreas

The normal architecture of the hepatopancreas was observed in all lobsters. It consisted of healthy hepatopancreatic tubules supported by a thin network of spongy connective tissue cells, interlaced rarely with reserve inclusion cells, with primary arterioles and fixed phagocytes interspersed sparingly between the tubules (Fig. 1). In most animals, the reserve inclusion cells were not abundant, and in postmolt and some intermolt animals, the reserve inclusion cells were depleted, giving the appearance of their absence from the hepatopancreas. The B cells ("blasenzellen," a cell that secretes digestive enzymes) in the hepatopancreas were observed for vacuoles as a possible sign of feeding or metabolic state, but they were too variable for this purpose (Fig. 1C–F). The other cell types—F, E, and R cells—did not appear to be altered in histological samples.

In 15% of the animals from Rhode Island, portions of the hepatopancreatic tubules exhibited pathological necroses that were rated as focal lesions (9%) or coalescent lesions (6%), and




Figure 2. Focal necrosis and coalescent lesions of necrotizing hepatopancreatitis in lobsters from Rhode Island. (A) Focal necrosis of a hepatopancreatic tubule in lobster RI026. Bar = 100 μ m. (B) Focal necrosis on the lateral margin of a tubule, with infiltrating hemocytes and rescaffolding of the connective tissues (*) occurring around the affected tubule. Bar = 100 μ m. (C) Coalescent lesion of necrotizing hepatopancreatitis involving several hepatopancreatic tubules in lobster RI048. Bar = 100 μ m. Note the intensive proliferation or rescaffolding of spongy connective tissues in the affected region (*). (D) Infiltration of granulocytes (arrows) within the spongy connective tissues of a coalescent lesion in lobster RI048. Bar = 50 μ m.

were identified as necrotizing hepatopancreatitis. The focal condition consisted of the necrosis of an entire hepatopancreatic tubule (Fig. 2). The surrounding tissues showed signs of nodulation and encapsulation, and in some cases, the rescaffolding of adjacent spongy connective tissues. Infiltration of granulocytes, the primary cells involved in nodulation and encapsulation, was rarely observed in the focal condition. In animals with coalescent hepatopancreatitis, the condition was more extensive than the focal form, involving at least four or more tubules (Figs. 2–4). The affected tubules showed extensive nodulation and encapsulation of the necrotic area, with the involvement of the basement membranes, extensive infiltration of hemocytes, particularly granulocytes into the surrounding connective tissues, and proliferative rescaffolding of the spongy connective tissue adjacent to necrotic areas (Figs. 3 and 4). Apoptotic cells were present in areas with extensive rescaffolding of the spongy connective tissues (Fig. 3D). In some cases, tubules adjacent to those undergoing nodulation possessed epithelial cells that had clearly undergone metaplasia, with cuboidal epithelial cells replacing the normally squamous E and R cells (Fig. 4). Hepatopancreatic tubules adjacent to necrotic tubules occasionally possessed sloughed cells from the tubules (Fig. 4A and D).

Idiopathic Granulomas

Nonspecific, or idiopathic, granulomas were present in significantly more lobsters from Rhode Island (44%) than in those from Maine (5%; chi-square = 10.263, $df = 1$, $P = 0.001$) (Fig. 4E, F). One animal from Maine and 6 animals from Rhode Island had granulomas present in large numbers in a number of different organs. The granulomas were focal in nature, never presenting as coalescent lesions. Calcinosis was




Figure 3. Coalescent lesions of necrotizing hepatopancreatitis in lobsters from Rhode Island. (A) Coalescent lesion with extensive rescaffolding of the spongy connective tissues (*) in lobster RI044. (B) Necrotic tubule with nodulation of the tubule wall (arrows) and infiltration of hemocytes (*) around the exterior of the nodule in lobster RI044. (C) Nodulation of a necrotic tubule showing extensive encapsulation (arrows) of the necrotic region, with adjacent unaffected tubule (h) in lobster RI044. (D) Rescaffolding of the spongy connective tissues in a coalescent lesion showing the presence of apoptotic cells (arrows) in lobster RI048. (E) Higher magnification of an encapsulated necrotic tubule in lobster RI044. Note the extensive zone of encapsulation (*). (F) Infiltration of granulocytes and encapsulation (arrows) of an affected hepatopancreatic tubule in lobster RI044. Bars in A–C = 100 μ m, and in D–F = 50 μ m.

not ruled out because histological stains for calcium were not done on these animals. Bacterial infections may be associated with the presence of granulomas in the tissues, but there was no association between the presence of *Vibrio*-like bacteria in the hemolymph and the presence or intensity of granulomas in the tissues of lobsters from Rhode Island (chi-square = 1.598, $df = 1$, $P = 0.206$), nor was there a difference when stratified by presence or absence of shell disease (chi-square = 0.804, $df = 1$, $P = 0.370$). In a few cases, the granulomas showed evidence of melanization by the host response (Fig. 4F). Bacteria were not observed in the tissues using the Leaver adaptation of the Brown and Brenn method (Humason 1979).

Epizootic Shell Disease

The pathology of animals infected with ESD was virtually identical to that published by Smolowitz et al. (2005a, 2005b). In the histology, deep, penetrating pits occurred on the cuticle of animals with the syndrome (Fig. 5). Within the pits were friable remnants of the cuticle that appeared as pillars of material in histological sections. Extensive melanization of the




Figure 4. Coalescent lesions of necrotizing hepatopancreatitis in lobster RI001 from Rhode Island. (A) Affected hepatopancreatic tubule showing metaplasia of cuboidal epithelial cells (arrows). Note the apparent sloughed cells of the tubules (h). Bar = 100 μm . (B) Extensive proliferation and rescaffolding of spongy connective tissue (*) adjacent to a necrotic tubule. Bar = 100 μm . (C) Detail of affected lesion showing extensive infiltration around the basement layer of an affected tubule. Note the apparent sloughed cells of the tubules (h). Bar = 50 μm . (D) Intensive nodulation and encapsulation of an affected nodule. Note the sloughed cells of the tubules within (h). The intensely eosinophilic substance may be melanin deposits. Bar = 50 μm . (E) Granuloma (g) at the base of a gill filament in lobster RI071. The entire filament was affected. Bar = 100 μm . Inset: A necrotic granuloma in lobster RI023 causing complete occlusion of the gill filament. Bar = 100 μm . (F) A large, melanized granuloma (g) in the epidermis of lobster RI020. Reserve inclusion cells (arrows) are abundant in the epidermis. Bar = 100 μm .

endocuticle was evident in the larger lesions. Lateral erosion of the endocuticle was not observed in the shell-disease lesions. In several histological sections, large areas of the endocuticle appeared to delaminate from their normal lamellar appearance, but this was likely an artifact of the processing of diseased tissues, which were more friable than healthy ones. The membranous layer of the cuticle was affected by nodulation and the infiltration of hemocytes into the adjacent epidermal area in some animals. Pseudomembranes underlying the affected cuticle were observed in 10 lobsters with severe ESD. In these animals, the medial portion of the membranous layer had moderate nodulation with pseudomembranes occurring between the affected cuticle and the underlying epidermis. The underlying musculature was not affected.

Idiopathic Blindness

Idiopathic lesions in the eye indicated the presence of a nonspecific blindness (hereafter called idiopathic blindness).




Figure 5. Epizootic shell disease in lobsters from Rhode Island. (A) Normal cuticle with underlying epidermis (lobster RI054). Bar = 300 μm . (B) Light infection of epizootic shell disease with an eroded pit that has penetrated through the outer portion of the endocuticle in lobster RI068. Note the lack of undermining laterally into the endocuticle. Bar = 100 μm . (C) More extensive erosion of the outer portion of the endocuticle with the typical “pillarlike” formations in the exo- and endocuticle typical of epizootic shell disease. Bar = 100 μm . (D) Extensive erosion of the outer portion of the endocuticle in lobster RI069. Note the melanization (m) of the inner endocuticle, and intensive nodulation (arrows) of the membranous layer. Note the lack of undermining of the outer endocuticle, as is typical in classic shell disease. Bar = 300 μm . (E) Detail of a lesion from lobster RI019 showing extensive melanization (m) of the endocuticle and intensive nodulation (arrows) of the underlying membranous layer. Bar = 100 μm . (F) Extensive erosion of the cuticle in lobster RI019. A pseudomembrane (p) has formed under the nodulated portion of the membranous layer. The separation of the endocuticle is an artifact. Bar = 100 μm . (G) Pseudomembrane (p) adherent to the nodulated membranous layer in lobster RI042. Note the intensive melanization (m) of the endocuticle. Bar = 100 μm . Endo, endocuticle; Epi, epicuticle; Exo, exocuticle; Mb, membranous layer.

The pathology in animals with idiopathic blindness was virtually identical to that published by Maniscalco and Shields (2006) and Magel et al. (2009). Lesions in the ommatidia of the eye were present in animals from both Rhode Island and Maine (Fig. 6). Only 3 lobsters from Maine exhibited idiopathic lesions, and 2 of these animals showed the disruption of only 1–2 ommatidia, perhaps an early response to the syndrome (Fig. 6C, E, F). The affected ommatidia in these 2 animals showed a posterior shift in their screening pigments medially into the underlying optic nerves. The third animal had a moderate case of the syndrome, with ~20% of the ommatidia affected in a centroid lesion (Fig. 6F). A large number of animals (54%, $n = 48$) from Rhode Island exhibited idiopathic blindness, with




Figure 6. Idiopathic blindness in lobsters from Rhode Island and Maine. (A) Normal, healthy eye from Lobster ME008 from Maine. Note the arrangement of the basal portions of the ommatidia (o) along the basement layer (b). Bar = 300 μ m. (B) Lobster from Narragansett Bay, RI, with obvious discoloration of several ommatidia within the compound eye. This animal was effectively blind. Bar \approx 30 mm. (C) Small idiopathic lesion (arrows) affecting only 1–2 centrally located ommatidia in lobster ME012 from Maine. Note the disruption (*) to the optic nerve basal to the basement layer. Bar = 300 μ m. (D) Moderate-size centroid lesion (*) in lobster RI077 from Rhode Island. There is little damage to the ommatidia in this animal, which is indicative of a mild case of the syndrome. The section plane does not show the effect on the entire affected ommatidia. Bar = 300 μ m. (E) Small, aberrant idiopathic lesion (arrow) in the eye of lobster ME003 from Maine. Bar = 300 μ m. (F) Moderate-size oblique lesion (*) in lobster ME011 from Maine. Note the extensive displacement of screening pigment (arrows) into the optic nerve region. The section plane does not show the effect on the whole ommatidia. The space between the ommatidia and the basement layer is an artifact. Bar = 300 μ m.

6 cases having a complete or nearly complete loss of the ommatidia, and 4 animals with light cases, less than 10% of the ommatidia affected.

The prevalence of idiopathic blindness was significantly higher in animals from Rhode Island (54%) than in those from Maine (16%). Moreover, the severity of idiopathic blindness was much higher and more variable in animals from Rhode Island (mean, $21.5\% \pm 26.5\%$ of the lineal area; $n = 89$; see Magel et al. 2009, for lineal area) compared with those from Maine (mean, $1.5\% \pm 4.7\%$ of the lineal area; $n = 19$) (untransformed t -test, $t = 3.263$, $df = 106$, $P = 0.001$). The severity data were calculated as a percentage of the lineal area of the cross-section of the eye; hence, the data were arcsin-square root-transformed to reduce the variance. Whereas the transformed data showed significant differences as above, they also

had a very large variance structure (arcsin-square root-transformed data, t -test, $t = 3.388$, $df = 106$, $P = 0.001$). In the lobsters from Rhode Island, the prevalence of idiopathic blindness was not significantly different between animals with shell disease (53%) and those without (54%; chi-square = 0.022, $df = 1$, $P = 0.882$), indicating no relationship between ESD and idiopathic blindness.

DISCUSSION

The histological assessment of lobsters in the 100 Lobsters Project revealed several important findings. First, lobsters from Rhode Island have considerably more idiopathic disease syndromes and significantly greater intensities of disease than their counterparts from Maine. This finding highlights the importance of incorporating animals from a control site from outside the endemic area of disease, because the conditions that give rise to ESD likely lie along a continuum of exposures within eastern LIS and Narragansett Bay, RI. That is, lobsters from Rhode Island exhibit similar levels of several disease syndromes regardless of whether they have ESD. These disease syndromes are indicative of exposure to stressors or etiological agents that do not occur off Maine.

Second, at least 3 of the idiopathic syndromes in lobsters from Rhode Island can be quite severe. The prevalence of necrotizing hepatopancreatitis in lobsters is startlingly high, and requires further examination. Necrotizing hepatopancreatitis in clawed lobsters is a new discovery. Its etiology in lobsters is unknown, but a similar disease, also known as necrotizing hepatopancreatitis, occurs in penaeid shrimp, and is caused by a rickettsial agent (Freiler et al. 1992, Lightner & Redman 1994). The pathology in the lobsters showed no obvious signs of intracellular bacteria. The presence of necrotic hepatopancreatic tubules has been described in clawed lobsters from Canada (Comeau & Benhalima 2009), but the pathology was not described in sufficient detail to compare it with necrotizing hepatopancreas from the current study. The consequences of necrotizing hepatopancreatitis to lobster health are unknown, but the damage to the affected parts of the hepatopancreas is certainly a complete loss of function. However, the histological samples represented only a small portion of the hepatopancreas; thus, the extent of the disease in individuals was not fully assessed.

The second syndrome with severe effects is idiopathic blindness. Using physiological assessments, Magel et al. (2009) showed that animals with light to moderate cases of this syndrome ($\sim 20\%$ affected areas) have nearly complete loss of vision in the affected eye. Given the prevalence of the syndrome in lobsters in the current study, and in those by Maniscalco and Shields (2006) and Magel et al. (2009), we estimate that nearly half of all adult lobsters in LIS have some form of blindness. The current study extends the range of idiopathic blindness eastward into Narragansett Bay, Block Island Sound, and potentially into the Gulf of Maine, albeit at low levels in the latter. The cause of the idiopathic blindness remains unknown. Maniscalco and Shields (2006) speculated that environmental contaminants released from sediments or multiple stressors during hypoxic events might be involved in blindness.

The third syndrome with severe effects is ESD. Several of the animals in the current study had severe ESD, with extensive erosion of the cuticle covering up to 50% of the body surface

(e.g., Fig. 1B). The host response to the syndrome involved infiltration of hemocytes, intensive nodulation, and incipient melanization of the affected cuticle. In extreme cases, an extensive pseudomembrane developed adjacent to the membranous layer of the cuticle. The latter may impinge on the ability of the lobster to molt, particularly if the pseudomembrane causes adherence to the molt (Smolowitz et al. 2005a, 2005b, Stevens 2009).

Third, 2 syndromes—idiopathic blindness and necrotizing hepatopancreatitis—indicate that lobsters from Rhode Island are likely being exposed to environmental contaminants. In fact, several metals such as cadmium, chromium, arsenic, and copper have been associated with shell disease in other crustaceans (Doughtie et al. 1983, Andersen et al. 2000). Similar syndromes (cataract, fin rot, liver cirrhosis, and hepatic cancers) have been reported in estuarine fishes living in sites heavily contaminated by polycyclic aromatic hydrocarbons (PAHs) and metals (Vogelbein et al. 1990, Huggett et al. 1992). Several metals have been analyzed in tissues from the 100 lobsters as well as from other lobsters with and without ESD (LeBlanc & Prince 2012). Arsenic, cadmium, cobalt, and copper were present in the hepatopancreas in relatively higher concentrations in lobsters from contaminated sites, but there was no relationship with ESD (LeBlanc & Prince 2012). Copper was notably higher in lobsters from Narragansett Bay and eastern LIS than in animals from other regions, but there was no apparent relationship between it and signs of ESD. However, several metals, including those mentioned, were found in sediments from the RI collection site, and their concentrations were similar to or exceeded sediment quality guidelines (LeBlanc & Prince 2012). Although PAHs were not examined in the current study, LIS and Buzzards Bay are known to have high levels of environmental contaminants, including PCBs, pesticides, metals, and PAHs (Turgeon & O'Conner 1991, Hartmann et al. 2004, Hartmann et al. 2005, Calabretta & Oviatt 2008, Mitch & Anisfeld 2010, Morgan & Lohmann 2010). In addition, a point source contaminant may be involved because in January 1996, there was a major spill of no. 2 fuel oil off Rhode Island during a storm that caused an estimated 9 million lobster mortalities (*North Cape* spill, NOAA 2009a, NOAA 2009b), with circumstantial emergence of ESD in animals in Narragansett Bay and Block Island Sound in summer 1996, and thereafter. However, it is not clear whether oil contaminants from no. 2 fuel oil, a relatively light oil, was swept into Narragansett Bay, nor is it clear that it would remain in the sediments for more than 13 y.

Pesticides are also of concern, given their annual use against mosquitoes and crop pests. Accordingly, Biggers and Laufer (2004) and Laufer et al. (2005) found that lobsters within LIS and nearby areas had high levels of alkyl phenols, and that bottom sediments had higher than normal levels of these compounds in areas where diseased lobsters resided. Jacobs et al. (2012) found significant association between alkyl phenol contamination and shell disease for lobsters from central and eastern LIS, but not in animals from Rhode Island. More recently Laufer et al. (2012) have shown that alkyl phenols interfere with tyrosine cross-linking, a process important for sclerotization, in the newly molted cuticle of lobsters. Thus, there is a sound basis for multiple stressors interacting at several different levels in the etiology and severity of ESD. The role of multiple stressors and host susceptibility to ESD has been

examined in a hypothetical model that provides a solid framework for the further exploration of causality and etiology (Tlusty et al. 2007).

Contaminants may not play as important a role in shell disease as some have suggested. Boston Harbor is a heavily contaminated embayment, yet there is a low prevalence of ESD in that region (Glenn & Pugh 2006), albeit endemic, or classic, shell disease is relatively common in lobsters harvested there (Estrella 1984). Interestingly, ESD does not occur in western LIS, an anthropogenically affected area that experiences high summer temperatures (>25°C) and environmental contaminants. These discrepancies indicate that the link between contaminants and shell disease is not consistent, or that focal contaminants through oil spills or other sources may be important to the etiology of the syndrome or that selection pressures have already operated on lobsters in that heavily impacted area. Regardless, the nature of the internal idiopathic syndromes observed from lobsters from Narragansett Bay, RI, is more like that observed in other animals exposed to contaminants (e.g., granulomatous lesions without microbial involvement) rather than to infectious agents. Indeed, some contaminants can cause subtle effects in marine animals, such as immunosuppression (Arkoosh et al. 1998) and reduced metabolic responses (Baldwin et al. 2009), that can appear as infectious etiologies in their hosts. Clearly, many factors, either alone or in combination, may lead to the development of ESD in American lobsters (Castro et al. 2006, Tlusty et al. 2007).

Idiopathic blindness represents another indicator of anthropogenic exposure in lobsters harvested from Rhode Island and eastern LIS. Because of the important role of the endocrine glands in the eyestalk of lobsters, and their role in several physiological processes such as molting and growth, we strongly suggest that this syndrome be included as part of a continued lobster health assessment management program so that this disease and its possible northward advancement into the Gulf of Maine can be monitored. Additional sampling from locations within Block Island Sound, central LIS, and elsewhere should be undertaken to determine the extent and range of the syndromes reported here. Clearly, more research is needed on lobster health to determine the factors and thresholds responsible for tipping the balance from normal host conditions to disease.

ACKNOWLEDGMENTS

We thank Tom Angell, Dan Costa, and Steve Seymour for their able collection of lobsters from Rhode Island, and Melissa Smith and Kathleen Reardon, Maine Department of Marine Resources, for obtaining lobsters for us. Barbara Somers and Kathy Castro provided technical assistance. Three reviewers provided valuable comments. This work was supported by the National Marine Fisheries Service as the New England Lobster Research Initiative: Lobster Shell Disease under NOAA grant NA06NMF4720100 to the University of Rhode Island Fisheries Center. The views expressed herein are those of the authors and do not necessarily reflect the views of NOAA or any of its subagencies. The U.S. government is authorized to produce and distribute reprints for government purposes, notwithstanding any copyright notation that may appear hereon. This is contribution no. 3172 from the Virginia Institute of Marine Science.

LITERATURE CITED

- Andersen, L. E., J. Norton & N. Levy. 2000. A new shell disease in the mud crab *Scylla serrata* from Port Curtis, Queensland (Australia). *Dis. Aquat. Organ.* 43:233–239.
- Arkoosh, A. M., E. Casillas, P. Huffman, E. Clemons, J. Evered, J. E. Stein & U. Varanasi. 1998. Increased susceptibility of juvenile Chinook salmon from a contaminated estuary to *Vibrio anguillarum*. *Trans. Am. Fish. Soc.* 127:360–374.
- Baldwin, D. H., J. A. Spromberg, T. K. Collier & N. Scholz. 2009. A fish of many scales: extrapolating sublethal pesticide exposures to the productivity of wild salmon populations. *Ecol. Appl.* 19:2004–2015.
- Battison, A., R. MacMillan, A. MacKenzie, P. Rose, R. Cawthorn & B. Horney. 2000. Use of injectable potassium chloride for euthanasia of American lobsters (*Homarus americanus*). *Comp. Med.* 50:545–550.
- Bell, S. L., B. Allam, A. McElroy, A. Dove & G. T. Taylor. 2012. Investigation of epizootic shell disease in American lobsters (*Homarus americanus*) from Long Island Sound: I. Characterization of associated microbial communities. *J. Shellfish Res.* 31:473–484.
- Biggers, W. J. & H. Laufer. 2004. Identification of juvenile hormone-active alkylphenols in the lobster *Homarus americanus* and in marine sediments. *Biol. Bull.* 206:13–24.
- Calabretta, C. J. & C. A. Oviatt. 2008. The response of benthic macrofauna to anthropogenic stress in Narragansett Bay, Rhode Island: a review of human stressors and assessment of community conditions. *Mar. Pollut. Bull.* 56:1680–1695.
- Castro, K. M. & T. E. Angell. 2000. Prevalence and progression of shell disease in American lobster, *Homarus americanus*, from Rhode Island waters and the offshore canyons. *J. Shellfish Res.* 19:691–700.
- Castro, K., T. Angell & B. Somers. 2005. Lobster shell disease in southern New England: monitoring and research. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 165–172.
- Castro, K. M., J. R. Factor, T. Angell & D. F. Landers, Jr. 2006. The conceptual model of shell disease revisited. *J. Crustac. Biol.* 26:646–660.
- Chistoserdov, A. Y., R. A. Quinn, S. L. Gubbala & R. Smolowitz. 2012. Bacterial communities associated with lesions of shell disease in the American lobster, *Homarus americanus* Milne-Edwards. *J. Shellfish Res.* 31:449–462.
- Chistoserdov, A., R. Smolowitz & A. Hsu. 2003. Bacterial assemblages involved in the development and progression of shell disease in the American lobster. Presented at the Third Long Island Sound Lobster Health Symposium, Bridgeport, Connecticut, March 7, 2003.
- Chistoserdov, A., S. L. Gubbala, R. Smolowitz, F. Mirazol & A. Hsu. 2005a. A microbiological assessment of epizootic shell disease in the American lobster indicates its strictly dermal etiology. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 12–20.
- Chistoserdov, A., R. Smolowitz, F. Mirazol & A. Hsu. 2005b. Culture-dependent characterization of the microbial community associated with epizootic shell disease in American lobster, *Homarus americanus*. *J. Shellfish Res.* 24:741–748.
- Cobb, J. S. & K. M. Castro. 2006. Shell disease in lobsters: a synthesis. Report prepared for the New England Lobster Research Initiative, Kingston, RI: Rhode Island Sea Grant. http://seagrant.gso.uri.edu/lobster_initiative/documents/SD_SynthesisFinal.pdf. 18 pp.
- Comeau, M. & K. Benhalima. 2009. Internal organ pathology of wild American lobster (*Homarus americanus*) from eastern Canada affected with shell disease. *N. Z. J. Mar. Freshw. Res.* 43:257–269.
- Doughtie, D. G., P. J. Conklin & K. R. Rao. 1983. Cuticular lesions induced in grass shrimp exposed to hexavalent chromium. *J. Invertebr. Pathol.* 42:249–258.
- Dove, A. D. M., C. LoBue, P. Bowser & M. Powell. 2004. Excretory calcinosis: a new fatal disease of wild American lobsters *Homarus americanus*. *Dis. Aquat. Organ.* 58:215–221.
- Estrella, B. T. 1984. Black gill and shell disease in the American lobster (*Homarus americanus*) as indicators of pollution in Massachusetts Bay and Buzzards Bay. Boston, MA: Massachusetts Division of Marine Fisheries Publication no. 14049-19-125-5-85-CR. 17 pp.
- Freiler, P. F., R. F. Sis, T. A. Bell & D. H. Lewis. 1992. Microscopic and ultrastructural studies of necrotizing hepatopancreatitis in Pacific white shrimp (*Penaeus vannamei*) cultured in Texas. *Vet. Pathol.* 29:269–277.
- Glenn, R. & T. L. Pugh. 2005. Observations on the chronology and distribution of lobster shell disease in Massachusetts coastal waters. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 141–155.
- Glenn, R. & T. L. Pugh. 2006. Epizootic shell disease in American lobsters (*Homarus Americanus*) in Massachusetts coastal waters: interactions of temperature, maturity, and intermolt duration. *J. Crustac. Biol.* 26:639–645.
- Hartmann, P. C., J. G. Quinn, R. W. Cairns & J. W. King. 2004. The distribution and sources of polycyclic aromatic hydrocarbons in Narragansett Bay surface sediments. *Mar. Pollut. Bull.* 48:351–358.
- Hartmann, P. C., J. G. Quinn, R. W. Cairns & J. W. King. 2005. Depositional history of organic contaminants in Narragansett Bay, Rhode Island, USA. *Mar. Pollut. Bull.* 50:388–395.
- Howell, P., C. Giannini & J. Benway. 2005. Status of shell disease in Long Island Sound. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 106–114.
- Huggett, R. J., P. A. Van Veld, C. L. Smith, W. J. Hargis & W. K. Vogelbein. 1992. The effects of contaminated sediments in the Elizabeth River. In: Burton, G. A. (ed.). Sediment toxicity assessment. Boca Raton, FL: Lewis Publishers. pp. 403–430.
- Humason, G. L. 1979. Animal tissue techniques, 4th ed. San Francisco, CA: W.H. Freeman. 661 pp.
- Jacobs, M., H. Laufer, J. Stuart, M. Chen & X. Pan. 2012. Endocrine disrupting alkyl phenols are widespread in the blood of lobsters from southern New England and adjacent offshore areas. *J. Shellfish Res.* 31:563–571.
- Kunkel, J. G., W. Nagel & M. J. Jercinovic. 2012. An apatite for the American lobster. *J. Shellfish Res.* 31:515–526.
- Landers, D. F. 2005. Prevalence and severity of shell disease in American lobster *Homarus americanus* from eastern Long Island Sound, Connecticut. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 94–97.
- Laufer, H., M. Chen, M. Johnson, N. Demir & J. M. Bobbitt. 2012. The effect of alkylphenols on lobster shell hardening. *J. Shellfish Res.* 31:555–562.
- Laufer, H., N. Demir & X. Pan. 2005. Shell disease in the American lobster and its possible relation to alkylphenols. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 72–25.
- LeBlanc, L. A. & D. Prince. 2012. Metal concentrations in tissues of American lobsters (*Homarus americanus*, Milne-Edwards) with epizootic shell disease. *J. Shellfish Res.* 31:543–553.
- Lightner, D. V. & R. M. Redman. 1994. An epizootic of necrotizing hepatopancreatitis in cultured penaeid shrimp (Crustacea: Decapoda) in northwestern Peru. *Aquaculture* 122:9–18.
- Luna, L. G. 1968. Manual of histologic staining methods of the Armed Forces Institute of Pathology. New York: McGraw-Hill Publications. 200 pp.
- Magel, D. R., J. D. Shields & R. Brill. 2009. Idiopathic lesions and visual deficits in the American lobster (*Homarus americanus*) from Long Island Sound, NY. *Biol. Bull.* 217:95–101.

- Maniscalco, A. M. & J. D. Shields. 2006. Histopathology of idiopathic lesions in the eyes of *Homarus americanus* from Long Island Sound. *J. Invertebr. Pathol.* 91:88–97.
- Meres, N. J., C. C. Ajuzie, M. Sikaroodi, M. Vemulapalli, J. D. Shields & P. M. Gillevet. 2012. Dysbiosis in epizootic shell disease of the American lobster (*Homarus americanus*). *J. Shellfish Res.* 31:463–472.
- Mitch, A. A. & S. C. Anisfeld. 2010. Contaminants in Long Island Sound: data synthesis and analysis. *Estuaries Coasts* 33:609–628.
- Morgan, E. J. & R. Lohmann. 2010. Dietary uptake from historically contaminated sediments as a source of PCBs to migratory fish and invertebrates in an urban estuary. *Environ. Sci. Technol.* 44:5444–5449.
- Mullen, T. E., R. Russell, M. T. Tucker, J. L. Maratea, C. Koerting, L. Hinckley, S. De Guise, S. J. Frasca & R. A. French. 2004. Paramebiasis associated with mass mortality of American lobster *Homarus americanus* in Long Island Sound, USA. *J. Aquat. Anim. Health* 16:29–38.
- NOAA. 2009a. North Cape oil spill. Damage assessment, remediation and restoration program. http://www.darrp.noaa.gov/northeast/north_cape/.
- NOAA. 2009b. North Cape oil spill restoration fact sheet. <http://www.fws.gov/contaminants/restorationplans/NorthCape/NorthCapeFactSheetGeneral.pdf>.
- Pearce, J. & N. Balcom. 2005. The 1999 Long Island Sound lobster mortality event: Findings of the Comprehensive Research Initiative. *J. Shellfish Res.* 24:691–698.
- Powell, P., C. Giannini & J. Benway. 2005. Status of shell disease in Long Island Sound. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 106–114.
- Quinn, R. A., A. Metzler, R. M. Smolowitz, M. Tlusty & A. Y. Chistoserdov. 2012. Exposures of *Homarus americanus* shell to three bacteria isolated from naturally occurring epizootic shell disease lesions. *J. Shellfish Res.* 31:485–493.
- Shields, J. D., K. N. Wheeler, J. Moss, B. Somers & K. Castro. 2012. The “100 Lobsters” Project: a cooperative demonstration project for health assessments of lobsters from Rhode Island. *J. Shellfish Res.* 31:431–438.
- Smolowitz, R., A. Y. Chistoserdov & A. Hsu. 2005a. A description of the pathology of epizootic shell disease in the American lobster, *Homarus americanus* H. Milne Edwards 1837. *J. Shellfish Res.* 24:49–756.
- Smolowitz, R., A. Y. Chistoserdov & A. Hsu. 2005b. Epizootic shell disease in the American lobster, *Homarus americanus*. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 2–11.
- Stevens, B. G. 2009. Effects of epizootic shell disease in American lobster *Homarus americanus* determined using a quantitative disease index. *Dis. Aquat. Organ.* 88:25–34.
- Tall, B. D., S. Fall, M. R. Pereira, M. Ramos-Valle, S. K. Curtis, M. H. Kothary, D. M. Chu, S. R. Monday, L. Kornegay, T. Donkar, D. Prince, R. L. Thunberg, K. A. Shangraw, D. E. Hanes, F. M. Khambaty, K. A. Lampel, J. W. Bier & R. C. Bayer. 2003. Characterization of *Vibrio fluvialis*-like strains implicated in limp lobster disease. *Appl. Environ. Microbiol.* 69:7435–7446.
- Tlusty, M. F., R. M. Smolowitz, H. O. Halvorson & S. E. DeVito. 2007. Host susceptibility hypothesis for shell disease in American lobsters. *J. Aquat. Anim. Health* 19:215–225.
- Turgeon, D. D. & T. P. O’Conner. 1991. Long Island Sound: distributions, trends and effects of chemical contamination. *Estuaries* 14:279–289.
- Vogelbein, W. K., J. W. Fournie, P. A. Van Veld & R. J. Huggett. 1990. Hepatic neoplasms in the mummichog *Fundulus heteroclitus* from a creosote-contaminated site. *Cancer Res.* 50:5978–5986.
- Wahle, R. A., M. Gibson & M. J. Fogarty. 2009. Distinguishing disease impacts from larval supply effects in a lobster fishery collapse. *Mar. Ecol. Prog. Ser.* 376:185–192.
- Wilson, C. 2005. Observations of shell disease in coastal Maine Waters: 2003 and 2004. In: M. F. Tlusty, H. O. Halvorson, R. Smolowitz & U. Sharma, editors. Lobster shell disease workshop. Aquatic Forum Series 05-1. Boston, MA: New England Aquarium. pp. 156–160.
- Zulkowsky, A. M., J. P. Ruggieri, S. A. Terracciano, B. J. Brownawell & A. E. McElroy. 2005. Acute toxicity of resmethrin, malathion and methoprene to larval and juvenile American lobsters (*Homarus americanus*) and analysis of pesticide levels in surface waters after Scourge, Anvil and Altosid application. *J. Shellfish Res.* 24:495–804.