

2004

## Cape Charles Seabird Watch, Fall 2003

J. W. Williams

*The Center for Conservation Biology*, [jwwil2@wm.edu](mailto:jwwil2@wm.edu)

B. J. Paxton

*The Center for Conservation Biology*, [bjpaxt@wm.edu](mailto:bjpaxt@wm.edu)

Follow this and additional works at: [https://scholarworks.wm.edu/ccb\\_reports](https://scholarworks.wm.edu/ccb_reports)

---

### Recommended Citation

Williams, J. W. and B. J. Paxton. 2004. Cape Charles Seabird Watch, Fall 2003. CCBTR-04-16. Center for Conservation Biology Technical Report Series. College of William and Mary, Williamsburg, VA. 19 pp.

This Report is brought to you for free and open access by the Center for Conservation Biology (CCB) at W&M ScholarWorks. It has been accepted for inclusion in CCB Technical Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).

# SEABIRD WATCH

FALL 2003

Northampton County, Virginia

**Bill Williams**

*center for conservation biology  
education director*

**Bryan Watts, PhD**

*center for conservation biology  
director*

**Bart Paxton**

*center for conservation biology  
biologist*

**Jason Wade**

**Elizabeth Rogan**

*seabird watch technicians*

Virginia's first full-time systematic seabird watch was conducted 9 November through 20 December 2003. The project was a collaborative effort among The Center for Conservation Biology of the College of William and Mary, the Virginia Coast Reserve of The Nature Conservancy, the Virginia Department of Mines, Minerals and Energy, the Virginia Department of Environmental Quality, the United States Coast Guard, and the Eastern Shore of Virginia National Wildlife Refuge.

The watch site was the 191 foot tall Cape Charles Lighthouse, located 2 kilometers north of the southern tip of Smith Island, an Atlantic coastal barrier island in Northampton County on Virginia's Eastern Shore. Two full-time technicians, Elizabeth Rogan and Jason Wade, manned the site 32 out of a possible 42 days, logging 184.25 observation hours. They documented the passage of 75,138 birds of 57 species of which 71,298 were seabirds, that is, species that utilize the ocean as a migration corridor and/or as a primary foraging resource. Birds which fall within this designation include Red-throated and Common loons, Horned and Red-necked grebes, Northern Gannet, Great and Double-crested cormorants, 17 duck species, Parasitic and Pomarine jaegers, 5 gull and 5 tern species, Razorbill and Dovekie. The additional 3,840 birds included songbirds, marsh dwelling waterfowl (ducks, geese, swans, and herons), and shorebirds.

Watch data, including metrological conditions such as wind speed and direction, ambient air temperature, percent cloud cover, visibility, relative humidity, and barometric pressure were recorded hourly following a prescribed protocol. Meteorological information was obtained with a Davis Vantage Pro Wireless Weather station. Also recorded were minutes of observation, species flight distance from shore, and flight altitude above the water surface. Reference markers demarcating distances up to 2 kilometers were placed off shore to aid in determining flight distance from shore. Visual observations were made with Leica Televid 20-60 power spotting scopes and Leica 10 X 42 Trinovid binoculars.



EASTERN SHORE OF VIRGINIA  
NATIONAL WILDLIFE REFUGE

Boat transportation for the technicians to and from Smith Island was provided daily by James Clark, a local contractor.

Six volunteers participated in the 2003 project including Virginia residents John Spahr of Waynesboro, Elisa Enders of Portsmouth, Robert Eggleston of Harrisonburg, and Alaina Thomas of Cape Charles. Fletcher Smith from Pecatonica, Illinois, and Jethro Runco from Newton, Kansas, both seasonal ornithologists, also assisted.



## Results

A total of 75,138 birds of 57 species was recorded during 184.25 hours of observation, a rate of 408 birds/hour. Sea ducks accounted for 49 % (34,687) of the total 71,298 seabirds counted. Dark-winged scoter species comprised 85 % (29,590) of the sea duck total and 42 % of the total number of seabirds. Northern Gannets constituted 25 % of the total seabirds with 18,100. Gull species (Laughing, Herring, Ring-billed, Great Black-backed, and Bonaparte's) made up 10% of the total with 7150 birds and terns (Caspian, Royal, Common, and Forster's) comprised another 5 %. Loons, primarily Red-throated, and cormorants represented 5 % and 4 % respectively of the seabirds total.

The highest count-day totals for all species were recorded 16 November (13,877 birds) and 1 (10,192 birds) and 2 (14,931 birds) December. The peak movement of sea ducks occurred 22 November - 4 December, with a maximum count of 2,927 25 November. A significant movement of Northern Gannets took place 16 November and 1-4 December, with a count period high of 4564 on 2 December.

Staff also conducted a parallel hawk count 9-30 November. Approximately, 100 hours were logged during 17 observation days. 137 diurnal raptors of 9 species were recorded of which 34 % (46) were Black Vultures, 24 % (33) were Northern Harriers, 14 % (20) were Bald Eagles and 9 % (12) were Sharp-shinned Hawks. Additionally, 1 Osprey, 6 Turkey Vultures, 2 Cooper's Hawks, 6 Red-tailed Hawks, 6 American Kestrels, and 5 unidentified raptors were noted.

**Table 1. Summary of Seabird Counts for the 2003 Cape Charles Seabird Watch**

<b>Species</b>	<b>Total</b>	<b>% Total</b>	<b>Peak Count</b>	<b>Date</b>
Red-throated Loon	2049	3	307	12/2
Common Loon	392	1	47	12/4
Loon sp.	902	1	131	11/22
Horned Grebe	27	<1	7	12/9
Red-necked Grebe	3	<1	3	11/9
Northern Gannet	18,100	25	4564	12/2
Brown Pelican	878	1	237	11/16
Great Cormorant	44	<1	26	12/4
Double-crested Cormorant	2351	3	295	11/22
Cormorant sp.	181	<1	60	11/15
Great Blue Heron	45	<1	5	11/11 and 12/9
Tundra Swan	133	<1	30	12/9
Snow Goose	335	<1	100	12/16
Brant	558	1	402	12/8
Canada Goose	18	<1	18	12/12
Canvasback	1	<1	1	12/2
Redhead	22	<1	22	12/12
Scaup sp.	2	<1	2	12/18
Long-tailed Duck	7	<1	2	12/9 and 12/12
Black Scoter	3729	5	958	11/10
Surf Scoter	714	1	267	11/16
"Dark-winged" Scoter	29,590	42	2851	11/25
White-winged Scoter	188	<1	39	11/16
Bufflehead	44	<1	12	11/25
Red-breasted Merganser	390	1	131	12/19
Pomarine Jaeger	1	<1	1	12/19
Laughing Gull	153	<1	30	11/22
Bonaparte's Gull	11	<1	3	11/9 and 11/11
Ring-billed Gull	2379	3	641	11/16
Herring Gull	868	1	107	11/15
Great Black-backed Gull	476	1	91	11/16
Gull sp.	3264	5	605	11/15
Caspian Tern	47	<1	18	11/16
Royal Tern	164	<1	54	11/16
Common Tern	39	<1	25	11/15
Forster's Tern	667	1	248	11/16
Tern sp.	2522	4	750	12/3
Dovekie	1	<1	1	11/9
Razorbill	5	<1	3	11/16
Alcid sp.	35	<1	12	11/15