

11-1-2001

Water Hemlock, Beaver Poison *Cicuta maculata* L.

Gene Silberhorn

Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Plant Sciences Commons](#)

Recommended Citation

Silberhorn, G. (2001) Water Hemlock, Beaver Poison *Cicuta maculata* L.. Wetland Flora Technical Reports, Wetlands Program, Virginia Institute of Marine Science. Virginia Institute of Marine Science, College of William and Mary. <http://dx.doi.org/doi:10.21220/m2-z1em-gh75>

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Technical Report



Wetland Flora

No. 01-11 / November 2001

Gene Silberhorn

Water Hemlock Beaver Poison

Cicuta maculata L.

Growth Habit and Diagnostic Characteristics

Water hemlock is a leafy, branching, perennial herb, 3 to 6 feet (1 to 2 m) tall, with pinnately compound, alternate leaves, with toothed margins. The leaves are divided into leaflets arranged along a central axis; those immediately subtending the flower head are usually divided into three parts. Farther down the stem, the compound leaves may be divided into 15 or more leaflets. Stems are erect, hollow, with longitudinal ridges and often mottled purple near the base.

In summer, the plant produces an unusual, radially branched inflorescence (compound umbel) with a multitude of tiny white flowers. The flowering head generally resembles that of Queen Anne's Lace or wild carrot (*Daucus carota*). The common names of this plant are appropriate. There are documented accounts of this plant poisoning animals and humans. A closely related Eurasian plant, poison hemlock (*Conium maculatum*), was introduced to the New World, and was very likely the cause of Socrates' death.

Distribution

Water hemlock is often found in freshwater wetlands throughout much of North America, as well as the Mid-Atlantic Coastal Plain.

Habitat

Cicuta maculata grows in various wetland habitats such as ditches, swales, tidal freshwater marshes, and the edges of swamps. Because it is a perennial, the plant is commonly found in the same location year after year. The author has consistently found it growing in one spot of Sweet Hall Marsh on the Pamunkey River for over 25 years. It is associated with northern wild rice (*Zizania aquatica*) at this location.

Ecological Value/Benefits

Beaver poison has little wildlife value and is a dangerous poison to wetland browsers such as beavers and muskrats.

Wetland Indicator Status

According to the *Revision of the National List of Plant Species that Occur in Wetlands, 1997*, *Cicuta maculata* is classified as an **obligate wetland plant (OBL)**. OBLs are plants that almost always occur in wetlands (> 99% probability).

Cicuta maculata L. -----



Wetlands Program
School of Marine Science
Virginia Institute of Marine Science
College of William and Mary
Gloucester Point, Virginia 23062
Dr. Carl Hershner, Program Director

This report was funded, in part, by the Department of
Environmental Quality's Coastal Resources Management
Program through Grant No. NA97020181-01 of the
National Oceanic and Atmospheric Administration,
Office of Ocean and Coastal Resource Management,
under the Zone Management Act of 1972, as amended.



Illustration by
Kent Forrest

Printed on
recycled
paper: 