

2014

World Health Organization infant and young child feeding indicators and their associations with child anthropometry: a synthesis of recent findings

Andrew D. Jones

Laura E. Smith

Amanda A. Zongrone

Scott B. Ickes

College of William and Mary

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Jones, A. D., Ickes, S. B., Smith, L. E., Mbuya, M. N., Chasekwa, B., Heidkamp, R. A., ... & Stoltzfus, R. J. (2014). World Health Organization infant and young child feeding indicators and their associations with child anthropometry: a synthesis of recent findings. *Maternal & child nutrition*, 10(1), 1-17.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Review Article

World Health Organization infant and young child feeding indicators and their associations with child anthropometry: a synthesis of recent findings

Andrew D. Jones*, Scott B. Ickes[†], Laura E. Smith*, Mduduzi N.N. Mbuya[‡], Bernard Chasekwa[‡], Rebecca A. Heidkamp[§], Purnima Menon[¶], Amanda A. Zongrone* and Rebecca J. Stoltzfus*

*Division of Nutritional Sciences, Cornell University, Ithaca, New York, USA, [†]Department of Kinesiology and Health Sciences, College of William and Mary, Williamsburg, Virginia, USA, [‡]ZVITAMBO Project, Harare, Zimbabwe, [§]The Johns Hopkins University, Baltimore, Maryland, USA, and [¶]Poverty, Health and Nutrition Division, International Food Policy Research Institute, New Delhi, India

Abstract

As the World Health Organization (WHO) infant and young child feeding (IYCF) indicators are increasingly adopted, a comparison of country-specific analyses of the indicators' associations with child growth is needed to examine the consistency of these relationships across contexts and to assess the strengths and potential limitations of the indicators. This study aims to determine cross-country patterns of associations of each of these indicators with child stunting, wasting, height-for-age *z*-score (HAZ) and weight-for-height *z*-score (WHZ). Eight studies using recent Demographic and Health Surveys data from a total of nine countries in sub-Saharan Africa (nine), Asia (three) and the Caribbean (one) were identified. The WHO indicators showed mixed associations with child anthropometric indicators across countries. Breastfeeding indicators demonstrated negative associations with HAZ, while indicators of diet diversity and overall diet quality were positively associated with HAZ in Bangladesh, Ethiopia, India and Zambia ($P < 0.05$). These same complementary feeding indicators did not show consistent relationships with child stunting. Exclusive breastfeeding under 6 months of age was associated with greater WHZ in Bangladesh and Zambia ($P < 0.05$), although CF indicators did not show strong associations with WHZ or wasting. The lack of sensitivity and specificity of many of the IYCF indicators may contribute to the inconsistent associations observed. The WHO indicators are clearly valuable tools for broadly assessing the quality of child diets and for monitoring population trends in IYCF practices over time. However, additional measures of dietary quality and quantity may be necessary to understand how specific IYCF behaviours relate to child growth faltering.

Keywords: infant and young child feeding, WHO feeding indicators, diet diversity, child growth, stunting, wasting.

Correspondence: Dr Andrew D. Jones, 313 Savage Hall, Ithaca, NY 14853, USA. E-mail: adj23@cornell.edu

Introduction

Children who are poorly nourished in the first 2 years of life are at increased risk of mortality and impaired cognitive development, as well as diminished work capacity and chronic disease later in life (Dewey & Begum 2011). Proper care for children is an essential element needed for the healthy growth and develop-

ment of a child (Engle *et al.* 1996). A major component of care is the set of practices caregivers employ to provide breast milk and complementary foods to children in their first years of life (PAHO/WHO 2003). Relatively simple indicators for assessing breastfeeding (BF) practices have been in widespread use since the early 1990s (World Health Organization 1991). The ability to measure and monitor these

Table 1. World Health Organization infant and young child feeding core indicators

Indicator	Description
Breastfeeding indicators	
1. Early initiation of breastfeeding	Proportion of children born in the last 24 months who were put to the breast within 1 h of birth
2. Exclusive breastfeeding under 6 months	Proportion of infants 0–5 months of age who were fed exclusively breast milk during the previous day
3. Continued breastfeeding at 1 year	Proportion of children 12–15 months of age who were fed any breast milk during the previous day
Complementary feeding indicators	
4. Introduction of solid, semi-solid or soft foods	Proportion of infants 6–8 months of age who received solid, semi-solid or soft foods during the previous day
5. Minimum dietary diversity	Proportion of children 6–23 months of age who received foods from 4 or more food groups during the previous day
6. Minimum meal frequency	Proportion of breastfed and non-breastfed children 6–23 months of age who received solid, semi-solid or soft foods (but also including milk feeds for non-breastfed children) the minimum number of times or more during the previous day
7. Minimum acceptable diet	Proportion of children 6–23 months of age who had at least the minimum dietary diversity and minimum meal frequency (apart from breast milk) during the previous day
8. Consumption of iron-rich or iron-fortified foods	Proportion of children 6–23 months of age who received an iron-rich food or iron-fortified food that is specially designed for infants and young children, or that is fortified in the home during the previous day

practices has helped raise awareness of the importance of BF and has facilitated progress in achieving improvements in BF practices worldwide (UNICEF 2006). Yet, despite an ever-growing evidence base supporting the importance of complementary feeding (CF) beginning at 6 months of age, defining simple CF indicators has proven challenging because of the multiple dimensions of CF, the variation in these practices across contexts and the changes in recommended practices that occur from 6 to 23 months of age (Arimond *et al.* 2008; World Health Organization 2008).

In 2008, the World Health Organization (WHO) published a set of population-level IYCF indicators developed in response to the need for simple, practi-

cal indicators of appropriate feeding practices in children aged 6–23 months that could be developed from large-scale survey data to describe trends over time and allow for national and sub-national comparisons not only of BF practices but also CF practices (World Health Organization 2002, 2008). A core set of eight indicators (i.e. three BF and five CF) include measures of dietary diversity, feeding frequency, consumption of iron-rich or iron-fortified foods as well as indicators of appropriate BF practices (Table 1). All of these indicators can be constructed from the existing data collected in surveys such as the Demographic and Health Surveys (DHS) (Arimond *et al.* 2008; ICF Macro 2011). An examination of these practices in 46 countries for which DHS data were available between

Key messages

- The World Health Organization (WHO) infant and young child feeding (IYCF) indicators are simple measures of diet quality with many applications.
- WHO breastfeeding indicators are negatively associated with child height-for-age z-score (HAZ) in many countries, suggesting that these relationships may be prone to reverse causality.
- Indicators of diet quality were positively associated with HAZ in some countries, although child stunting was not consistently associated with these indicators.
- The WHO IYCF indicators lack specificity, and therefore, additional measures of diet quality may be useful for understanding how specific behaviours are associated with child growth.

2002 and 2008 revealed inadequate CF practices among large proportions of respondents, particularly in sub-Saharan Africa (SSA) and South Asia (World Health Organization 2010a). In 11 of the 27 SSA countries analysed, fewer than 30% of children aged 6–23 months achieved a minimally diverse diet, while in no SSA country did greater than 30% of children achieve a minimally acceptable diet according to the WHO indicators.

Previous dietary diversity scores and child feeding indices have been used to assess the quality of IYCF practices in different country contexts and to examine the association of these practices with child anthropometry (Hatløy *et al.* 2000; Arimond & Ruel 2002; Ruel & Menon 2002; Savy *et al.* 2005). These studies have generally found that proper child feeding practices are associated with a lower probability of child stunting (i.e. low height-for-age), an indicator of chronic undernutrition. One recent study pooled data from 14 low-income countries to assess the overall average effect of the WHO IYCF indicators on child anthropometry across all countries rather than measuring the impacts for individual countries (Marriott *et al.* 2012). The authors found that all of the WHO CF indicators, with the exception of the indicator defining minimum meal frequency (MMF), were associated with a significantly lower probability of a child being underweight (i.e. low weight-for-age) and stunted.

Understanding the extent to which indicators of dietary quality predict anthropometric outcomes is important for interpreting the meaning of the measurements arising from these indicators. Especially because the WHO IYCF indicators are meant to be used not only for describing trends in IYCF practices over time but also for identifying populations at risk and evaluating the impacts of interventions, the indicators may not achieve their intended purpose if they do not predict the ultimate outcomes of interest for assessing nutritional risk or impact, namely child growth and development. However, the strength and relevance of an indicator depend on the purpose for which it is intended (Habicht & Pelletier 1990). Individual-level associations with child anthropometry may be unimportant for assessing the relevance of an indicator if the inferences that need to be drawn from the data are not at the individual level, but

rather are at the national or sub-national level (e.g. comparing trends across countries or regions), or are not directly concerned with child anthropometric outcomes (e.g. drawing attention to disparities in child feeding practices for advocacy purposes).

It is only recently that analyses have emerged examining associations between the gamut of the WHO recommended IYCF indicators and child anthropometry. As these indicators are increasingly used in programmatic and research settings to inform assessment, targeting, and monitoring and evaluation efforts, analyses of the associations between these indicators and child anthropometric outcomes are needed to develop an understanding of the strengths and potential limitations of the indicators when used for different purposes. This understanding then may lead to a more informed process for selecting indicators and metrics of child diets and feeding practices in different contexts. Furthermore, country-specific analyses of the relationships between these IYCF indicators and child anthropometry are also needed. Pooling data across countries increases statistical power and improves precision for estimating average effects; however, it may also mask important variability across countries (e.g. feeding trends in a subgroup of countries may disproportionately drive the average effects observed).

We review here the current literature examining the eight core WHO IYCF indicators and their relationships with child anthropometry using country-level data. Our objectives are (1) to examine the patterns of associations across countries of each of the indicators with child stunting and wasting as well as child height-for-age z-score (HAZ) and weight-for-height z-score (WHZ); (2) to assess the consistency of the relationships observed across countries; and based on these analyses, (3) to develop an understanding of the strengths and potential limitations of the indicators when used for different purposes.

Materials and methods

Study selection

We conducted a literature search to identify peer-reviewed journal articles that examined associations

between at least one of the WHO IYCF indicators and at least one of the following child anthropometric outcomes: stunting, wasting, HAZ or WHZ. We searched Google Scholar, PubMed Central and Web of Science for relevant articles from 2008 to 2013, reviewing the titles and abstracts of all identified studies. All citations from relevant articles were examined for additional references and we identified several yet-to-be-published articles through personal contacts with expert colleagues. Articles that examined relationships between IYCF practices and child anthropometry, but did not construct measures of child feeding using the definitions recommended by the WHO (World Health Organization 2008), were not included.

Measurement of variables

All of the studies reviewed used DHS data as their data source. The DHS model survey instruments changed in many countries in 2008 from phase 5 questionnaires (used from 2003 to 2008) to phase 6 questionnaires (used from 2008 to 2013). Only the Kenya and Zimbabwe data presented here were collected using phase 6 questionnaires. Data from all other countries were collected using phase 5 questionnaires.

We review data only on the eight core WHO indicators (Table 1). Early initiation of breastfeeding (EIB) is based on historic recall by mothers of children born in the last 24 months who were put to the breast within 1 h of birth. All other indicators are based on recall of practices from the previous day. For example, exclusive breastfeeding (EBF) among children 0–5 months of age and continued breastfeeding at 1 year (CBF) among children 12–15 months of age are calculated from recall of exclusive and any BF, respectively, the previous day only. Introduction of solid, semi-solid or soft foods (ISF) is the proportion of infants 6–8 months of age who received solid, semi-solid or soft foods the previous day. The minimum dietary diversity (MDD) indicator is calculated based on consumption of at least four of the following seven food groups: (1) grains, roots and tubers; (2) legumes and nuts; (3) dairy products; (4) flesh foods; (5) eggs; (6) vitamin A-rich fruits and vegetables; and (7) other fruits and vegetables. Breastfed infants aged 6–8 and

9–23 months must have received solid, semi-solid or soft foods at least two times and three times, respectively, in the previous day to achieve the MMF. Non-breastfed children aged 6–23 months must receive solid, semi-solid or soft foods or milk feeds at least four times in the previous day to achieve the MMF. The minimum acceptable diet (MAD) indicator is a composite of the MDD and MMF indicators. Breastfed children aged 6–23 months must have had the MDD and MMF the previous day, while non-breastfed children aged 6–23 months must have received at least two milk feedings and had at least the MDD (not including milk feeds) and the MMF during the previous day. The consumption of iron-rich or iron-fortified foods (IRF) indicator is defined as children 6–23 months of age who received, during the previous day, an iron-rich food or a food that was specially designed for infants and young children and was fortified with iron.

All identified studies calculated child anthropometric measures based on the WHO Child Growth Standards (World Health Organization 2010b) and among those studies reporting stunting and wasting data, *z*-score cut-off points of ≤ 2 SD were used to classify low height-for-age (i.e. stunting) and low weight-for-height (i.e. wasting).

Heterogeneity in model specification

We report on regression coefficients, *P*-values and 95% confidence intervals around point estimates from multiple linear regression models that regressed the IYCF indicators on HAZ and WHZ in separate models for each indicator. In models using stunting and wasting as the dependent variable, we present odds ratios, 95% confidence intervals and *P*-values from logistic regression models. In the Cambodia analysis, stunting was the only outcome variable assessed, and in the Ethiopia and Zambia analyses, only HAZ and WHZ were assessed. In all other countries, data were reported from analyses using all four outcome variables. Analyses from all countries used sample weights to adjust standard errors for the complex DHS survey design.

Data are presented only for models that adjusted for potentially confounding variables of the relation-

ship between IYCF practices and child anthropometry. The included covariates differed across country data sets. All studies included child age, child sex and an asset-based wealth measure in models, and all studies, but Cambodia, included maternal education. The DHS does not measure household income or expenditures, but rather calculates an asset-based measure of wealth status using data on household assets, utility services, ownership of agricultural land and employment of domestic servants (Rutstein & Johnson 2004). Household assets are assigned weights generated through principal component analysis, and the resulting asset scores are standardised and used to define population wealth quintiles.

All studies, but Cambodia and Uganda, also included maternal height, maternal body mass index, and maternal antenatal care attendance. Additional covariates used in models by country include the square of child age (Bangladesh, Kenya, Zimbabwe, India), current BF status (Bangladesh, Haiti, India, Kenya, Zimbabwe), urban vs. rural residence (Bangladesh, Haiti, Kenya, Zimbabwe), regional dummy variables (Bangladesh, Ethiopia, Haiti, India, Zambia), number of children (Bangladesh, India, Kenya, Zimbabwe), age of mother at first birth (Bangladesh, India, Kenya, Zimbabwe), current age of mother (Cambodia), birth type (Haiti) and child diarrhoea in the past 2 weeks (Kenya, Uganda, Zimbabwe). The India analysis also included the following covariates not included in other studies: size of child at birth, maternal employment status in the 12 months preceding the survey date, frequency of reading newspaper or magazine, frequency of listening to radio, frequency of watching television, household head's social identity (caste/tribe) and religion, and maternal diet diversity.

We did not conduct a meta-analysis of the results from the identified studies because of the small number of studies identified and their focus on different child anthropometric measures. Previous authors have examined pooled results of these relationships across countries (Marriott *et al.* 2012). This study aims to examine country-specific relationships between the IYCF indicators and child anthropometry and conduct cross-country comparisons. Therefore, we present results as summary tables and make qualita-

tive comparisons across studies. *P*-values <0.05 were considered statistically significant for associations between the IYCF indicators and anthropometric outcomes within each country.

Results

Characteristics of the samples

We identified eight relevant studies that reported data from a total of nine countries (five from sub-Saharan Africa, three from Asia and one from the Caribbean region); one study examined data from both Ethiopia and Zambia (Ali *et al.* 2012) and another examined two consecutive DHS data sets from Cambodia (Marriott *et al.* 2010), although only analyses from the more recent 2005 DHS data set examined in that study are presented. Table 2 reports select sample characteristics from the sample populations in each of the studies.

The sample populations in all countries were predominantly rural. The Ethiopian sample had the highest percentage of rural households (93%) and the lowest percentage was in Haiti (66%). More than three-quarters of the mothers in Ethiopia received no formal education, while in all other countries except India, fewer than one-third of sample mothers never attended school. The proportion of women attending at least some secondary schooling was extremely low in Kenya and Ethiopia at less than 5%, while in Zimbabwe and Bangladesh, this percentage was much larger (64% and 47%, respectively).

In all countries, approximately one-quarter or more of children aged less than 24 months were stunted, with nearly half of all children stunted in Ethiopia and Zambia (47% and 45%, respectively) (Table 3). In all studies that disaggregated data by age grouping, the percentage of stunted children increased with increasing age group: the proportion of stunted children aged 18–23 months was two to three times greater than that of children aged 6–11 months. In Ethiopia and Haiti, 17% and 13% of children aged 0–23 months, respectively, were wasted, as were 21% of children aged 6–23 months in Bangladesh. This percentage was lower among children 0–23 months in Kenya, Zambia and Zimbabwe at 8%, 7% and 5%, respectively. The

Table 2. Select child, maternal and household sample characteristics, by country

Characteristic	Africa				Asia				Caribbean		
	Ethiopia <i>Ali et al.</i> (2012) 2005 DHS	Kenya <i>Smith,</i> <i>Chasekwa</i> 2008–2009 DHS	Uganda <i>Ikes</i> 2006 DHS	Zambia <i>Ali et al.</i> (2012) 2007 DHS	Zimbabwe <i>Mbuya, Smith,</i> <i>Chasekwa</i> 2011 DHS	Bangladesh <i>Zongrone</i> <i>et al.</i> (2012) 2007 DHS	Cambodia <i>Marriott</i> <i>et al.</i> (2010) 2005 DHS	India <i>Menon et al.</i> (2013) 2005–2006 DHS	Haiti <i>Heidkamp</i> 2005–2006 DHS		
Country											
Authors											
Data source											
Sample size	1810	2498	1011	2512	2526	1508	3112	14 257	2286		
Proportion of sample living in rural area (%)	92.6	79.8	89.6	71.5	70.9	–	–	74.7	66.3		
Child											
Age range of children (months) for reported data on demographic characteristics of sample	0–23	0–23	0–23	0–23	0–23	6–23	0–23	0–23	0–23		
Female (%)	–	47.9	49.6	–	49.7	–	49.9	47.9	49.4		
Maternal											
Height (cm)	157 (–)	159 (0.24)	160 (1.2)	166 (–)	160 (0.14)	151 (0.14)	–	152 (–)	158 (0.25)		
BMI (kg m ⁻²)	–	22.32 (0.13)	21.9 (0.15)	–	23.3 (0.10)	–	20.7 (0.07)	19.7 (–)	22.7 (0.25)		
	20.1	9.9	11.2	7.6	6.4	–	–	–	12.7		
	3.5	23.1	12.5	16.7	29.5	–	–	–	17.8		
	–	–	–	–	–	–	–	–	–		
Education	77.2	12.3	21.3	13.7	1.1	22.8	22.5	47.1	29.0		
Primary (%)	18.1	63.1	63.4	63.7	31.6	30.4	59.6	–	44.8		
≥Secondary (%)	4.6	4.8	15.3	22.6	63.9	46.8	17.8	–	26.2		
Mothers attending any antenatal care visit (%)	29.3	92.5	92.5	97.8	87.4	87.4	–	–	50.8		
Household											
Electricity (%)	7.2	15.8	5.5	13.8	32.1	–	–	–	27.4		
Wealth quintiles	22.6	24.0	19.7	24.1	22.2	18.50	26.4	24.9	24.8		
Poorest (%)	21.0	19.8	22.6	22.8	21.0	19.36	21.7	22.3	20.9		
Poorer (%)	21.3	18.0	22.2	21.2	19.3	19.43	18.4	19.6	19.8		
Middle (%)	20.6	18.3	18.0	18.4	21.3	18.24	16.2	18.1	20.1		
Richer (%)	14.3	20.0	17.6	13.5	16.1	24.47	17.3	15.0	14.5		
Richest (%)											

BMI, body mass index; DHS; Demographic and Health Surveys. ‘–’ indicates that data were not reported. In the Haiti data set, $n = 1099$ for maternal height variable and $n = 1106$ for BMI and education calculations. In some data sets, data were reported on multiple children aged 0–23 months per household. In Haiti, there were 180 children aged 0–23 months from the same households (i.e. 2286 children and 2106 households). Data were also reported on multiple children per household in Kenya and Zimbabwe where multiple families commonly constituted a single household (i.e. 2498 children and 2365 households in Kenya, and 2526 children and 1812 households in Zimbabwe).

Table 3. Prevalence of child stunting and wasting, by country

Country	Stunting (%)		Wasting (%)										
	Child age range (months)												
	0–5	6–11	12–17	18–23	6–23	0–23	0–5	6–11	12–17	18–23	6–23	0–23	
Bangladesh	–	–	–	–	33.7 (n = 1508)	–	–	–	–	–	–	–	–
Cambodia	17.2 (n = 291)	17.5 (n = 367)	37.2 (n = 359)	52.7 (n = 378)	–	–	–	–	–	–	–	–	–
Ethiopia	–	–	–	–	–	46.6 (n = 1 079)	–	–	–	–	–	–	17.1
Haiti	15.5 (n = 260)	17.3 (n = 256)	26.7 (n = 312)	38.1 (n = 230)	–	24.1 (n = 1 058)	10.5	17.6	11.1	12.1	–	–	12.7
India	–	–	–	–	–	38.8 (n = 14 257)	–	–	–	–	–	–	26.4
Kenya	10.6 (n = 492)	26.6 (n = 591)	42.9 (n = 510)	45.3 (n = 502)	37.9 (n = 1603)	31.9 (n = 2 095)	9.2	10.8	6.2	4.1	–	–	7.6
Uganda	15.9 (n = 235)	19.4 (n = 273)	37.5 (n = 267)	44.5 (n = 236)	33.2 (n = 777)	29.2 (n = 1 011)	7.9	16.2	13.7	7.5	–	–	11.6
Zambia	–	–	–	–	–	45.4 (n = 1653)	–	–	–	–	–	–	7.4
Zimbabwe	10.0 (n = 553)	15.5 (n = 600)	30.6 (n = 546)	47.3 (n = 347)	28.4 (n = 1493)	23.4 (n = 2046)	4.9	6.1	3.3	5.0	–	–	4.9

Age group-specific sample sizes for the percentage of wasted children are the same as for stunting.

prevalence of wasting in children under 24 months was highest in India at 26%.

Breastfeeding and complementary feeding practices

Approximately 40% of mothers in Bangladesh, Haiti and Uganda initiated BF of their youngest child within 1 h of birth (Table 4a). More than half of mothers reported early BF initiation in Kenya and Zambia (58% and 56%, respectively), and this percentage reached 67% in Ethiopia and Zimbabwe. Only 23% of mothers reported initiating BF early in India. EBF under 6 months was lowest in Bangladesh at 36% and highest in Kenya at 78%. At least four of five children aged 12–15 months in all countries received breast milk during the previous day, and approximately this same proportion of children aged 6–8 months received solid, semi-solid or soft foods during the previous day in all countries, except Ethiopia and India where fewer than two-thirds of children had received these foods (i.e. 61% and 64%, respectively).

All studies, except for Cambodia, reported data on CF practices for children 6–23 months of age combined. Ethiopia demonstrated the poorest achievement of the WHO CF indicators, with only 7% of children receiving a minimally diverse diet during the previous day, 5% achieving the MAD indicator and 9% achieving the IRF indicator (Table 4b). Haiti, India, Uganda and Zimbabwe also showed a low proportion of children meeting these indicators (i.e. in India, Uganda and Zimbabwe, less than a quarter of children met the MDD indicator, and in all four countries, fewer than one in five children received a minimally acceptable diet) with somewhat higher percentages in Zambia, Bangladesh and Cambodia. Only in Bangladesh did the percentage of children aged 6–23 months that achieved the MAD indicator exceed 30%.

The percent of children achieving the MMF indicator was considerably higher in all countries as compared with the MDD indicator, with more than half or nearly half of all children meeting this indicator in most countries.

Table 4a. Percentage of children 0–23 months, disaggregated by age grouping, meeting the eight core WHO IYCF indicators, by country

Age range (months) and sample size	Early initiation of breastfeeding				Exclusive breastfeeding under 6 months				Continued breastfeeding at 1 year		Introduction of solid, semi-solid or soft foods					
	0–11	12–23	0–23	n	0–1	n	2–3	n	4–5	n	0–5	n	12–15	n	6–8	n
	Bangladesh	–	–	42.8	2 096	–	–	–	–	–	–	36.1	–	96.0	–	79.7
Cambodia	34.8	1542	33.8	1529	73.8	226	66.4	275	48.7	245	–	746	89.6	532	82.4	393
Ethiopia	–	–	66.9	1 810	–	–	–	–	–	–	43.0	585	94.8	430	60.7	244
Haiti	44.6	1050	43.5	1035	60.8	166	41.6	214	24.3	198	41.2	577	82.3	408	87.3	296
India	–	–	23	14 257	–	–	–	–	–	–	42	3740	87	3151	64	2064
Kenya	57.2	1267	58.0	1160	89.2	193	83.9	210	63.6	216	78.0	619	79.4	377	85.6	362
Uganda	40.6	508	43.1	503	95.6	72	72.2	84	45.3	78	70.5	235	93.1	202	77.9	139
Zambia	–	–	56.2	2 512	–	–	–	–	–	–	51.4	719	93.1	516	90	344
Zimbabwe	66.2	1358	68.9	1076	85.1	182	70.6	245	53.1	293	67.1	720	81.9	438	87.1	359

Data for breastfed and non-breastfed children are presented together.

Associations between WHO IYCF indicators and stunting/HAZ

EIB was significantly positively associated with HAZ in Haiti ($P < 0.05$). EBF under 6 months was negatively associated with HAZ in seven of the data sets assessed; however, this association was significant only in Ethiopia and Kenya ($P < 0.01$ and $P < 0.05$, respectively) (Table 6). Continued BF at 1 year was also significantly negatively associated with HAZ in Ethiopia as well as Zimbabwe ($P < 0.01$ and $P < 0.05$, respectively), and demonstrated similar, although non-significant, trends in all other countries.

The ISF indicator was positively associated with HAZ in Bangladesh and Zambia ($P < 0.01$), and the odds of stunting were significantly lower for children 6–8 months of age in Bangladesh who achieved this indicator ($P < 0.01$) (Table 5). The MDD indicator was positively associated with HAZ in Bangladesh, India and Zambia ($P < 0.05$), and the odds of stunting were lower in children in India who achieved the MDD indicator ($P < 0.001$). Achieving the MAD indicator was similarly associated with a higher HAZ in these same three countries as well as in Ethiopia ($P < 0.05$). In Zimbabwe, the odds of stunting were significantly lower for children aged 6–23 months who met the MAD indicator ($P < 0.05$). The MMF and IRF indicators were not associated with HAZ or stunting in any of the data sets that assessed these relationships.

Associations between WHO IYCF indicators and wasting/WHZ

EIB, CBF at 1 year, and the MDD and IRF indicators were not associated with wasting or WHZ in any of the data sets. Children in Bangladesh aged 0–5 months who received only breast milk the previous day were significantly less likely to be wasted ($P < 0.05$). The EBF indicator was also associated with higher WHZ in Bangladesh ($P < 0.05$) as well as in Zambia ($P < 0.001$) (Tables 5,6). Zambian and Zimbabwean children 6–8 months of age who were fed solid, semi-solid or soft foods the previous day showed significantly lower WHZ scores ($P < 0.05$). In Uganda, the odds of being wasted were significantly

Table 4b. Percentage of children 0–23 months, disaggregated by age grouping, meeting the eight core WHO IYCF indicators, by country (breastfed and non-breastfed children combined)

Country	Child age (months)	Sample size	Minimum dietary diversity	Minimum meal frequency	Minimum acceptable diet	Consumption of iron-rich/iron-fortified foods
Bangladesh	6–11	451	–	–	–	–
	12–17	610	–	–	–	–
	18–23	447	–	–	–	–
Cambodia	6–23	1 508	45.2	84.5	42.7	51.5
	6–11	790	27.4	62.2	20.9	59.4
	12–17	794	26.2	70.8	20.7	51.5
	18–23	758	56.6	64.8	41.5	91.0
	6–23	2 342	–	–	–	–
	6–11	–	–	–	–	–
Ethiopia	12–17	–	–	–	–	–
	18–23	–	–	–	–	–
	6–23	Variable	7.1 (<i>n</i> = 1227)	54.7 (<i>n</i> = 1116)	5.2 (<i>n</i> = 1116)	8.9 (<i>n</i> = 1227)
Haiti	6–11	–	–	–	–	–
	12–17	Variable	32.4 (<i>n</i> = 600)	42.6 (<i>n</i> = 557)	19.6 (<i>n</i> = 456)	–
	18–23	Variable	35.1 (<i>n</i> = 518)	36.5 (<i>n</i> = 434)	23.9 (<i>n</i> = 204)	–
	6–23	Variable	29.2 (<i>n</i> = 1687)	45.3 (<i>n</i> = 1534)	17.1 (<i>n</i> = 1187)	–
India	6–11	–	–	–	–	–
	12–17	–	–	–	–	–
	18–23	–	–	–	–	–
Kenya	6–23	10 517	16.0	45.0	9.0	22.0
	6–11	Variable	18.6 (<i>n</i> = 679)	74.8 (<i>n</i> = 630)	15.9 (<i>n</i> = 630)	–
	12–17	Variable	30.6 (<i>n</i> = 592)	74.7 (<i>n</i> = 566)	25.7 (<i>n</i> = 566)	–
	18–23	Variable	37.5 (<i>n</i> = 608)	81.3 (<i>n</i> = 592)	33.1 (<i>n</i> = 591)	–
	6–23	Variable	28.7 (<i>n</i> = 1879)	76.9 (<i>n</i> = 1788)	24.7 (<i>n</i> = 1788)	–
	6–11	273	14.9	37.5	9.0	20.2
Uganda	12–17	267	29.4	32.3	10.3	36.4
	18–23	236	25.6	30.8	7.1	39.4
	6–23	777	23.1	33.7	8.9	31.6
	6–11	–	–	–	–	–
Zambia	12–17	–	–	–	–	–
	18–23	–	–	–	–	–
	6–23	Variable	37.4 (<i>n</i> = 1793)	56.3 (<i>n</i> = 1445)	25.1 (<i>n</i> = 1445)	59.5 (<i>n</i> = 1793)
	12–17	Variable	3.3 (<i>n</i> = 600)	55.3 (<i>n</i> = 589)	16.6 (<i>n</i> = 565)	–
	18–23	Variable	24.9 (<i>n</i> = 399)	86.0 (<i>n</i> = 378)	23.2 (<i>n</i> = 295)	–
	6–23	Variable	18.9 (<i>n</i> = 1655)	62.0 (<i>n</i> = 1664)	14.5 (<i>n</i> = 1511)	–

MAD, minimum acceptable diet; MDD, minimum dietary diversity; MMF, minimum meal frequency. In the Kenya data set, data on 122 non-breastfed children who did not achieve the MMF indicator were excluded in calculations of MMF and MAD because data on milk feeds were not available, and therefore, these indicators could not be properly constructed for these children. In the Zimbabwe data set, data on 170 such children were excluded for these two indicators for the same reason. Data for the MAD indicator are presented only for breastfed children in the Bangladesh and Haiti data sets. Data for the MMF indicator in Bangladesh are also presented only for breastfed children. Data were not available in the Haiti, Kenya or Zimbabwe DHS data sets to create the iron-fortified foods indicator. In Bangladesh, disaggregated food group data for flesh foods and eggs were not available, so only six food groups were analysed for the creation of the MDD.

Table 5. Adjusted logistic regression results for the association of each of the eight core WHO YCF indicators with stunting and wasting, by country

Country	Stunting										Wasting									
	Early initiation of breastfeeding (0–24 months)	Exclusive breastfeeding under 6 months (0–5 months)	Continued breastfeeding at 1 year (12–15 months)	Introduction of solid, semi-solid, or soft foods (6–8 months)	Minimum dietary diversity (6–23 months)	Minimum meal frequency (6–23 months)	Minimum acceptable diet (6–23 months)	Consumption of iron-rich/iron-fortified foods (6–23 months)	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n		
Bangladesh	1.23 (–, –)	2 096	1.05 (–, –)	–	0.26** (–, –)	–	0.88 (–, –)	1 508	0.70 (–, –)	1 508	0.88 (–, –)	1 508	0.93 (–, –)	1 508	0.80 (0.52, 1.24)	291	–	–		
Cambodia	–	–	1.00 (0.49, 2.02)	1104	–	–	0.75 (0.56, 1.02)	–	–	–	–	–	–	–	–	–	–	–		
Haiti	0.83 (0.57, 1.20)	996	1.50 (0.64, 3.48)	253	0.98	–	1.21 (0.79, 1.32)	121	0.79 (0.47, 1.32)	753	1.23 (0.79, 1.92)	726	1.12 (0.49, 2.54)	578	–	–	–	–		
India	1.10 (0.98, 1.25)	14 257	1.08 (0.84, 1.40)	3740	1.05 (0.77, 1.45)	3151	0.78 (0.58, 1.06)	2064	0.76*** (0.65, 0.89)	10 517	0.94 (0.83, 1.07)	8802	0.88 (0.71, 1.08)	8802	0.95 (0.83, 1.10)	10 517	–	–		
Kenya	1.13 (0.80, 1.60)	1 944	1.67 (0.54, 5.13)	433	0.76 (0.28, 2.09)	319	1.23 (0.46, 3.27)	300	0.94	1 517	1.09 (0.79, 1.51)	1234	1.07 (0.69, 1.67)	1234	–	–	–	–		
Uganda	0.76 (0.57, 1.02)	1 011	0.55 (0.26, 1.15)	235	0.42 (0.14, 1.28)	202	1.89 (0.44, 8.03)	139	0.88 (0.60, 1.29)	777	0.75 (0.53, 1.06)	777	0.69 (0.38, 1.23)	777	0.99 (0.71, 1.38)	777	–	–		
Zimbabwe	1.06 (0.83, 1.35)	1 980	1.17 (0.51, 2.68)	544	2.13 (0.79, 5.75)	376	1.18 (0.30, 4.57)	289	0.78 (0.57, 1.06)	1 445	0.97 (0.72, 1.33)	1 118	0.58* (0.31, 0.97)	1112	–	–	–	–		
	Stunting										Wasting									
	Early initiation of breastfeeding (0–24 months)	Exclusive breastfeeding under 6 months (0–5 months)	Continued breastfeeding at 1 year (12–15 months)	Introduction of solid, semi-solid, or soft foods (6–8 months)	Minimum dietary diversity (6–23 months)	Minimum meal frequency (6–23 months)	Minimum acceptable diet (6–23 months)	Consumption of iron-rich/iron-fortified foods (6–23 months)	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n	OR (95% CI)	n		
Bangladesh	1.11 (–, –)	2 096	0.52* (–, –)	–	1.17 (–, –)	–	1.58 (–, –)	–	1.15 (–, –)	1 508	0.90 (–, –)	1 508	1.14 (–, –)	1 508	1.02 (–, –)	1 508	–	–		
Haiti	0.75 (0.47, 1.19)	996	1.07 (0.21, 5.45)	253	0.41	–	211	1.39	1.13 (0.59, 2.16)	753	0.77 (0.43, 1.38)	726	0.93 (0.36, 2.39)	578	–	–	–	–		
India	0.93 (0.82, 1.06)	14 257	0.96 (0.76, 1.20)	3740	0.89 (0.61, 1.29)	3151	1.11 (0.82, 1.49)	2064	0.91 (0.75, 1.10)	10 517	0.96 (0.84, 1.10)	8802	0.86 (0.68, 1.10)	8802	1.07 (0.92, 1.26)	10 517	–	–		
Kenya	1.08 (0.69, 1.70)	1 944	1.55 (0.55, 4.33)	437	N/A	N/A	0.58 (0.20, 1.75)	300	0.73	1 517	1.06 (0.57, 1.97)	1234	0.80 (0.35, 1.85)	1234	–	–	–	–		
Uganda	1.10 (0.74, 1.63)	1 011	1.71 (0.54, 5.45)	235	2.07 (0.31, 14.07)	202	0.60 (0.18, 1.95)	139	1.22 (0.72, 2.06)	777	0.61* (0.38, 0.99)	777	0.96 (0.45, 2.02)	777	0.85 (0.52, 1.38)	777	–	–		
Zimbabwe	1.09 (0.66, 1.80)	1 980	N/A	N/A	1.52 (0.22, 10.46)	226	N/A	1.55 (0.71, 3.38)	1.445	N/A	N/A	N/A	2.85* (1.11, 7.3)	1080	–	–	–	–		

OR, odds ratio from adjusted logistic regression models regressing either stunting or wasting on the YCF indicator; CI, confidence interval. N/A, odds ratios could not be calculated because of complete separation of data, or perfect correlation, between the feeding indicator and child wasting. In Kenya, no children complying with the 'continued breastfeeding at 1 year' indicator were wasted. In Zimbabwe, no children complying with the 'EBF under 6 months', 'introduction of solid, semi-solid, or soft foods' and 'minimum meal frequency' indicators were wasted. * <0.05, **<0.01, ***<0.001. Values set in boldface are those for which the observed relationship had a *P*-value <0.05.

Table 6. Adjusted linear regression results for the association of each of the 8 core WHO IYCF indicators with mean HAZ and WHZ, by country

Country	Height-for-age z-score (HAZ)				Weight-for-height z-score (WHZ)										
	Est (95% CI)	n	Est (95% CI)	n	Est (95% CI)	n	Est (95% CI)	n							
	Early initiation of breastfeeding (0–24 months)		Exclusive breastfeeding under 6 months (0–5 months)		Continued breastfeeding at 1 year (12–15 months)		Introduction of solid, semi-solid, or soft foods (6–8 months)		Minimum dietary diversity (6–23 months)		Minimum meal frequency (6–23 months)		Minimum acceptable diet (6–23 months)		Consumption of iron-rich/iron-fortified foods (6–23 months)
Bangladesh	0.07 (–, –)		2.096 (–0.08, –)		–	–	–	–	–	–	–	–	–	–	
Ethiopia	–0.12 (–, –)		1.436 (–0.80, –)		391 (–1.66, –)		374 (0.43, –)		216 (0.40, –)		1508 (0.16, –)		1508 (0.19, –)		1508 (0.06, –)
Haiti	0.20* (0.01, 0.40)		996 (–0.09, –0.52, 0.33)		253 (–0.64, –)		211 (–0.92, –)		121 (0.01, –0.26, 0.27)		753 (0.26, –)		961 (0.43, –)		961 (0.52, –)
India	–0.01 (–0.10, 0.07)		14 257 (–0.12, –0.28, 0.04)		3740 (–0.14, –0.37, 0.08)		3151 (0.13, –0.07, 0.33)		2064 (0.19, –)		10 517 (0.08, –0.004, 0.16)		8802 (0.15, –)		8802 (0.03, –0.06, 0.11)
Kenya	0.04 (–0.15, 0.23)		1 944 (–0.45, –0.89, 0.00)		437 (–0.40, –1.35, 0.55)		319 (0.01, –0.73, 0.75)		300 (0.11, –0.16, 0.37)		1 518 (–0.14, –0.37, 0.08)		1234 (–0.45, –0.33, 0.24)		1234 (–)
Uganda	0.04 (–0.14, 0.23)		1 011 (0.43, –0.02, 0.87)		234 (–0.02, –0.78, 0.73)		201 (0.24, –0.42, 0.92)		138 (0.15, –0.10, 0.41)		776 (0.21, –0.01, 0.43)		776 (0.25, –0.11, 0.61)		776 (–0.08, –0.30, 0.15)
Zambia	0.00 (–, –)		2 133 (–0.25, –)		553 (–0.59, –)		464 (1.10, –)		300 (0.23, –)		1 580 (0.15, –)		1274 (0.27, –)		1274 (0.13, –)
Zimbabwe	0.07 (–0.06, 0.21)		1 980 (–0.23, –0.57, 0.10)		544 (–0.53, –1.06, 0.01)		376 (0.05, –0.44, 0.54)		297 (0.11, –0.10, 0.32)		1 445 (0.02, –0.15, 0.19)		1 118 (0.27, –0.04, 0.57)		1 112 (–)
	Early initiation of breastfeeding (0–24 months)		Exclusive breastfeeding under 6 months (0–5 months)		Continued breastfeeding at 1 year (12–15 months)		Introduction of solid, semi-solid, or soft foods (6–8 months)		Minimum dietary diversity (6–23 months)		Minimum meal frequency (6–23 months)		Minimum acceptable diet (6–23 months)		Consumption of iron-rich/iron-fortified foods (6–23 months)
Bangladesh	–0.01 (–, –)		2 096 (0.29, –)		–	–	–	–	–	–	–	–	–	–	
Ethiopia	–0.20 (–, –)		1 436 (0.53, –)		391 (–0.06, –)		374 (0.18, –)		216 (0.23, –)		1 045 (0.05, –)		961 (0.34, –)		961 (0.07, –)
Haiti	0.16 (–0.05, 0.38)		996 (0.25, –0.12, 0.61)		253 (–0.004, –)		211 (–0.207, –)		121 (–0.02, –0.28, 0.24)		753 (–0.09, –0.34, 0.16)		726 (–0.03, –0.45, 0.39)		578 (–)
India	0.04 (–0.03, 0.11)		14 257 (0.05, –0.11, 0.20)		3740 (0.09, –0.09, 0.27)		3151 (0.03, –0.15, 0.21)		2064 (0.05, –0.03, 0.14)		10 517 (0.02, –0.05, 0.09)		8802 (0.07, –0.03, 0.18)		8802 (0.02, –0.06, 0.10)
Kenya	0.02 (–0.14, 0.19)		1 944 (0.01, –0.33, 0.34)		437 (–0.15, –0.73, 0.42)		319 (–0.05, –0.61, 0.51)		300 (–0.00, –0.21, 0.21)		1 518 (0.10, –0.12, 0.33)		1234 (0.10, –0.16, 0.35)		1234 (–)
Uganda	–0.01 (–0.18, 0.17)		1 011 (0.06, –0.37, 0.50)		234 (0.21, –0.49, 0.90)		201 (0.13, –0.44, 0.70)		138 (0.11, –0.12, 0.34)		776 (0.21, –0.01, 0.41)		776 (0.17, –0.16, 0.51)		776 (0.07, –0.13, 0.27)
Zambia	0.10 (–, –)		2 133 (0.65, –)		553 (0.03, –)		464 (0.94, –)		300 (–0.04, –)		1 580 (0.08, –)		1274 (0.04, –)		1274 (0.07, –)
Zimbabwe	–0.6 (–1.96, 0.07)		1 980 (0.12, –0.17, 0.43)		544 (–0.14, –0.48, 0.20)		376 (–0.39, –0.75, 0.03)		297 (–0.11, –0.32, 0.10)		1 445 (–0.10, –0.24, 0.04)		1 118 (–0.12, –0.45, 0.21)		1 112 (–)

Est, regression coefficient from adjusted multiple linear regression models regressing either HAZ or WHZ on the IYCF indicator. * <0.05 ; ** <0.01 ; *** <0.001 . Values set in boldface are those for which the observed relationship had a P -value <0.05 .

lower among children who met the MMF indicator, and there was a positive association observed between achievement of this indicator and WHZ ($P < 0.05$). In Zimbabwe, the odds of being wasted were significantly higher among children who met the MAD indicator ($P < 0.05$).

Discussion

The patterns of associations between the WHO IYCF indicators and child anthropometry varied widely across the country data sets examined. Among the BF indicators, EBF under 6 months was associated with lower odds of wasting in Bangladesh and higher WHZ in Bangladesh and Zambia. EBF under 6 months and CBF at 1 year were consistently negatively associated with HAZ. Of the CF indicators, the MAD indicator showed the strongest associations with HAZ (i.e. demonstrating significant positive relationships in four of the data sets) and the MDD indicator was positively associated with HAZ in Bangladesh, India and Zambia. The MMF and IRF indicators were not strongly associated with HAZ in any of the data sets. Child stunting was positively associated with the ISF indicator in Bangladesh, the MDD indicator in India and the MAD indicator in Zimbabwe, but was not associated with any other indicator in any country.

Child growth is determined by multiple factors, each is necessary but not sufficient. While the quality of child diets is certainly one of these determinants (Ruel & Menon 2002), proper hygiene practices and access to adequate water, proper sanitation and reliable health services may be even more important determinants in some contexts. Yet, one might expect many of the IYCF practices that the WHO indicators aim to measure to be associated with child anthropometry. Considering first the CF indicators, dietary diversity is associated with child linear growth in many different contexts (Ruel & Menon 2002; Arimond & Ruel 2004; Rah *et al.* 2010). We would therefore expect the MDD and MAD indicators to show positive relationships with stunting and HAZ. To the extent that consumption of iron-rich foods is associated with consumption of animal-source foods or lipid-based nutrient supplements that are fortified

with iron (e.g. Nutributter), we might also expect the IRF indicator to be associated with child linear growth (Rivera *et al.* 2003; Abu-Afarwuh *et al.* 2007). The MMF indicator is intended as a proxy for energy intake from non-breast-milk foods (World Health Organization 2008). If greater energy intakes result in more adequate nutrient intakes generally (Willett 2013), we might expect the MMF indicator to show positive associations with HAZ. However, adequate energy intakes alone are likely not sufficient to improve child linear growth (Allen 1994). The expected relationships between the BF indicators and child anthropometry are not as straightforward. Infants exclusively fed breast milk to 6 months do not show deficits in weight or length gain compared with infants exposed to mixed BF beginning at 3–4 months (Kramer & Kakuma 2004). However, infants achieving the EBF indicator may not necessarily exhibit improved growth compared with non-EBF infants (Kramer *et al.* 2001). Longer duration of any BF has been shown to be associated with greater linear growth (Onyango *et al.* 1999; Simondon *et al.* 2001), although some studies have presented evidence to the contrary (Caulfield *et al.* 1996; Habicht 2000). Therefore, the relationship between the CBF indicator and child HAZ may depend, in part, on the quality of complementary foods fed at 1 year.

Therefore, there is evidence to suggest that the IYCF practices measured by the WHO indicators, especially the CF practices related to the overall quality of child diets, should demonstrate associations with the growth of young children. The WHO IYCF indicators though were designed as simple indicators that could be incorporated into large-scale surveys. These indicators are especially well suited for monitoring trends in diet quality in large-scale data sets wherein detailed dietary data cannot be collected; however, they may not be highly sensitive or specific measures of dietary quality in the analysis of the causal pathways to child growth. Although the relevance of these indicators does not depend in all cases on how well they are associated with child anthropometry at the individual level, understanding these associations is essential for selecting appropriate metrics for different purposes. We examine in more depth below the design of the individual

indicators and their potential strengths and limitations in predicting anthropometric outcomes.

Associations of WHO breastfeeding indicators with child anthropometry

EBF under 6 months was negatively associated with HAZ in nearly all of the studies examined, and the association was statistically significant in Ethiopia and Kenya. The WHO EBF indicator, however, lacks sensitivity (i.e. it may commonly classify children as exclusively breastfed who may have received non-breast-milk liquids or foods prior to the survey) and therefore overestimates the proportion of exclusively breastfed infants (World Health Organization 2008). This overestimation of EBF rates by 1-day recall measures has been observed previously (Zohoori *et al.* 1993; Piwoz *et al.* 1995; Arimond & Ruel 2002). The lack of sensitivity of the EBF indicator could lead to the diminution of a potential positive association between EBF and HAZ. Alternatively, if the misclassification of infants aged 0–5 months who are given non-breast-milk liquids and are growing more poorly is substantial, this misclassification could, in fact, lead to an observed negative association.

Socio-economic status (SES) is known to correlate with both child anthropometric outcomes (Adler & Ostrove 1999; Ruel & Menon 2002) and the practice of EBF (Lawoyin *et al.* 2001; Aidam *et al.* 2005). If not properly controlled for in analyses examining the association of EBF with child anthropometry, SES may confound the observed relationship. The DHS does not provide a stand-alone measure of SES, but rather an asset-based wealth measure. Aside from wealth, type of occupation and level of education are normally associated with SES. These two variables are not included in the DHS wealth measure so as to retain a pure measure of economic status and because these variables may independently influence health status (Rutstein & Johnson 2004). All of the studies included in this synthesis included this same asset-based wealth measure as a covariate in regression analyses, and all but one study (i.e. Cambodia) included maternal education. Although it is possible that uncontrolled domains of SES (e.g. type of occupation) account for the negative associations

observed between EBF and HAZ, this is unlikely given that a carefully developed measure of household wealth was included in models across all countries, maternal education was included as a covariate, and a consistent relationship was observed between the two variables across contexts.¹

While EBF was negatively associated with HAZ, it was positively associated with WHZ in every country, with the relationship achieving statistical significance in Bangladesh and Zambia. This reverse trend has been observed previously (Trowbridge *et al.* 1987) and may be due, in part, to the statistical relationship between child height-for-age and child weight-for-height. As height is in the numerator and denominator of the two metrics, respectively, child linear growth spurts may lead to increases in height-for-age with simultaneous declines in weight-for-height (Golden 2009). Alternatively, WHZ may simply be a more responsive indicator of growth in this early infancy period as compared with HAZ – an indicator of chronic undernourishment for which cumulative growth deficits will be most pronounced later in infancy.

Consistent with previous studies (Caulfield *et al.* 1996; Marquis *et al.* 1997; Simondon *et al.* 2001; Sawadogo *et al.* 2006), findings from all eight data sets demonstrated a negative association between continued BF at 12–15 months and HAZ. This relationship reached statistical significance in Zimbabwe and Ethiopia where continued BF was associated with 0.53 and 1.66 SD lower HAZ, respectively. The authors have suggested that reverse causality underlies the negative associations observed between continued BF and HAZ, such that smaller, malnourished children, often seen as more vulnerable, are weaned later (Simondon *et al.* 2001; Habicht 2002). While the findings from the nine data sets reviewed here seem to support this reverse causality hypothesis, the cross-sectional nature of these studies does not allow for causal inferences to be drawn from the observed relationships.

¹Even in the India models where maternal employment was controlled for, EBF under 6 months showed a negative association with HAZ.

Association of WHO complementary feeding indicators with child anthropometry

Introduction of solid, semi-solid or soft foods was positively associated with HAZ in Zambia and Bangladesh and reduced odds of stunting in Bangladesh. It was also associated, however, with lower child WHZ in Zambia and Zimbabwe. Because this indicator relies on a 24-h recall period, similar to the BF indicators, it is plausible that children who received complementary foods prior to 6 months are, in fact, not excluded from the calculation of the indicator. On the other hand, children who are introduced to solid, semi-solid or soft foods in the 6–8 month window that the indicator examines may not have received those foods the day prior to the survey and therefore may also be misclassified. If so, the indicator will lack specificity and will not accurately reflect the timing of the initial introduction of solid, semi-solid or soft foods.

The MMF indicator, intended as a proxy for dietary energy intake, was positively associated with WHZ in Uganda, but was not strongly associated with any other anthropometric outcome. An appropriate number of daily feedings depends on the energy density of foods fed to children and the amount consumed at each feeding (PAHO/WHO 2003). The cut-offs as defined by the MMF indicator, therefore, would tend to underestimate the needs of children fed complementary foods of low energy density (i.e. approximately less than 0.8 kcal g^{-1} ; Dewey & Brown 2003). On the other hand, as noted earlier, a high number of daily feedings of complementary foods may excessively displace breast milk and deleteriously affect child growth. Capturing these complex dynamics in a single indicator, amenable to large-scale surveys may not be feasible and may contribute to the lack of association observed between the MMF indicator and child anthropometry.

In at least three of the countries where the MAD indicator was positively associated with HAZ, Bangladesh, India and Zambia, the relationship appears to be driven largely by the dietary diversity component of the MAD indicator. Previous studies have similarly observed the important role of dietary diversity in shaping relationships with child anthropometry when

using summary child feeding measures (Arimond & Ruel 2002; Sawadogo *et al.* 2006; Moursi *et al.* 2008).

The MDD indicator was significantly positively associated with HAZ in only three of the studies examined and was significantly negatively associated with stunting only in India. This finding was unexpected given the considerable evidence noted earlier, demonstrating the positive association between more diverse diets and better child growth. It is possible that the small percentage of children achieving the MDD indicator in some countries, Ethiopia in particular (Ali *et al.* 2012), did not allow for sufficient power to detect differences in child anthropometry. Alternatively, the consistent positive relationship observed between dietary diversity and household SES in different contexts (Ferguson *et al.* 1993; Hatløy *et al.* 2000; Hoddinott & Yohannes 2002) may mean that controlling for SES in regression models that estimate the effects of dietary diversity on child anthropometry dampens the strength of the association observed between the two variables. Removing SES from models entirely, however, may lead to biased estimates if SES does not influence child anthropometry entirely through dietary diversity. Assessing multicollinearity and analysing regression models with and without controlling for household SES would help identify this as a potential complicating factor of such analyses.

An analysis of 10 data sets from Africa, Asia and Latin America that served as one input to the development of the WHO IYCF indicators found dietary diversity to be a useful indicator of dietary quality, with the mean micronutrient density adequacy (MMDA)² of diets increasing with increasing food group diversity (FANTA 2006). However, at a cut-off of four food groups, the dietary diversity indicator had a high sensitivity and low specificity for detecting a MMDA of less than 50% (i.e. the indicator would rarely misclassify an inadequate diet as adequate, but

²Mean micronutrient density adequacy is a measure of overall dietary quality, calculated as the mean of individual micronutrient density adequacies in the diet, each capped at 100%. Individual micronutrient density adequacies are calculated as the percentage of the desired nutrient density for that age and breastfeeding status.

would commonly misclassify adequate diets as inadequate). Such an indicator may be appropriate for identifying a population's potential to benefit from a dietary diversification intervention relative to other groups or relative to interventions emphasising other CF practices. The indicator may also be appropriate for assessing the adequacy of programs (Habicht *et al.* 1999) by providing a measuring stick to assess how well programme activities meet expected objectives. However, its frequent misclassification of adequate diets may make it less well suited in evaluation research settings where the impact of interventions on dietary quality or the causal pathway from child diet to growth must be rigorously assessed.

Limitations

This synthesis of findings has several limitations. Although the studies reviewed do not make causal claims, the data presented are cross sectional and the directions of the relationships observed are only hypothesised to be in the stated directions (i.e. feeding practices influence child anthropometry). The causal sequencing of these relationships cannot be determined, however, from the available data. In addition, assessing the associations between model covariates and child anthropometric outcomes across studies was beyond the scope of this synthesis. A fuller treatment of the associations of these potentially confounding variables with child anthropometric outcomes would be useful in future synthesis work. Finally, the studies reviewed were remarkably similar in both their standardised construction of the WHO indicators and their analysis approach. Yet, the different DHS questionnaires used in Kenya and Zimbabwe, differences in model covariates and the minor differences in construction of the CF indicators between countries (Table 4b) may limit, to some extent, the comparability of findings across countries.

Conclusion

The WHO IYCF indicators are clearly valuable tools in both programme and research settings for broadly assessing the quality of child diets and for monitoring population trends in IYCF practices over time. Their development has also served to remove a barrier to

scaling up nutrition programming centred on child feeding by providing simple metrics for monitoring such programmes. This simplicity, however, means that the indicators necessarily simplify a great deal of the complexity inherent to child feeding, especially CF. For research and programmes that require analysis of how the quality of child diets is influenced by upstream determinants or how it affects downstream outcomes such as child growth, stronger proxies of dietary quality might be needed to discern these relationships (e.g. feeding scores or measures based on 7-day food frequency recall data). Further research, in diverse contexts, examining the relationship of these indicators to other metrics of dietary quality and child nutritional status may be beneficial to programme planners, researchers and policy-makers in determining how best to apply them in specific contexts.

Acknowledgements

We would like to acknowledge Tawanda W. Ndapwadza for his contributions to the preparation and analysis of the Zimbabwe data set. We are also grateful to Jean-Pierre Habicht for his thoughtful comments on earlier drafts of this manuscript.

Source of funding

None.

Conflicts of interest

The authors declare that they have no conflicts of interest.

Contributions

SBI conducted the analysis of the Uganda data set. LES, MNNM and BC conducted the analysis of the Zimbabwe data set. LES also conducted the analysis of the Kenya data set. RAH conducted the analysis of the Haiti data set. PM conducted the analyses of the India, Zambia and Ethiopia data sets. AAZ conducted the analysis of the Bangladesh data set. ADJ synthesized the analyses. RJS and PM conceptualized the manuscript. The manuscript was written by ADJ

and edited by SBI, LES, MNNM, RAH, PM, RJS, AAZ, and ADJ.

References

- Abu-Afarwuah S., Lartey A., Brown K.H., Zlotkin S., Briend A. & Dewey K.G. (2007) Randomized comparison of 3 types of micronutrient supplements for home fortification of complementary foods in Ghana: effects on growth and motor development. *American Journal of Epidemiology* **166**, 412–420.
- Adler N.E. & Ostrove J.M. (1999) Socioeconomic status and health: what we know and what we don't. *Annals of the New York Academy of Sciences* **896**, 3–15.
- Aidam B.A., Pérez-Escamilla R., Lartey A. & Aidam J. (2005) Factors associated with exclusive breastfeeding in Accra, Ghana. *European Journal of Clinical Nutrition* **59**, 786–796.
- Ali D., Rawat R., Subandoro A. & Menon P. (2012) Infant and young child feeding (IYCF) practices in Ethiopia and Zambia and their association with child nutrition: analysis of demographic and health survey data. *African Journal of Food, Agriculture, Nutrition and Development* **12**, 5895–5914.
- Allen L.H. (1994) Nutritional influences on linear growth: a general review. *European Journal of Clinical Nutrition* **48** (Suppl. 1), S75–S89.
- Arimond M. & Ruel M.T. (2002) *Progress in Developing an Infant and Child Feeding Index: An Example Using the Ethiopia Demographic and Health Survey 2000*. International Food Policy Research Institute: Washington, DC.
- Arimond M. & Ruel M.T. (2004) Dietary diversity is associated with child nutritional status: evidence from 11 demographic and health surveys. *The Journal of Nutrition* **134**, 2579–2585.
- Arimond M., Daelmans B. & Dewey K. (2008) Indicators for feeding practices in children. *Lancet* **371**, 541–542.
- Caulfield L.E., Bentley M.E. & Ahmed S. (1996) Is prolonged breastfeeding associated with malnutrition? Evidence from nineteen demographic and health surveys. *International Journal of Epidemiology* **25**, 693–703.
- Dewey K.G. & Begum K. (2011) Long-term consequences of stunting in early life. *Maternal & Child Nutrition* **7**, 5–18.
- Dewey K.G. & Brown K.H. (2003) Update on technical issues concerning complementary feeding of young children in developing countries and implications for intervention programs. *Food and Nutrition Bulletin* **24**, 5–28.
- Engle P., Menon P. & Haddad L. (1996) *Care and Nutrition: Concepts and Measurement*. International Food Policy Research Institute: Washington, DC.
- Ferguson E.L., Gibson R.S., Opore Obisaw C., Osei Opore F., Lamba C. & Ounpuu S. (1993) Seasonal food consumption patterns and dietary diversity of rural preschool Ghanaian and Malawian children. *Ecology of Food and Nutrition* **29**, 219–234.
- Food and Nutrition Technical Assistance Project (FANTA) (2006) *Developing and Validating Simple Indicators of Dietary Quality and Energy Intake of Infants and Young Children in Developing Countries: Summary of Findings from Analysis of 10 Data Sets*. Academy for Educational Development: Washington, DC.
- Golden M.H. (2009) Proposed recommended nutrient densities for moderately malnourished children. *Food and Nutrition Bulletin* **30**, S267–S342.
- Habicht J.P. (2000) The association between prolonged breastfeeding and poor growth. In: *Short and Long Term Effects of Breast Feeding on Child Health* (eds B. Koletzko, K.F. Michaelsen & O. Hernell), pp 193–200. Kluwer Academic/Plenum Publishers: New York.
- Habicht J.P. (2002) The association between prolonged breastfeeding and poor growth – what are the implications? *Advances in Experimental Medicine and Biology* **478**, 193–200.
- Habicht J.P. & Pelletier D.L. (1990) The importance of context in choosing nutritional indicators. *The Journal of Nutrition* **120** (Suppl. 11), 1519–1524.
- Habicht J.P., Victora C.G. & Vaughan J.P. (1999) Evaluation designs for adequacy, plausibility and probability of public health programme performance and impact. *International Journal of Epidemiology* **28**, 10–18.
- Hatloy A., Hallund J., Diarra M.M. & Oshaug A. (2000) Food variety, socioeconomic status and nutritional status in urban and rural areas in Koutiala (Mali). *Public Health Nutrition* **3**, 57–65.
- Hoddinott J. & Yohannes Y. (2002) *Dietary Diversity as a Food Security Indicator*. International Food Policy Research Institute: Washington, DC.
- ICF Macro (2011) *Demographic and Health Surveys Methodology: Interviewer's Manual*. ICF Macro: Calverton, MD.
- Kramer M.S. & Kakuma R. (2004) The optimal duration of exclusive breastfeeding: a systematic review. *Advances in Experimental Medicine and Biology* **554**, 63–77.
- Kramer M.S., Chalmers B., Hodnett E.D., Sevkovskaya Z., Dzikovitch I., Shapiro S. *et al.* (2001) Promotion of breastfeeding intervention trial (PROBIT): a randomized trial in the Republic of Belarus. *Journal of the American Medical Association* **285**, 413–420.
- Lawoyin T.O., Olawuyi J.F. & Onadeko M.O. (2001) Factors associated with exclusive breastfeeding in Ibadan, Nigeria. *Journal of Human Lactation* **17**, 321–325.
- Marquis G.S., Habicht J.P., Lanata C.F., Black R.E. & Rasmussen K.M. (1997) Association of breastfeeding and stunting in Peruvian toddlers: an example of

- reverse causality. *International Journal of Epidemiology* **26**, 349–356.
- Marriott B.P., White A.J., Hadden L., Davies J.C. & Wallingford J.C. (2010) How well are infant and young child World Health Organization (WHO) feeding indicators associated with growth outcomes? An example from Cambodia. *Maternal & Child Nutrition* **6**, 358–373.
- Marriott B.P., White A., Hadden L., Davies J.C. & Wallingford J.C. (2012) World Health Organization (WHO) infant and young child feeding indicators: associations with growth measures in 14 low-income countries. *Maternal & Child Nutrition* **8**, 354–370.
- Menon P., Bamezai A., Subandoro A., Ayoya M.A. & Aguayo V. (2013) Age-appropriate infant and young child feeding practices are associated with child nutrition in India: insights from nationally-representative data. *Maternal & Child Nutrition* doi: 10.1111/mcn.12036.
- Moursi M.M., Martin-Prevel Y., Eymard-Duvernay S., Capon G., Treche S., Maire B. & Delpeuch F. (2008) Assessment of child feeding practices using a summary index: stability over time and association with child growth in urban Madagascar. *The American Journal of Clinical Nutrition* **87**, 1472–1479.
- Onyango A.W., Esrey S.A. & Kramer M.S. (1999) Continued breastfeeding and child growth in the second year of life: a prospective cohort study in western Kenya. *Lancet* **354**, 2041–2045.
- PAHO/WHO (2003) *Guiding Principles for Complementary Feeding of the Breastfed Child*. Pan American Health Organization/World Health Organization: Washington, DC.
- Piwoz E.G., Creed De Kanashiro H., Lopez De Romana G., Black R.E. & Brown K.H. (1995) Potential for misclassification of infants' usual feeding practices using 24-hour dietary assessment methods. *The Journal of Nutrition* **125**, 57–65.
- Rah J.H., Akhter N., Semba R.D., de Pee S., Bloem M.W., Campbell A.A. *et al.* (2010) Low dietary diversity is a predictor of child stunting in rural Bangladesh. *European Journal of Clinical Nutrition* **64**, 1393–1398.
- Rivera J.A., Hotz C., González-Cossío T., Neufeld L. & García-Guerra A. (2003) The effect of micronutrient deficiencies on child growth: a review of results from community-based supplementation trials. *Journal of Nutrition* **133**, 4010S–4020S.
- Ruel M.T. & Menon P. (2002) Child feeding practices are associated with child nutritional status in Latin America: innovative uses of the demographic and health surveys. *The Journal of Nutrition* **132**, 1180–1187.
- Rutstein S.O. & Johnson K. (2004) *The DHS Wealth Index. DHS Comparative Reports No. 6*. ORC Macro: Calverton, MD.
- Savy M., Martin-Prevel Y., Sawadogo P., Kameli Y. & Delpeuch F. (2005) Use of variety/diversity scores for diet quality measurement: relation with nutritional status of women in a rural area in Burkina Faso. *European Journal of Clinical Nutrition* **59**, 703–716.
- Sawadogo P.S., Martin-Prevel Y., Savy M., Kameli Y., Traisac P., Traore A.S. *et al.* (2006) An infant and child feeding index is associated with the nutritional status of 6- to 23-month-old children in rural Burkina Faso. *The Journal of Nutrition* **136**, 656–663.
- Simondon K.B., Costes R., Delaunay V., Diallo A. & Simondon F. (2001) Children's height, health and appetite influence mothers' weaning decisions in rural Senegal. *International Journal of Epidemiology* **30**, 476–481.
- Trowbridge F.L., Marks J.S., Lopez De Romana G., Madrid S., Boutton T.W. & Klein P.D. (1987) Body composition of Peruvian children with short stature and high weight-for-height. II. Implications for the interpretation for weight-for-height as an indicator of nutritional status. *The American Journal of Clinical Nutrition* **46**, 411–418.
- UNICEF (2006) *Nutrition Indicators: Exclusive Breastfeeding. A Report Card on Nutrition*. <http://www.unicef.org>. (Number 4). [Online]. Available at: http://www.unicef.org/progressforchildren/2006n4/index_breastfeeding.html (Accessed 18 September 2012).
- Willett W.C. (2013) *Nutritional Epidemiology*. Oxford University Press: New York, NY.
- World Health Organization (1991) *Indicators for Assessing Breastfeeding Practices*. Division of Child Health and Development, World Health Organization: Geneva, Switzerland.
- World Health Organization (2002) *Report of Informal Meeting to Review and Develop Indicators for Complementary Feeding, 3–5 December*. World Health Organization. Department of Child and Adolescent Health and Development and Department of Nutrition for Health and Development: Geneva, Switzerland.
- World Health Organization (2008) *Indicators for Assessing Infant and Young Child Feeding Practices: Part I Definitions*. World Health Organization: Geneva, Switzerland.
- World Health Organization (2010a) *Indicators for Assessing Infant and Young Child Feeding Practices: Part 3 Country Profiles*. World Health Organization: Geneva, Switzerland.
- World Health Organization (2010b) *The WHO Child Growth Standards*. World Health Organization: Geneva, Switzerland.
- Zohoori N., Popkin B.M. & Fernandez M.E. (1993) Breastfeeding patterns in the Philippines: a prospective analysis. *Journal of Biosocial Science* **25**, 127–138.
- Zongrone A., Winkell K. & Menon P. (2012) Infant and young child feeding practices and child undernutrition in Bangladesh: insights from nationally representative data. *Public Health Nutrition* **15**, 1697–1704.