

7-2006

## Nitrogen fixation and release of fixed nitrogen by *Trichodesmium* spp. in the Gulf of Mexico

Margaret R. Mulholland

Peter W. Bernhardt

Cynthia A. Heil

Deborah A. Bronk

*Virginia Institute of Marine Science*

Judith M. O'Neil

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Environmental Sciences Commons](#)

---

### Recommended Citation

Mulholland, Margaret R.; Bernhardt, Peter W.; Heil, Cynthia A.; Bronk, Deborah A.; and O'Neil, Judith M., Nitrogen fixation and release of fixed nitrogen by *Trichodesmium* spp. in the Gulf of Mexico (2006). *Limnology and Oceanography*, 51(4), 1762-1776.  
<https://doi.org/10.4319/lo.2006.51.4.1762>

This Article is brought to you for free and open access by the Virginia Institute of Marine Science at W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).

## Nitrogen fixation and release of fixed nitrogen by *Trichodesmium* spp. in the Gulf of Mexico

Margaret R. Mulholland<sup>1</sup> and Peter W. Bernhardt

Ocean, Earth, and Atmospheric Sciences, Old Dominion University, Norfolk, Virginia 23529-0276

Cynthia A. Heil<sup>2</sup>

College of Marine Science, University of South Florida, St. Petersburg, Florida 33701

Deborah A. Bronk

Virginia Institute of Marine Science, the College of William and Mary, Gloucester Point, Virginia 23062

Judith M. O'Neil

University of Maryland Center for Environmental Science, Horn Point Laboratory, Cambridge, Maryland 21613

### Abstract

During a 3-yr study in the Gulf of Mexico, we measured dinitrogen (N<sub>2</sub>) fixation and nitrogen (N) release by *Trichodesmium* and compared these rates with water column N demand and the estimated N necessary to support blooms of *Karenia brevis*, a toxic dinoflagellate that severely affects the West Florida shelf. Net and gross N<sub>2</sub> fixation rates were compared in simultaneous incubations using <sup>15</sup>N<sub>2</sub> uptake and acetylene reduction, respectively. The difference between net and gross N<sub>2</sub> fixation is assumed to be an approximation of the rate of N release. Results demonstrate that *Trichodesmium* in the Gulf of Mexico are fixing N<sub>2</sub> at high rates and that an average of 52% of this recently fixed N<sub>2</sub> is rapidly released. Calculations suggest that observed densities of *Trichodesmium* can provide enough N to support moderately sized *K. brevis* blooms. Based on other studies where <sup>15</sup>N<sub>2</sub> and acetylene reduction were compared directly, it appears that N release from *Trichodesmium* is common but varies in magnitude among environments. In addition, carbon (C) and N-based doubling times for *Trichodesmium* vary among studies and environments. Comparing gross N<sub>2</sub> fixation and C fixation directly, C-based doubling times exceeded N-based doubling times, which suggests an imbalance in elemental turnover or a failure to fully quantify *Trichodesmium* N use.

Blooms of the toxic dinoflagellate *Karenia brevis* occur in the oligotrophic waters of the eastern Gulf of Mexico where known nitrogen (N) sources are insufficient to support the observed biomass accumulations (Steidinger et al. 1998; Walsh and Steidinger 2001; Vargo et al. 2004). Estuarine transport can supply only an estimated 5% to 20% of the N required for the daily growth needs of a moderate population ( $3 \times 10^5$  cells L<sup>-1</sup>) of *K. brevis* (Vargo et al.

2004). However, large *K. brevis* blooms frequently co-occur or occur subsequent to blooms of the N<sub>2</sub>-fixing cyanobacteria, *Trichodesmium* spp. (Walsh and Steidinger 2001). *Trichodesmium* spp. have long been recognized to be both abundant and common in the eastern Gulf of Mexico, forming dense blooms from February to August, often in shallower coastal waters (King 1950). Correlations between the timing and magnitude of blooms of *Trichodesmium* spp. and *K. brevis* in the Gulf of Mexico and the coastal Atlantic have been made based on historical red tide monitoring data, unpublished research results, and anecdotal information (Lenes et al. 2001; Walsh and Steidinger 2001). *Trichodesmium* was originally thought to have a role in *K. brevis* blooms by providing a biological mechanism for concentrating phosphorus in surface waters (Graham et al. 1953), which could be made available to *K. brevis* via decomposition and regeneration (Chew 1953). More recently, it has been hypothesized that *Trichodesmium* provide a source of new N to fuel growth of *K. brevis* (Walsh and Steidinger 2001).

*Trichodesmium* spp. can fix dinitrogen (N<sub>2</sub>) at high rates and are therefore not N limited. Further, *Trichodesmium* release fixed N as dissolved organic N (DON) (Glibert and Bronk 1994), amino acids (Capone et al. 1994), and NH<sub>4</sub><sup>+</sup> (Mulholland and Capone 2001; Mulholland et al. 2004), and the accumulation of NH<sub>4</sub><sup>+</sup> and/or DON has been observed in and around *Trichodesmium* blooms in the

<sup>1</sup> Corresponding author (mmulholl@odu.edu).

<sup>2</sup> Present address: Florida Fish and Wildlife Conservation Commission, Fish and Wildlife Research Institute, St. Petersburg, Florida 33701.

### Acknowledgments

We thank the captains and crews of the R/V *Walton Smith*, R/V *Suncoaster*, and R/V *Pelican* for help with sample collection. We also thank George Boneillo, Esther Cornfeld, Lloyd Godson, Ian Hewson, Sue Marasko, Ryan Morse, Sue Reynolds, Marta Sanderson, and Michelle Watson for their help in sample collection and analysis. We thank the Florida ECOHAB program for sharing data and ship time and Gabe Vargo for use of laboratory and facilities at the University of South Florida. We also thank two anonymous reviewers and Mary Scranton for their helpful comments. Funding for this project was provided by the National Science Foundation with grants to M.R.M. (OCE-0095923), C.A.H. (OCE-0095970), and D.A.B. (OCE-0095940). This is UMCES contribution 3948.

Pacific (Devassy et al. 1978; Karl et al. 1992, 1997) and the Gulf of Mexico (Lenes et al. 2001; Heil et al. unpubl. data), along the coast of Australia (Glibert and O'Neil 1999), and in aging *Trichodesmium* cultures (Mulholland and Capone 2001). At other times, nutrient concentrations within and around blooms may be depleted because the released N is rapidly taken up by organisms growing on and around colonies (e.g., see Sellner 1992) or co-occurring in the water column.

In some coastal systems, blooms of dinoflagellates and diatoms have been observed during and subsequent to *Trichodesmium* blooms (Devassy et al. 1978; Revelante et al. 1982; Furnas and Mitchell 1996). On the West Florida shelf, dense *K. brevis* blooms occur during and subsequent to *Trichodesmium* blooms, and it has been hypothesized that they provide a source of new N to fuel growth of *K. brevis* (Walsh and Steidinger 2001). Here we report the first estimates of N<sub>2</sub> fixation from the Gulf of Mexico and compare them with rates reported for other regions. Simultaneous N<sub>2</sub> and carbon (C) fixation rates are also compared to determine biomass-specific doubling times of *Trichodesmium* in this region. In addition to direct estimates of N release, gross N<sub>2</sub> fixation, estimated using the acetylene reduction method, was compared with net N<sub>2</sub> uptake as an independent measure of N release in the field (Mulholland et al. 2004; Mulholland and Berardt 2005). These rates are then compared to estimates of C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> reduction ratios observed in other systems and water column N demand in the Gulf of Mexico in areas where *Trichodesmium* are seasonally abundant and where *K. brevis* blooms routinely occur.

## Methods

During cruises on the West Florida shelf in the Gulf of Mexico (Fig. 1) a suite of parameters was measured including (1) N<sub>2</sub> fixation by *Trichodesmium* using the acetylene reduction method and by <sup>15</sup>N<sub>2</sub> uptake, (2) the production of <sup>15</sup>NH<sub>4</sub><sup>+</sup> from <sup>15</sup>N<sub>2</sub> by *Trichodesmium*, (3) regeneration of NH<sub>4</sub><sup>+</sup> by isotope dilution, (4) the distribution of *Trichodesmium* in the water column, and (5) water column uptake of NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, urea, and a mixture of amino acids. During 2001, cruises were aboard the R/V *Walton Smith* (July) and the R/V *Suncoaster* (October); in 2002, cruises were aboard the R/V *Walton Smith* (July and October); and, in 2003, cruises were aboard the R/V *Pelican* (June and November). On the June and July cruises, a surface radio-beacon drogue was deployed and used to follow surface populations of *Trichodesmium* spp. for periods of up to 5 d.

For N<sub>2</sub> fixation measurements, *Trichodesmium* colonies were collected from the upper 20 m using a 64- $\mu$ m, 0.5-m plankton net towed at speeds <0.5 m s<sup>-1</sup>. Colonies were transferred to filtered (Whatman GF/F, nominal pore size of 0.7  $\mu$ m) seawater using inoculating loops to rinse the colonies and then transferred to experimental incubation bottles filled with 0.2- $\mu$ m filtered seawater.

Rates of N<sub>2</sub> fixation were estimated using the acetylene reduction method (Capone 1993), which estimates gross N<sub>2</sub> fixation (Mulholland et al. 2004), and using <sup>15</sup>N-labeled N<sub>2</sub>,




Fig. 1. Map of the study area along the West Florida shelf in the Gulf of Mexico.

which estimates net <sup>15</sup>N<sub>2</sub> uptake into particulate organic nitrogen (PON) (Montoya et al. 1996; Mulholland et al. 2004). Filtrate from the <sup>15</sup>N<sub>2</sub> incubations was collected to determine rates of production of dissolved <sup>15</sup>NH<sub>4</sub><sup>+</sup> from <sup>15</sup>N<sub>2</sub> (see following). Acetylene (C<sub>2</sub>H<sub>2</sub>) reduction assays were initiated by removing 2 mL of headspace from 24-mL serum vials containing 10 mL of 0.2- $\mu$ m filtered seawater and 10 colonies of *Trichodesmium* and then adding 2 mL of C<sub>2</sub>H<sub>2</sub>. Immediately after the C<sub>2</sub>H<sub>2</sub> addition, and at increments over the course of the day, 100  $\mu$ L of headspace was removed and the production of ethylene was measured using a Shimadzu gas chromatograph equipped with a flame ionization detector.

Nitrogenase preferentially reduces C<sub>2</sub>H<sub>2</sub> as opposed to its natural substrate N<sub>2</sub>, and the stoichiometry of these two reactions is different. As a result, one must use a conversion factor to convert acetylene reduction rates to N<sub>2</sub> fixation rates. The theoretical molar ratio of C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> reduction is 3 : 1 (mol : mol) (Montoya et al. 1996; Postgate 1998). However, nitrogenase-dependent H<sub>2</sub> release, which is inhibited by C<sub>2</sub>H<sub>2</sub>, results in deviations from this theoretical stoichiometry (see Discussion following for details). We used a 3 : 1 conversion factor and evaluated deviations between N<sub>2</sub> fixation rate estimates using the C<sub>2</sub>H<sub>2</sub> reduction method and <sup>15</sup>N<sub>2</sub> uptake. These are compared with previous intercalibrations between the two methods to estimate N release by difference (Mulholland et al. 2004; Mulholland and Bernhardt 2005).

Uptake experiments were initiated by putting tracer additions (<10%) of highly enriched (99%) <sup>15</sup>N<sub>2</sub> into gas-tight bottles (315 or 727 mL) filled with whole water or <0.2- $\mu$ m filtered seawater containing 50 or 100 colonies of *Trichodesmium* (Mulholland and Capone 2001). Carbon fixation was measured simultaneously by putting tracer additions (<10% or the calculated dissolved inorganic C pool) of H<sup>13</sup>CO<sub>3</sub><sup>-</sup> into the <sup>15</sup>N incubations (Mulholland and Capone 2001). After 2 to 6 h, incubations were

terminated by filtration through combusted (450°C for 2 h) GF/F filters. Filtrate was reserved to measure the production of  $^{15}\text{NH}_4^+$  from recently fixed  $^{15}\text{N}_2$ . Parallel incubations were conducted to measure  $\text{NH}_4^+$  uptake and independently measure  $\text{NH}_4^+$  regeneration by the isotope dilution method (Glibert et al. 1982). All  $^{15}\text{N}$  and  $^{13}\text{C}$  samples were analyzed on a Europa 20/20 mass spectrometer equipped with an automated nitrogen and carbon analysis for gas, solids, and liquids (ANCA-GSL) preparation module. Rates of uptake were calculated based on the equations of Montoya et al. (1996) and Orcutt et al. (2001) using a mixing model

N uptake =

$$\frac{(\text{atom \% PON})_{\text{final}} - (\text{atom \% PON})_{\text{initial}}}{(\text{atom \% enrichment N source pool} - \text{atom \% PON})_{\text{initial}} \times \text{time}} \times [\text{PON}] \quad (1)$$

C uptake =

$$\frac{(\text{atom \% POC})_{\text{final}} - (\text{atom \% POC})_{\text{initial}}}{(\text{atom \% enrichment C source pool} - \text{atom \% POC})_{\text{initial}} \times \text{time}} \times [\text{POC}] \quad (2)$$

Daily rates of  $\text{N}_2$  fixation were calculated from hourly measurements of acetylene reduction or from  $^{15}\text{N}_2$  incubations lasting 4 h or more. Ten hours of C and  $\text{N}_2$  fixation per day were assumed. N- and C-based doubling times were calculated as in Orcutt et al. (2001).

Production of  $\text{NH}_4^+$  was measured with two techniques: (1) by adding  $^{15}\text{N}$ -labeled  $\text{N}_2$  gas and quantifying the appearance of  $^{15}\text{N}$  in the  $\text{NH}_4^+$  pool (Mulholland et al. 2004) and (2) by adding  $^{15}\text{N}$ -labeled  $\text{NH}_4^+$  and measuring isotope dilution over time (Glibert et al. 1982; Mulholland and Bernhardt 2005). The  $\text{NH}_4^+$  pool was isolated with solid phase extraction (Dudek et al. 1986). Rates of  $^{15}\text{NH}_4^+$  production from  $^{15}\text{N}_2$  were calculated using  $\text{N}_2$  as the source pool and assuming that intracellular pools of  $\text{NH}_4^+$  and DON were minimal so that release of these compounds occurred prior to assimilation into PON; mixing is between the enriched  $\text{N}_2$  and  $\text{NH}_4^+$  pools (Glibert and Bronk 1994; Mulholland et al. 2004). Rates of  $\text{NH}_4^+$  regeneration determined by isotope dilution were calculated assuming that both isotopic signatures and  $\text{NH}_4^+$  concentrations changed over the course of incubations (Eq. 2; Mulholland and Bernhardt 2005). This is a modification of the equations outlined by Glibert et al. (1982) but is consistent with the original model of Caperon et al. (1979).

Water column N uptake was estimated in 1-liter incubations (4–6 h in duration) using additions ( $0.05 \mu\text{mol L}^{-1}$  N) of highly enriched (96–99%) tracer compounds:  $^{15}\text{NH}_4^+$ ,  $^{15}\text{NO}_3^-$ , and  $^{15}\text{N}$ -labeled urea or dissolved primary amines. Uptake rates were calculated using Eq. 1 above.

Nutrient concentrations were measured using an Astoria-Pacific automated nutrient analyzer. Dissolved free amino acid concentrations were measured by high performance liquid chromatography (Cowie and Hedges 1992). Particulate organic C and PON concentrations were measured with the mass spectrometer described above.

Table 1. Maximum hourly rates of  $\text{N}_2$  fixation estimated using the  $\text{C}_2\text{H}_2$  reduction technique and a conversion factor of  $3 \pm \text{SD}$  from triplicate incubations.

Date	Time of maximum $\text{N}_2$ fixation (h)	Gross $\text{N}_2$ fixation $\text{C}_2\text{H}_2$ reduction (nmol N col $^{-1}$ h $^{-1}$ )
08 Jul 2001	16:00	1.17±0.02
09 Jul 2001	16:30	1.27±0.31
10 Jul 2001	16:30	1.96±0.37
11 Jul 2001	12:30	1.15±0.31
12 Jul 2001	15:00	1.20±0.25
11 Jul 2002	12:30	2.41±1.77
12 Jul 2002	13:00	0.93±0.54
13 Jul 2002	12:00	0.48±0.20
14 Jul 2002	12:30	1.89±0.57
05 Jun 2003	15:30	0.44±0.08
06 Jun 2003	14:00	0.93±0.10
07 Jun 2003	13:00	0.85±0.22
08 Jun 2003	15:00	0.94±0.24
09 Jun 2003	17:00	0.77±0.26
10 Jun 2003	14:00	1.50±0.12
11 Jun 2003	12:30	2.56±0.12
06 Nov 2003	15:30	0.72±0.15
07 Nov 2003	12:00	0.92±0.45
08 Nov 2003	13:00	1.52±0.46
09 Nov 2003	15:00	1.17±0.49

Water column *Trichodesmium* spp. abundance was estimated by filtering the entire contents of 20-liter Niskin bottles collected along depth profiles onto filters and then counting colonies microscopically.

## Results

*N<sub>2</sub> fixation, C fixation, and N release by Trichodesmium*—During daily  $\text{C}_2\text{H}_2$  reduction experiments, average maximum hourly rates of gross  $\text{N}_2$  fixation ranged from 0.44 to 2.56 nmol N colony $^{-1}$  (col $^{-1}$ ) h $^{-1}$  over the entire study period (six cruises over 3 yr; Table 1). The time of day that the maximum  $\text{N}_2$  fixation rate was observed varied but was always in the afternoon. The daily pattern in rates, maximum hourly rates, and integrated daily  $\text{N}_2$  fixation rate estimates also varied from day to day over a drogue study in 2001, but there was no pattern of increasing or decreasing colony-specific rates over the course of the drogue deployment (Fig. 2A). When we compared rates of  $\text{N}_2$  fixation made over the course of the same day using the  $\text{C}_2\text{H}_2$  reduction technique (gross  $\text{N}_2$  fixation) versus  $^{15}\text{N}_2$  uptake (net  $\text{N}_2$  uptake), the pattern and rates of  $\text{N}_2$  fixation varied between the two methods (Fig. 2B). Similarly, the timing of the daily  $\text{N}_2$  fixation maxima, estimated using the  $\text{C}_2\text{H}_2$  reduction technique, varied daily during cruises in July 2002 (Fig. 3A), June 2003 (Fig. 3B), and November 2003 (Fig. 3C).

Daily integrated rates of  $\text{N}_2$  and C fixation were calculated based on estimates of net  $^{15}\text{N}_2$  uptake, gross  $\text{N}_2$  fixation ( $\text{C}_2\text{H}_2$  reduction), and  $\text{H}^{13}\text{CO}_3^-$  uptake (Table 2). Net  $\text{N}_2$  uptake from  $^{15}\text{N}_2$  incubations ranged from 2.1 to 9.2 nmol N col $^{-1}$  d $^{-1}$ , while gross  $\text{N}_2$  fixation rates ranged from 3.3 to 20.5 nmol N col $^{-1}$  d $^{-1}$  based on






Fig. 2. (A) Rates of gross N<sub>2</sub> fixation (estimated using the C<sub>2</sub>H<sub>2</sub> reduction method and a conversion factor of 3) over the course of several days during a cruise on the West Florida shelf in the Gulf of Mexico in July 2001. (B) Paired comparisons of gross N<sub>2</sub> fixation (C<sub>2</sub>H<sub>2</sub> reduction) and net N<sub>2</sub> uptake (<sup>15</sup>N<sub>2</sub>) rates on 11 July 2001. Error bars are standard deviations from triplicate incubations.

C<sub>2</sub>H<sub>2</sub> reduction. Bicarbonate uptake was only measured in 2003, and daily rates ranged from 82 to 281.6 nmol C col<sup>-1</sup> d<sup>-1</sup>. Molar ratios of C fixation to gross N<sub>2</sub> fixation (estimated using the C<sub>2</sub>H<sub>2</sub> reduction method) ranged from 10.4 to 27.4, in excess of the molar C : N ratio of *Trichodesmium* (average of about five during 2003). Based on daily N<sub>2</sub> and C fixation, N turnover time for *Trichodesmium* ranged from 0.03 to 0.15 d<sup>-1</sup> over the 3-yr study period, while C turnover in 2003 was estimated to be between 0.16 and 0.24 d<sup>-1</sup>. This translates into N- and C-specific doubling times for *Trichodesmium* of 4.7 to 27.1 d and 2.9 and 4.2 d, respectively.

Surface abundance of *Trichodesmium* spp. varied from 0.2 to 7.8 colonies L<sup>-1</sup> during our cruises in July 2001 and 2002. However, these values are closer to background *Trichodesmium* concentrations (0.75 colonies L<sup>-1</sup>) and low compared with summer averages of 20 colonies L<sup>-1</sup> (Lenes et al. 2001) and bloom concentrations in excess of 1,000 colonies L<sup>-1</sup> (Vargo et al. 2004). Surface concentrations of *Trichodesmium* in excess of 6,500 colonies L<sup>-1</sup> were observed during the ecology and oceanography of harmful algal blooms (ECOHAB) Florida program from 1998 to 2002 (Vargo pers. comm.).




Fig. 3. Rates of gross N<sub>2</sub> fixation (estimated using C<sub>2</sub>H<sub>2</sub> reduction and a conversion factor of 3) over daily cycles during (A) July 2002, (B) June 2003, and (C) November 2003.

We also compared net and gross N<sub>2</sub> fixation using <sup>15</sup>N<sub>2</sub> uptake (net) and acetylene reduction (gross) methods, with C fixation over discrete intervals in order to examine variability in the ratio of net : gross N<sub>2</sub> fixation and the ratio of C : N<sub>2</sub> fixation. Rates of gross N<sub>2</sub> fixation always exceeded net N<sub>2</sub> uptake from <sup>15</sup>N<sub>2</sub> incubations, with net N<sub>2</sub> uptake ranging from 19% to 91% of the gross N<sub>2</sub> fixation (Table 3). This suggests substantial loss or release of recently fixed N<sub>2</sub> from cells (9% to 81% release, average

Table 2. Integrated daily estimates of C and N<sub>2</sub> fixation by *Trichodesmium*, estimated turnover of particulate nitrogen (PN) and particulate carbon (PC), and *Trichodesmium* doubling times  $\pm$  SD. C : N uptake is the ratio of C fixation to gross N<sub>2</sub> fixation measured using the acetylene reduction technique and a conversion factor of 3.

Date	Net <sup>15</sup> N <sub>2</sub> uptake (nmol N col <sup>-1</sup> d <sup>-1</sup> )	Gross N <sub>2</sub> fixation (nmol N col <sup>-1</sup> d <sup>-1</sup> )	C fixation (nmol C col <sup>-1</sup> d <sup>-1</sup> )	C : N uptake molar ratio	PN ( $\mu$ mol N col <sup>-1</sup> )	PC ( $\mu$ mol C col <sup>-1</sup> )	PC : PN molar ratio	N turnover (d <sup>-1</sup> )	N doubling time (d)	C turnover (d <sup>-1</sup> )	C doubling time (d)
08 Jul 2001		7.5			0.15 $\pm$ 0.01			0.05	13.8		
09 Jul 2001		10.6			0.14 $\pm$ 0.01			0.08	9.2		
10 Jul 2001	2.61	20.5			0.14 $\pm$ 0.03			0.15	4.7		
11 Jul 2001	5.68	9.8			0.11 $\pm$ 0.01			0.09	7.8		
12 Jul 2001	2.56	8.4			0.10 $\pm$ 0.01			0.08	8.3		
11 Jul 2002		11.6						0.13	5.4		
12 Jul 2002	2.06	6.0			0.09 $\pm$ 0.01			0.07	10.5		
13 Jul 2002		5.6						0.08	8.7		
14 Jul 2002	3.72	10.0			0.07 $\pm$ 0.01			0.14	4.9		
28 Oct 2002	2.86				0.07 $\pm$ 0.004						
05 Jun 2003		3.3			0.13 $\pm$ 0.01	0.74 $\pm$ 0.04	5.9 $\pm$ 0.07	0.03	27.1		
06 Jun 2003		7.7			0.10 $\pm$ 0.01	0.55 $\pm$ 0.05	5.8 $\pm$ 0.28	0.08	9.0		
07 Jun 2003		7.3	156.8	21.5	0.16 $\pm$ 0.02	0.72 $\pm$ 0.04	4.8 $\pm$ 0.13	0.05	15.2	0.22	3.2
08 Jun 2003		5.8	116.2	20.0	0.15 $\pm$ 0.04	0.70 $\pm$ 0.02	4.8 $\pm$ 0.06	0.04	17.9	0.17	4.2
09 Jun 2003	5.61	5.7	151.0	26.7	0.16 $\pm$ 0.01	0.80 $\pm$ 0.03	5.1 $\pm$ 0.13	0.04	19.6	0.19	3.7
10 Jun 2003	9.21	11.6	120.2	10.4	0.14 $\pm$ 0.01	0.69 $\pm$ 0.07	5.0 $\pm$ 0.09	0.08	8.4	0.17	4.0
11 Jun 2003	8.60	15.8	281.6	17.9	0.23 $\pm$ 0.08	1.16 $\pm$ 0.01	4.8 $\pm$ 0.01	0.07	10.1	0.24	2.9
06 Nov 2003		4.4	119.7	27.4	0.14 $\pm$ 0.01	0.73 $\pm$ 0.02	5.1 $\pm$ 0.18	0.03	22.2	0.16	4.2
07 Nov 2003	5.48	6.9	129.8	18.8	0.13 $\pm$ 0.06	0.63 $\pm$ 0.01	4.9 $\pm$ 0.05	0.05	13.1	0.21	3.4
08 Nov 2003	2.31	7.2	82.0	11.4	0.09 $\pm$ 0.02	0.44 $\pm$ 0.04	5.0 $\pm$ 0.10	0.08	8.7	0.19	3.7
09 Nov 2003		10.1	125.3	12.4	0.11 $\pm$ 0.02	0.64 $\pm$ 0.04	4.9 $\pm$ 0.10	0.09	7.5	0.20	3.5
Study average									11.6		3.6

Table 3. Direct comparisons of N<sub>2</sub> fixation estimates made using C<sub>2</sub>H<sub>2</sub> (gross) and <sup>15</sup>N<sub>2</sub> uptake (net), estimates of N release on a per-colony basis and as a percentage of the gross N<sub>2</sub> fixation, molar ratios of C<sub>2</sub>H<sub>2</sub> reduction : <sup>15</sup>N<sub>2</sub> uptake, and C fixation over discrete time intervals.

Date	Time interval (h)	N <sub>2</sub> fixation			N release (% of N <sub>2</sub> fixation)	C <sub>2</sub> H <sub>2</sub> : <sup>15</sup> N <sub>2</sub> molar ratio	C fixation (nmol C col <sup>-1</sup> )	C : N uptake molar ratio
		C <sub>2</sub> H <sub>2</sub> reduction (nmol N col <sup>-1</sup> )	<sup>15</sup> N <sub>2</sub> uptake (nmol N col <sup>-1</sup> )	N release (nmol N col <sup>-1</sup> )				
08 Jul 2001	12:00–14:00	1.62	1.14	0.48	29.6	4.3		
09 Jul 2001	10:00–12:00	2.10	0.40	1.70	81.0	15.8		
10 Jul 2001	11:00–17:00	9.54	1.98	7.56	79.2	14.5		
11 Jul 2001	09:00–16:00	6.30	1.54	4.76	75.6	12.3		
	10:30–14:30	3.56	2.84	0.72	20.2	3.8		
	10:30–12:30	2.30	0.80	1.50	65.2	8.6		
12 Jul 2001	10:00–17:00	4.83	2.24	2.59	53.6	6.5		
	12:30–17:30	4.45	2.30	2.15	48.3	5.8		
	12:30–14:30	1.86	0.74	1.12	60.2	7.5		
12 Jul 2002	14:00–18:00	1.40	1.04	0.36	25.7	4.0		
14 Jul 2002	14:00–20:00	4.38	2.82	1.56	35.6	4.7		
07 Jun 2003	15:00–16:30	0.55	0.22	0.33	60.1	7.5	23.6	42.7
08 Jun 2003	14:30–16:30	1.52	0.62	0.90	59.2	7.4	23.2	15.3
09 Jun 2003	14:30–17:00	1.50	0.93	0.58	38.3	4.9	37.8	25.2
	15:00–17:00	1.54	1.40	0.14	9.1	3.3	26.8	17.4
10 Jun 2003	10:30–11:30	1.35	0.66	0.69	51.1	6.1	10.2	7.6
	10:30–12:30	2.64	1.62	1.02	38.6	4.9	16.2	6.1
	10:30–14:30	5.64	4.60	1.04	18.4	3.7	48.0	8.5
11 Jun 2003	11:00–13:00	3.38	1.94	1.44	42.6	5.2	65.0	19.2
	14:00–16:00	3.92	2.36	1.56	39.8	5.0	56.4	14.4
06 Nov 2003	14:00–16:00	1.44	0.32	1.12	77.8	13.5	24.0	16.7
07 Nov 2003	12:00–16:00	3.32	2.72	0.60	18.1	3.7	52.0	15.7
08 Nov 2003	11:00–13:00	3.04	0.58	2.46	80.9	15.7	16.4	5.4
Study average					48	7.3		13.8

of 52%; Fig. 4A). Based on differences in gross N<sub>2</sub> fixation and net N<sub>2</sub> uptake, we estimate release rates ranging from 0.14 to 7.56 nmol N col<sup>-1</sup> (average of 0.59 nmol N col<sup>-1</sup> h<sup>-1</sup>; Fig. 4B) over intervals ranging from 1.5 to 7 h. Release was not related to the incubation duration. The molar ratio of C<sub>2</sub>H<sub>2</sub> reduction : <sup>15</sup>N<sub>2</sub> uptake varied between 3.3 and 15.8 with an overall average of 7.3 (Table 3).

During 2003, we conducted parallel experiments wherein we measured production of <sup>15</sup>NH<sub>4</sub><sup>+</sup> from <sup>15</sup>N<sub>2</sub> and NH<sub>4</sub><sup>+</sup> regeneration by isotope dilution to compare the two estimates of N regeneration. During these experiments, net <sup>15</sup>N<sub>2</sub> uptake ranged from 0.01 to 2.15 nmol col<sup>-1</sup> h<sup>-1</sup>, production of <sup>15</sup>NH<sub>4</sub><sup>+</sup> from <sup>15</sup>N<sub>2</sub> was at the limit of detection, and NH<sub>4</sub><sup>+</sup> regeneration estimated from isotope dilution ranged from 0.01 to 0.72 nmol N col<sup>-1</sup> h<sup>-1</sup> (Table 4). Production of NH<sub>4</sub><sup>+</sup> within incubation bottles could be in excess of net <sup>15</sup>N<sub>2</sub> uptake, which suggests substantial release of NH<sub>4</sub><sup>+</sup> during incubations. This analysis does not consider the probability that DON is also being released (e.g., Capone et al. 1994; Glibert and Bronk 1994; Mulholland et al. 2004). Pooling the data, there was no clear pattern suggesting higher release rates at any particular part of the day during these paired comparisons.

*Water column N<sub>2</sub> fixation*—In addition to *Trichodesmium*, it is now known that there are other unicellular cyanobacterial N<sub>2</sub> fixers that occur in the water column in marine environments (Zehr et al. 2001; Montoya et al. 2004). In this study, we estimated that water column N<sub>2</sub> uptake ranged from 11.1 to 234 pmol L<sup>-1</sup> h<sup>-1</sup> (Table 5). If we consider that unicells might fix for 24 h a day (e.g., Montoya et al. 2004), then daily fixation rates range from 0.27 to 5.62 nmol N L<sup>-1</sup> d<sup>-1</sup>. This estimate is comparable to that which might be expected for *Trichodesmium* under background conditions if there are 0.75 colonies L<sup>-1</sup> and using our study averages of 9.4 nmol N col<sup>-1</sup> d<sup>-1</sup> (gross N<sub>2</sub> fixation) or 4.8 nmol N col<sup>-1</sup> d<sup>-1</sup> (net N<sub>2</sub> uptake).

*Water column N uptake*—To put these measurements of N<sub>2</sub> fixation in surface waters in context, we compared water column uptake rates of combined N sources (NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub><sup>-</sup>, urea, and amino acids) during cruises in 2001 and 2002. Total rates of combined N uptake were lowest in July 2001 (30.6 and 53.7 nmol N L<sup>-1</sup> h<sup>-1</sup>), and the bulk of the N that was taken up was as NH<sub>4</sub><sup>+</sup> or as NH<sub>4</sub><sup>+</sup> and urea (Table 6). Total measured N uptake was within the range of N demand that could be supported by an average (20 colonies L<sup>-1</sup>) population of *Trichodesmium* fixing N<sub>2</sub> at the maximal rates we observed (2.56 nmol N col<sup>-1</sup> h<sup>-1</sup>). In




Fig. 4. Pooled estimates of N release estimated by the difference between gross  $N_2$  fixation ( $C_2H_2$  reduction) and net  $N_2$  uptake made over the entire study period (2001 to 2003) (A) expressed as a percentage of recently fixed  $N_2$  and (B) expressed as a colony-specific rate.

contrast, in the fall of 2001, there was a *K. brevis* bloom and total surface N uptake was two orders of magnitude higher than in July 2001 (2,330 and 1,540  $nmol\ N\ L^{-1}\ h^{-1}$ ; Table 6). The bulk of that N taken up, 69% to 84%, was as urea. During the summer of 2002, total N uptake was again reduced, and  $NH_4^+$  and urea satisfied the bulk of the N demand. In the fall, N demand increased, and urea was the primary source of N taken up in surface waters.

## Discussion

**$N_2$  fixation**—These are the first reported measurements of  $N_2$  fixation from the eastern Gulf of Mexico. Results from  $N_2$  fixation studies indicate that the  $N_2$  fixation rates we report for *Trichodesmium* in the Gulf of Mexico (Table 1) are within the range of those reported from studies in other basins (Table 7) and have similar variability. Our calculated N-based doubling times (4.7 to 27 d; Table 2) were also within the ranges reported elsewhere (Table 7). Despite the number of studies now reporting rates of  $N_2$  fixation by *Trichodesmium*, measured per-colony rates can vary by nearly two orders of magnitude, as can N-based doubling times. There are fewer but equally variable (e.g., two orders of magnitude) estimates of C

fixation and C-based doubling times for *Trichodesmium* (Table 8). In this study, N-based doubling times (average 11.6 d) exceeded those estimated from C fixation (average 3.6 d), but N-based doubling times varied only by a factor of about three within individual cruises, and C fixation times were within a factor of 1.5. *Trichodesmium* have also been shown to use combined N (see table 5 in Mulholland and Capone 1999), and this may result in underestimates of total N turnover. In addition, high N release rates could also result in underestimates of total N turnover. As such, we assert that C-based doubling times are likely to be more robust than those based on estimates of  $N_2$  fixation.

One likely reason for the highly variable rates and doubling times is that the physiological status of natural populations is unknown and the environmental variables controlling rates of  $N_2$  fixation and growth by these organisms are poorly constrained. Additionally, species- or strain-specific differences in the field are poorly described in general, and differences in  $N_2$  fixation rates in field incubations may depend on which species or strains of *Trichodesmium* are present (Carpenter et al. 1993).

Similarly, there is very little information on factors controlling growth rates for these organisms under oligotrophic conditions. Doubling times of hundreds of days are unlikely to produce the surface accumulations observed in nature, and even doubling times on the order of 3 d may be insufficient for producing the large ephemeral aggregations that are often observed. It is therefore likely that physical mechanisms are also involved in producing large surface blooms.

Oceanic concentrations of both phosphorus (P) and iron (Fe) have been proposed to limit growth of *Trichodesmium* and phytoplankton in nature (Karl et al. 1995; Sañudo-Wilhelmy et al. 2001; Mills et al. 2004), but there is little experimental evidence demonstrating relationships between P and Fe turnover and  $N_2$  fixation and doubling times for *Trichodesmium*. During culture studies,  $N_2$  fixation rates are higher in cultures grown at higher P concentrations (Fu and Bell 2003; Mulholland and Bernhardt 2005). In order to better evaluate global rates of  $N_2$  fixation, it will be important to constrain this variability and make models of  $N_2$  fixation regionally specific.

In addition to implications for growth rates, paired comparisons of  $N_2$  and C fixation suggest some stoichiometric imbalance in the cycling of these elements through particulate material. While *Trichodesmium* particles are very close to Redfield stoichiometry with respect to C and N, C is fixed at much higher rates than  $N_2$  (Table 9). This may be because C turnover is faster than N turnover, because N is released at high rates prior to its assimilation, or because cells are rapidly turning over C to achieve other cellular goals such as reducing intracellular oxygen through Mehler reactions (e.g., Kana 1992).

High N release rates would seem to argue for high N turnover; however, if N is released prior to its assimilation into biomass, this would contribute to gross  $N_2$  fixation (e.g., reduction of  $N_2$  to  $NH_4^+$ ) but not net uptake. In numerous paired comparisons between acetylene reduction and  $^{15}N_2$  uptake, ratios of  $C_2H_2$  reduced to  $N_2$  taken up have varied by at least an order of magnitude (Table 10).



Table 4. Simultaneous estimates of <sup>15</sup>N<sub>2</sub> uptake and NH<sub>4</sub><sup>+</sup> regeneration by isotope dilution and <sup>15</sup>NH<sub>4</sub><sup>+</sup> production in 2003, ±SD (NA indicates that there was none available). <sup>15</sup>NH<sub>4</sub><sup>+</sup> production during <sup>15</sup>N<sub>2</sub> incubations was also measured but was always below detection (data not shown).

Date	N <sub>2</sub> fixation <sup>15</sup> N <sub>2</sub> uptake (nmol N col <sup>-1</sup> h <sup>-1</sup> )	NH <sub>4</sub> <sup>+</sup> production isotope dilution (nmol N col <sup>-1</sup> h <sup>-1</sup> )	Incubation time (h)
05 Jun 2003	0.01±0.005		13:00–14:30
06 Jun 2003	0.05±0.002		09:30–11:00
07 Jun 2003	0.17±0.06		09:45–11:15
	0.37±0.09	0.11±NA	14:40–16:10
08 Jun 2003	0.01±0.01	0.06±NA	09:40–11:20
	0.31±NA		14:30–16:00
09 Jun 2003	0.63±0.40	0.72±NA	14:50–15:20
	0.37±0.14	0.11±0.02	14:50–16:00
	2.15±0.30	0.02±NA	14:50–17:00
	0.70±0.01	0.04±0.001	14:50–19:00
10 Jun 2003	0.20±NA	0.36±0.34	10:30–11:20
	0.66±0.19	0.07±0.002	10:30–12:00
	0.81±0.52	0.01±NA	10:30–13:00
	1.15±0.08	0.20±0.14	10:30–15:00
11 Jun 2003	0.97±0.21	0.07±0.01	10:45–12:00
	1.18±0.88	0.08±0.02	14:00–15:30
06 Nov 2003	0.290±NA	0.28±0.07	13:15–14:45
07 Nov 2003	0.685±0.205	0.20±0.01	12:00–14:15
08 Nov 2003	0.170±0.08	0.12±0.05	12:30–17:00
09 Nov 2003	0.019±0.01	0.02±0.01	11:30–13:45

Table 5. Water column N<sub>2</sub> fixation rates ± SD. NA indicates there were none available.

Date	Depth	N <sub>2</sub> fixation (pmol L <sup>-1</sup> h <sup>-1</sup> )
10 Jul 2001	surface	38.4±7.0
11 Jul 2001	surface	11.1±NA
12 Jul 2001	surface	74.0±NA
09 Jun 2003	surface	233.7±105.4
07 Nov 2003	surface	39.3±NA
	Chl max	63.0±22.4
08 Nov 2003	surface	174.9±20.7
	surface	158±NA
09 Nov 2003	surface	57.6±1.2
	Chl max	44.2±27.5

Table 6. Rates of combined N uptake (nmol N L<sup>-1</sup> h<sup>-1</sup>) ± SD in surface waters of the Gulf of Mexico.

Date	NH <sub>4</sub> <sup>+</sup>	NO <sub>3</sub> <sup>-</sup>	Urea	Amino acids	Total
08 Jul 2001	22.7±<0.1	1.5±0.1	3.9±<0.1	2.5±0.1	30.6
09 Jul 2001	25.1±1.6	1.5±0.2	21.6±1.9	5.5±3.9	53.7
30 Sep 2001	163±11.5	26.5±1.6	1,950±52.3	189±1.2	2,329
01 Oct 2001	190±1.0	44.2±0.3	1,060±56.3	246±15.4	1,540
11 Jul 2002	46.7±0.6	1.3±0.4	41.7±4.8	2.1±0.2	91.8
13 Jul 2002	66.6±3.7	1.7±0.3	60.8±5.7	1.1±0.1	130.2
14 Jul 2002	66.2±11.6	1.7±0.1	141±0.4	4.3±<0.1	212.8
27 Oct 2002	86.2±13.1	246±4.5	638±13.5	56.6±1.5	1,027
29 Oct 2002	9.5±4.0	24.9±0.9	214±13.4	5.2±0.3	253.7
30 Oct 2002	6.0±0.4	3.4±0.3	30.5±5.4		39.9

Table 7. N-based turnover and doubling times based on 10 h of N<sub>2</sub> fixation per day for *Trichodesmium* collected from different environments. Times were calculated based on study-specific estimates of N biomass unless noted. Rates were recalculated using molar ratios of 3 : 1 moles C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> reduced where noted. References are found in Web Appendix 1, [http://www.aslo.org/lo/toc/vol\\_51/issue\\_4/1762a1.pdf](http://www.aslo.org/lo/toc/vol_51/issue_4/1762a1.pdf).

Location	Reported as	N <sub>2</sub> fixation		PN turnover time (d)	N-based doubling time (d)	Reference
		(nmol N col <sup>-1</sup> h <sup>-1</sup> )	(nmol N col <sup>-1</sup> d <sup>-1</sup> )			
Subtropics						
Sargasso Sea	0.2 ng N (μg algal N) <sup>-1</sup> h <sup>-1</sup>	0.30	3.0	42	35	Dugdale et al. (1961)*
Sargasso Sea	0.044 μg N col <sup>-1</sup> d <sup>-1</sup>	0.26	2.6	80	56	Carpenter (1973)
W. Sargasso Sea	0.069 pg N cell <sup>-1</sup> h <sup>-1</sup>	0.15	1.5	73	51	Carpenter and McCarthy (1975)
N. Pacific	0.27 μg N μg Chl a <sup>-1</sup> h <sup>-1</sup>	0.42	4.2	41	29	Mague et al. (1977)†
Sargasso Sea	0.033 pg N cell <sup>-1</sup> h <sup>-1</sup>	0.15	1.5	103	71	Carpenter and Price (1977)‡§
N. Pacific	504 pmol C <sub>2</sub> H <sub>4</sub> μmol N <sup>-1</sup> h <sup>-1</sup>	0.05	0.5	298	206	Saino and Hattori (1978)*
Sargasso Sea	0.022 pg N cell <sup>-1</sup> h <sup>-1</sup>	0.10	1.0	155	107	McCarthy and Carpenter (1979)‡
East China Sea	0.7 pmol C <sub>2</sub> H <sub>4</sub> trichome <sup>-1</sup> h <sup>-1</sup>	0.09	0.9	161	111	Saino and Hattori (1980)¶
Tanzania Coast	69 nmol L <sup>-1</sup> C <sub>2</sub> H <sub>2</sub> (10 <sup>3</sup> trichome) <sup>-1</sup> d <sup>-1</sup> (Fig. 5)	0.77	7.7	20	14	Bryceson and Fay (1981)*¶#
Sargasso Sea	0.266 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup> (min.)	0.18	1.8	86	59	Scranton (1984)*
Sargasso Sea	1.47 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup> (max.)	0.98	9.8	16	11	Scranton (1984)*
Sargasso Sea	0.94 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup> (avg.)	0.63	6.3	24	17	Scranton et al. (1987)*††
Sargasso Sea	4.97 pmol C <sub>2</sub> H <sub>4</sub> trichome <sup>-1</sup> h <sup>-1</sup>	0.73	7.3	21	14	Carpenter et al. (1987)¶
N. Atlantic	4.04 pmol C <sub>2</sub> H <sub>4</sub> trichome <sup>-1</sup> h <sup>-1</sup>	0.59	5.9	26	18	Carpenter et al. (1987)*
N. Atlantic	32.6 nmol C <sub>2</sub> H <sub>4</sub> μg Chl a <sup>-1</sup> d <sup>-1</sup> (Fig. 1)	0.09	0.9	175	121	Paerl (1994)*§
N. Atlantic	4.3 nmol C <sub>2</sub> H <sub>4</sub> μg Chl a <sup>-1</sup> h <sup>-1</sup>	0.14	1.4	110	77	Paerl (1994)*§
N. Atlantic	305 pmol N col <sup>-1</sup> h <sup>-1</sup>	0.31	3.1	50	35	Capone et al. (1994)*
N. Atlantic (BATS)**	0.89 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup> (net tows)	0.59	5.9	35	25	Orcutt et al. (2001)
N. Atlantic (BATS)	1.00 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup> (SCUBA)	0.67	6.7	32	22	Orcutt et al. (2001)
N. Atlantic (BATS)	0.18 nmol N col <sup>-1</sup> h <sup>-1</sup> (net tows)	0.18	1.8	100	69	Orcutt et al. (2001)
N. Atlantic (BATS)	0.70 nmol N col <sup>-1</sup> h <sup>-1</sup> (SCUBA)	0.70	7.0	26	18	Orcutt et al. (2001)
N. Atlantic (BATS)	0.39 nmol N col <sup>-1</sup> h <sup>-1</sup>	0.39	3.9	51	36	Orcutt et al. (2001)
N. Atlantic (BATS)	32.2 pmol N col <sup>-1</sup> h <sup>-1</sup> (min.)	0.03	0.3	559	387	Orcutt et al. (2001)
N. Atlantic (BATS)	796.2 pmol N col <sup>-1</sup> h <sup>-1</sup> (max.)	0.80	8.0	23	16	Orcutt et al. (2001)
Gulf of Mexico	3.3 nmol N col <sup>-1</sup> h <sup>-1</sup> (min.)	0.28	2.8	51	35	Mulholland et al. (this study)
Gulf of Mexico	20.5 nmol N col <sup>-1</sup> h <sup>-1</sup> (max.)	1.71	17.1	8.2	5.7	Mulholland et al. (this study)

Table 7. Continued.

Location	Reported as	N <sub>2</sub> fixation		PN turnover time (d)	N-based doubling time (d)	Reference
		(nmol N col <sup>-1</sup> h <sup>-1</sup> )	(nmol N col <sup>-1</sup> d <sup>-1</sup> )			
Tropics						
Atlantic	1.9 ng N (μg algal N) <sup>-1</sup> h <sup>-1</sup>	0.29	2.9	53	36	Goering et al. (1966)‡
Caribbean Sea	0.077 pg N cell <sup>-1</sup> h <sup>-1</sup>	0.34	3.4	44	31	Carpenter and Price (1977)‡§
Caribbean Sea	0.090 pg N cell <sup>-1</sup> h <sup>-1</sup>	0.40	4.0	38	26	Carpenter and Price (1977)‡§
Caribbean Sea	2 fg N cell <sup>-1</sup> h <sup>-1</sup> (min.)	0.01	0.1	1,700	1,179	Scranton (1983)‡‡
Caribbean Sea	120 fg N cell <sup>-1</sup> h <sup>-1</sup> (max.)	0.54	5.4	28	20	Scranton (1983)‡‡
Caribbean Sea	1.3 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup>	0.87	8.7	18	12	Scranton et al. (1987)‡††
Caribbean Sea	6.45 pmol C <sub>2</sub> H <sub>4</sub> trichome <sup>-1</sup> h <sup>-1</sup>	0.95	9.5	16	11	Carpenter et al. (1987)*¶
Virgin Islands	99 pg N trichome <sup>-1</sup> h <sup>-1</sup>	1.41	14.1	10	7.0	Carpenter and Capone (1992)¶
Blooms worldwide	150 pg N trichome <sup>-1</sup> h <sup>-1</sup>	2.14	21.4	6.7	4.6	Carpenter and Capone (1992)¶
Caribbean Sea	0.45 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup> ( <i>Trichodesmium thiebautii</i> )	0.30	3.0	55	38	Carpenter et al. (1993)
Caribbean Sea	0.28 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup> ( <i>Trichodesmium erythraeum</i> )	0.19	1.9	83	58	Carpenter et al. (1993)
Caribbean Sea	156 pmol N col <sup>-1</sup> h <sup>-1</sup>	0.16	1.6	97	68	Capone et al. (1994)*
Caribbean Sea	182 pmol N col <sup>-1</sup> h <sup>-1</sup>	0.18	1.8	84	58	Glibert and Bronk (1994)*
Arabian Sea	1.45 pmol trichome <sup>-1</sup> h <sup>-1</sup>	0.29	2.9	52	36	Capone et al. (1998)¶‡‡
N. Atlantic (May1994)	0.44 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup>	0.29	2.9	34	23	Capone et al. (2005)§§
N. Atlantic (Apr 1996)	0.18 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup>	0.12	1.2	70	48	Capone et al. (2005)§§
N. Atlantic (Oct 1996)	0.44 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup>	0.29	2.9	20	14	Capone et al. (2005)§§
N. Atlantic (Feb 2001)	0.24 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup>	0.16	1.6	62	43	Capone et al. (2005)§§
N. Atlantic (Jul 2001)	0.17 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup>	0.11	1.1	87	60	Capone et al. (2005)§§
N. Atlantic (Apr 2003)	0.14 nmol C <sub>2</sub> H <sub>4</sub> col <sup>-1</sup> h <sup>-1</sup>	0.09	0.9	106	73	Capone et al. (2005)§§

\* Assumed an average N biomass of 2.13 μg N col<sup>-1</sup> (McCarthy and Carpenter 1979).  
 † Original paper used a molar ratio of 1.9 : 1 moles C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> reduced (Mague et al. 1977).  
 ‡ Assumed an average of 29,800 cells col<sup>-1</sup> (Carpenter and McCarthy 1975).  
 § Assumed an average of 48 ng Chl *a* col<sup>-1</sup> for *T. thiebautii* (Carpenter et al. 1993).  
 || Original paper used a molar ratio of 6.3 : 1 moles C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> reduced (Carpenter and Price 1977).  
 ¶ Assumed an average of 200 trichomes col<sup>-1</sup> (Carpenter 1983; Carpenter and Capone 1992).  
 # Colony size was 2 to 45 trichomes bundle<sup>-1</sup> (Bryceson and Fay 1981).  
 \*\* BATS, Bermuda Atlantic Time-series Study.  
 †† Colony size was 10 to 30 trichomes col<sup>-1</sup> (Capone et al. 1998).  
 ‡‡ Original paper used a molar ratio of 4.1 : 1 moles C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> reduced (Scranton et al. 1987).  
 §§ We used PN values from same cruises reported in Carpenter et al. (2004); for the last three cruises, we used the average N biomass of 1.38 μg N col<sup>-1</sup> from May 1994.

Table 8. C-based turnover and doubling times based on 10 h of C fixation per day for *Trichodesmium* collected from different environments. Times were calculated based on study-specific estimates of C biomass unless noted. References are found in Web Appendix 1.

Location	Reported as	C fixation		PC turnover time (d)	C-based doubling time (d)	Reference
		(nmol C col <sup>-1</sup> h <sup>-1</sup> )	(nmol C col <sup>-1</sup> d <sup>-1</sup> )			
Subtropics						
N. Pacific	0.38 $\mu\text{g C } \mu\text{g Chl } a^{-1} \text{ h}^{-1}$	1.1	10.8	75	52	Mague et al. (1977)
Sargasso Sea	1 $\mu\text{g C } 10^{-6} \text{ cell}^{-1} \text{ h}^{-1}$	2.5	24.8	38	26	Carpenter and Price (1977)*†
Sargasso Sea	4.2 ng C col <sup>-1</sup> h <sup>-1</sup>	0.4	3.5	269	186	McCarthy and Carpenter (1979)
N. Atlantic	11.3 $\mu\text{g C } \mu\text{g Chl } a^{-1} \text{ h}^{-1}$ (Fig. 1)	3.8	37.7	25	17	Paerl (1994)‡
N. Atlantic	1.63 $\mu\text{g C col}^{-1} \text{ d}^{-1}$	13.6	135.8	6.1	4.3	Carpenter and Roenneberg (1995)
N. Atlantic (BATS)	17.54 nmol C col <sup>-1</sup> h <sup>-1</sup> (net tows)	17.5	175.4	5.4	3.7	Orcutt et al. (2001)*
N. Atlantic (BATS)	13.91 nmol C col <sup>-1</sup> h <sup>-1</sup> (SCUBA)	13.9	139.1	6.8	4.7	Orcutt et al. (2001)*
N. Atlantic (BATS)	17.4 nmol C col <sup>-1</sup> h <sup>-1</sup> (simultaneous N and C)	17.4	174.0	5.4	3.8	Orcutt et al. (2001)*
Gulf of Mexico	82 nmol C col <sup>-1</sup> h <sup>-1</sup> (min.)	6.8	68.3	7.6	5.3	Mulholland et al. (this study)
Gulf of Mexico	281.6 nmol C col <sup>-1</sup> h <sup>-1</sup> (max.)	23.5	234.7	5.0	3.5	Mulholland et al. (this study)
Tropics						
Caribbean Sea	2 $\mu\text{g C } 10^{-6} \text{ cell}^{-1} \text{ h}^{-1}$	5.0	49.7	19	13	Carpenter and Price (1977)*†
Kuroshio	3.0 mg C mg Chl <i>a</i> <sup>-1</sup> h <sup>-1</sup>	8.5	85	11	7.7	Shimura et al. (1978)§
Kuroshio	5.2 mg C mg Chl <i>a</i> <sup>-1</sup> h <sup>-1</sup>	14.7	147.3	6.4	4.4	Shimura et al. (1978)§
Caribbean Sea	0.606 $\mu\text{g C col}^{-1} \text{ d}^{-1}$ (gross)	4.2	42.1	22	16	Li et al. (1980)
Caribbean Sea	0.426 $\mu\text{g C col}^{-1} \text{ d}^{-1}$ (net)	3.0	29.6	32	22	Li et al. (1980)
Caribbean Sea	3.9 $\mu\text{g C col}^{-1} \text{ d}^{-1}$ ( <i>T. thiebautii</i> )	32.5	325	3.1	2.1	Carpenter et al. (1993)
Caribbean Sea	3.0 $\mu\text{g C col}^{-1} \text{ d}^{-1}$ ( <i>T. erythraeum</i> )	25.0	250	3.7	2.6	Carpenter et al. (1993)
Caribbean Sea	10.8 $\mu\text{g C col}^{-1} \text{ d}^{-1}$	90.0	900.0	0.9	0.6	Carpenter and Roenneberg (1995)
Caribbean Sea	1.29 $\mu\text{g C col}^{-1} \text{ d}^{-1}$	10.8	107.5	7.8	5.4	Carpenter and Roenneberg (1995)
N. Atlantic (May 1994)	0.22 $\mu\text{g C col}^{-1} \text{ h}^{-1}$	18.3	183.3	3.5	2.4	Carpenter et al. (2004)
N. Atlantic (Apr 1996)	0.07 $\mu\text{g C col}^{-1} \text{ h}^{-1}$	5.8	58.3	8.1	5.6	Carpenter et al. (2004)
N. Atlantic (Oct 1996)	0.09 $\mu\text{g C col}^{-1} \text{ h}^{-1}$	7.5	75.0	5.3	3.7	Carpenter et al. (2004)

\* Assumed an average C biomass of 11.3  $\mu\text{g C col}^{-1}$  (McCarthy and Carpenter 1979).

† Assumed an average of 29,800 cells col<sup>-1</sup> (Carpenter and McCarthy 1975).

‡ Assumed an average of 48 ng Chl *a* col<sup>-1</sup> for *T. thiebautii* (Carpenter et al. 1993).

§ Assumed an average of 34 ng Chl *a* col<sup>-1</sup> for the central Pacific (Mague et al. 1977).

Because C<sub>2</sub>H<sub>2</sub> reduction measures just the reduction step, it is a measure of gross N<sub>2</sub> fixation, while movement of <sup>15</sup>N<sub>2</sub> from the dissolved to the particulate pool measures net N assimilation (see Mulholland et al. 2004; Mulholland and Bernhardt 2005). Release of recently reduced N<sub>2</sub> and the

difficulty in chemically recovering all possible dissolved pools into which products of N<sub>2</sub> fixation might be released may make intercalibration between the two methods impossible. However, the difference between N<sub>2</sub> reduction (gross N<sub>2</sub> fixation) and net N<sub>2</sub> assimilation has proved

Table 9. Paired comparisons of C and N<sub>2</sub> fixation. References are found in Web Appendix 1.

Location	C : N <sub>2</sub> fixation		Reference
	Range	Average	
Gulf of Mexico	5.4–42.7	13.1	Mulholland et al. (this study)
New Caledonia (lagoon)	9.2–77.7		Mulholland et al. (unpubl. data)
North Atlantic (latitudinal gradient)	13.6–33.3	21.9	Mulholland et al. (unpubl. data)
North Pacific	1.2–2.1		Mague et al. (1977)
Sargasso Sea		16	Carpenter and Price (1977)
Sargasso Sea	1.5–87		McCarthy and Carpenter (1979)
BATS (puffs)	13–437	128	Orcutt et al. (2001)
BATS (tufts)	15–703	198	Orcutt et al. (2001)
N. Atlantic		63	Carpenter et al. (2004); Capone et al. (2005)
N. Atlantic		63	Carpenter et al. (2004); Capone et al. (2005)
N. Atlantic		63	Carpenter et al. (2004); Capone et al. (2005)
<i>Trichodesmium</i> IMS101 (batch)	4.6–132.5	28	Mulholland and Capone (2001)*
<i>Trichodesmium</i> IMS101 (batch)	6.5–15.2	9.6	Mulholland and Capone (2001)†
<i>Trichodesmium</i> IMS101 (continuous)	13.4–20		Mulholland and Bernhardt (2005)‡

\* Midday estimate during exponential growth.

† Cumulative estimate over a growth or diel cycle. Ratio increased during stationary phase growth.

‡ Lower at faster growth rates.

promising in culture systems as an index of the release of recently fixed N<sub>2</sub> (Mulholland et al. 2004; Mulholland and Bernhardt 2005). What this means is that measuring an average C<sub>2</sub>H<sub>2</sub> : <sup>15</sup>N<sub>2</sub> uptake ratio of six would translate into a release rate of about 50%.

There has long been debate about the “correct” calibration factor to use in translating C<sub>2</sub>H<sub>2</sub> reduction to N<sub>2</sub> fixation. Most of this discussion has revolved around the role of nitrogenase in H<sub>2</sub> production. Although the theoretical molar ratio of C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> reduction is 3 : 1

Table 10. Results from paired comparisons of C<sub>2</sub>H<sub>2</sub> : N<sub>2</sub> measurements. Numbers are reported as molar ratios, and N release is estimated as the observed molar ratio minus the theoretical ratio (3) divided by the observed molar ratio. References are found in Web Appendix 1.

Location	C <sub>2</sub> H <sub>2</sub> : N <sub>2</sub> (molar ratio)		Estimated N release (%)	Reference
	Range	Average		
Gulf of Mexico	3.3–15.8	7.3	59	Mulholland et al. (this study)
N. Atlantic (latitudinal gradient)	2.1–7.4	4.2	29	Mulholland et al. (unpubl. data)
Sargasso Sea		6.0	50	Carpenter and McCarthy (1975)
N. Pacific		1.9		Mague et al. (1977)
N. Pacific	3–10			Saino (1977)
Sargasso Sea		6.3	52	Carpenter and Price (1977)
Sargasso Sea		2.9		Scranton (1984)
Caribbean and Sargasso Seas		4.1	27	Scranton et al. (1987)
Caribbean Sea		3.4	12	Glibert and Bronk (1994)
BATS (net tows)		4.9	39	Orcutt et al. (2001)
BATS (SCUBA)		1.4		Orcutt et al. (2001)
BATS		3.0		Orcutt et al. (2001)
N. Atlantic	0.9–7.3	3.6	17	Capone et al. (2005)
<i>Trichodesmium</i> IMS101 (batch)	1.7–9.8	5.6	46	Mulholland et al. (2004)
<i>Trichodesmium</i> IMS101 (continuous)	3.0–22.2	11.4	74	Mulholland and Bernhardt (2005)



(mol:mol) (Montoya et al. 1996; Postgate 1998), because nitrogenase catalyzes the reduction of other compounds, including  $H^+$ , in conjunction with  $N_2$  fixation, higher ratios are often assumed (e.g.,  $>3:1$ ). Acetylene blocks  $H_2$  production by nitrogenase, but  $N_2$  does not, and so nitrogenase-mediated  $H_2$  production and  $N_2$  fixation can proceed simultaneously in the absence of  $C_2H_2$ , thereby altering the stoichiometry of  $N_2$  reduction. For example, if a mole of  $H_2$  is produced with each mole of  $N_2$  reduced, the molar ratio of  $C_2H_2:N_2$  reduction becomes 4:1 (mol:mol). However, little net  $H_2$  production was observed under environmental conditions (e.g., rates of  $H_2$  production were two to three orders of magnitude lower than  $N_2$  fixation rates in incubations containing *Trichodesmium* and could not account for a substantial deviation from the 3:1 theoretical ratio; Scranton 1984; Scranton et al. 1987), which suggests either (1) that hydrogen production from nitrogenase is low in nature, justifying a 3:1 conversion factor, or (2) that cells efficiently recoup  $H_2$  produced during  $N_2$  fixation, arguing for higher ratios and a thorough evaluation of hydrogen cycling within cells. We did not measure  $H_2$  production as part of this study. Despite low rates of  $H_2$  production, Scranton et al. (1987) measured a  $C_2H_2$  reduction: $^{15}N_2$  uptake ratio of 4.1:1. We contend that in the absence of  $H_2$  production, deviations from the theoretical 3:1 ratio of  $C_2H_2:N_2$  reduction might also arise from N release. Alternatively, measurements of  $H_2$  production may be underestimated if  $N_2$  fixers recycle or recoup  $H_2$  intracellularly rather than release it into the environment or if production and consumption of  $H_2$  are tightly coupled in the environment (e.g., co-occurring taxa such as *K. brevis* take up  $H_2$ ). Physiological studies of  $H_2$  production catalyzed by nitrogenase in *Trichodesmium* are hampered by the lack of in vitro assays, and we are unaware of studies examining  $H_2$  uptake by *K. brevis* or other co-occurring taxa.

While there have been observations that *Trichodesmium* release recently fixed  $N_2$  as DON in natural populations (Capone et al. 1994; Glibert and Bronk 1994) and as  $NH_4^+$  in cultures of *Trichodesmium* IMS101 (Mulholland et al. 2004; Mulholland and Bernhardt 2005), it is unclear why cells do this. Previous speculation suggested that this is a mechanism for the extracellular transfer of fixed N between cells that fix  $N_2$  and those that do not have that capability (Mulholland and Capone 1999, 2000). Another possible fate for regenerated N is co-occurring organisms (O'Neil et al. 1996). In the current study, release rates averaged about 52% of the recently fixed  $N_2$  or 0.29 nmol  $col^{-1} h^{-1}$ , and much of this was probably  $NH_4^+$  and DON (including amino acids; Capone et al. 1994). So far in this discussion of  $N_2$  fixation, we have not considered unicellular diazotrophs, which fixed between 0.01 and 0.23 nmol N  $L^{-1} h^{-1}$  during this study (Table 5). These rates are within the range reported in water collected from 25 m from Sta. ALOHA in the Pacific (0.01 to 0.15 nmol  $L^{-1} h^{-1}$ ) but lower than that reported for the pigment maximum (about 120 m) at the same site (0.047 to 1.85 nmol  $L^{-1} h^{-1}$ ), during a transect in the North Pacific (0.31 to 1.3 nmol  $L^{-1} h^{-1}$ ), or along the north coast of Australia (up to 62 nmol  $L^{-1} h^{-1}$ ) (Montoya et al. 2004).

The fate of  $N_2$  fixed by unicellular cyanobacteria is unknown, and so we cannot speculate on its contribution to the dissolved pool or to higher trophic levels.

*Water column N uptake*—To put the observed  $N_2$  fixation and release rates into the context of water column N demand, *Trichodesmium* abundance was between 0.2 and 7.8  $col L^{-1}$  during 2001 and 2002, and this would result in a release of about 0.12 to 4.6 nmol N  $L^{-1} h^{-1}$  (using an average release rate of 0.59 nmol N  $L^{-1} h^{-1}$ ). While background concentrations of *Trichodesmium* during November to April in the eastern Gulf are about 0.75  $col L^{-1}$  (Vargo pers. comm.), summer *Trichodesmium* concentrations are generally much higher, averaging 20  $col L^{-1}$  (11.8 nmol N  $L^{-1} h^{-1}$  released based on our average release rates). During blooms, densities along the West Florida shelf can reach  $>1,000 col L^{-1}$  (average release rate of 590 nmol N  $L^{-1} h^{-1}$ ). Based on in situ rates of N uptake by *K. brevis* (Bronk et al. 2004), we calculate an N demand for the large 2001 *K. brevis* bloom off St. Petersburg (up to  $8.5 \times 10^6 cells L^{-1}$ ) to be on the order of 0.15 to 1.15 nmol N  $L^{-1} h^{-1}$ . This is well within the range of N release rates measured during our July study. Independently, Vargo et al. (2004) calculated a total N demand of 56 to 267 nmol N  $L^{-1} d^{-1}$  for a moderate ( $3 \times 10^5 cells L^{-1}$ ) *K. brevis* population undergoing 0.2 divisions  $d^{-1}$ . Based on the current study, if we assume 10 h of  $N_2$  fixation per day and modest *Trichodesmium* populations of 20  $col L^{-1}$ , we calculate an average release of 118 nmol N  $L^{-1} d^{-1}$ , within the range necessary to support a moderate *K. brevis* population. If we invoke the higher *Trichodesmium* densities, *Trichodesmium* can produce ample dissolved N to fuel *K. brevis* population growth.

When we compare these estimates with estimates of surface water N demand during this study (Table 6), *Trichodesmium* can account for a substantial fraction of the water column N demand. During September and October 2001, *K. brevis* was abundant in the surface waters and the N demand was quite high as compared with summer sampling periods when *Trichodesmium* was more abundant. The temporal separation between populations of *K. brevis* and *Trichodesmium* may indicate indirect trophic coupling as well. Urea was the primary source of N during the fall, when *K. brevis* blooms reach their peak. The source of this urea is not known, but we are currently examining the possibility that urea may be regenerated from *Trichodesmium* biomass during its decay (Mulholland et al. unpubl. data). In addition,  $NH_4^+$  and amino acids, including glutamate, were important components of the bulk water column uptake observed in this study and during the *K. brevis* bloom during 2001 (Bronk et al. 2004). These compounds are also common, identifiable release products during *Trichodesmium* growth (Capone et al. 1994; Mulholland et al. 2004; Mulholland and Bernhardt 2005).

While these results do not establish a direct trophic link between the two populations, we conclude that  $N_2$  fixation is quantitatively significant in the eastern Gulf of Mexico and that recently fixed  $N_2$  from *Trichodesmium* is released as dissolved N in forms that can be taken up by natural

populations of phytoplankton that include *K. brevis*. This is the first step toward demonstrating a direct link between N<sub>2</sub> fixation by *Trichodesmium* and the harmful bloom-former *K. brevis*.

### References

- BRONK, D. A., M. P. SANDERSON, M. R. MULHOLLAND, C. A. HEIL, AND J. M. O'NEIL. 2004. Organic and inorganic nitrogen uptake kinetics in field populations dominated by *Karenia brevis*, p. 80–82. In K. A. Steidinger, J. H. Lansberg, C. R. Tomas, and G. A. Vargo [eds.], *Harmful algae 2002*. Florida Fish and Wildlife Conservation Commission, Florida Institute of Oceanography, and Intergovernmental Oceanographic Commission of UNESCO.
- CAPERON, J., D. SCHELL, J. HIROTA, AND E. LAWS. 1979. Ammonium excretion rates in Kaneohe Bay, Hawaii, measured by a <sup>15</sup>N isotope dilution technique. *Mar. Biol.* **54**: 33–40.
- CAPONE, D. G. 1993. Determination of nitrogenase activity in aquatic samples using the acetylene reduction procedure, p. 621–631. In P. F. Kemp, B. F. Sherr, E. B. Sherr, and J. J. Cole [eds.], *Handbook of methods in aquatic microbial ecology*. Lewis.
- , M. D. FERRIER, AND E. J. CARPENTER. 1994. Amino acid cycling in colonies of the planktonic marine cyanobacterium, *Trichodesmium thiebautii*. *Appl. Environ. Microbiol.* **60**: 3989–3995.
- CARPENTER, E. J., J. M. O'NEIL, R. DAWSON, D. G. CAPONE, P. J. A. SIDDIQUI, T. ROENNEBERG, AND B. BERGMAN. 1993. The tropical diazotrophic phytoplankton *Trichodesmium*: Biological characteristics of two common species. *Mar. Ecol. Prog. Ser.* **95**: 295–304.
- CHEW, F. 1953. Results of hydrographic and chemical investigations in the region of the 'Red Tide' bloom on the west coast of Florida in November 1952. *Bull. Mar. Sci. Gulf Caribb.* **2**: 610–625.
- COWIE, G. L., AND J. I. HEDGES. 1992. Improved amino acid quantification in environmental samples: Charge-matched recovery standards and reduced analysis time. *Mar. Chem.* **37**: 223–238.
- DEVASSY, V. P., P. M. A. BHATTATHIRI, AND S. Z. QASIM. 1978. *Trichodesmium* phenomenon. *Ind. J. Mar. Sci.* **7**: 168–186.
- DUDEK, N., M. A. BRZEZINSKI, AND P. A. WHEELER. 1986. Recovery of ammonium nitrogen by solvent extraction for the determination of relative <sup>15</sup>N abundance in regeneration experiments. *Mar. Chem.* **18**: 59–69.
- FU, F.-X., AND P. R. F. BELL. 2003. Factors affecting N<sub>2</sub> fixation by the cyanobacterium *Trichodesmium* sp. GBRTRL1101. *FEMS Microbiol. Ecol.* **45**: 203–209.
- FURNAS, M. J., AND A. W. MITCHELL. 1996. Pelagic primary production in the Coral and southern Solomon Seas. *Mar. Freshw. Res.* **47**: 695–705.
- GLIBERT, P. M., AND D. A. BRONK. 1994. Release of dissolved organic nitrogen by marine diazotrophic cyanobacteria, *Trichodesmium* spp. *Appl. Environ. Microbiol.* **60**: 3996–4000.
- , F. LIPSCHULTZ, J. J. MCCARTHY, AND M. A. ALTABET. 1982. Isotope dilution models of uptake and remineralization of ammonium by marine plankton. *Limnol. Oceanogr.* **27**: 639–650.
- , AND J. M. O'NEIL. 1999. Dissolved organic nitrogen release and amino acid oxidase activity by *Trichodesmium* spp., p. 265–271. In L. Charpy and A. W. D. Larkum [eds.], *Marine cyanobacteria*. Bull. De l'Institut océanographique, Monaco, Numéro spécial 19.
- GRAHAM, H. W., J. M. AMISON, AND K. T. MARVIN. 1953. Phosphorus content of waters along the west coast of Florida. U.S. Fish. Wildl. Serv., Spec. Sci. Rep. Fish. 122.
- KANA, T. M. 1992. Oxygen cycling in cyanobacteria with specific reference to oxygen protection in *Trichodesmium* spp., p. 29–41. In E. J. Carpenter, D. G. Capone, and J. G. Rueter [eds.], *Marine pelagic cyanobacteria: Trichodesmium and other diazotrophs*. Kluwer.
- KARL, D. M., R. LETELIER, D. V. HEBEL, D. F. BIRD, AND C. D. WINN. 1992. *Trichodesmium* blooms and new nitrogen in the North Pacific Gyre, p. 219–237. In E. J. Carpenter, D. G. Capone, and J. G. Rueter [eds.], *Marine pelagic cyanobacteria: Trichodesmium and other diazotrophs*. Kluwer.
- , ———, ———, L. TUPAS, J. DORE, J. CHRISTIAN, AND C. WINN. 1995. Ecosystem changes in the North Pacific subtropical gyre attributed to the 1991–1992 El Niño. *Nature* **373**: 230–234.
- , ———, L. TUPAS, J. DORE, J. CHRISTIAN, AND D. HEBEL. 1997. The role of nitrogen fixation in biogeochemical cycling in the subtropical North Pacific. *Nature* **388**: 533–538.
- KING, J. E. 1950. A preliminary report on the plankton of the west coast of Florida. *J. Fla. Acad. Sci.* **12**: 109–137.
- LENES, J. M., AND OTHERS. 2001. Iron fertilization and the *Trichodesmium* response on the West Florida shelf. *Limnol. Oceanogr.* **46**: 1261–1277.
- MILLS, M. M., C. RIDAME, M. DAVEY, J. LAROCHE, AND R. J. GEIDER. 2004. Iron and phosphorus co-limit nitrogen fixation in the eastern tropical North Atlantic. *Nature* **429**: 292–294.
- MONTONA, J. P., C. M. HOLL, J. P. ZEHR, A. HANSEN, T. A. VILLAREAL, AND D. G. CAPONE. 2004. High rates of N<sub>2</sub> fixation by unicellular diazotrophs in the oligotrophic Pacific Ocean. *Nature* **430**: 1027–1032.
- , M. VOSS, P. KÄHLER, AND D. G. CAPONE. 1996. A simple, high precision tracer assay for dinitrogen fixation. *Appl. Environ. Microbiol.* **62**: 986–993.
- MULHOLLAND, M. R., AND P. W. BERNHARDT. 2005. The effect of growth rate, phosphorus concentration, and temperature on N<sub>2</sub> fixation and N regeneration rates in continuous cultures of *Trichodesmium* IMS101. *Limnol. Oceanogr.* **50**: 839–849.
- , D. A. BRONK, AND D. G. CAPONE. 2004. N<sub>2</sub> fixation and regeneration of NH<sub>4</sub><sup>+</sup> and dissolved organic N by *Trichodesmium* IMS101. *Aquat. Microb. Ecol.* **37**: 85–94.
- , AND D. G. CAPONE. 1999. Nitrogen utilization and metabolism relative to patterns of N<sub>2</sub> fixation in populations of *Trichodesmium* from the North Atlantic Ocean and Caribbean Sea. *Mar. Ecol. Prog. Ser.* **188**: 33–49.
- , AND ———. 2000. The physiology of the marine N<sub>2</sub> fixing cyanobacteria *Trichodesmium*. *Trends Plant Sci* **5**: 148–153.
- , AND ———. 2001. The stoichiometry of N and C utilization in cultured populations of *Trichodesmium* IMS101. *Limnol. Oceanogr.* **46**: 436–443.
- O'NEIL, J. M., P. M. METZLER, AND P. M. GLIBERT. 1996. Ingestion of <sup>15</sup>N-labelled *Trichodesmium* spp. and ammonium regeneration by the harpacticoid copepod *Macrosetella gracilis*. *Mar. Biol.* **125**: 89–96.
- ORCUTT, K. M., F. LIPSCHULTZ, K. GUNDERSEN, R. ARIMOTO, A. F. MICHAELS, A. H. KNAP, AND J. R. GALLON. 2001. A seasonal study of the significance of N<sub>2</sub> fixation by *Trichodesmium* spp. at the Bermuda Atlantic Time-series Study (BATS) site. *Deep-Sea Res. II* **48**: 1583–1608.
- POSTGATE, J. R. 1998. *The fundamentals of nitrogen fixation*. Cambridge Univ. Press.
- REVELANTE, N., W. T. WILLIAMS, AND J. S. BUNT. 1982. Temporal and spatial distribution of diatoms, dinoflagellates and *Trichodesmium* in waters of the Great Barrier Reef. *J. Exp. Mar. Biol. Ecol.* **63**: 27–45.

- SANUDO-WILHELMY, S. A., AND OTHERS. 2001. Phosphorus limitation of nitrogen fixation by *Trichodesmium* in the central North Atlantic Ocean. *Nature* **411**: 66–69.
- SCRANTON, M. I. 1984. Hydrogen cycling in the waters near Bermuda: The role of the nitrogen fixer, *Oscillatoria thiebautii*. *Deep-Sea Res* **31**: 133–143.
- , P. C. NOVELLI, A. MICHAELS, S. G. HERRIGAN, AND E. J. CARPENTER. 1987. Hydrogen production and nitrogen fixation by *Oscillatoria thiebautii* during in situ incubations. *Limnol. Oceanogr.* **32**: 998–1006.
- SELLNER, K. G. 1992. Trophodynamics of marine cyanobacteria blooms, p. 75–94. In E. J. Carpenter, D. G. Capone, and J. G. Rueter [eds.], *Marine pelagic cyanobacteria: Trichodesmium and other diazotrophs*. Kluwer.
- STEIDINGER, K. A., G. A. VARGO, P. A. TESTER, AND C. R. TOMAS. 1998. Bloom dynamics and physiology of *Gymnodinium breve* with emphasis on the Gulf of Mexico, p. 133–153. In D. M. Anderson, A. D. Cembella, and G. M. Hallegraeff [eds.], *Physiological ecology of harmful algal blooms*. Springer-Verlag.
- VARGO, G. A., AND OTHERS. 2004. Four *Karenia brevis* blooms: A comparative analysis, p. 14–16. In K. A. Steidinger, J. H. Lansberg, C. R. Tomas, and G. A. Vargo [eds.], *Harmful algae 2002*. Florida Fish and Wildlife Conservation Commission, Florida Institute of Oceanography, and Intergovernmental Oceanographic Commission of UNESCO.
- WALSH, J. J., AND K. A. STEIDINGER. 2001. Saharan dust and red tides: The cyanophyte connection. *J. Geophys. Res.* **106**: 11597–11612.
- ZEHR, J. P., AND OTHERS. 2001. Unicellular cyanobacteria fix N<sub>2</sub> in the subtropical North Pacific Ocean. *Nature* **412**: 635–638.

Received: 14 September 2005

Amended: 6 February 2006

Accepted: 5 March 2006