

W&M ScholarWorks

Arts & Sciences Articles

Arts and Sciences

2016

General Conditions for Proximity-Induced Odd-Frequency Superconductivity in Two-Dimensional Electronic Systems

Christopher Triola

Alexander V. Balatsky

Driss M. Badiane *William & Mary*

E. Rossi *William & Mary*

Alexander V. Balatsky

Follow this and additional works at: https://scholarworks.wm.edu/aspubs

Recommended Citation

Triola, Christopher; Balatsky, Alexander V.; Badiane, Driss M.; Rossi, E.; and Balatsky, Alexander V., General Conditions for Proximity-Induced Odd-Frequency Superconductivity in Two-Dimensional Electronic Systems (2016). *Physical Review Letters*, 116(25). 10.1103/PhysRevLett.116.257001

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

General conditions for proximity-induced odd-frequency superconductivity in two-dimensional electronic systems

Christopher Triola,^{1,2} Driss M. Badiane,³ Alexander V. Balatsky,^{1,2,4} and E. Rossi³

¹Nordita, Roslagstullsbacken 23, SE-106 91 Stockholm, Sweden

²Center for Quantum Materials (CQM), KTH and Nordita, Stockholm, Sweden

³Department of Physics, College of William and Mary, Williamsburg, Virginia 23187, USA

⁴Institute for Materials Science, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

We obtain the general conditions for the emergence of odd-frequency superconducting pairing in a two-dimensional (2D) electronic system proximity-coupled to a superconductor, making minimal assumptions about both the 2D system and the superconductor. Using our general results we show that a simple heterostructure formed by a monolayer of a group VI transition metal dichalcogenide, such as molybdenum disulfide, and an s-wave superconductor with Rashba spin-orbit coupling will exhibit odd-frequency superconducting pairing. Our results allow the identification of a new class of systems among van der Waals heterostructures in which odd-frequency superconductivity should be present.

Low-dimensional heterostructures hold the promise for new technologies^{1–5} as well as granting us access to many unconventional quantum states including: novel forms of superfluidity⁶, manipulation of spin textures^{7,8}, and unconventional superconductivity^{4,9–15}. In addition, theoretical analyses have shown that Majorana bound states can appear in heterostructures incorporating superconducting materials^{16–28}. Given the variety of possible exotic states in low dimensional heterostructures has rapidly advanced in recent years⁵ it is important to continue developing our understanding of their electronic properties. One important facet of this understanding is the classification of the possible symmetries of the proximityinduced superconductivity in these structures.

The symmetries of a superconductor can be characterized by investigating the properties of the anomalous Green's function $F_{\alpha\beta}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2)$ $\langle Tc_{\alpha}(\mathbf{r}_1, t_1)c_{\beta}(\mathbf{r}_2, t_2)\rangle$, where $c_{\sigma}(\mathbf{r}_i, t_i)$ is the fermionic annihilation operator for an electron at position \mathbf{r}_i time t_i with spin σ , T is time ordering operator, and the angle brackets denote the expectation value. Given the fermionic nature of the quasiparticles $F_{\alpha\beta}(\mathbf{r}_1, t_1; \mathbf{r}_2, t_2) = -F_{\beta\alpha}(\mathbf{r}_2, t_2; \mathbf{r}_1, t_1).$ Conventionally this is taken to imply that if the quasiparticle pair is in a spin singlet state then the pairing amplitude is even in parity while if it is a spin triplet the pairing amplitude is odd in parity. However, if the pairing amplitude is odd in time, or, equivalently, odd in frequency, spin triplet pairs can be even in parity and spin singlet pairs can be odd in parity as was originally proposed for superfluid ³He by Berezinskii²⁹ and later for superconductivity by Balatsky and Abrahams³⁰.

The study of odd-frequency superconductivity (SC) has been hindered by the scarcity of experimental systems in which it can be realized. Soon after the original suggestion that in general an odd-frequency pairing term could be present it was realized that it would be challenging to get such a term via electron-phonon interactions and that a spin-dependent electron-electron

interaction would be necessary 31 . This fact greatly restricts the number of systems in which odd-frequency SC could be realized. However, in recent years it has become apparent that odd-frequency SC can be obtained in heterostructures^{9,12,14,15,32–39}. Each of this works considered a different type of heterostructure. The recent impressive explosion of the types of heterostructures that can be realized has made this piecemeal approach unfeasible: a theoretical treatment able to provide the general conditions in which odd-frequency SC should be present in heterostructures has become necessary. In this work we present such a general treatment. Our general treatment also makes possible the identification of novel, somehow unexpected, engineered systems in which such pairing should be present, as exemplified by the heterostructure formed by one monolayer of MoS_2 placed on superconducting Pb, that we discuss in the second part of the manuscript. In particular by showing what are the necessary elements that a van der Waals heterostructure must have to exhibit odd-frequency SC it adds this important class of systems to the odd-frequency playbook. Our work also makes possible to select among such systems, the ones in which a direct observation – for example via scanning tunneling microscopy (STS) and angle resolved photoemission spectroscopy (ARPES) – of the signatures due to odd-frequency SC is more readily achievable.

The Hamiltonian (H) describing the most general heterostructure formed by a 2D electron gas (2DEG) and a superconductor can be written as $H = H_{2D} + H_{SC} + H_t$ where

$$H_{2D} = \sum_{\mathbf{k},\sigma,\sigma'} c^{\dagger}_{\mathbf{k},\sigma} \left[h_0(\mathbf{k})\sigma_0 + \mathbf{h}(\mathbf{k}) \cdot \boldsymbol{\sigma} \right]_{\sigma,\sigma'} c_{\mathbf{k},\sigma'}$$
(1)

$$H_{SC} = \sum_{\mathbf{k}\sigma\sigma'} d^{\dagger}_{\mathbf{k}\sigma} h^{SC}_{\sigma\sigma'}(\mathbf{k}) d_{\mathbf{k}\sigma'} + \sum_{\mathbf{k}\sigma\sigma'} d^{\dagger}_{\mathbf{k}\sigma} \Delta_{\mathbf{k}\sigma\sigma'} d^{\dagger}_{-\mathbf{k}\sigma'} + \text{h.c.}$$
(2)

$$H_t = t \sum_{\mathbf{k},\sigma} d^{\dagger}_{\mathbf{k},\sigma} c_{\mathbf{k},\sigma} + \text{h.c.}$$
(3)

are the Hamiltonians describing the 2DEG, the superconductor, and the tunneling between the two systems, respectively. In Eqs. (1)-(3) σ_0 is the identity matrix in spin space, $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)$ is the vector of Pauli matrices in spin space, $c_{\mathbf{k},\sigma}^{\dagger}$ $(d_{\mathbf{k},\sigma}^{\dagger})$ and $c_{\mathbf{k},\sigma}$ $(d_{\mathbf{k},\sigma})$ are the creation and annihilation operators, respectively, acting on the fermionic states in the 2DEG (SC) layer with momentum **k** and spin σ , $h_0(\mathbf{k})$ is the spin-independent part of H_{2D} and $\mathbf{h}(\mathbf{k})$ is the field that describes its spin-dependent part due to an exchange field and/or spin-orbit coupling, $h^{SC}_{\sigma,\sigma'}(\mathbf{k})$ describes the quasiparticle spectrum of the normal state of the superconductor, $\Delta_{\mathbf{k};\sigma,\sigma'}$ is the superconducting gap, and t is the tunneling between the 2D system and the SC. We assume the tunneling to conserve both spin and momentum given that this is the most common situation and to be able to identify the most general condition to realize odd-frequency SC without having to resort to spin-active interfaces that are often difficult to realize experimentally. To keep the treatment general we make no assumptions on the form of $\mathbf{h}(\mathbf{k})$, $h_{\sigma,\sigma'}^{SC}(\mathbf{k})$, and $\Delta_{\mathbf{k};\sigma,\sigma'}$.

The anomalous Green's function associated with the superconductor described by Eq. (2) is given by $\hat{F}_{\mathbf{k};i\omega_n}^{SC} = \left[\hat{\Delta}_{-\mathbf{k}}^{\dagger} - \left(i\omega_n + \hat{h}^{SC}(-\mathbf{k})^*\right)\hat{\Delta}_{\mathbf{k}}^{-1}\left(i\omega_n - \hat{h}^{SC}(\mathbf{k})\right)\right]^{-1}$. We can parameterize this matrix in terms of singlet and triplet parts:

$$\hat{F}_{\mathbf{k};i\omega_n}^{SC} = \left(s_{\mathbf{k},i\omega_n}^{SC}\sigma_0 + \mathbf{d}_{\mathbf{k},i\omega_n} \cdot \boldsymbol{\sigma}\right)i\sigma_2 \tag{4}$$

where ω_n is the Matsubara frequency, and $s_{\mathbf{k},i\omega_n}^{SC}$ and the three-component complex vector $\mathbf{d}_{\mathbf{k},i\omega_n}^{40}$ give the singlet and triplet superconducting amplitudes, respectively. The leading order contributions to the proximityinduced superconducting pairing in the 2DEG are given by:

$$\hat{F}_{\mathbf{k};i\omega_n}^{2D} = t^2 \; \hat{G}_{\mathbf{k};i\omega_n}^{2D} \; \hat{F}_{\mathbf{k};i\omega_n}^{SC} \; \left(\hat{G}_{-\mathbf{k};-i\omega_n}^{2D} \right)^T \tag{5}$$

where

$$\hat{G}_{\mathbf{k};i\omega_n}^{2D} = \frac{(i\omega_n - h_0(\mathbf{k}))\sigma_0 + \mathbf{h}(\mathbf{k}) \cdot \boldsymbol{\sigma}}{(i\omega_n - h_0(\mathbf{k}))^2 - |\mathbf{h}(\mathbf{k})|^2}$$
(6)

is the Green's function associated with the 2DEG.

It is convenient to separate the anomalous Green's function $\hat{F}^{2D}_{\mathbf{k};i\omega_n}$ into two parts $\hat{F}^{2D}_{\mathbf{k};i\omega_n} = A_{\mathbf{k};i\omega_n} \left(F^{odd}_{\mathbf{k};i\omega_n} + F^{even}_{\mathbf{k};i\omega_n}\right)$ where $A_{\mathbf{k};i\omega_n}$ is generally a function even in ω_n^{41} , and $F^{odd}_{\mathbf{k};i\omega_n}$ and $F^{even}_{\mathbf{k};i\omega_n}$ are the odd- and even- frequency 2×2 matrices describing the spin structure of the induced superconducting pairs respectively.

Let $\mathbf{h}_{\pm}(\mathbf{k}) \equiv \mathbf{h}(\mathbf{k}) \pm \mathbf{h}(-\mathbf{k})$. Then for $F_{\mathbf{k};i\omega_n}^{even}$ we find:

$$F^{even}_{\mathbf{k};i\omega_n} = \left(S^{even}_{\mathbf{k};i\omega_n}\sigma_0 + \mathbf{D}^{even}_{\mathbf{k};i\omega_n} \cdot \boldsymbol{\sigma}\right)i\sigma_2$$

where $S_{\mathbf{k};i\omega_n}^{even}$, $\mathbf{D}_{\mathbf{k};i\omega_n}^{even}$ are the singlet and triplet compo-

nents, respectively, given by:

$$S_{\mathbf{k};i\omega_{n}}^{even} = \left[\omega_{n}^{2} + h_{0}^{2}(\mathbf{k}) - \frac{1}{4}(|\mathbf{h}_{+}(\mathbf{k})|^{2} - |\mathbf{h}_{-}(\mathbf{k})|^{2})\right]s_{\mathbf{k};i\omega_{n}}^{SC}$$
$$- \left[h_{0}(\mathbf{k})\mathbf{h}_{-}(\mathbf{k}) + \frac{i}{2}\mathbf{h}_{+}(\mathbf{k}) \times \mathbf{h}_{-}(\mathbf{k})\right] \cdot \mathbf{d}_{\mathbf{k};i\omega_{n}}$$
$$\mathbf{D}_{\mathbf{k};i\omega}^{even} = \left[\omega_{n}^{2} + h_{0}^{2}(\mathbf{k}) + \frac{1}{4}(|\mathbf{h}_{+}(\mathbf{k})|^{2} - |\mathbf{h}_{-}(\mathbf{k})|^{2})\right]\mathbf{d}_{\mathbf{k};i\omega_{n}}$$
$$-ih_{0}(\mathbf{k})\mathbf{h}_{+}(\mathbf{k}) \times \mathbf{d}_{\mathbf{k};i\omega_{n}} - \frac{1}{2}\mathbf{h}_{+}(\mathbf{k})\left(\mathbf{h}_{+}(\mathbf{k}) \cdot \mathbf{d}_{\mathbf{k};i\omega_{n}}\right)$$
$$+ \frac{1}{2}\mathbf{h}_{-}(\mathbf{k})\left(\mathbf{h}_{-}(\mathbf{k}) \cdot \mathbf{d}_{\mathbf{k};i\omega_{n}}\right)$$
$$- \left[h_{0}(\mathbf{k})\mathbf{h}_{-}(\mathbf{k}) - \frac{i}{2}\mathbf{h}_{+}(\mathbf{k}) \times \mathbf{h}_{-}(\mathbf{k})\right]s_{\mathbf{k};i\omega_{n}}^{SC}.$$
$$\tag{7}$$

The first line (three lines) of the expression for $S^{even}_{\mathbf{k};i\omega_n}$ $(\mathbf{D}_{\mathbf{k}:i\omega_n}^{even})$ show that, as expected a singlet (triplet) pairing is induced, via the proximity effect, in the 2DEG by a singlet (triplet) superconductor, regardless of the value of **h**. The last line for the expression of $S^{even}_{\mathbf{k};i\omega_n}$ ($\mathbf{D}^{even}_{\mathbf{k};i\omega_n}$) shows that if $\mathbf{h}_{-} \neq 0$, by proximity effect, in the 2DEG we will have even-frequency superconductivity with both singlet and triplet pairing even if the substrate superconductor only has singlet or triplet pairing. It also shows that the strength of the pairing in the 2DEG with spin-structure different from the one of the substrate is proportional to $\mathbf{h}_{-}(\mathbf{k})$ and is augmented when $\mathbf{h}_{-} \times \mathbf{h}_{+} \neq 0$. This result shows how the presence of spin-orbit coupling, that gives rise to $\mathbf{h}_{-} \neq 0$, qualitatively affects the nature of the conventional (even-frequency) superconducting pairing induced by proximity. We then find that the interplay of the field **h** in the 2DEG, and the superconducting pairing in the substrate gives rise to an odd-frequency pairing term:

$$F_{\mathbf{k};i\omega_n}^{odd} = i\omega_n \left(S_{\mathbf{k};i\omega_n}^{odd} \sigma_0 + \mathbf{D}_{\mathbf{k};i\omega_n}^{odd} \cdot \boldsymbol{\sigma} \right) i\sigma_2$$

with $S^{odd}_{\mathbf{k};i\omega_n}$, $\mathbf{D}^{odd}_{\mathbf{k};i\omega_n}$ the odd-frequency singlet and triplet components, respectively, given by:

$$\begin{aligned} S^{odd}_{\mathbf{k};i\omega_n} &= -\mathbf{h}_+(\mathbf{k}) \cdot \mathbf{d}_{\mathbf{k};i\omega_n} \\ \mathbf{D}^{odd}_{\mathbf{k};i\omega} &= -\mathbf{h}_+(\mathbf{k}) s^{SC}_{\mathbf{k};i\omega_n} - i\mathbf{h}_-(\mathbf{k}) \times \mathbf{d}_{\mathbf{k};i\omega_n}. \end{aligned} \tag{8}$$

This result clearly shows that it *is* possible to get an oddfrequency singlet term provided the substrate is a triplet superconductor with a **d** vector that is not perpendicular to the even component of \mathbf{h} , \mathbf{h}_+ . Notice that because \mathbf{h} and **d** belong to different layers they are not constrained to be in any specific relation. Eq. (8) also shows that an odd-frequency triplet term will be present if both \mathbf{h}_+ and the singlet pairing in the substrate s^{SC} are not zero, as shown previously 9,32,33 . Eq. (8) therefore shows that when $\mathbf{h}_{+} \neq 0$, and $\mathbf{h}_{-} = 0$, by proximity effect, we will have odd-frequency superconductivity in the 2DEG that has the "opposite" spin structure from the superconductivity in the substrate: triplet if the substrate is a singlet superconductor, singlet if the substrate is a triplet superconductor (with d not ortogonal to h_{+}). A very interesting and novel result is that even when $\mathbf{h}_{+} = 0$, i.e. no



FIG. 1. (Color online) a) Unit cell for a monolayer TMD. A single monolayer is composed of three covalently bonded layers trigonally coordinated with a layer of transition metal sandwiched between two layers of chalcogen. b) Schematic of a heterostructure formed by exfoliating a TMD monolayer onto a superconductor. c) Sketch of the band structure of a TMD monolayer with the *d*-electron bands appearing at the K and K' points with a band gap of 1.8eV separating a pair of spin-degenerate conduction bands from a pair of spin-polarized bands. Notice that the polarization is different in the two inequivalent valleys (K and K').

ferromagnetism is present in the 2DEG, we can have oddfrequency superconductivity in the 2DEG, without having to assume the presence of a spin-active interface, if the 2DEG has spin-orbit coupling, so that $\mathbf{h}_{-} \neq 0$, and the substrate is a triplet superconductor with **d** not parallel to \mathbf{h}_{-} (again, we emphasize that because \mathbf{h}_{-} and **d** belong to different layers they are not locked to each other). This is a result that significantly enlarges the set of engineered structures in which to look for odd-frequency superconductivity by adding a whole new class of heterostructures. As we show below, a system that falls into this class is a heterostructure formed by a group-VI dichalcogenide monolayer and a superconductor's surface with Rashba spin-orbit coupling.

Transition metal dichalcogenides (TMDs), such as molybdenum disulfide (MoS₂) have recently received a lot of attention due to their unusual electronic properties and their potential for applications in electronics. MoS₂ can be exfoliated down to monolayer 2D crystals^{4,42–44}. These monolayers have been shown to possess a direct band gap of $1.8 \text{eV}^{4,45}$, they can be gated⁴, and have exhibited electron mobilities as high as 200 cm²V⁻¹s⁻¹⁴. Furthermore, the *d*-electron states exhibit a valley degree of freedom that is coupled to the electron spin^{46–48}. In the context of our problem, this material is of great interest not only because it is a two-dimensional material that is readily available, easily manufactured and incorporated into heterostructures, but also because of its strong spin-orbit coupling. Consider the heterostructure shown in Fig. 1 composed of a transition metal dichalcogenide (TMD) monolayer on top of a superconductor. The low-energy electronic states of an TMD monolayer are well described by the following valley-dependent Hamiltonian⁴⁶:

$$\hat{H}_{\mathbf{k},\lambda}^{TMD} = \left[a\gamma \left(\lambda k_x \tau_1 + k_y \tau_2 \right) + \frac{u}{2} \tau_3 - \mu \tau_0 \right] \otimes \sigma_0 - \frac{\lambda \alpha}{2} \left(\tau_3 - \tau_0 \right) \otimes \sigma_3$$
(9)

where τ_i are Pauli matrices acting on the orbital space of the TMD monolayer, a is the lattice constant, γ is the effective hopping integral, u is the energy gap between the valence and conduction bands, α is the strength of the spin-orbit coupling, $\lambda = \pm 1$ is the valley index ($\lambda = 1$ denotes the K valley, $\lambda = -1$ denotes the K' valley, see Fig. 1), $\mathbf{k} = (k_x, k_y, 0)$ is a vector describing small deviations from the K or K' point in k-space, and μ is the chemical potential. For MoS₂: a = 3.193 Å, $\gamma = 1.10$ eV, u = 1.66 eV, and $2\alpha = 0.15$ eV⁴⁶.

The Hamiltonian in Eq. (9) possesses four eigenstates at the K and K' points; two spin-degenerate conduction states separated by an eV-scale gap from two spin polarized valence states, as shown in Fig. 1. For our analysis the most interesting case is when MoS₂ is hole doped. For this reason in the following we will use an effective 2band model in which we include only the valence bands. Considering the large gap between the valence and the conduction bands this does not introduce any inaccuracy. For small k the valence band Hamiltonian can be written in spin space as:

$$\hat{H}_{\mathbf{k},\lambda}^{TMD} = -\left(\frac{a^2\gamma^2}{u}k^2 + \frac{u}{2} + \mu\right)\sigma_0 + \lambda\alpha\sigma_3.$$
(10)

Notice that, taking into account the valley index λ , for the parity operator, \mathcal{P} , acting on a function, $f(\mathbf{k}, \lambda)$ we have $\mathcal{P}f(\mathbf{k}, \lambda) = f(-\mathbf{k}, -\lambda)$. Using the notation used in Eqs (7) and (8) we then find that in this case $h_0(\mathbf{k}) = -\left(\frac{a^2\gamma^2}{u}k^2 + \frac{u}{2} + \mu\right)$, $\mathbf{h}_+(\mathbf{k}, \lambda) = 0$, and $\mathbf{h}_-(\mathbf{k}, \lambda) = 2\lambda\alpha\hat{z}$, where \hat{z} is the unit vector normal to the TMD monolayer. Starting from the general Eqs (7) and (8) we then find:

$$S_{\mathbf{k},\lambda;i\omega_{n}}^{even} = \left(\omega_{n}^{2} + \xi_{\mathbf{k}}^{2} + \alpha^{2}\right) s_{\mathbf{k},\lambda;i\omega_{n}}^{SC} - 2\lambda\alpha\xi_{\mathbf{k}}\hat{z}\cdot\mathbf{d}_{\mathbf{k},\lambda;i\omega_{n}}$$
$$\mathbf{D}_{\mathbf{d},\lambda;i\omega}^{even} = \left(\omega_{n}^{2} + \xi_{\mathbf{k}}^{2} - \alpha^{2}\right) \mathbf{d}_{\mathbf{k},\lambda;i\omega_{n}} + 2\alpha^{2} \left(\hat{z}\cdot\mathbf{d}_{\mathbf{k},\lambda;i\omega_{n}}\right)\hat{z} - 2\lambda\alpha\xi_{\mathbf{k}}s_{\mathbf{k},\lambda;i\omega_{n}}^{SC}\hat{z}$$
(11)

and

$$S^{odd}_{\mathbf{k},\lambda;i\omega_n} = 0$$

$$\mathbf{D}^{odd}_{\mathbf{k},\lambda;i\omega} = -i2\lambda\alpha\hat{z} \times \mathbf{d}_{\mathbf{k},\lambda;i\omega_n}$$
(12)

In accordance with Eq. (8) we find that, given that $\mathbf{h}_{+} = 0$, to get odd frequency superconductivity in the TMD we need a substrate with non-zero triplet superconducting pairing. In general this situation is realized in non-centrosymmetric superconductors. Additionally, this condition can be realized at the surface of centrosymmetric singlet superconductors with spin-orbit coupling since the surface breaks inversion symmetry leading to the appearance of a Rashba spin-orbit term that in turn induces a superconducting triplet component⁴⁹. This is expected to be the case for the surface of superconducting Pb.

Considering the case in which the superconductor in Fig. 1 (b) has Rashba spin-orbit coupling, the Hamiltonian matrix describing the single particle spectrum of the superconductor is $\hat{h}^{SC}(\mathbf{k}) = \epsilon_{\overline{\mathbf{k}}} \hat{\sigma}_0 + \eta \hat{z} \cdot (\boldsymbol{\sigma} \times \overline{\mathbf{k}})$ where η is the Rashba spin-orbit coupling in the superconductor surface, $\epsilon_{\overline{\mathbf{k}}}$ is the dispersion of the normal state quasiparticles in the absence of spin-orbit coupling, and $\overline{\mathbf{k}}$ is the momentum measured from the Brillouin zone center. Considering that the dominant pairing is intraband we obtain^{41,49}

$$\hat{F}_{\overline{\mathbf{k}};i\omega_n}^{SC} = \frac{\Delta}{(s_{\overline{\mathbf{k}};i\omega_n}^{SC})^2 - |\mathbf{d}_{\overline{\mathbf{k}}}|^2} (s_{\overline{\mathbf{k}};i\omega_n}^{SC} \sigma_0 + \mathbf{d}_{\overline{\mathbf{k}}} \cdot \boldsymbol{\sigma}) i\sigma_2 \quad (13)$$

where Δ is the substrate's superconducting gap, $s_{\mathbf{k};i\omega_n}^{SC} = \Delta^2 + \omega_n^2 + \epsilon_{\mathbf{k}}^2 + \eta^2 \overline{\mathbf{k}}^2$ and $\mathbf{d}_{\mathbf{k}} = 2\epsilon_{\mathbf{k}}\eta(-\overline{k}_y,\overline{k}_x,0)$. The key point of Eq. (13) is that thanks to the Rashba spin-orbit coupling induced by the breaking of the inversion symmetry at the surface of the Pb substrate a triplet term appears in the \hat{F}^{SC} and that in addition the **d** vector for such triplet component is perpendicular to the field \mathbf{h}_- in the TMD monolayer. The interplay of such triplet component with the spin-orbit cupling of the TMD monolayer gives rise to odd-frequency SC in the TMD.

With the above definitions we can follow the same steps leading to Eqs 8 and 7 and obtain the leading order contribution to the proximity-induced anomalous Green's function in the TMD layer as $\hat{F}_{\mathbf{k},\lambda;i\omega_n}^{TMD} = A_{\mathbf{k},\lambda;i\omega_n}^{TMD} \left(F_{\mathbf{k},\lambda;i\omega_n}^{odd} + F_{\mathbf{k},\lambda;i\omega_n}^{even}\right)$ where

$$A_{\mathbf{k},\lambda;i\omega_n}^{TMD} = \frac{\Delta t^2}{[(i\omega_n - \xi_{\mathbf{k}})^2 - \alpha^2]^2 [(s_{\mathbf{k}+\mathbf{K}_\lambda}^{SC};i\omega_n)^2 - |\mathbf{d}_{\mathbf{k}+\mathbf{K}_\lambda}|^2]}$$

For the even-frequency singlet and triplet components of \hat{F}^{TMD} we find:

$$S_{\mathbf{k},\lambda;i\omega_{n}}^{even} = \left(\omega_{n}^{2} + \xi_{\mathbf{k}}^{2} + \alpha^{2}\right) s_{\mathbf{k}+\mathbf{K}_{\lambda};i\omega_{n}}^{SC} \mathbf{D}_{\mathbf{k},\lambda;i\omega}^{even} = -\left(\omega_{n}^{2} + \xi_{\mathbf{k}}^{2} - \alpha^{2}\right) \mathbf{d}_{\mathbf{k}+\mathbf{K}_{\lambda}} - 2\lambda\alpha\xi_{\mathbf{k}}s_{\mathbf{k}+\mathbf{K}_{\lambda};i\omega_{n}}^{SC}$$
(14)

Given that $\mathbf{h}_{+} = 0$, see Eq. (12), the odd-frequency singlet component vanishes whereas for the triplet component we find:

$$\mathbf{D}_{\mathbf{k},\lambda;i\omega}^{odd} = i4\lambda\alpha\eta\epsilon_{\mathbf{k}+\mathbf{K}_{\lambda}}(\mathbf{k}+\mathbf{K}_{\lambda}) \tag{15}$$

where \mathbf{K}_{λ} is the momentum vector at the K(K') point for $\lambda = 1$ ($\lambda = -1$). Eq. (15) shows that in the TMD the odd-frequency triplet component has a d-vector pointing in the direction of the momentum. One can verify that this corresponds to an equal-spin spin triplet amplitude given by $F_{\uparrow\uparrow/\downarrow\downarrow}^{TMD} \sim i\omega_n \eta \alpha \epsilon_{\overline{\mathbf{k}}} \lambda \left(\overline{k}_y \pm i \overline{k}_x \right)$ which is proportional to the product of the spin-orbit couplings in the two materials. Consistent with the general case, we see that the emergence of this term requires the spin-orbit couplings in the two media to be non parallel.

Our results add a new class of systems, Van der Waals (VdW) heterostructures, to the odd-frequency playbook. Van der Waals systems have many advantages: i) the 2DEG in which odd-frequency pairing is present lives in a layer with an exposed surface, a fact allows for ideal STS and ARPES measurements; ii) as shown by the example of the MoS_2/Pb heterostructure, it is possible to realize VdW systems with no ferromagnetic layers, or spin-active interfaces that exhibit odd-frequency SC; iii) the 2DEG in which odd-frequency pairing is present can be just one atom thick, this fact removes many of the complications associated with the interpretation of STS and ARPES data done in heterostructures in which each layer is several nanometers thick; iv) because the top layer is just one atom thick the electrons are truly confined in 2D, this fact, combined with the fact that according to our results the top layer can be a semiconductor, rather than a ferromagnetic metal as in previous proposals, ensures that the DOS of the normal state is quite low and therefore allows for an easier observation of the features in the DOS due to the presence of odd-frequency pairing.

In conclusion, in this work we investigated the symmetries of proximity-induced superconducting pairing amplitudes in a 2DEG coupled to a superconductor. We arrived at a general expression relating the induced pairing amplitudes to the components of the anomalous Green's function of the superconducting substrate and the elements of the 2DEG Hamiltonian matrix, $\hat{h}(\mathbf{k}) =$ $h_0(\mathbf{k})\hat{\sigma}_0 + \mathbf{h}(\mathbf{k}) \cdot \boldsymbol{\sigma}$. We have shown that the interplay of the spin-orbit coupling in the 2DEG and the superconducting pairing of the substrate can give rise, via proximity effect, to unusual superconducting pairings in the 2DEG. We find that even when no ferromagnetism is present in the 2DEG, and there is no spin-active interface, odd-frequency superconductivity can be induced in the 2DEG provided the 2DEG has spin-orbit coupling and the substrate has some triplet superconductivity. We then showed that this condition can be realized in a MoS_2/Pb heterostructure. This result, combined with the general equations that we obtain, adds a new class of systems, Van der Waals (VdW) heterostructures, to the odd-frequency playbook.

Acknowledgements: We wish to thank Annica Black-Schaffer, Matthias Eschrig, Satrio Gani, Martin Rodriguez-Vega, Yudistira Virgus, and Junhua Zhang for useful discussions. The work of C.T. and A.V.B. was supported by the European Research Council (ERC) DM-321031 and US DOE BES E304, DMB and ER have been supported by ONR-N00014-13-1-0321 and NSF-DMR-1455233.

- ¹ J. Xiang, W. Lu, Y. Hu, Y. Wu, H. Yan, and C. M. Lieber, Nature **441**, 489 (2006).
- ² J. R. Miller, R. Outlaw, and B. Holloway, Science **329**, 1637 (2010).
- ³ Y. Zhu, S. Murali, M. D. Stoller, K. Ganesh, W. Cai, P. J. Ferreira, A. Pirkle, R. M. Wallace, K. A. Cychosz, M. Thommes, et al., Science **332**, 1537 (2011).
- ⁴ B. Radisavljevic, A. Radenovic, J. Brivio, V. Giacometti, and A. Kis, Nature nanotechnology **6**, 147 (2011).
- ⁵ A. Geim and I. Grigorieva, Nature **499**, 419 (2013).
- ⁶ J. Zhang and E. Rossi, Phys. Rev. Lett. **111**, 086804 (2013).
- ⁷ K.-H. Jin and S.-H. Jhi, Phys. Rev. B 87, 075442 (2013).
- ⁸ J. Zhang, C. Triola, and E. Rossi, Phys. Rev. Lett. **112**, 096802 (2014).
- ⁹ C. Triola, E. Rossi, and A. V. Balatsky, Phys. Rev. B 89, 165309 (2014).
- ¹⁰ E. Demler, G. Arnold, and M. Beasley, Phys. Rev. B 55, 15174 (1997).
- ¹¹ T. Tokuyasu, J. Sauls, and D. Rainer, Phys. Rev. B 38, 8823 (1988).
- ¹² A. Black-Schaffer and A. Balatsky, Phys. Rev. B 87, 220506(R) (2013).
- ¹³ J. Linder, Y. Tanaka, T. Yokoyama, A. Sudbø, and N. Nagaosa, Phys. Rev. B **81**, 184525 (2010).
- ¹⁴ J. Linder, T. Yokoyama, A. Sudbø, and M. Eschrig, Phys. Rev. Lett. **102**, 107008 (2009).
- ¹⁵ F. Parhizgar and A. M. Black-Schaffer, Physical Review B 90, 184517 (2014).
- ¹⁶ L. Fu and C. L. Kane, Physical Review Letters **100**, 096407 (2008).
- ¹⁷ J. D. Sau, R. M. Lutchyn, S. Tewari, and S. D. Sarma, Phys. Rev. Lett. **104**, 040502 (2010).
- ¹⁸ R. M. Lutchyn, J. D. Sau, and S. D. Sarma, Phys. Rev. Lett. **105**, 077001 (2010).
- ¹⁹ Y. Oreg, G. Refael, and F. von Oppen, Phys. Rev. Lett. 105, 177002 (2010).
- ²⁰ M. Z. Hasan and C. L. Kane, Reviews of Modern Physics 82, 3045 (2010).
- ²¹ X.-L. Qi and S.-C. Zhang, Reviews of Modern Physics 83, 1057 (2011).
- ²² J. Alicea, Nature nanotechnology **8**, 623 (2013).
- ²³ V. Mourik, K. Zuo, S. Frolov, S. Plissard, E. Bakkers, and L. Kouwenhoven, Science **336**, 1003 (2012).
- ²⁴ L. P. Rokhinson, X. Liu, and J. K. Furdyna, Nature Physics 8, 795 (2012).
- ²⁵ A. Das, Y. Ronen, Y. Most, Y. Oreg, M. Heiblum, and H. Shtrikman, Nature Physics 8, 887 (2012).
- ²⁶ M. Deng, C. Yu, G. Huang, M. Larsson, P. Caroff, and H. Xu, Nano letters **12**, 6414 (2012).

- ²⁷ A. Finck, D. Van Harlingen, P. Mohseni, K. Jung, and X. Li, Physical review letters **110**, 126406 (2013).
- ²⁸ H. Churchill, V. Fatemi, K. Grove-Rasmussen, M. Deng, P. Caroff, H. Xu, and C. M. Marcus, Physical Review B 87, 241401 (2013).
- ²⁹ V. L. Berezinskii, Pis' ma Zh. Eksp. Teor. Fiz. **20**, 628 (1974).
- ³⁰ A. Balatsky and E. Abrahams, Phys. Rev. B 45, 13125 (1992).
- ³¹ E. Abrahams, A. Balatsky, J. R. Schrieffer, and P. B. Allen, Phys. Rev. B 47, 513 (1993), URL http://link.aps.org/ doi/10.1103/PhysRevB.47.513.
- ³² M. Eschrig and T. Löfwander, Nature Physics 4, 138 (2008).
- ³³ J. Linder, T. Yokoyama, and A. Sudbø, Phys. Rev. B 77, 174514 (2008).
- ³⁴ T. Yokoyama, Phys. Rev. B 86, 075410 (2012).
- ³⁵ A. Black-Schaffer and A. Balatsky, Phys. Rev. B 86, 144506 (2012).
- ³⁶ J. Linder, A. Sudbø, T. Yokoyama, R. Grein, and M. Eschrig, Phys. Rev. B **81**, 214504 (2010).
- ³⁷ Y. Tanaka, Y. Tanuma, and A. Golubov, Phys. Rev. B 76, 054522 (2007).
- ³⁸ Y. Tanaka, M. Sato, and N. Nagaosa, J. Phys. Soc. Jpn. 81, 011013 (2012).
- ³⁹ A. Di Bernardo, S. Diesch, Y. Gu, J. Linder, G. Divitini, C. Ducati, E. Scheer, M. G. Blamire, and J. W. Robinson, Nature communications 6 (2015).
- ⁴⁰ A. P. Mackenzie and Y. Maeno, Rev. Mod. Phys. **75**, 657 (2003).
- ⁴¹ Supplementary material.
- ⁴² K. Novoselov, D. Jiang, F. Schedin, T. Booth, V. Khotkevich, S. Morozov, and A. Geim, Proceedings of the National Academy of Sciences of the United States of America **102**, 10451 (2005).
- ⁴³ K. F. Mak, C. Lee, J. Hone, J. Shan, and T. F. Heinz, Physical Review Letters **105**, 136805 (2010).
- ⁴⁴ J. S. Ross, S. Wu, H. Yu, N. J. Ghimire, A. M. Jones, G. Aivazian, J. Yan, D. G. Mandrus, D. Xiao, W. Yao, et al., Nature communications 4, 1474 (2013).
- ⁴⁵ A. Splendiani, L. Sun, Y. Zhang, T. Li, J. Kim, C.-Y. Chim, G. Galli, and F. Wang, Nano letters **10**, 1271 (2010).
- ⁴⁶ D. Xiao, G.-B. Liu, W. Feng, X. Xu, and W. Yao, Physical Review Letters **108**, 196802 (2012).
- ⁴⁷ C. Mai, A. Barrette, Y. Yu, Y. G. Semenov, K. W. Kim, L. Cao, and K. Gundogdu, Nano letters **14**, 202 (2013).
- ⁴⁸ X. Xu, W. Yao, D. Xiao, and T. F. Heinz, Nature Physics 10, 343 (2014).
- ⁴⁹ L. P. Gor'kov and E. I. Rashba, Physical review letters 87, 037004 (2001).