

2016

Preservers of unitary similarity functions on Lie products of matrices

Jianlian Cui

Tsinghua Univ, Dept Math, Beijing 100084, Peoples R China;

Chi-Kwong Li

College of William & Mary, Dept Math, Williamsburg, VA 23187 USA

Yiu-Tung Poon

Iowa State Univ, Dept Math, Ames, IA 50011 USA

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Cui, J., Li, C. K., & Poon, Y. T. (2016). Preservers of unitary similarity functions on Lie products of matrices. *Linear Algebra and its Applications*, 498, 160-180.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

PRESERVERS OF UNITARY SIMILARITY FUNCTIONS ON LIE PRODUCTS OF MATRICES

JIANLIAN CUI, CHI-KWONG LI, AND YIU-TUNG POON

In memory of Professor Hans Schneider.

ABSTRACT. Denote by M_n the set of $n \times n$ complex matrices. Let $f : M_n \rightarrow [0, \infty)$ be a continuous map such that $f(\mu U A U^*) = f(A)$ for any complex unit μ , $A \in M_n$ and unitary $U \in M_n$, $f(X) = 0$ if and only if $X = 0$ and the induced map $t \mapsto f(tX)$ is monotonically increasing on $[0, \infty)$ for any rank 1 nilpotent $X \in M_n$. Characterization is given for surjective maps ϕ on M_n satisfying $f(AB - BA) = f(\phi(A)\phi(B) - \phi(B)\phi(A))$. The general theorem are then used to deduce results on special cases when the function is the pseudo spectrum and the pseudo spectral radius.

AMS Subject Classification Primary 15A60, 46B04

Keywords Lie product, unitary similarity invariant function, pseudo spectrum.

1. INTRODUCTION

Let M_n be the set of $n \times n$ matrices. A function $f : M_n \rightarrow \mathbb{R}$ is a *radial unitary similarity invariant function* if

$$(P1) \quad f(\mu U A U^*) = f(A) \text{ for a complex unit } \mu, A \in M_n \text{ and unitary } U \in M_n.$$

In [11], the authors studied unitary similarity invariant functions that are norms on M_n , and determine the structure of maps $\phi : M_n \rightarrow M_n$ satisfying

$$(1.1) \quad f(AB - BA) = f(\phi(A)\phi(B) - \phi(B)\phi(A)) \quad \text{for all } A, B \in M_n.$$

In [11, Remark 2.7], it was pointed out that the result actually holds for more general unitary similarity invariant functions. However, no detail was given, and it is not straightforward to apply the results to a specific problem. For instance, it is unclear how one can apply the result to study preservers of pseudo spectrum of Lie product of matrices;¹ see the definition in Section 3. To fill this gap, we extend the result in [11] to continuous radial unitary similarity invariant functions $f : M_n \rightarrow \mathbb{R}$ satisfying the following properties.

(P2) For any $X \in M_n$ we have $f(X) = f(0_n)$ if and only if $X = 0_n$, the $n \times n$ zero matrix.

(P3) For any rank 1 nilpotent $X \in M_n$, the map $t \mapsto f(tX)$ on $[0, \infty)$ is strictly increasing.

For a function $f : M_n \rightarrow [0, \infty)$ satisfying (P1) – (P3), we show that if $\phi : M_n \rightarrow M_n$ is a surjective map satisfying (1.1), then there is a unitary $U \in M_n$ and a subset \mathcal{N}_n of normal matrices in M_n such that ϕ has the form

$$\phi(A) = \begin{cases} \mu_A U A^\dagger U^* + \nu_A I_n & A \in M_n \setminus \mathcal{N}_n \\ \mu_A U (A^\dagger)^* U^* + \nu_A I_n & A \in \mathcal{N}_n, \end{cases}$$

¹This is a question raised by Professor Molnar to the second and third author at the 2014 Summer Conference of the Canadian Mathematics Society.

where

- (a) $\mu_A, \nu_A \in \mathbb{C}$ with $|\mu_A| = 1$, depending on A ,
- (b) $A^\dagger = A, \bar{A}, A^t$ or A^* , and
- (c) \mathcal{N}_n depends on the given unitarily invariant function f .

The proof of this result will be given in Section 2. In Section 3, we apply the main result to the case when f is the pseudo spectral radius, and then obtain the result for the case when f is the pseudo spectrum.

For other preserver problems on different types of products on matrices and operators, one may see [1, 2, 3, 7, 11, 12] and their references.

2. MAIN THEOREM

In this section, we prove Theorem 2.1 extending the result in [11]. We use similar ideas in [11] with some intricate arguments to make the extension possible.

Theorem 2.1. *Let $f : M_n \rightarrow [0, \infty)$ be a function on M_n satisfying (P1) – (P3). Suppose $n \geq 3$, and $\phi : M_n \rightarrow M_n$ is a surjective map satisfying*

$$f([\phi(A), \phi(B)]) = f([A, B]).$$

Then there is a unitary matrix U and a subset \mathcal{N}_n of normal matrices with non-collinear eigenvalues such that ϕ has the form

$$\phi(A) = \begin{cases} \mu_A U \psi(A) U^* + \nu_A I_n & A \in M_n \setminus \mathcal{N}_n \\ \mu_A U \psi(A)^* U^* + \nu_A I_n & A \in \mathcal{N}_n, \end{cases}$$

where $\mu_A, \nu_A \in \mathbb{C}$ with $|\mu_A| = 1$ depending on A , and ψ is one of the maps: $A \mapsto A, A \mapsto \bar{A}, A \mapsto A^t$ or $A \mapsto A^$.*

A bijective map P on M_n is said to be a *locally regular polynomial map* [14] if for every $A \in M_n$, there exists a polynomial $p_A(t)$ such that $P(A) = p_A(A)$ and A have the same commutant. To prove the above theorem, we need the following result from Šemrl [14].

Theorem 2.2. *Suppose $n \geq 3$, and $\phi : M_n \rightarrow M_n$ is a bijective map satisfying*

$$[A, B] = 0_n \iff [\phi(A), \phi(B)] = 0_n.$$

Let Γ be the set of matrices A such that the Jordan form of A only has Jordan blocks of sizes 1 or 2. Then there are an invertible matrix S , an automorphism τ of the complex field and a regular locally polynomial map $A \rightarrow p_A(A)$ such that

$$(2.1) \quad \phi(A) = S(p_A(A_\tau^\dagger))S^{-1} \quad \text{for all } A \in \Gamma.$$

Here, X_τ is the matrix whose (i, j) -entry is $\tau(X_{ij})$, and $A^\dagger = A$ or A^t .

Our proof strategy is to show that $\phi(A)$ has the asserted form described in the theorem for a special class \mathcal{C}_1 of matrices A . Then we modify the map ϕ to ϕ_1 so that it will satisfy the same

hypothesis of ϕ with the additional assumption that $\phi(X) = X$ for every $X \in \mathcal{C}_1$. Then we can set $B = \phi(A)$ for a certain matrix A not in \mathcal{C}_1 and use the condition that

$$f([A, X]) = f([\phi_1(A), \phi_1(X)]) = f([B, X]) \quad \text{for all } X \in \mathcal{C}_1$$

to show that $B = \phi_1(A)$ also has the asserted form. Thus, ϕ_1 has the asserted form for a larger class \mathcal{C}_2 of matrices, and so on and so forth until we show that the modified map will fix every matrices after a finite number of steps.

In the next few lemmas, we will focus on the relations between a pair of matrices A and B such that

$$f([A, X]) = f([B, X]) \quad \text{for all } X \in \mathcal{C}$$

for a certain subset \mathcal{C} of matrices.

Lemma 2.3. *Suppose $A \in M_n$ is a rank one nilpotent matrix. Then $A = xy^*$ for some non-zero orthogonal vectors, x and y . Furthermore, A is unitarily similar to $\|x\|\|y\|E_{21}$.*

Proof. Suppose $A \in M_n$ is a rank one matrix. Then $A = xy^*$ for some non-zero column vectors, x and $y \in \mathbb{C}^n$. If A is nilpotent, then $A^k = 0$ for some integer $k > 1$. Then we have

$$0 = \text{tr } A^k = \text{tr } (xy^*)^k = (y^*x)^k.$$

Therefore, x and y are orthogonal. Let U be a unitary matrix with $\frac{y}{\|y\|}$ and $\frac{x}{\|x\|}$ as the first and second columns respectively. Then $U^*AU = \|x\|\|y\|E_{21}$ \square

Denote by $\sigma(A)$ the spectrum of A and $N(A)$ the null space of A .

Lemma 2.4. *For any two matrices A and B , if*

$$(2.2) \quad f([A, X]) = f([B, X]) \quad \text{for all rank one } X \in M_n,$$

then there are $\mu, \nu \in \mathbb{C}$ with $|\mu| = 1$ such that one of the following holds with $\hat{A} = \mu A + \nu I_n$.

(a) $\sigma(B) = \sigma(\hat{A})$ and for any $\lambda \in \sigma(\hat{A})$,

$$N(B - \lambda I_n) = N(\hat{A} - \lambda I_n) \quad \text{and} \quad N(B^t - \lambda I_n) = N(\hat{A}^t - \lambda I_n).$$

(b) The eigenvalues of A are not collinear, $\sigma(B) = \overline{\sigma(\hat{A})}$ and for any $\lambda \in \sigma(\hat{A})$,

$$N(B - \bar{\lambda} I_n) = N(\hat{A} - \lambda I_n) \quad \text{and} \quad N(B^t - \bar{\lambda} I_n) = N(\hat{A}^t - \lambda I_n).$$

Proof. Note that for any rank one matrix $X = xy^t$, $[C, X] = 0$ if and only if x and y^t are the right and left eigenvectors of C corresponding to the same eigenvalue. To see this, as $[C, X] = (Cx)y^t - x(y^tC)$, then $[C, X] = 0$ if and only if $Cx = \lambda x$ and $y^tC = \lambda y^t$ for some $\lambda \in \mathbb{C}$.

Suppose A and B satisfy (2.2). By the above observation on rank one matrices and property (P2) of f , A and B must have the same set of left and right eigenvectors. Furthermore, x_1 and x_2 are the right eigenvectors of A corresponding to the same eigenvalue if and only if the two eigenvectors correspond to the same eigenvalue of B . Thus, the eigenvalues of A and B have the same geometric multiplicity.

Let $\lambda_1, \dots, \lambda_k$ be the distinct eigenvalues of A with x_1, \dots, x_k and y_1, \dots, y_k being the right and left eigenvectors. Also for each pair of eigenvectors x_i and y_i^t , let γ_i be the corresponding eigenvalue

of B . Take $X_{ij} = x_i y_j^t$. Then $A X_{ij} = \lambda_i X_{ij}$ and $X_{ij} A = \lambda_j X_{ij}$. Using (P1), we see that for any $1 \leq i, j \leq n$,

$$f([A, X_{ij}]) = f(\lambda_i X_{ij} - \lambda_j X_{ij}) = f((\lambda_i - \lambda_j) X_{ij}) = f(|\lambda_i - \lambda_j| X_{ij}).$$

Similarly, $f([B, X_{ij}]) = f((\gamma_i - \gamma_j) X_{ij}) = f(|\gamma_i - \gamma_j| X_{ij})$.

By the fact that $f([A, X_{ij}]) = f([B, X_{ij}])$ and Property (P3),

$$|\lambda_i - \lambda_j| = |\gamma_i - \gamma_j| \quad \text{for all } 1 \leq i, j \leq k.$$

As a result, there are $\mu, \nu \in \mathbb{C}$ with $|\mu| = 1$ such that either

- (1) $\gamma_i = \mu \lambda_i + \nu$ for all $1 \leq i \leq k$; or
- (2) the eigenvalues of A are non-collinear and $\bar{\gamma}_i = \mu \lambda_i + \nu$ for all $1 \leq i \leq k$.

Then the result follows with $\hat{A} = \mu A + \nu I_n$. □

Lemma 2.5. *Suppose A and B commute and satisfy (2.2). If A has at least two distinct eigenvalues, then there are $\mu, \nu \in \mathbb{C}$ with $|\mu| = 1$ such that either*

- (a) $B = \mu A + \nu I_n$, or
- (b) A is normal with non-collinear eigenvalues and $B = \mu A^* + \nu I_n$.

Proof. As A and B commute, there is a unitary matrix U such that both $U^* A U$ and $U^* B U$ are upper triangular, see [9, Theorem 2.3.3]. Replacing (A, B) with $(U^* A U, U^* B U)$, we may assume that A and B are upper triangular.

As A and B satisfy (2.2), Lemma 2.4 holds. Suppose Lemma 2.4(a) holds with $\hat{A} = \mu A + \nu I_n$. Notice that $\sigma(B) = \sigma(\hat{A})$ and

$$f([\hat{A}, X]) = f([\mu A + \nu I_n, X]) = f([B, X]) \quad \text{for all rank one } X \in M_n.$$

Suppose λ is an eigenvalue of \hat{A} and $y \in N(\hat{A}^t - \lambda I_n)$. For any $z \in \mathbb{C}^n$, let $Z = z y^t$. Then $Z \hat{A} = \lambda Z$ and $[\hat{A}, Z] = (\hat{A} - \lambda I_n) Z$. Note that $(\hat{A} - \lambda I_n) Z$ has rank at most one and $\text{tr}((\hat{A} - \lambda I_n) Z) = \text{tr}([\hat{A}, Z]) = 0$, so $(\hat{A} - \lambda I_n) Z$ is unitarily similar to $\|(\hat{A} - \lambda I_n) z\| \|y^t\| E_{12}$. Thus,

$$f([\hat{A}, Z]) = f(\|(\hat{A} - \lambda I_n) z\| \|y^t\| E_{12}).$$

Similarly, $f([B, Z]) = f(\|(B - \lambda I_n) z\| \|y^t\| E_{12})$. Hence, by (P1) and (P3),

$$\|(\hat{A} - \lambda I_n) z\| = \|(B - \lambda I_n) z\| \quad \text{for all } z \in \mathbb{C}^n \text{ and } \lambda \in \sigma(\hat{A}).$$

As a result,

$$\begin{aligned} z^* \hat{A}^* \hat{A} z - 2\text{Re}(\bar{\lambda} z^* \hat{A} z) + |\lambda|^2 z^* z &= \|(\hat{A} - \lambda I_n) z\|^2 \\ &= \|(B - \lambda I_n) z\|^2 = z^* B^* B z - 2\text{Re}(\bar{\lambda} z^* B z) + |\lambda|^2 z^* z. \end{aligned}$$

This implies that

$$2\text{Re}(\bar{\lambda} z^* (\hat{A} - B) z) = z^* (\hat{A}^* \hat{A} - B^* B) z \quad \text{for all } z \in \mathbb{C}^n \text{ and } \lambda \in \sigma(\hat{A}).$$

As A has at least two distinct eigenvalues, so does \hat{A} . Taking any $\lambda, \gamma \in \sigma(\hat{A})$ with $\lambda \neq \gamma$, we have

$$2\text{Re}(\bar{\lambda} z^* (\hat{A} - B) z) = z^* (\hat{A}^* \hat{A} - B^* B) z = 2\text{Re}(\bar{\gamma} z^* (\hat{A} - B) z).$$

Thus, $W((\bar{\lambda} - \bar{\gamma})(\hat{A} - B)) \subseteq i\mathbb{R}$, where $W(X)$ is the numerical range of X .

Then $(\overline{\lambda - \gamma})(\hat{A} - B)$ is a skew-Hermitian matrix [8]. Since both \hat{A} and B are upper triangular, they must be diagonal matrices. Now for any $1 \leq i \leq n$, $b_{ii} \in \sigma(B) = \sigma(\hat{A})$. Then

$$0 = \|(B - b_{ii}I_n)e_i\| = \|(\hat{A} - b_{ii}I_n)e_i\| = \|(B - b_{ii}I_n)e_i + (\hat{A} - B)e_i\| \|(\hat{A} - B)e_i\|$$

Thus, $(\hat{A} - B)e_i = 0$ for all $1 \leq i \leq n$ and hence $B = \hat{A}$.

Now suppose Lemma 2.4(b) holds. Then by a similar argument, we can show that

$$(2.3) \quad \|(\hat{A} - \lambda I_n)z\| = \|(B - \bar{\lambda}I_n)z\| \quad \text{for all } \lambda \in \sigma(\hat{A}) \text{ and } z \in \mathbb{C}^n$$

and so $(\overline{\lambda - \gamma})\hat{A} - (\lambda - \gamma)B$ is a skew-Hermitian matrix. It follows that $(\overline{\lambda - \gamma})T_A - (\lambda - \gamma)T_B = 0$, or equivalently, $T_B = \frac{\overline{\lambda - \gamma}}{\lambda - \gamma}T_A$, where T_A and T_B are the strictly upper triangular parts of A and B .

Now as the eigenvalues of A and hence \hat{A} are not collinear, we can always find another $\omega \in \sigma(\hat{A})$ such that $\frac{\overline{\lambda - \omega}}{\lambda - \omega} \neq \frac{\overline{\lambda - \gamma}}{\lambda - \gamma}$. Then the above equation is possible only if $T_A = T_B = 0$. In this case, A and B are both diagonal and hence normal. Then (2.3) implies that $\hat{A} = \bar{B}$. \square

From Lemma 2.5, we have the following consequence for diagonalizable matrices.

Corollary 2.6. *Suppose A and B satisfy (2.2) and A is diagonalizable. Then there are $\mu, \nu \in \mathbb{C}$ with $|\mu| = 1$ such that*

- (a) $B = \mu A + \nu I_n$, or
- (b) A is normal with non-collinear eigenvalues and $B = \mu A^* + \nu I_n$.

Proof. Suppose A is diagonalizable. Then $A = SDS^{-1}$ for some invertible S and diagonal D . By Lemma 2.4, $B = S(\mu D + \nu I_n)S^{-1}$ or $B = S(\mu \bar{D} + \nu I_n)S^{-1}$. If A has only one eigenvalue, then A is a scalar matrix and so is B . Then the result follows. Suppose A has at least two eigenvalues. As A and B commute, the result now follows by Lemma 2.5. \square

Lemma 2.7. *For any two matrices A and B , if*

$$(2.4) \quad f([A, X]) = f([B, X]) \quad \text{for all } X \in M_n,$$

then there are $\mu, \nu \in \mathbb{C}$ with $|\mu| = 1$ such that either

- (a) $B = \mu A + \nu I_n$, or
- (b) A is normal with non-collinear eigenvalues and $B = \mu A^* + \nu I_n$.

Proof. Suppose A and B satisfy (2.4). Then, putting $X = B$ in (2.4), it follows from (P2) that A and B commute. If A has at least two eigenvalues, then the result follows from Lemma 2.5.

Suppose A has only one eigenvalue, say λ . Then by Lemma 2.4, B has one eigenvalue only, say γ . Write $A = SJS^{-1} + \lambda I_n$, where S is invertible and $J = J_{n_1} \oplus \cdots \oplus J_{n_s}$ is the Jordan form of A with $n_1 \geq \cdots \geq n_s$. Now as A and B satisfy (2.4), A and B have the same set of commuting matrices. Then $B = Sp(J)S^{-1} + \gamma I_n$ for some polynomial p of degree at most $m = n_1 - 1$ with $p(0) = 0$.

By a similar argument as in Lemma 2.5, we can show that

$$\|(B - \gamma I_n)z\| = \|(A - \lambda I_n)z\| \quad \text{for all } z \in \mathbb{C}^n.$$

Then there is a unitary matrix W such that

$$Sp(J)S^{-1} = (B - \gamma I_n) = W(A - \lambda I_n) = WSJS^{-1}.$$

Write $S = UT$ for unitary U and upper triangular T , $V = U^*WU$ and $p(x) = \sum_{i=1}^m c_i x^i$. Then we have

$$(2.5) \quad Tp(J)T^{-1} = VTJT^{-1}.$$

Notice that both $Tp(J)T^{-1}$ and TJT^{-1} are strictly upper triangular. Furthermore, the first $n_1 - 1$ entries in the super-diagonal of $Tp(J)T^{-1}$ are c_1 times the corresponding $n_1 - 1$ super-diagonal entries of TJT^{-1} .

As V is unitary, we must have $|c_1| = 1$ and $V = c_1 I_{n_1-1} \oplus V_1$ for some unitary $V_1 \in M_{n-n_1+1}$. Now comparing the first $n_1 \times n_1$ principal submatrices in (2.5), we have

$$T_1 p(J_{n_1}) T_1^{-1} = (c_1 I_{n_1-1} \oplus [v_{n_1, n_1}]) T_1 J_{n_1} T_1^{-1} = c_1 T_1 J_{n_1} T_1^{-1},$$

where T_1 is the $n_1 \times n_1$ principal submatrix of T . Therefore, $T_1 (\sum_{i=2}^m c_i J_{n_1}^i) T_1^{-1} = 0$ and so $\sum_{i=2}^m c_i J_{n_1}^i = 0$. Hence, $c_2 = \dots = c_m = 0$. Then $p(x) = c_1 x$ and so $B = c_1 A + (\gamma - c_1 \lambda) I_n$. \square

We are now ready to present the following.

Proof of Theorem 2.1.

First we assume that ϕ is bijective. Suppose ϕ is a bijective map satisfying

$$f([A, B]) = f([\phi(A), \phi(B)]) \quad \text{for all } A, B \in M_n.$$

Because $f(X) = f(0)$ if and only if $X = 0$ by (P2), we see that $[A, B] = 0$ if and only if $[\phi(A), \phi(B)] = 0$. We can apply Theorem 2.2 and conclude that ϕ has the form (2.1) with $A^\dagger = A$ or A^t . In particular, for any rank one matrix $R \in M_n$, there are $\mu_R, \nu_R \in \mathbb{C}$ such that

$$\phi(R) = S(\mu_R R_\tau^\dagger + \nu_R I_n) S^{-1}.$$

Suppose $\mu_R = |\mu_R| e^{i\theta_R}$. By replacing $\phi(R)$ with $e^{-i\theta_R} (\phi(R) - \nu_R I_n)$, we may assume that $\mu_R > 0$ and $\nu_R = 0$.

Here we consider only the case when $A^\dagger = A$. The case when $A^\dagger = A^t$ is similar. Fixed an orthonormal basis $\{x_1, \dots, x_n\}$ and define $X_{ij} = x_i x_j^*$. Take $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{C}^n$ and let $A = \sum_{j=1}^n \alpha_j X_{j1}$. For $k = 2, \dots, n$,

$$(2.6) \quad f(\mu_A \mu_{X_{kk}} \tau(\alpha_k) S(X_{k1})_\tau S^{-1}) = f([\phi(A), \phi(X_{kk})]) = f([A, X_{kk}]) = f(\alpha_k X_{k1}).$$

In particular, if $Z = \mu_A \mu_{X_{22}} S(X_{21})_\tau S^{-1}$, then

$$f(\tau(\alpha) Z) = f(\alpha X_{21}) \quad \text{for all } \alpha \in \mathbb{C}.$$

Suppose τ is neither the identity map $\lambda \rightarrow \lambda$ nor the conjugate map $\lambda \rightarrow \bar{\lambda}$. By [10, Theorem 1], the set $\tau([0, 1])$ is an unbounded subset of \mathbb{C} . Thus, there exists $\alpha \in [0, 1]$ such that $|\tau(\alpha)| > |\tau(2)|$. But then by (P1) and (P3), we have

$$f(2X_{21}) = f(\tau(2)Z) = f(|\tau(2)|Z) < f(|\tau(\alpha)|Z) = f(\tau(\alpha)Z) = f(\alpha X_{21}) < f(2X_{21}),$$

which is a contradiction. Thus, τ is either the identity map or the conjugate map.

Furthermore, as $f([X_{32}, X_{22}]) = f(X_{32}) = f([X_{32}, X_{33}])$,

$$\begin{aligned} & f(\mu_{X_{32}}\mu_{X_{22}}S(X_{32})_{\tau}S^{-1}) = f([\phi(X_{32}), \phi(X_{22})]) \\ & = f([\phi(X_{32}), \phi(X_{33})]) = f(\mu_{X_{32}}\mu_{X_{33}}S(X_{32})_{\tau}S^{-1}). \end{aligned}$$

Thus, $\mu_{X_{22}} = \mu_{X_{33}}$ by (P3). By (2.6) and the fact that $f(\xi X_{21}) = f(\xi X_{31})$ for all $\xi \in \mathbb{C}$, we have

$$f(S(X_{21})_{\tau}S^{-1}) = f(S(X_{31})_{\tau}S^{-1}).$$

We now claim that S is a multiple of some unitary matrix. If not, then there is a pair of orthonormal vectors y_2, y_3 such that $\|Sy_2\| \neq \|Sy_3\|$. Extend y_2, y_3 to an orthonormal basis $\{y_1, y_2, y_3, \dots, y_n\}$ and let $x_j = (y_j)_{\tau^{-1}}$. Then $\{x_1, \dots, x_n\}$ also forms an orthonormal basis. By the above study, we have

$$f(\|Sy_2\|\|y_1^*S^{-1}\|E_{12}) = f(S(X_{21})_{\tau}S^{-1}) = f(S(X_{31})_{\tau}S^{-1}) = f(\|Sy_3\|\|y_1^*S^{-1}\|E_{12}),$$

which contradicts that $\|Sy_2\| \neq \|Sy_3\|$. Thus, S is a multiple of some unitary matrix. By absorbing the constant term, we may assume that S is unitary. Now for any rank one matrices R and S ,

$$f([R, S]) = f([\phi(R), \phi(S)]) = f(\mu_R\mu_S[R_{\tau}, S_{\tau}]).$$

By (P1), $f([R, S]) = f([R_{\tau}, S_{\tau}])$ whenever $[R, S]$ is a rank one nilpotent matrix, and hence $\mu_R\mu_S = 1$ in this case.

Now for any rank one matrix A , we can always find two other rank one matrices B and C such that $[A, B]$, $[A, C]$ and $[B, C]$ are all rank one nilpotents. Then we must have $\mu_A\mu_B = \mu_A\mu_C = \mu_B\mu_C = 1$. As all μ_A, μ_B, μ_C are positive real numbers, the equality is possible only when $\mu_A = \mu_B = \mu_C = 1$. Then we have $\phi(A) = SA_{\tau}S^{-1} = SA_{\tau}S^*$ for all rank one A .

By replacing ϕ with the map $A \mapsto S^*\phi(A)S$, we may assume that $\phi(X) = X^+$ for all rank one matrices X , where $X^+ = X, \bar{X}, X^t$ or X^* . Then

$$f([A, B]) = f([\phi(A), \phi(B)]) = f([A^+, B^+]) = f([A, B]^+)$$

for all rank one $A, B \in M_n$. Notice that the set

$$\{X : X = [A, B] \text{ for some rank one } A \text{ and } B\}$$

contains the set of trace zero non-nilpotent matrices with rank at most two and so is dense in the set of trace zero matrices with rank at most two. Thus, by continuity of f we see that

$$f(X) = f(X^+) \quad \text{for all trace zero matrices } X \text{ with rank at most two.}$$

Now define $\Phi : M_n \rightarrow M_n$ by $A \mapsto \phi(A)^+$. Then $\Phi(X) = X$ for all rank one matrices X . For any $A \in M_n$ and rank one matrix $X \in M_n$, as $[A, X]$ is a trace zero matrix with rank at most two,

$$f([A, X]) = f([\phi(A), \phi(X)]) = f([\phi(A), X^+]) = f([\phi(A)^+, X]) = f([\Phi(A), X]).$$

Thus, $f([A, X]) = f([\Phi(A), X])$ for all rank one X . Then Corollary 2.6 implies that $\Phi(A) = \mu_A A + \nu_A I_n$ or $\Phi(A) = \mu_A A^* + \nu_A I_n$ for all diagonalizable matrices A and the latter case happens only when A is normal with non-collinear eigenvalues.

After absorbing the constants μ_A and ν_A , we may assume that $\Phi(X) = X$ for all non-normal diagonalizable matrices X . Then

$$f([A, B]) = f([\phi(A), \phi(B)]) = f([\Phi(A), \Phi(B)]^+) = f([A, B]^+)$$

for all non-normal diagonalizable matrices A and B . Since the set of all non-normal diagonalizable matrices is dense in M_n , we see that $f([A, B]) = f([A, B]^+)$ for all $A, B \in M_n$. Then for any $A \in M_n$,

$$f([A, X]) = f([\phi(A), \phi(X)]) = f([\Phi(A), \Phi(X)]^+) = f([\Phi(A), X])$$

for all non-normal diagonalizable matrices X , and so $f([A, X]) = f([\Phi(A), X])$ for all $X \in M_n$ by the continuity of f . Now the result follows by Lemma 2.7.

Finally, we show that one only needs the surjective assumption on ϕ . For any $A, B \in M_n$, we say $A \sim B$ if

$$f([A, X]) = f([B, X]) \quad \text{for all } X \in M_n.$$

Clearly, \sim is an equivalence relation and for each $A \in M_n$, denote by $S_A = \{B : B \sim A\}$ the equivalence class of A . By Lemma 2.7, either

- (I) S_A is the set of matrices of the form $\mu A + \nu I$ for some $\mu, \nu \in \mathbb{C}$ with $|\mu| = 1$, or
- (II) A is normal and $A \sim A^*$, S_A is the set of matrices of the form $\mu A + \nu I$ or $\mu A^* + \nu I$ for some $\mu, \nu \in \mathbb{C}$ with $|\mu| = 1$.

Pick a representative for each equivalence class and write \mathcal{A} for the set of these representatives. Since ϕ is surjective, S_A and $\phi^{-1}(S_A)$ have the same cardinality c for every $A \in \mathcal{A}$. Thus there exists a map $\psi : M_n \rightarrow M_n$ which maps $\phi^{-1}(S_A)$ bijectively onto S_A for each $A \in \mathcal{A}$. Clearly ψ is bijective and $\psi(A) \sim \phi(A)$ for all $A \in M_n$. Then, for any $A, B \in M_n$,

$$f([A, B]) = f([\phi(A), \phi(B)]) = f([\psi(A), \phi(B)]) = f([\psi(A), \psi(B)]).$$

That is, ψ is bijective map satisfying (2.2). By the proof of Theorem 2.1 with bijective ϕ in the previous paragraphs, ψ has the desired form and hence so does ϕ , as $\psi(A) \sim \phi(A)$ implies $\phi(A) = \mu\psi(A) + \nu I$ or $\phi(A) = \mu\psi(A)^* + \nu I$ when $\psi(A)^*$ is normal and $\psi(A)^* \sim \psi(A)$. \square

Remark Using the argument in the last part of the proof on the replacement of the bijective assumption by the surjective assumption on ϕ , one may further weaken the surjective assumption on ϕ by any one of the following (weaker) assumptions on the following modified map $\tilde{\phi}$ defined by

$$\tilde{\phi}(X) = \phi(X) - \text{tr}(\phi(X))I/n$$

on the set M_n^0 of trace zero matrices in M_n .

- (a) The map $\tilde{\phi} : M_n^0 \rightarrow M_n^0$ surjective.
- (b) For any $A \in M_n^0$ the range of $\tilde{\phi}$ contains a matrix of the form $e^{it}A$ for some $t \in [0, 2\pi)$.

3. PSEUDO SPECTRUM AND PSEUDO SPECTRAL RADIUS

In this section, we use Theorem 2.1 to study maps preserving the pseudo spectral radius (see the definitions below) of the Lie product of matrices. Then we further deduce the result for maps preserving the pseudo spectrum. As one shall see, with considerable effort, one will be able to get more specific structure of the preserving maps.

For $\varepsilon > 0$, define the ε -pseudospectrum $\sigma_\varepsilon(A)$ of $A \in M_n$ as

$$\sigma_\varepsilon(A) = \{z \in \sigma(A + E) : E \in M_n, \|E\| < \varepsilon\} = \{z \in \mathbb{C} : s_n(A - zI_n) < \varepsilon\},$$

where $s_1(X) \geq \dots \geq s_n(X)$ denote the singular values of $X \in M_n$, and the ε -pseudospectral radius $r_\varepsilon(A)$ of $A \in M_n$ as

$$r_\varepsilon(A) = \sup\{|\mu| : \mu \in \sigma_\varepsilon(A)\}.$$

Note that the pseudo spectral radius is useful in studying the stability of matrices under perturbations, and there are efficient algorithm for its computation; see for example, [6] and its references. Preservers of pseudo spectrum has been considered for several types of products in [4] (see also [5]). Here we characterize the preservers of pseudo spectral radius and pseudo spectrum for Lie products. We first prove the following.

Theorem 3.1. *Suppose $n \geq 3$ and $\varepsilon > 0$. Then a surjective map $\phi : M_n \rightarrow M_n$ satisfies*

$$r_\varepsilon([A, B]) = r_\varepsilon([\phi(A), \phi(B)]) \quad \text{for all } A, B \in M_n$$

if and only if there is a unitary $U \in M_n$ such that

$$\phi(A) = \mu_A U \psi(A) U^* + \nu_A I_n \quad \text{for all } A \in M_n,$$

where $\mu_A, \nu_A \in \mathbb{C}$ with $|\mu_A| = 1$, depending on A , and ψ is one of the following maps: $A \mapsto A$, $A \mapsto \bar{A}$, $A \mapsto A^t$ or $A \mapsto A^$.*

Proof. The sufficiency can be readily checked. To prove the necessity, let $f(A) = r_\varepsilon(A)$ for $A \in M_n$. It is clear that f is a continuous map satisfying (P1) and (P2). Suppose X is a rank one nilpotent matrix. It follows from Proposition 2.4 in [5] that $r_\varepsilon(X) = \sqrt{\varepsilon^2 + \|X\|\varepsilon}$. Hence, (P3) is also satisfied. So, we can apply Theorem 2.1 and conclude that ϕ has the form in Theorem 2.1. To get the desired conclusion, we need to show that the set \mathcal{N} is empty. Assume not, and there is $A \in \mathcal{N}$. Since A is normal with non-collinear eigenvalues, there is a unitary V and $\gamma, \xi \in \mathbb{C}$ such that

$$V(\psi(A) - \xi I)V^* = \gamma \text{diag}(1, \mu, 0, \mu_4, \dots, \mu_n),$$

where $\mu \notin \mathbb{R}$. Let $B \in M_n$ be such that

$$\tilde{B} = V\psi(B)V^* = \begin{bmatrix} 0 & 1 & 0 \\ a & 0 & b \\ 0 & c & 0 \end{bmatrix} \oplus O_{n-3},$$

where $a = (1 - \bar{\mu})/(1 - \mu)$, $b > 0$ and $c = b\bar{\mu}/\mu$. Then

$$\tilde{B}\tilde{B}^* = \begin{bmatrix} 1 & 0 & \bar{c} \\ 0 & |a|^2 + |b|^2 & 0 \\ c & 0 & |c|^2 \end{bmatrix} \quad \text{and} \quad \tilde{B}^*\tilde{B} = \begin{bmatrix} |a|^2 & 0 & \bar{a}b \\ 0 & 1 + |c|^2 & 0 \\ \bar{b}a & 0 & |b|^2 \end{bmatrix}$$

and we can choose $b > 0$ so that \tilde{B} is not normal, and neither is B . As a result, $\phi(B) = \mu_B U \psi(B) U^* + \nu_B I$.

Now,

$$C_1 = V[\psi(A), \psi(B)]V^* = \gamma \begin{bmatrix} 0 & 1 - \mu & 0 \\ \bar{\mu} - 1 & 0 & b\mu \\ 0 & -b\bar{\mu} & 0 \end{bmatrix} \oplus O_{n-3}$$

is normal with eigenvalues $s_\pm = \pm\gamma\sqrt{|1 - \mu|^2 + b^2|\mu|^2}$ so that

$$r_\varepsilon([A, B]) = r_\varepsilon([\psi(A), \psi(B)]) = |\gamma|\sqrt{|1 - \mu|^2 + b^2|\mu|^2} + \varepsilon.$$

However, $[\phi(A), \phi(B)]$ is unitarily similar to

$$C_2 = \mu_A \mu_B \bar{\gamma} \begin{bmatrix} 0 & 1 - \mu & 0 \\ (1 - \bar{\mu})^2 / (\mu - 1) & 0 & b\bar{\mu} \\ 0 & -b\bar{\mu}^2 / \mu & 0 \end{bmatrix} \oplus O_{n-3}.$$

One readily checks that the matrix C_2 is normal if and only if μ is pure imaginary. In all other cases, there is a unitary $R \in M_n$ obtained from I_n by changing the $(1, 1), (1, 3), (3, 1), (3, 3)$ entries so that

$$RC_2R^* = \bar{\gamma} \begin{bmatrix} 0 & c_1 & 0 \\ c_2 & 0 & c_3 \\ 0 & 0 & 0 \end{bmatrix} \oplus O_{n-3}.$$

If C_2 has singular values $s_1 \geq s_2$, then

$$|\gamma|^2(|c_1|^2 + |c_2|^2 + |c_3|^2) = \text{tr}(C_2C_2^*) = \text{tr}(C_1C_1^*) = |\gamma|^2(s_+^2 + s_-^2).$$

Because C_2 is not normal, $s_1 < s_+$, we see that $s_2 > s_-$. Then for any $z \in \mathbb{C}$, if $\tilde{C} - zI$ has singular values $s_1(z) \geq s_2(z)$, then

$$s_1(z)^2 + s_2(z)^2 = 2|z|^2 + |c_1|^2 + |c_2|^2 + |c_3|^2 2|z|^2 + s_+^2 + s_-^2 = s_+(z)^2 + s_-(z)^2,$$

where $s_+(z) \geq s_-(z)$ are the singular values of $C_1 - zI$. Again, because $C_2 - zI$ is not normal, we see that $s_+(z) > s_1(z) \geq s_2(z) > s_-(z)$. It follows that $s_+(z) > s_-(z)$ for any $z \in \mathbb{C}$ with $|z| \leq |\gamma| \sqrt{|1 + \mu|^2 + b^2|\mu|^2} + \varepsilon$. Thus,

$$\max\{z \in \mathbb{C} : s_2(C_2 - zI) \leq \varepsilon\} < \max\{z \in \mathbb{C} : s_2(C_1 - zI) \leq \varepsilon\}.$$

So, if a normal matrix A with three collinear eigenvalues $\gamma + \nu, \gamma\mu + \nu, \nu$, where μ is not real and $\mu \neq \pm i$, then $A \notin \mathcal{N}$. Clearly, if $A \in \mathcal{N}$ has eigenvalues of the form $\gamma + \nu, \gamma + i\nu, \gamma$, then $\psi(A)^*$ can be viewed as a multiple of $\psi(A)$. Thus, we may assume that $A \notin \mathcal{N}$ by adjusting μ_A and ν_A . The result follows. \square

We will use the above theorem to determine the structure of preservers of the pseudo spectrum of Lie product of matrices. To achieve this, we need a characterization of normal matrices A with two distinct eigenvalues, i.e., $A - bI$ is a nonzero multiple of a rank k orthogonal projection P with $1 \leq k < n$; see Proposition 3.3 below. The proof depends on the following lemma.

Lemma 3.2. *Suppose $C = C_1 \oplus O_{n-3}$, where $C_1 \in M_3$ has rank ≤ 2 and $\text{tr } C_1 = 0$. Then for every $\varepsilon > 0$, $\sigma_\varepsilon(C) = \sigma_\varepsilon(C_1)$. Furthermore, suppose for $t \in \mathbb{R}$,*

$$f(\lambda, t) = \det(\lambda I_3 - (C_1 - tI_3)^*(C_1 - tI_3)) = \lambda^3 + p_2(t)\lambda^2 + p_1(t)\lambda + p_0(t)$$

where $p_1(t) = q_1(t) + at$ with $a \neq 0$ and $p_0(t), q_1(t), p_2(t)$ contains only even powers of t . Then $\sigma_\varepsilon(C) \neq -\sigma_\varepsilon(C)$.

Proof. Since $\text{rank } C_1 \leq 2, 0 \in \sigma(C_1)$. Therefore, $\sigma_\varepsilon(C) = \sigma_\varepsilon(C_1) \cup \sigma_\varepsilon(0_{n-k}) = \sigma_\varepsilon(C_1)$.

Note that for each $t \in \mathbb{R}$, $f(\lambda, t)$ is a cubic polynomial in λ with three non-negative real roots $\lambda_1(t) \geq \lambda_2(t) \geq \lambda_3(t) \geq 0$ and $s_{\min}(C_1 - tI_3) = \sqrt{\lambda_3(t)}$.

Without loss of generality, we may assume that $a < 0$. Given $\varepsilon > 0$, $t \in \sigma_\varepsilon(C_1) \cap \mathbb{R}$ if and only if $\lambda_3(t) < \varepsilon^2$. Since $\lambda_3(0) = 0$ and $\lim_{t \rightarrow \infty} \lambda_3(t) = \infty$, there exists $t_0 > 0$ such that $\lambda_3(t_0) = \varepsilon^2$. We have $t_0 \notin \sigma_\varepsilon(C)$ and $f(\varepsilon^2, t_0) = 0$. But then

$$f(\varepsilon^2, -t_0) = f(\varepsilon^2, t_0) - 2at_0\varepsilon^2 > 0$$

Thus, $\lambda_3(-t_0) < \varepsilon^2$ implying that $-t_0 \in \sigma_\varepsilon(C)$. So, $t_0 \in -\sigma_\varepsilon(C)$, and thus $\sigma_\varepsilon(C) \neq -\sigma_\varepsilon(C)$. \square

Proposition 3.3. *Let $n \geq 3$ and $A \in M_n$. The following conditions are equivalent.*

- (a) *A is a normal matrix with at most two distinct eigenvalues.*
- (b) *$\sigma_\varepsilon([A, B]) = -\sigma_\varepsilon([A, B])$ for all $B \in M_n$.*
- (c) *$\sigma_\varepsilon([A, B]) = -\sigma_\varepsilon([A, B])$ for all rank one nilpotent $B \in M_n$.*

Proof. Suppose (a) holds. Then there is a unitary V and $\nu \in \mathbb{C}$ such that $VAV^* - \nu I = \lambda J$ with $J = I_k \oplus -I_{n-k}$. Then for any $B \in M_n$ such that $VBV^* = (B_{ij})_{1 \leq i, j \leq 2}$ with $B_{11} \in M_k, B_{22} \in B_{22}$, we have

$$C = V[A, B]V^* = 2\lambda \begin{bmatrix} O_k & B_{12} \\ -B_{21} & O_{n-k} \end{bmatrix}$$

satisfies $-C = JCJ^*$. Thus,

$$\sigma_\varepsilon([A, B]) = \sigma_\varepsilon([VAV^*, VBV^*]) = \sigma_\varepsilon(-J[A, B]J^*) = \sigma_\varepsilon(-[A, B]).$$

So, condition (b) holds.

The implication (b) \Rightarrow (c) is clear. To prove (c) \Rightarrow (a), we consider the contra-positive. Assume (a) is not true. We consider 2 cases.

Case 1. Suppose A is normal with more than two distinct eigenvalues. We may assume that $A = \text{diag}(a, b, c) \oplus A_2$ such that a, b and c are distinct. If $\text{Re}((b-a)\overline{(c-a)}) \leq 0$, then we have $\text{Re}((b-c)\overline{a-c}) = \text{Re}((b-a+a-c)\overline{a-c}) = |a-c|^2 - \text{Re}((b-a)\overline{(c-a)}) > 0$. Thus, we may assume that $\text{Re}((b-a)\overline{(c-a)}) > 0$ which implies that

$$|2a - (b+c)|^2 = |(b-a) + (c-a)|^2 > |b-a|^2 + |c-a|^2 > |b-c|^2 \Rightarrow \left| a - \frac{b+c}{2} \right| > \frac{|b-c|}{2}.$$

Thus, by replacing A with $\frac{2}{(b-c)} \left(A - \frac{(b+c)}{2} I \right)$, we may assume that $A = \text{diag}(a, 1, -1) \oplus A_2$

such that $|a| > 1$. Consider the rank one nilpotent $X = \begin{bmatrix} 0 & -\sqrt{2} & \sqrt{2} \\ 0 & -1 & 1 \\ 0 & -1 & 1 \end{bmatrix} \oplus 0_{n-3}$. We have $[A, X] =$

$C \oplus 0_{n-3}$, where $C = \begin{bmatrix} 0 & \sqrt{2}(1-a) & \sqrt{2}(1+a) \\ 0 & 0 & 2 \\ 0 & 2 & 0 \end{bmatrix}$. Then

$$\det(\lambda I_3 - (C - tI_3)^*(C - tI_3)) = \lambda^3 + p_2(t)\lambda^2 + p_1(t)\lambda + p_0(t),$$

where

$$p_2(t) = -3t^2 - 4|a|^2 - 12,$$

$$p_1(t) = 3t^4 + 4(1 + |a|^2)t^2 + 16(1 - |a|^2)t + 16(2 + |a|^2),$$

$$p_0(t) = -t^6 + 8t^4 - 16t^2.$$

Since $|a| > 1$, the condition in Lemma 3.2 is satisfied. Therefore, $\sigma_\varepsilon(C) \neq -\sigma_\varepsilon(C)$.

Case 2. Assume that A is not normal. We may assume that $A = (a_{ij})$ is in upper triangular form such that the $(1, 2)$ entry is nonzero; see [13, Lemma 1]. We may replace A by $A - a_{33}I$ and assume that $A = (A_{ij})$ with $A_{22} \in M_{n-3}$, $A_{21} = O$, and

$$A_{11} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22} & a_{23} \\ 0 & 0 & 0 \end{bmatrix}.$$

Subcase (2.a) Suppose not both $[a_{13}, \dots, a_{1n}]$ and $[a_{23}, \dots, a_{2n}]$ are zero. Then there is a unitary $U = U_1 \oplus U_2$ with $U_1 \in M_2$ such that $UAU^* = \tilde{A} = (\tilde{a}_{ij})$, where the second row of \tilde{A} equals $[\tilde{a}_{21}, \tilde{a}_{22}, \tilde{a}_{23}, 0, \dots, 0]$ with $\tilde{a}_{21} \in \mathbb{R}$ and $\tilde{a}_{21} \neq 0$ and $\tilde{a}_{23} \neq 0$. Let $B = E_{12}$. Then

$$C = [\tilde{A}, B] = \begin{bmatrix} -\tilde{a}_{21} & \tilde{a}_{11} - \tilde{a}_{22} & -\tilde{a}_{23} \\ 0 & \tilde{a}_{21} & 0 \\ 0 & 0 & 0 \end{bmatrix} \oplus O_{n-3}.$$

Then

$$\det(\lambda I_3 - (C - tI_3)^*(C - tI_3)) = \lambda^3 + p_2(t)\lambda^2 + p_1(t)\lambda + p_0(t),$$

where

$$p_2(t) = -3t^2 - |\tilde{a}_{22} - \tilde{a}_{11}|^2 - |\tilde{a}_{23}|^2 - 2\tilde{a}_{21}^2,$$

$$p_1(t) = 3t^4 + (|\tilde{a}_{22} - \tilde{a}_{11}|^2 + |\tilde{a}_{23}|^2)t^2 - 2\tilde{a}_{21}|\tilde{a}_{23}|^2t + \tilde{a}_{21}^2(\tilde{a}_{21}^2 + |\tilde{a}_{23}|^2),$$

$$p_0(t) = -t^6 + 2\tilde{a}_{21}^2t^4 - \tilde{a}_{21}^4t^2.$$

Since \tilde{a}_{21} and $\tilde{a}_{23} \neq 0$, the condition in Lemma 3.2 is satisfied. Therefore, $\sigma_\varepsilon(C) \neq -\sigma_\varepsilon(C)$.

Subcase (2.b) Suppose both $[a_{13}, \dots, a_{1n}]$ and $[a_{23}, \dots, a_{2n}]$ are zero.

i) If $a_{11} = a_{22} = 0$, then we may assume that $a_{12} = 1$. Let

$$B = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 0 & 1 \\ -1 & 0 & -1 \end{bmatrix} \oplus O_{n-3} \text{ so that } C = [A, B] = \begin{bmatrix} 1 & -1 & 1 \\ 0 & -1 & 0 \\ 0 & 1 & 0 \end{bmatrix} \oplus O_{n-3}.$$

Then

$$\det(\lambda I_3 - (C - tI_3)^*(C - tI_3)) = \lambda^3 + p_2(t)\lambda^2 + p_1(t)\lambda + p_0(t),$$

where

$$p_2(t) = -3t^2 - 5,$$

$$p_1(t) = 3t^4 + 3t^2 - 2t + 4,$$

$$p_0(t) = -t^6 + 2t^4 - t^2.$$

Therefore, the condition in Lemma 3.2 is satisfied and $\sigma_\varepsilon(C) \neq -\sigma_\varepsilon(C)$.

ii) If either a_{11} or $a_{22} \neq 0$, then, applying a unitary similarity, we may assume that $a_{11} \neq 0$. Replacing A by $e^{i\theta}A$, we may assume that $a_{11} \in \mathbb{R}$. Then we may further assume that $a_{12} = 1$. Let

$B = \begin{bmatrix} 1 & 0 & 1 \\ 0 & 0 & 0 \\ -1 & 0 & -1 \end{bmatrix} \oplus O_{n-3}$ and $C = [A, B]$. Then $C = C_1 \oplus 0_{n-3}$, where $C_1 = \begin{bmatrix} 0 & -1 & a_{11} \\ 0 & 0 & 0 \\ a_{11} & 1 & 0 \end{bmatrix}$. Then

$$\det(\lambda I_3 - (C_1 - tI_3)^*(C_1 - tI_3)) = \lambda^3 + p_2(t)\lambda^2 + p_1(t)\lambda + p_0(t),$$

where

$$p_2(t) = -3t^2 - 2 - 2a_{11}^2,$$

$$p_1(t) = 3t^4 + 2t^2 - 4a_{11}t + 2a_{11}^2 + a_{11}^4,$$

$$p_0(t) = -t^6 + 2a_{11}^2t^4 - a_{11}^4t^2.$$

Therefore, the condition in Lemma 3.2 is satisfied and $\sigma_\varepsilon(C) \neq -\sigma_\varepsilon(C)$.

The proof is complete. \square

Theorem 3.4. *Suppose $n \geq 3$ and $\varepsilon > 0$. Then a surjective map $\phi : M_n \rightarrow M_n$ satisfies*

$$\sigma_\varepsilon([A, B]) = \sigma_\varepsilon([\phi(A), \phi(B)]) \quad \text{for all } A, B \in M_n$$

if and only if there exist $\mu \in \{1, -1\}$, a unitary matrix $U \in M_n$, and a set \mathcal{T} of normal matrices with at most two distinct eigenvalues such that

$$(3.1) \quad \phi(A) = \begin{cases} \mu U \psi(A) U^* + \nu_A I & \text{if } A \in M_n \setminus \mathcal{T}, \\ -\mu U \psi(A) U^* + \nu_A I & \text{if } A \in \mathcal{T}, \end{cases}$$

where $\nu_A \in \mathbb{C}$ depends on A , and ψ is one of the maps: $A \mapsto A, A \mapsto iA^t$.

Proof. To prove the sufficiency, if ψ has the first form, then $\sigma_\varepsilon([A, B]) = \sigma_\varepsilon([\phi(A), \phi(B)]) = \mu_A \mu_B \sigma_\varepsilon([A, B])$ if none, one, or both of $A, B \in \mathcal{T}$ by Proposition 3.3. If ψ has the second form, then $\sigma_\varepsilon([A, B]) = \sigma_\varepsilon([\phi(A), \phi(B)]) = -\mu_A \mu_B \sigma_\varepsilon([A^t, B^t]) = \mu_A \mu_B \sigma_\varepsilon([A, B])$ if none, one, or both of $A, B \in \mathcal{T}$ by Proposition 3.3.

To prove the necessity, we may compose ϕ by a map of the form $X \mapsto VXV^*$ and adjust ν_X if necessary so that ϕ has the form $A \mapsto \mu_A \psi(A)$, where ψ is one of the maps $A \mapsto A, A \mapsto A^t, A \mapsto \bar{A}, A \mapsto A^*$. Focusing on rank one Hermitian matrices, we see that one of the following happens.

- (1) For any rank one $A = xx^*$, $\phi(A) = \mu_A A$. (2) For any rank one $A = xx^*$, $\phi(A) = \mu_A A^t$.

Suppose (2) holds. We may replace ϕ by the map $X \mapsto i\phi(X)^t$. Then the modified map will satisfy condition (1). Thus, we can focus on the case when (1) holds, and prove that ϕ has the asserted form with $\psi(X) = X$ for all $X \in M_n$.

In the rest of the proof, we assume that (1) holds. Then we have either

- i) $\phi(A) = \mu_A A$ for all $A \in M_n$, or ii) $\phi(A) = \mu_A A^*$ for all $A \in M_n$.

We will show that for some μ , we have $\mu_A = \mu$ for all $A \in M_n \setminus \mathcal{T}$ and $\mu_A = -\mu$ for all $A \in \mathcal{T}$.

satisfying (3.1). Clearly, we need only consider non-scalar matrices.

Assertion 1 For every non-scalar matrix $A \in M_n$, $\mu_A \in \{-1, 1\}$.

To prove Assertion 1, let $A = xx^*$. If $B = yy^*$ such that $0 \neq [A, B]$, then $[A, B]$ is unitarily similar to $\text{diag}(ai, -ai) \oplus O_{n-2}$ with $a = \sqrt{-\text{tr}([A, B]^2)/2} > 0$ so that

$$\sigma_\varepsilon([A, B]) = D(-ai, \varepsilon) \cup D(0, \varepsilon) \cup D(ai, \varepsilon).$$

Because $\sigma_\varepsilon([\phi(A), \phi(B)]) = \mu_A \mu_B \sigma_\varepsilon([A, B])$, we see that $\mu_A \mu_B = \pm 1$.

Let $\mu = \mu_{E_{11}}$. Suppose $B = xx^*$ for a nonzero $x \in \mathbb{C}^n$. We can find $C = yy^*$ such that $[E_{11}, C] \neq 0$ and $[B, C] \neq 0$. Then $\mu \mu_C, \mu_B \mu_C \in \{1, -1\}$ so that $\mu \mu_C = \pm \mu_B \mu_C$. It follows that $\mu_B \in \{\mu, -\mu\}$.

Choose $B_j = x_j x_j^*$, $j = 1, 2$ so that $[E_{11}, B_1]$, $[E_{11}, B_2]$ and $[B_1, B_2] \neq 0$. Then

$$\mu \mu_{B_1}, \mu \mu_{B_2}, \mu_{B_2} \mu_{B_1} \in \{1, -1\}.$$

Hence, $\mu^2 \in \{-1, 1\}$. So we have either

$$(a) \mu^2 = -1 \Rightarrow \mu_B \in \{-i, i\} \text{ for all } B = xx^*, \quad \text{or} \quad (b) \mu^2 = 1 \Rightarrow \mu_B \in \{-1, 1\} \text{ for all } B = xx^*.$$

Next we will show that $\phi(A) = \mu_A A$ for all $A \in M_n$. Assume the contrary that $\phi(A) = \mu_A A^*$ for all $A \in M_n$. Let $B_1 = E_{11} + E_{13} + E_{31} + E_{33}$, $B_2 = E_{22} + E_{23} + E_{32} + E_{33}$ and $C = E_{11} + e^{i\pi/6} E_{22}$.

Then

$$\sigma_\varepsilon([B_1, C]) = D(-i, \varepsilon) \cup D(i, \varepsilon) \cup D(0, \varepsilon)$$

and

$$\sigma_\varepsilon([\phi(B_1), \phi(C)]) = \mu_{B_1} \mu_C D(-i, \varepsilon) \cup D(i, \varepsilon) \cup D(0, \varepsilon).$$

Hence, $\mu_{B_1} \mu_C \in \{-1, 1\}$. By a direct computation,

$$\sigma_\varepsilon([B_2, C]) = D(-e^{-2\pi i/3}, \varepsilon) \cup D(e^{-2\pi i/3}, \varepsilon) \cup D(0, \varepsilon)$$

and

$$\sigma_\varepsilon([\phi(B_2), \phi(C)]) = \mu_{B_2} \mu_C \left(D(-e^{-\pi i/3}, \varepsilon) \cup D(e^{-\pi i/3}, \varepsilon) \cup D(0, \varepsilon) \right).$$

Since $\mu_{B_1} = \pm \mu_{B_2}$ and $\mu_{B_1} \mu_C \in \{-1, 1\}$, we have $\mu_{B_2} \mu_C \in \{-1, 1\}$. Hence, $\sigma_\varepsilon([\phi(B_2), \phi(C)]) \neq \sigma_\varepsilon([B_2, C])$, a contradiction. Therefore, we have $\phi(A) = \mu_A A$ for all $A \in M_n$.

For any non-scalar normal matrix B with spectral decomposition $\sum_{j=1}^n b_j x_j x_j^*$ with $b_1 \neq b_2$, let $C = yy^*$ with $y = x_1 + x_2$. Then $[B, C]$ is unitarily similar to $\text{diag}(a, -a) \oplus O_{n-2}$. It follows that $\mu_B \mu_C \in \{1, -1\}$. Because $\mu_C \mu \in \{1, -1\}$, we see that $\mu_B \in \{\mu, -\mu\}$. Suppose B is non-normal. There is a unitary U such that $UBU^* = H + iG$, where $G = G^*$ is in diagonal form and $H = H^*$ has a nonzero $(1, 2)$ entry. Then for $C = UE_{11}U^*$, the matrix $[B, C]$ is unitarily similar to $\text{diag}(a, -a) \oplus O_{n-2}$. Again, we can conclude that $\mu_B = \pm \mu$. So, $\mu_B \in \{\mu, -\mu\}$ for every $B \in M_n$. Consequently, we have

$$(c) \mu_B \in \{-i, i\} \text{ for all } X \in M_n, \quad \text{or} \quad (d) \mu_B \in \{-1, 1\} \text{ for all } X \in M_n.$$

We claim that the condition (d) holds. To this end, let $D = \text{diag}(1, -1) \oplus O_{n-2}$ and $B = E_{12}/2 + E_{23} + E_{31}$. Then $[D, B] = E_{12} - E_{23} - E_{31}$ is a unitary matrix with eigenvalues $\lambda_1 = 1, \lambda_2 = e^{i2\pi/3}, \lambda_3 = e^{i4\pi/3}$. Thus,

$$\sigma_\varepsilon([D, B]) = D(\lambda_1, \varepsilon) \cup D(\lambda_2, \varepsilon) \cup D(\lambda_3, \varepsilon).$$

We see that $\mu_B \mu_D = 1$ for such a matrix B . Similarly, if $C = -E_{21}/2 + iE_{32} - iE_{13}$, then $\mu_C \mu_D = 1$. Thus, $\mu_B = \mu_C$. Now, $[B, C] = (1 + i/4)E_{11} + (1 - i/4)E_{22} - 2E_{33}$. Then $\mu_B \mu_C \sigma_\varepsilon([B, C]) = \sigma_\varepsilon([B, C])$ will imply that $\mu_B \mu_C = 1$. Because, $\mu_B = \mu_C$, we see that $\mu_B = \mu_C \in \{-1, 1\}$. Hence, the condition (d) holds.

Assertion 2 There is $\mu \in \{1, -1\}$ such that $\mu_A = \mu$ if A is not a normal matrix with at most two distinct eigenvalues.

Proof. First we show that for any nonzero vectors x, y, f such that 1) $y, f \in x^\perp$, 2) $\{y, f\}$ linearly independent and 3) $\operatorname{Re}(f^*y) \neq 0$, then the following holds.

$$(3.2) \quad \mu_{xf^*} = \mu_{yx^*}$$

Note that $C = [xf^*, yx^*] = (f^*y)xx^* - \|x\|^2yf^*$ which has a matrix representation of the form

$$C = \begin{bmatrix} \alpha & 0 & 0 \\ 0 & -\alpha & 0 \\ 0 & \beta & 0 \end{bmatrix} \oplus 0 = X \oplus 0$$

with $\alpha = f^*y\|x\|^2$, $\beta = \|x\|^2\sqrt{\|f\|^2\|y\|^2 - |f^*y|^2} \neq 0$. Then

$$\det(\lambda I_3 - (X - tI_3)^*(X - tI_3)) = \lambda^3 + p_2(t)\lambda^2 + p_1(t)\lambda + p_0(t),$$

where

$$p_2(t) = -3t^2 - (2|\alpha|^2 + |\beta|^2),$$

$$p_1(t) = 3t^4 + (4(\operatorname{Im}(\alpha))^2 + \beta^2)t^2 - 2\operatorname{Re}(\alpha)\beta^2t + |\alpha|^2(|\alpha|^2 + \beta^2),$$

$$p_0(t) = -t^6 + (\alpha^2 + \bar{\alpha}^2)t^4 - |\alpha|^4t^2.$$

Since $\operatorname{Re}(\alpha)$ and $\beta \neq 0$, the condition in Lemma 3.2 is satisfied. Therefore, $\sigma_\varepsilon(C) \neq -\sigma_\varepsilon(C)$. Since $\sigma_\varepsilon(C) = \mu_{xf^*}\mu_{yx^*}\sigma_\varepsilon(C)$, we have $\mu_{xf^*}\mu_{yx^*} = 1$, and thus $\mu_{xf^*} = \mu_{yx^*}$.

If xf^* and xu^* are rank-1 nilpotent and if $u \in f^\perp$, then (3.2) ensures that

$$\mu_{xf^*} = \mu_{(f+u)x^*} = \mu_{xu^*} = \mu_{u(x+f)^*} = \mu_{fu^*} = \mu_{(x+u)f^*} = \mu_{fx^*}.$$

So we have

$$(3.3) \quad \mu_{xf^*} = \mu_{xu^*} = \mu_{fx^*}$$

whenever the vectors x, f, u are pairwise orthogonal.

Next we show that

$$(3.4) \quad \mu_{xf^*} = \mu_{xu^*} \quad \text{for any nonzero vectors } f, u \in x^\perp.$$

Suppose f, u are nonzero vectors in x^\perp . If $u \in f^\perp$, the equality follows from (3.3). If $u = \lambda f$ for some nonzero scalar λ , taking $v \in \{x, f\}^\perp$ we have

$$\mu_{xf^*} = \mu_{xv^*} = \mu_{xu^*}.$$

If $u \notin f^\perp$ and the vectors u, f are linearly independent, then let $v = u - cf$, where $c = \frac{f^*u}{f^*f}$. Then

$v \in \{x, f\}^\perp$ and $u^*v = u^*u - \frac{|f^*u|^2}{f^*f} \neq 0$. By (3.2) and (3.3), we have

$$\mu_{xu^*} = \mu_{v^*} = \mu_{xf^*}.$$

Next, we show that $\mu_A = \mu_B$ for any rank one nilpotent matrices A, B . To this end, $A = xf^*$ and $B = yg^*$, taking unit vector $u \in \{x, y\}^\perp$ and using (3.4), we have

$$\mu_{xf^*} = \mu_{xu^*} = \mu_{yu^*} = \mu_{yg^*}.$$

By Proposition 3.3, if A is not a normal matrix with at most two distinct eigenvalues, then there is a rank one nilpotent B such that

$$-\sigma_\varepsilon([A, B]) \neq \sigma_\varepsilon([B, A]) = \mu_B \mu_A \sigma_\varepsilon([B, A]).$$

Thus, $\mu_A \mu_B = 1$, which implies $\mu_A = \mu_B$. The desired conclusion follows. \square

Acknowledgment

The research of the first author was partially supported by National Natural Science Foundation of China (No.11271217). The research of the second and third authors was partially supported by USA NSF and HK RCG. The second author is an honorary professor of the Shanghai University and an honorary professor of the University of Hong Kong.

REFERENCES

1. J.T. Chan, C.K. Li, and N.S. Sze, Isometries for unitarily invariant norms, *Linear Algebra Appl.* 399 (2005), 53-70.
2. J.T. Chan, C.K. Li, and N.S. Sze, Mappings on matrices: Invariance of functional values of matrix products, *J. of Aust. Math. Soc.*, 81 (2006), 165-184.
3. W.S. Cheung, S. Fallat and C.K. Li, Multiplicative preservers on semigroups of matrices, *Linear Algebra Appl.* 355 (2002), 173-186.
4. J. Cui, V. Forstall, C.K. Li and V. Yannello, Properties and Preservers of the Pseudospectrum, *Linear Algebra and its Applications* 436 (2012), 316-325.
5. J. Cui, C.K. Li and Y.T. Poon, Pseudospectra of special operators and Pseudospectrum preservers, *J. Math. Anal. Appl.* (2014), 1261-1273.
6. N. Guglielmi and M.L. Overton, Fast algorithms for the approximation of the pseudospectral abscissa and pseudospectral radius of a matrix, *SIAM J. Matrix Anal. appl.* 32 (2011), 1166-1192.
7. R. Guralnick and C.K. Li, Invertible preservers and algebraic groups III: preservers of unitary similarity (congruence) invariants and overgroups of some unitary subgroups, *Linear and Multilinear Algebra* 43 (1997), 257-282.
8. K. E. Gustafson and D. K. M. Rao, *Numerical range: The Field of values of linear operators and matrices*, Universitext, Springer-Verlag, New York (1997).
9. R. Horn and C. Johnson, *Matrix analysis*. Cambridge University Press, Cambridge, 1985.
10. H. Kestelman, Automorphisms of the field of complex numbers, *Proc. London Math. Soc.* 53 (1951), 1-12.
11. C.K. Li, E. Poon, and N.S. Sze, Preservers for norms of Lie product, *Operators and Matrices* 3 (2009), 187-203.
12. C.K. Li and N.K. Tsing, Linear operators preserving certain classes of functions on singular values of matrices, *Linear and Multilinear Algebra* 26 (1990), 133-143.
13. M. Marcus and M. Sandy, Conditions for the generalized numerical range to be real, *Linear Algebra Appl.* 71 (1985). 219-239.
14. P. Šemrl, Non-linear commutativity preserving maps. *Acta Sci. Math. (Szeged)* 71 (2005), 781-819.

DEPARTMENT OF MATHEMATICS, TSINGHUA UNIVERSITY, BEIJING 100084, P.R. CHINA

E-mail address: jcui@math.tsinghua.edu.cn

DEPARTMENT OF MATHEMATICS, COLLEGE OF WILLIAM & MARY, WILLIAMSBURG, VA 23187, USA

E-mail address: ckli@math.wm.edu

DEPARTMENT OF MATHEMATICS, IOWA STATE UNIVERSITY, AMES, IA 50011, USA

E-mail address: ytpoon@iastate.edu

Response to reviewers

We have revised the paper following the reviewer's comments