

2016

Bounds on probability of state transfer with respect to readout time and edge weight

Whitney Gordon

Brandon Univ, Dept Math & Comp Sci, Brandon, MB R7A 6A9, Canada;

Sarah Plosker

Brandon Univ, Dept Math & Comp Sci, Brandon, MB R7A 6A9, Canada;

Steve Kirkland

Univ Manitoba, Dept Math, Winnipeg, MB R3T 2N2, Canada;

Chi-Kwong Li

Coll William & Mary, Dept Math, Williamsburg, VA 23187 USA

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Gordon, W., Kirkland, S., Li, C. K., Plosker, S., & Zhang, X. (2016). Bounds on probability of state transfer with respect to readout time and edge weight. *Physical Review A*, 93(2), 022309.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Bounds on probability of state transfer with respect to readout time and edge weight

Whitney Gordon,¹ Steve Kirkland,² Chi-Kwong Li,³ Sarah Plosker,^{1,2,4} and Xiaohong Zhang²

¹*Department of Mathematics & Computer Science,
Brandon University, Brandon, MB, Canada R7A 6A9*

²*Department of Mathematics, University of Manitoba, Winnipeg, MB, Canada R3T 2N2*

³*Department of Mathematics, College of William and Mary, Williamsburg, VA, USA 23187*

⁴*Winnipeg Institute for Theoretical Physics, Winnipeg, Manitoba**

We analyse the sensitivity of a spin chain modelled by an undirected weighted connected graph exhibiting perfect state transfer to small perturbations in readout time and edge weight in order to obtain physically relevant bounds on the probability of state transfer. At the heart of our analysis is the concept of the numerical range of a matrix; our analysis of edge weight errors additionally makes use of the spectral and Frobenius norms.

PACS numbers: 02.10.Ud, 02.10.Yn, 03.65.Aa, 03.67.Hk, 03.67.Ac

Keywords: fidelity, transition probability, adjacency matrices, weighted graph, numerical range

I. INTRODUCTION

Transmitting a quantum state from one location to another is a critical task within a quantum computer. This task can be realised through the use of a spin chain (1D magnet). The seminal work by Bose [1] describes how a spin chain can be used as a quantum data bus for quantum communication within a quantum computer. This leads to the notion of perfect state transfer (PST, described in more detail below), a desirable property for quantum communication. Although the spin chain considered by Bose minimizes the amount of physical and technological resources required to transfer quantum states, it only exhibits PST for $n \leq 3$, where n is the number of spins in the chain. A spin chain can be represented by a graph, so many other types of graphs have been considered for quantum state transfer with $n > 3$.

The *fidelity* or *probability of state transfer* is a measure of the closeness between two quantum states and is used to determine the accuracy of state transfer through a quantum data bus between quantum registers and/or processors. Fidelity is a number between 0 and 1; when the fidelity between two quantum states is equal to 1 we have *perfect state transfer* (PST), and when the fidelity can be made arbitrarily close to 1 we have *pretty good state transfer* (PGST). Many families of graphs been shown to exhibit PST [2–8] or PGST [9].

In this work, we take a mathematical approach to perturbations which decrease the probability of state transfer. While our approach is similar in nature to that found in [10] and in [11], it should be noted that other authors take different approaches. In particular, there have been a number of numerical studies investigating the robustness of fidelity with respect to perturbations (e.g. [12–14]), while the consideration of a disordered XY model leads to discussions of the appearance of Anderson localization (see [15]). A recent paper [16] concerns the

use of error correcting codes as a strategy for dealing with imperfections (in contrast, we do not consider encoding/decoding schemes herein).

We consider state transfer probability as it applies to a weighted, undirected graph G , where vertices are labelled $1, \dots, n$ and the weight of the edge between vertices j and k is denoted $w(j, k)$. For any graph G , we consider its $n \times n$ adjacency matrix $A = [a_{jk}]$ defined via

$$a_{jk} = \begin{cases} w(j, k) & \text{if } j \text{ and } k \text{ are adjacent} \\ 0 & \text{otherwise} \end{cases}$$

as well as its Laplacian matrix $L = R - A$, where R is the diagonal matrix of row sums of A .

Depending on the dynamics of our system, the Hamiltonian H , representing the total energy of our system, is taken to be either A (in the case of XX dynamics) or L (in the case of Heisenberg (XXX) dynamics). Its spectrum represents the possible measurement outcomes when one measures the total energy of the system. Here we are not making full use of the Hamiltonian in that we are taking a snapshot in time—neither A nor L depends on t . We account for time by setting $U(t) = e^{itH}$. The fidelity of transfer from vertex s (sender) to vertex r (receiver) is then given by $p(t) = |u(t)_{sr}|^2$, where $u(t)_{sr}$ is the (s, r) -th entry of $U(t) = e^{itH}$.

Ideally, the fidelity is 1, representing perfect state transfer (PST) between the sender and receiver. In [10], the author discusses the very issue of tolerance of a spin chain with respect to timing errors and with respect to edge weight errors (so-called *manufacturing* errors). For timing errors, he derives a simple lower bound based on the squared difference between each eigenvalue and the smallest eigenvalue, noting that a Hamiltonian with minimal eigenvalue spread would optimize the bound for small perturbations in readout time. The bounds that we produce (for both the adjacency and Laplacian cases) look similar and in fact extend the lower bound given in [10]. Moreover, we give an example where our bound is attained for the adjacency matrix case, so it cannot be further improved in that setting. Sensitivity with respect

* ploskers@brandonu.ca

to perturbations in readout time is discussed in Section II.

For manufacturing errors, again in [10] the author finds that distances between eigenvalues are key, although no bound is given. This sensitivity analysis was continued in [11] through an analysis of the derivatives of the fidelity of state transfer with respect to either readout time or a fixed (j, k) -th edge weight. Again it was noted that minimizing the spectral spread optimizes the bound on the fidelity of state transfer for small perturbations in time. No explicit bound with respect to perturbations in edge weight were given in [11]; the edge weight results were more qualitative in nature. Here, we take several different approaches to give bounds on the probability of state transfer with respect to edge weight perturbation which involve both the spectral and Frobenius norms. Sensitivity with respect to perturbations in edge weights is discussed in Section III.

II. SENSITIVITY WITH RESPECT TO READOUT TIME

Suppose we have PST between vertices j and k at time t_0 . How sensitive is $p(t_0)$ to small changes in time? We would like $p(t_0 + h)$ to be close to $p(t_0)$ for small h .

Let us fix some notation now. Let $\{|1\rangle, |2\rangle, \dots, |n\rangle\}$ denote the standard ordered basis for \mathbb{C}^n . Let H be a real symmetric matrix that we decompose as $H = Q^T \Lambda Q$, where $\Lambda = \text{diag}(\lambda_1, \dots, \lambda_n)$ is a (real) diagonal matrix of eigenvalues and Q^T is an orthogonal matrix of corresponding eigenvectors. If we have PST between vertices j and k , we can permute the columns of Q so that we have PST between vertices 1 and 2. Thus we will always focus on the $(1, 2)$ entry of our matrix for simplicity and ease of notation. Let $|q_1\rangle = (q_{11}, \dots, q_{n1})^T$ and $|q_2\rangle = (q_{12}, \dots, q_{n2})^T$ be the first two columns of Q ; these vectors represent the first (and the second, respectively) entries of all the eigenvectors of H . We then have $e^{it_0 H} = Q^T e^{it_0 \Lambda} Q$. We are assuming PST between vertices 1 and 2 at time t_0 , and so

$$|\langle q_1 | e^{it_0 \Lambda} | q_2 \rangle| = 1,$$

giving $e^{i\theta} |q_1\rangle = e^{it_0 \Lambda} |q_2\rangle$ for some $\theta \in \mathbb{R}$, where (obviously) $|e^{i\theta}| = 1$. In particular, we have

$$\left| \langle q_1 | \tilde{M} e^{it_0 \Lambda} | q_2 \rangle \right| = \left| \langle q_1 | \tilde{M} e^{i\theta} | q_1 \rangle \right| = |\langle q_1 | M | q_1 \rangle|$$

for any matrix \tilde{M} (where we have set $M = \tilde{M} e^{i\theta}$). This innocuous observation will allow us to consider the $(1, 1)$ entry of $Q^T M Q$ rather than the $(1, 2)$ entry of $Q^T \tilde{M} e^{it_0 \Lambda} Q$ (see the proof of Theorem II.1 below).

The change from the $(1, 2)$ entry to the $(1, 1)$ entry will forge the link between the sensitivity of the fidelity with respect to readout time and the notion of the *numerical range* of an $n \times n$ matrix M , defined by

$$W(M) = \{ \langle x | M | x \rangle : |x\rangle \in \mathbb{C}^n, \langle x | x \rangle = 1 \}.$$

We now consider a small perturbation of readout time. The motivation for this is highly practical: even with lab equipment calibrated to an arbitrary amount of precision, if we want to readout at time t_0 , the readout time in practice will be $t_0 + h$ for small h (e.g. $h = \pm 0.0001$).

Theorem II.1 *Let H be either the adjacency matrix or the Laplacian associated with an undirected weighted connected graph with perfect state transfer at time t_0 ; that is, $p(t_0) = 1$. Suppose there is a small perturbation and the readout time is instead $t_0 + h$, where, denoting the smallest and largest eigenvalues of H by λ_1, λ_n , respectively, h satisfies $|h| < \frac{\pi}{\lambda_n - \lambda_1}$. Then the fidelity at the perturbed time $t_0 + h$ satisfies the following lower bound:*

$$p(t_0 + h) \geq \frac{1}{4} |e^{ih\lambda_1} + e^{ih\lambda_n}|^2.$$

Proof: Let $D = t_0 \Lambda = \text{diag}(t_0 \lambda_1, \dots, t_0 \lambda_n)$ and $\hat{D} = \text{diag}((t_0 + h)\lambda_1, \dots, (t_0 + h)\lambda_n)$. We find

$$\begin{aligned} \langle q_1 | e^{i\hat{D}} | q_2 \rangle &= \langle q_1 | (e^{i\hat{D}} e^{-iD}) e^{iD} | q_2 \rangle \\ &= \langle q_1 | (e^{i\hat{D}} e^{-iD}) e^{i\theta} | q_1 \rangle = \langle q_1 | M | q_1 \rangle \in W(M) \end{aligned}$$

with $M = \text{diag}(e^{ih\lambda_1}, \dots, e^{ih\lambda_n}) e^{i\theta}$. Now, $W(M)$ is the convex hull of $\{e^{i\theta} e^{ih\lambda_1}, \dots, e^{i\theta} e^{ih\lambda_n}\}$. Since $|h\lambda_n - h\lambda_1| < \pi$, there exists an $s \in [0, 2\pi)$ such that $e^{is} M$ has eigenvalues $e^{i\xi_1}, \dots, e^{i\xi_n}$ with $-\pi/2 < \xi_1 \leq \dots \leq \xi_n < \pi/2$ with $\xi_1 = -\xi_n$. Let $e^{is} M = M_1 + iM_2$ such that $M_1 = M_1^\dagger$ and $M_2 = M_2^\dagger$. Then M_1 has eigenvalues $\cos \xi_1, \dots, \cos \xi_n$ so that $0 \leq \cos \xi_1 = \cos \xi_n \leq \cos \xi_j$ for all $j = 2, \dots, n-1$. As a result, for every unit vector $|q\rangle \in \mathbb{C}^n$, we have

$$\begin{aligned} |\langle q | M | q \rangle| &= |\langle q | (M_1 + iM_2) | q \rangle| \geq |\langle q | M_1 | q \rangle| \geq \cos \xi_1 \\ &= |e^{i\xi_1} + e^{i\xi_n}|/2 = |e^{ih\lambda_1} + e^{ih\lambda_n}|/2. \end{aligned}$$

Thus, every point in $W(M)$ has a distance larger than $|e^{ih\lambda_1} + e^{ih\lambda_n}|/2$ from 0. Consequently,

$$\begin{aligned} p(t_0) - p(t_0 + h) &= |\langle q_1 | e^{iD} | q_2 \rangle|^2 - |\langle q_1 | e^{i\hat{D}} | q_2 \rangle|^2 \\ &\leq 1 - \frac{1}{4} |e^{ih\lambda_1} + e^{ih\lambda_n}|^2, \end{aligned}$$

and the result follows. \square

In fact, in the above proof, one can get a better estimate of $|\langle q_1 | M | q_1 \rangle|$ using the information of $|q_1\rangle = (q_{11}, \dots, q_{n1})^T$ and $M = \text{diag}(e^{ih\lambda_1}, \dots, e^{ih\lambda_n}) e^{i\theta}$; namely, for any $s \in \mathbb{R}$,

$$\begin{aligned} |\langle q_1 | M | q_1 \rangle| &= \left| \sum_{j=1}^n q_{j1}^2 e^{ih(\lambda_j - s)} \right| \geq \left| \sum_{j=1}^n q_{j1}^2 (\cos(h(\lambda_j - s))) \right| \\ &\geq \sum_{j=1}^n q_{j1}^2 - \frac{h^2}{2} \sum_{j=1}^n q_{j1}^2 (\lambda_j - s)^2 \\ &= 1 - \frac{h^2}{2} \sum_{j=1}^n q_{j1}^2 (\lambda_j - s)^2, \end{aligned}$$

the second inequality following from the fact that $\cos(x) \geq 1 - \frac{x^2}{2}$ for any $x \in \mathbb{R}$.

In particular, if we let $s = \lambda_1$ in the above, we obtain a result that is parallel to the bound in [10], without that paper's extra hypotheses on the Hamiltonian.

For general s , the above implies

$$\begin{aligned} p(t_0) - p(t_0 + h) &= 1 - |\langle q_1 | M | q_1 \rangle|^2 \\ &= (1 + |\langle q_1 | M | q_1 \rangle|)(1 - |\langle q_1 | M | q_1 \rangle|) \\ &\leq 2 \left(\frac{h^2}{2} \sum_{j=1}^n q_{j_1}^2 (\lambda_j - s)^2 \right) \\ &= h^2 \sum_{j=1}^n q_{j_1}^2 (\lambda_j - s)^2. \end{aligned}$$

We summarize these derivations in the following theorem.

Theorem II.2 *Let H be either the adjacency matrix or the Laplacian associated to an undirected weighted connected graph with perfect state transfer at time t_0 ; that is, $p(t_0) = 1$. Suppose there is a small perturbation and the readout time is instead $t_0 + h$, where, for $\lambda_1 \leq \dots \leq \lambda_n$, h satisfies $|h| < \frac{\pi}{\lambda_n - \lambda_1}$. Then, for any $s \in \mathbb{R}$, the transition probability for the perturbed time has the following lower bound:*

$$p(t_0 + h) \geq 1 - h^2 \sum_{j=1}^n q_{j_1}^2 (\lambda_j - s)^2.$$

Theorem II.2 is an improved bound compared to Theorem II.1. Yet direct use of Theorem II.2 requires one to find all eigenvalues $\lambda_1, \dots, \lambda_n$ of the Hamiltonian H , while Theorem II.1 requires only that the smallest and the largest eigenvalues are known. For large spin systems, Theorem II.1 would then be more practical. However, the following consequence of Theorem II.2 yields a lower bound on the fidelity that involves the physical parameters of the Hamiltonian itself, and does not require knowledge of any of the eigenvalues of H .

Corollary II.3 *Under the hypotheses of Theorem II.2, we have*

$$p(t_0 + h) \geq 1 - h^2 (\langle 1 | H^2 | 1 \rangle - (\langle 1 | H | 1 \rangle)^2). \quad (1)$$

Proof: Observe that the quantity $\sum_{j=1}^n q_{j_1}^2 (\lambda_j - s)^2$ is minimised when $s \sum_{j=1}^n q_{j_1}^2 = \sum_{j=1}^n q_{j_1}^2 \lambda_j$, i.e. when $s = \sum_{j=1}^n q_{j_1}^2 \lambda_j / \sum_{j=1}^n q_{j_1}^2 = \langle 1 | H | 1 \rangle$. The corresponding minimum value is then given by $\sum_{j=1}^n q_{j_1}^2 (\lambda_j - \langle 1 | H | 1 \rangle)^2 = \sum_{j=1}^n q_{j_1}^2 \lambda_j^2 - 2 \langle 1 | H | 1 \rangle \sum_{j=1}^n q_{j_1}^2 \lambda_j + (\langle 1 | H | 1 \rangle)^2 \sum_{j=1}^n q_{j_1}^2 = (\langle 1 | H^2 | 1 \rangle - (\langle 1 | H | 1 \rangle)^2)$. Inequality (1) now follows readily from Theorem II.2. \square

Inequality 1 of Corollary II.3 is fairly accurate in the following sense: In [11], the author considers the derivatives of p at time t_0 under the hypotheses of PST at t_0 . In [11, Theorem 2.2], it is shown that all odd order derivatives of p at t_0 are zero, while the second derivative is equal to $-2(\langle 1 | H^2 | 1 \rangle - (\langle 1 | H | 1 \rangle)^2)$. From [11, Theorem 2.4], it follows that the fourth derivative of p at t_0 is positive. It now follows that for all h

with $|h|$ sufficiently small, there is a $c > 0$ such that $p(t_0 + h) = 1 - h^2(\langle 1 | H^2 | 1 \rangle - (\langle 1 | H | 1 \rangle)^2) + ch^4 + O(h^6)$. In other words, for small h , the lower bound of Corollary II.3 is accurate to terms in h^3 .

We now consider the case where equality holds in the bound of Theorem II.1 when H is the adjacency matrix of a connected weighted graph. For concreteness, suppose that H is of order n and that there is perfect state transfer at time t_0 . Suppose further that for some h with $|h| < \frac{\pi}{\lambda_n - \lambda_1}$ we have $p(t_0) - p(t_0 + h) = 1 - \frac{1}{4}|e^{ih\lambda_1} + e^{ih\lambda_n}|^2$. Denote the multiplicity of λ_1 by k , and recall that λ_n , as the Perron value of H (that is, the unique maximal eigenvalue as per the Perron–Frobenius theorem), is necessarily simple. Examining the proof of Theorem II.1, it follows that the $|q_1\rangle$ can only have nonzero entries in positions corresponding to the eigenvalues λ_1 and λ_n , and that further the entry in the position corresponding to λ_n , the j -th position say, must be $\pm \frac{1}{\sqrt{2}}$. Since the entries of $|q_2\rangle$ can only differ from the corresponding entries of $|q_1\rangle$ by a sign, we deduce that the j -th entry of $|q_2\rangle$ must also be $\pm \frac{1}{\sqrt{2}}$. Observe that since every column of Q^T has 2-norm equal to 1, the j -th column of Q^T has nonzero entries only in its first two rows. But the j -th column of Q^T is a Perron vector for H —that is, an eigenvector corresponding to the (positive and dominant) Perron eigenvalue of H —and so it cannot have any zero entries (again by the Perron–Frobenius theorem). We thus deduce that n must be 2, and that H must be a positive scalar multiple of $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$.

Conversely, suppose that H is a positive scalar multiple of $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$, and without loss of generality we assume that $H = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$. Then

$$\begin{aligned} e^{itH} &= \sum_{j=0}^{\infty} \frac{i^j}{j!} H^j t^j = \sum_{j \text{ even}} \frac{(i)^j}{j!} t^j I + \sum_{j \text{ odd}} \frac{(i)^j}{j!} t^j H \\ &= \begin{pmatrix} \cos t & i \sin t \\ i \sin t & \cos t \end{pmatrix}. \end{aligned}$$

At time $\frac{\pi}{2}$ we have $\begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$ and so we have PST (since $|i| = 1$). At time $\frac{\pi}{2} + h$, the (1, 2) entry is $i \cos h = \frac{1}{2}(e^{ih} + e^{-ih})$, and so the bound in Theorem II.1 is attained.

Although Theorem II.1 is true for either adjacency matrices or Laplacians, we can adapt the technique of Theorem II.1 slightly to produce an improved bound on the fidelity in the setting of the Laplacian matrix, since we have more information at hand.

Theorem II.4 *Let L be the Laplacian matrix of a connected weighted graph on $n \geq 3$ vertices. Denote the eigenvalues of L by $0 \equiv \lambda_1 < \lambda_2 \leq \dots \leq \lambda_n$. Suppose that there is perfect state transfer at time t_0 ; that is, $p(t_0) = 1$. Suppose there is a small timing perturbation, and the readout time is instead $t_0 + h$, where h*

satisfies $|h| < \frac{\pi}{\lambda_n - \lambda_2}$. Then

$$p(t_0 + h) \geq 1 - \frac{(n-1)^2(1 - \cos((\lambda_n - \lambda_2)h))}{2n^2} - \frac{(n-1)(2 - \cos(\lambda_2 h) - \cos(\lambda_n h))}{n^2} - \frac{(\cos(\lambda_2 h) - \cos(\lambda_n h))^2}{2n^2(1 - \cos((\lambda_n - \lambda_2)h))}. \quad (2)$$

Proof: We note that the normalised all-ones vector $\frac{1}{\sqrt{n}} \sum_{i=1}^n |i\rangle$ is a null vector for L . Then $L = Q^T \text{diag}(\lambda_1, \dots, \lambda_n) Q$ such that $|q_1\rangle$ has the form $(1/\sqrt{n}, x_2, \dots, x_n)^T$ where $\sum_{j=2}^n x_j^2 = \frac{n-1}{n}$. Mimicking the proof of Theorem II.1, we find that $p(t_0 + h)$ is bounded below by

$$\min \left| \frac{1}{n} + \langle z | \text{diag}(e^{ih\lambda_2}, \dots, e^{ih\lambda_n}) | z \rangle \right|^2,$$

where the minimum is taken over all unnormalised $|z\rangle \in \mathbb{R}^{n-1}$ such that $\langle z | z \rangle = \frac{n-1}{n}$. From elementary geometric considerations (in short, the $e^{ih\lambda_j}$ are points on the unit circle and so the minimum will be attained by taking a convex combination of the smallest and largest values, namely $e^{ih\lambda_2}$ and $e^{ih\lambda_n}$), we find that in fact

$$\min \left| \frac{1}{n} + \langle z | \text{diag}(e^{ih\lambda_2}, \dots, e^{ih\lambda_n}) | z \rangle \right|^2 = \min_{0 \leq \alpha \leq \frac{n-1}{n}} \left| \frac{1}{n} + \alpha e^{ih\lambda_2} + \left(\frac{n-1}{n} - \alpha \right) e^{ih\lambda_n} \right|^2.$$

A routine calculus exercise (the minimum corresponds to $\alpha = \frac{n-1}{2n} + \frac{\cos(h\lambda_n) - \cos(h\lambda_2)}{2n(1 - \cos(h(\lambda_2 - \lambda_n)))}$) shows that this last quantity is given by the right hand side of (2). \square

Example II.5 Suppose that n is divisible by 4, and consider the unweighted graph on vertices $1, \dots, n$, say G , formed by deleting the edge between vertices 1 and 2 from the complete graph on n vertices. Let L denote the Laplacian matrix for G , and note that L has three eigenvalues: 0, with corresponding eigenprojection matrix $\frac{1}{n} J$ (where J is the all ones matrix), $n-2$ with eigenprojection matrix $\frac{1}{2}(|1\rangle - |2\rangle)(\langle 1| - \langle 2|)$, and n with eigenprojection matrix

$$\left[\begin{array}{c|c} \frac{n-2}{2n} J_2 & -\frac{1}{n} J_{2,n-2} \\ \hline -\frac{1}{n} J_{n-2,2} & I - \frac{1}{n} J_{n-2} \end{array} \right].$$

It is shown in [7] that, using L as the Hamiltonian, there is perfect state transfer from vertex 1 to vertex 2 at time $\frac{\pi}{2}$.

Using the eigenvalues and eigenprojection matrices above, we find that for any h , the fidelity at time $\frac{\pi}{2} + h$ is given by

$$p\left(\frac{\pi}{2} + h\right) = \left| \frac{1}{n} + \frac{1}{2} e^{ih(n-2)} + \frac{n-2}{2n} e^{ihn} \right|^2.$$

This last expression can be simplified to yield

$$p\left(\frac{\pi}{2} + h\right) = 1 - \frac{n-2}{2n}(1 - \cos(2h)) - \frac{1}{n}(1 - \cos((n-2)h)) - \frac{(n-2)}{n^2}(1 - \cos(nh)).$$

An uninteresting computation reveals that $p(\frac{\pi}{2} + h)$ exceeds the lower bound of (2) in the amount of

$$\frac{(\cos((n-2)h) - \cos(nh) + 1 - \cos(2h))^2}{2n^2(1 - \cos(2h))}.$$

We note in passing that $\frac{(\cos((n-2)h) - \cos(nh) + 1 - \cos(2h))^2}{2n^2(1 - \cos(2h))}$ is asymptotic to h^2 as $h \rightarrow 0$.

III. SENSITIVITY WITH RESPECT TO EDGE WEIGHTS

The motivation for studying edge weight perturbation is similar to that for readout time: although lab equipment can be calibrated to high precision, small errors (even machine epsilon) will affect the state transition probability. We want to bound this effect so that small perturbations in edge weight do not drastically reduce transition probability.

Suppose we have PST between vertices j and k at time t_0 . As in section II, without loss of generality we consider the $(1, 2)$ entry of our matrix under consideration. Here, we keep the time constant at t_0 , and perturb edge weights, although the techniques employed in section II for readout time can readily be combined with the techniques in this section to obtain a bound for the situation when both readout time and edge weights are perturbed.

Let $H \in M_n$ be a real symmetric matrix representing the Hamiltonian of our system. Suppose $\langle 1 | e^{it_0 H} | 2 \rangle$ has modulus 1; this is the case for PST with H either an adjacency or Laplacian matrix. Consider now $\hat{H} = t_0 H + H_0$ where H_0 is a matrix representing small perturbations of edge weights and we have absorbed the time component into H_0 since we are keeping time fixed. Mathematically, we would like to find a perturbation bound for $|\langle 1 | e^{i\hat{H}} | 2 \rangle|^2 - |\langle 1 | e^{i(t_0 H + H_0)} | 2 \rangle|^2$ or, when $t = t_0$,

$$1 - |\langle 1 | e^{i(t_0 H + H_0)} | 2 \rangle|^2$$

for a symmetric matrix H with sufficiently small H_0 , say, measured by the operator norm $\|H_0\|$ or the Frobenius norm $\|H_0\|_F$.

Note that the entries of the matrix H_0 represent individual edge weight errors, so our approach allows for individual edge weight perturbations rather than simply an overall (global) edge weight perturbation (where all edge weights are perturbed by e.g. 0.0001 in the same direction) or a single edge weight perturbation (where all other edge weights remain unperturbed); the latter case was the (rather restrictive) situation considered in [11].

We begin with the following.

Theorem III.1 Suppose a perfect state transfer occurs at time t_0 , and $\hat{H} = t_0 H + H_0$, with a small nonzero perturbation H_0 . Then

$$\|e^{i(Ht_0 + H_0)} - e^{iHt_0}\| \leq \|H_0\| e^{\|H_0\|}.$$

Consequently,

$$1 - |\langle 1 | e^{i(Ht_0 + H_0)} | 2 \rangle|^2 \leq 2\|H_0\| e^{\|H_0\|} - \|H_0\|^2 e^{2\|H_0\|} \leq 2\|H_0\| + \|H_0\|^2 - \|H_0\|^3. \quad (3)$$

Proof: Set $\Delta t_0 = H_0$. Using the result in [17, p.532] and the fact that H is Hermitian, we have

$$\|e^{i(H+\Delta)t_0} - e^{iHt_0}\| \leq t_0 \|\Delta\| e^{t_0 \|\Delta\|} = \|H_0\| e^{\|H_0\|}.$$

Consequently,

$$|\langle 1|e^{it_0H}|2\rangle| - |\langle 1|e^{i(t_0H+H_0)}|2\rangle| \leq \|H_0\| e^{\|H_0\|}$$

so that

$$1 - \|H_0\| e^{\|H_0\|} \leq |\langle 1|e^{i(t_0H+H_0)}|2\rangle|.$$

Squaring both sides and rearranging terms, we have

$$\begin{aligned} & 1 - |\langle 1|e^{i(t_0H+H_0)}|2\rangle|^2 \\ & \leq 2\|H_0\| e^{\|H_0\|} - \|H_0\|^2 e^{2\|H_0\|} \\ & = 2\|H_0\| \left(1 + \frac{\|H_0\|}{1!} + \frac{\|H_0\|^2}{2!} + \dots\right) \\ & \quad - \|H_0\|^2 \left(1 + \frac{2\|H_0\|}{1!} + \frac{(2\|H_0\|)^2}{2!} + \dots\right) \\ & \leq 2\|H_0\| + \|H_0\|^2 - \|H_0\|^3. \end{aligned}$$

so that (3) holds. \square

We note that the estimate $\|e^{i(Ht_0+H_0)} - e^{iHt_0}\| \leq \|H_0\| e^{\|H_0\|}$ of Theorem III.1 can be reasonably accurate. For example, suppose H is the adjacency matrix of a connected weighted graph yielding perfect state transfer at time t_0 . Let $|v\rangle$ denote the positive Perron vector of H with norm one, and suppose that H_0 has the form $\epsilon|v\rangle\langle v|$ for some small $\epsilon > 0$. Then $\|e^{i(Ht_0+H_0)} - e^{iHt_0}\| = |e^{i\epsilon} - 1|$ while $\|H_0\| e^{\|H_0\|} = \epsilon e^\epsilon$, so that

$$\frac{\|e^{i(Ht_0+H_0)} - e^{iHt_0}\|}{\|H_0\| e^{\|H_0\|}} \rightarrow 1$$

as $\epsilon \rightarrow 0^+$.

If we have additional information about the matrix t_0H , we may be able to produce some better bounds as shown in Theorem III.3 below. Before presenting the theorem, we require a preliminary proposition which is intuitively clear. Its proof consists of elementary linear algebra manipulation techniques; we give the proof for completeness.

Proposition III.2 *Suppose there is a perfect state transfer at time t_0 ; that is, $|\langle 1|e^{it_0H}|2\rangle| = 1$. Then for some $\theta \in \mathbb{R}$, $t_0H = Q^T D Q - \theta I$, where*

$$D = \pi \text{diag}(r_1, \dots, r_\ell, r_{\ell+1}, \dots, r_m, r_{m+1}, \dots, r_n)$$

such that $r_1 \geq \dots \geq r_\ell$ are positive even integers, and $r_{\ell+1} \geq \dots \geq r_m$ are positive odd integers. Further, the first two rows of Q^T can be taken to have the form $(x_1, \dots, x_m, 0, \dots, 0)$ and $(x_1, \dots, x_\ell, -x_{\ell+1}, \dots, -x_m, 0, \dots, 0)$ satisfying $x_1, \dots, x_m \geq 0$.

Proof: As in section II, let $H = Q^T \Lambda Q$ for a real diagonal matrix Λ and Q an orthogonal matrix. Since we are focusing here on edge weights rather than time,

let us consider $D = t_0 \Lambda = \text{diag}(t_0 \lambda_1, \dots, t_0 \lambda_n)$. Suppose the first two columns of Q are $|q_1\rangle$ and $|q_2\rangle$. Then $|\langle q_1|e^{iD}|q_2\rangle| = 1$ implies that $e^{iD}|q_2\rangle = e^{i\theta}|q_1\rangle$ or, equivalently, $e^{-i\theta}e^{iD}|q_2\rangle = |q_1\rangle$ for some $\theta \in \mathbb{R}$. Any zero entries of $|q_1\rangle$ show up correspondingly as zero entries of $|q_2\rangle$. For a suitable permutation matrix P_1 we can replace (D, Q) by $(P_1 D P_1^T, P_1 Q)$ so that the zero entries of $P_1|q_1\rangle$ all occur in the last $n - m$ entries, for some $0 < m \leq n$. In this way, we may assume that $P_1(D - \theta I)P_1^T$ is a diagonal matrix with diagonal entries of the form $s_1\pi, \dots, s_m\pi, *, \dots, *$ for some integers s_1, \dots, s_m . The asterisks in the $(m+1, m+1)$ up to (n, n) entries of the diagonal matrix $P_1(D - \theta I)P_1^T$ represent unknown constants, corresponding to the zero entries (if any) of $P_1|q_1\rangle$. We can replace θ by $\theta - 2s\pi$ for a sufficiently large integer s so that we may assume that s_1, \dots, s_m are positive integers.

Next, for a suitable permutation matrix P_2 we can replace the pair $(P_1 D P_1^T, P_1 Q)$ by $(P_2 P_1 D P_1^T P_2^T, P_2 P_1 Q)$, so that $P_2 P_1(D - \theta I)P_1^T P_2^T = \pi \text{diag}(r_1, \dots, r_n)$ with $r_1 \geq \dots \geq r_\ell$ even, $r_{\ell+1} \geq \dots \geq r_m$ odd, and r_{m+1}, \dots, r_n unknown constants; note that we still have $t_0H = (P_2 P_1 Q)^T (P_2 P_1 D P_1^T P_2^T) (P_2 P_1 Q)$. Further, we may replace the pair $(P_2 P_1 D P_1^T P_2^T, P_2 P_1 Q)$, by $(S P_2 P_1 D P_1^T P_2^T S, S P_2 P_1 Q)$, for some diagonal orthogonal matrix (i.e. a signature matrix) S such that the first column of $S P_2 P_1 Q$, namely $S P_2 P_1|q_1\rangle = (x_1, \dots, x_m, 0, \dots, 0)^T$, satisfies $x_1, \dots, x_m \geq 0$. Now, $S P_2 P_1 e^{-i\theta} e^{iD}|q_2\rangle = S P_2 P_1|q_1\rangle$ implies that

$$S P_2 P_1|q_2\rangle = (x_1, \dots, x_\ell, -x_{\ell+1}, \dots, -x_m, 0, \dots, 0)^T.$$

Relabelling for simplicity, the result now follows. \square

Theorem III.3 *Suppose a perfect state transfer occurs at time t_0 , and $\hat{H} = t_0H + H_0$, with a small nonzero perturbation H_0 . Furthermore, assume that value m in Proposition III.2 equals n . Then*

$$1 - |\langle 1|e^{i\hat{H}}|2\rangle|^2 \leq \frac{2\|H_0\|_F^2}{(\pi - \|H_0\|)^2} + \|H_0\|^2 + O(\|H_0\|^3).$$

Proof: Let $t_0H = Q^T D Q - \theta I$, where $D = \text{diag}(d_1, \dots, d_n)$ is such that the first ℓ entries are even multiples of π and the last $n - \ell$ entries are odd multiples of π , and θ is as in Proposition III.2. Let $J = e^{iD} = I_\ell \oplus -I_{n-\ell}$.

Suppose $\hat{H} = t_0H + H_0 = \hat{Q}^T \hat{D} \hat{Q}$. By a suitable choice of \hat{Q} , we may assume that there is a permutation matrix P such that both $P^T D P$ and $P^T \hat{D} P$ have diagonal entries arranged in descending order. Then (e.g., see [18, p.101,(IV.62)])

$$\|D - \hat{D}\| = \|P^T D P - P^T \hat{D} P\| \leq \|Q^T D Q - \hat{Q}^T \hat{D} \hat{Q}\| = \|H_0\|$$

and hence

$$\|e^{iD} - e^{i\hat{D}}\| \leq \|D - \hat{D}\| \leq \|H_0\|. \quad (4)$$

Let V be an orthogonal matrix close to I , and consider the power series $\log(V) = -\sum_{j=1}^{\infty} \frac{1}{j} (I - V)^j$. Setting

$K = \log(V)$, we have $e^K = V$. It follows that $I = VV^T = e^K e^{K^T} = e^{K+K^T}$, where the last equality comes from the fact that K commutes with K^T . We deduce that $K^T = -K$, i.e. K is skew-symmetric. We will use this idea in what follows.

If H_0 is small, we may assume that the differences between the eigenspaces of \hat{H} and H are small so that \hat{D} is close to D , and $\hat{Q}Q^T$ is close to I (see [18, Section VII.3]). As a result, we can write $\hat{D} - D = wD_1$ and $e^{wK} = \hat{Q}Q^T$ for a small positive number w , a diagonal matrix D_1 and a skew-symmetric matrix K such that $\max\{\|D_1\|, \|K\|\} = 1$ (the norm condition is required so that the terms like K^3, D_1^3 can be lumped into the $O(w^3)$ term below). Note that it is possible that $D_1 = 0$ or $K = 0$ but not both as $H_0 \neq 0$. We emphasize that $\log(\hat{Q}Q^T)$ is skew-symmetric, from the above remark about a matrix V .

Now $\hat{H} = Q_w^T(D + wD_1)Q_w$ where we write $Q_w = e^{wK}Q$, using the subscript w here to emphasize the dependence on some small positive number w . Using the power series expansion of H_w and the fact that $K = -K^T$, we get

$$\begin{aligned} \langle 1|e^{iH_w}|2\rangle &= \langle 1|Q_w^T e^{iD_w} Q_w|2\rangle \\ &= \langle 1|[Q^T(I + wK + \frac{1}{2}w^2K^2)^T e^{iD}(I + iwD_1 - w^2\frac{1}{2}D_1^2)] \\ &\quad \times (I + wK + \frac{1}{2}w^2K^2)]Q|2\rangle + O(w^3) \\ &= \langle 1|Q^T e^{iD} Q|2\rangle + w\langle 1|Q^T [K^T e^{iD} + e^{iD}iD_1 + e^{iD}K]Q|2\rangle \\ &\quad + \frac{1}{2}w^2\langle 1|Q^T [(K^T)^2 e^{iD} + e^{iD}K^2 - e^{iD}D_1^2]Q|2\rangle \\ &\quad + w^2\langle 1|Q^T [K^T e^{iD}iD_1 + K^T e^{iD}K + e^{iD}iD_1K]Q|2\rangle + O(w^3). \end{aligned}$$

By the facts that $J|q_2\rangle = |q_1\rangle$ and $\langle q|K|q\rangle = 0$ for any vector $|q\rangle$, the above expression becomes

$$\begin{aligned} \langle 1|Q_w^T e^{iD_w} Q_w|2\rangle &= 1 + w(\langle q_1|K^T|q_1\rangle + \langle q_2|K|q_2\rangle) \\ &\quad + \frac{1}{2}w^2\langle q_1|((K^T)^2 - D_1^2 + JK^2J + 2K^TJKJ)|q_1\rangle \\ &\quad + iw\langle q_1|D_1|q_1\rangle + iw^2\langle q_1|(K^T D_1 + JD_1KJ)|q_1\rangle + O(w^3) \\ &= 1 + \frac{1}{2}w^2\langle q_1|((K^T)^2 - D_1^2 + JK^2J + 2K^TJKJ)|q_1\rangle \\ &\quad + iw\langle q_1|D_1|q_1\rangle + iw^2\langle q_1|(K^T D_1 + D_1JKJ)|q_1\rangle + O(w^3). \end{aligned}$$

Let $x(w) = \Re(\langle 1|Q_w^T e^{iD_w} Q_w|2\rangle)$, and $y(w) = \Im(\langle 1|Q_w^T e^{iD_w} Q_w|2\rangle)$. Then

$$\begin{aligned} |\langle 1|e^{i(t_0H+H_0)}|2\rangle|^2 - |\langle 1|e^{it_0H}|2\rangle|^2 &= |\langle 1|e^{i(t_0H+H_0)}|2\rangle|^2 - 1 \\ &= x(w)^2 + y(w)^2 - 1 \\ &= w^2[\langle q_1|(K^2 + JK^2J - 2KJKJ - D_1^2)|q_1\rangle \\ &\quad + (\langle q_1|D_1|q_1\rangle)^2] + O(w)^3 \\ &= -w^2(\|(KJ - JK)|q_1\rangle\|^2 + \|D_1|q_1\rangle\|^2 \\ &\quad - (\langle q_1|D_1|q_1\rangle)^2) + O(w^3). \end{aligned}$$

For the last equality in the above expression, we use the fact that, although $JKJK \neq KJKJ$, it is true that

$\langle q_1|JKJK|q_1\rangle = \langle q_1|KJKJ|q_1\rangle$, which is all that is required here.

$$\begin{aligned} \text{Thus, if } wK &= \begin{bmatrix} K_{11} & K_{12} \\ -K_{12}^T & K_{22} \end{bmatrix}, \text{ then } wKJ - wJK = \\ &\begin{bmatrix} O & -2K_{12} \\ -2K_{12}^T & O \end{bmatrix} \text{ and hence} \\ &\|(wKJ - wJK)|q_1\rangle\|^2 \leq 4\|K_{12}\|^2. \end{aligned} \quad (5)$$

Now,

$$\hat{Q}Q^T = e^{wK} = I + wK + \frac{(wK)^2}{2!} + \frac{(wK)^3}{3!} + \dots$$

So, $wK \approx (\hat{Q}Q^T - Q\hat{Q}^T)/2$.

As a result, if $\hat{Q}Q^T \equiv V = \begin{bmatrix} V_{11} & V_{12} \\ V_{21} & V_{22} \end{bmatrix}$, then V_{12} and V_{21} have the same nonzero singular values (there are a number of ways of seeing this; perhaps the simplest is to note that since $\hat{Q}Q^T$ is orthogonal, it follows that $V_{11}V_{11}^T + V_{12}V_{12}^T = I$ and $V_{11}^TV_{11} + V_{21}^TV_{21} = I$; that is, $V_{11}V_{11}^T = I - V_{12}V_{12}^T$ and $V_{11}^TV_{11} = I - V_{21}^TV_{21}$, from which we see $V_{12}V_{12}^T$ and $V_{21}^TV_{21}$ have the same eigenvalues). Note that

$$\begin{aligned} \|H_0\|_F^2 &= \|Q^T DQ - \hat{Q}^T \hat{D} \hat{Q}\|_F^2 = \|VD - \hat{D}V\|_F^2 \\ &= \sum_{j,k} (d_k - \hat{d}_j)^2 v_{jk}^2. \end{aligned}$$

The reverse triangle inequality gives us

$$\begin{aligned} |d_k - \hat{d}_j| &= |d_k - d_j + d_j - \hat{d}_j| \geq |d_k - d_j| - |d_j - \hat{d}_j| \\ &\geq \pi - \|H_0\|. \end{aligned}$$

We now have

$$\begin{aligned} \|H_0\|_F^2 &= \sum_{j,k} (d_k - \hat{d}_j)^2 v_{jk}^2 \\ &= \sum_{1 \leq j \leq \ell < k \leq n} (d_k - \hat{d}_j)^2 v_{jk}^2 + \sum_{1 \leq k \leq \ell < j \leq n} (d_k - \hat{d}_j)^2 v_{jk}^2 \\ &\geq (\pi - \|H_0\|)^2 (\|V_{12}\|_F^2 + \|V_{21}\|_F^2). \end{aligned}$$

It follows that

$$\|V_{12}\|_F^2 + \|V_{21}\|_F^2 \leq \frac{\|H_0\|_F^2}{(\pi - \|H_0\|)^2}.$$

Hence,

$$\begin{aligned} \|K_{12}\| &\leq (\|V_{12}\| + \|V_{21}\|)/2 = \|V_{12}\| \leq \|V_{12}\|_F \\ &\leq \frac{\|H_0\|_F}{\sqrt{2}(\pi - \|H_0\|)}. \end{aligned} \quad (6)$$

As a result, from (5) and (6) we have

$$w^2\|(KJ - JK)|q_1\rangle\|^2 \leq 4\|K_{12}\|^2 \leq \frac{2\|H_0\|_F^2}{(\pi - \|H_0\|)^2}.$$

A result of Mirsky [19] states that for any Hermitian matrix M , the eigenvalue spread for M is equal

to $2 \max |\langle u|A|v \rangle|$, where the maximum is taken over all pairs of orthonormal vectors $|u\rangle$ and $|v\rangle$. Consequently, for any symmetric matrix A , if $\{|u\rangle, |v\rangle\}$ is an orthonormal set, then

$$2|\langle u|A|v \rangle| \leq \lambda_n(A) - \lambda_1(A),$$

where we recall λ_1 is the minimum eigenvalue and λ_n is the maximum eigenvalue. In particular, if we set $A = wD_1$, $|u\rangle = |q_1\rangle$, and $A|q_1\rangle = \mu_1|q_1\rangle + \mu_2|q\rangle$ such that $\{|q_1\rangle, |q\rangle\}$ is an orthonormal set, then

$$\begin{aligned} & \|wD_1|q_1\rangle\|^2 - (\langle q_1|wD_1|q_1\rangle)^2 \\ &= \mu_2^2 = |\langle q|(wD_1)|q_1\rangle|^2 \\ &\leq ((\lambda_n(wD_1) - \lambda_1(wD_1))/2)^2. \end{aligned}$$

By (4), and recalling that D_1 is diagonal, we have

$$((\lambda_n(wD_1) - \lambda_1(wD_1))/2)^2 \leq \|\hat{D} - D\|^2 \leq \|H_0\|^2.$$

□

Consider the bounds of Theorems III.1 and III.3 when the perturbing matrix H_0 is small. The upper bound in the former result is $2\|H_0\| + \|H_0\|^2 - \|H_0\|^3$, while the upper bound in the latter result is $\frac{2\|H_0\|_F^2}{(\pi - \|H_0\|)^2} + \|H_0\|^2 + O(\|H_0\|^3)$. Thus we find that, neglecting terms of order $\|H_0\|^3$, the bound of Theorem III.3 is sharper than that of Theorem III.1 provided that

$$\frac{\|H_0\|_F^2}{(\pi - \|H_0\|)^2} < \|H_0\|. \quad (7)$$

Suppose for concreteness that H_0 has rank r . Recalling that $\|H\|_F^2 \leq r\|H_0\|^2$, we find that in order for (7) to hold, it is sufficient that $r\|H_0\| < (\pi - \|H_0\|)^2$, or equivalently, that $\|H_0\| < \frac{2\pi + r - \sqrt{4\pi r + r^2}}{2}$. It now follows that for all sufficiently small H_0 , the bound of Theorem III.3 is improvement upon that of Theorem III.1. Thus, in the case that the more restrictive hypothesis of Theorem III.3 holds, we get a better estimate from that result than from Theorem III.1.

Example III.4 *Here we give a small numerical example illustrating the main result of Theorem III.3. Consider the 10×10 symmetric tridiagonal matrix H with $h_{j,j+1} = h_{j+1,j} = \sqrt{j(10-j)}$, $j = 1, \dots, 9$ and all other entries equal to 0. It is known that for this H , there is perfect state transfer from 1 to 10 at time $t_0 = \frac{\pi}{2}$, with the $(1, 10)$ entry of $e^{it_0 H}$ equal to i .*

Next, we consider the perturbing matrix

$$H_0 = 10^{-5} \times \begin{bmatrix} 0 & 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0.533 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & .533 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \end{bmatrix}.$$

Setting $\hat{H} = t_0 H + H_0$, a couple of MATLAB[©] computations yield $1 - |\langle 1|e^{i\hat{H}}|10\rangle|^2 \approx 0.02497 \times 10^{-9}$ and $\frac{2\|H_0\|_F^2}{(\pi - \|H_0\|)^2} + \|H_0\|^2 \approx 0.19257 \times 10^{-9}$. We note that the ratio of the latter to the former is approximately 7.7110.

IV. CONCLUSION

We have obtained bounds on the probability of state transfer for a perturbed system, where either readout time or edge weights have been perturbed. By considering such timing and manufacturing errors, our results are physically relevant and more consistent with reality. We worked in the most general setting where the adjacency matrix A (or, alternatively, the Laplacian L) was arbitrary, and the perturbations themselves were arbitrary. More precise bounds can be obtained by considering more structured perturbations. Furthermore, it would be of interest to combine readout time error with edge weight error to create one bound encompassing both types of perturbations. Finally, we note that our analysis assumed perfect state transfer (PST). While there are a number of classes of graphs exhibiting PST, it is of interest to allow for pretty good state transfer (PGST) and perform a similar analysis with respect to readout time and edge weight errors; note that the numerical evidence reported in Examples 3.16 and 3.17 in [11] suggests that the fidelity may not be so well-behaved under perturbation of edge weights in the PGST setting. Analysis in the PGST case would require alternate techniques, however, since our arguments hinged on the modulus of the $(1, 2)$ entry of our matrix e^{itH} being exactly 1, which facilitates the key observation that $e^{it_0} |q_1\rangle = e^{it_0 \Lambda} |q_2\rangle$. We leave these as open problems for further study.

ACKNOWLEDGMENTS

W.G. was supported through a NSERC Undergraduate Student Research Award. S.K. and S.P. are supported by NSERC Discovery Grants. X.Z. is supported by the University of Manitoba's Faculty of Science and Faculty of Graduate Studies. C.-K.L. is supported by USA NSF grant DMS 1331021, Simons Foundation Grant 351047,

and NNSF of China Grant 11571220. The authors wish to thank the anonymous referee for useful comments and

for pointing out additional relevant literature; in particular, for suggesting the result Corollary II.3.

-
- [1] S. Bose, *Quantum communication through an unmodulated spin chain*, Phys. Rev. Lett., **91** (2003), 207901.
- [2] M. Christandl, N. Datta, A. Ekert, and A. J. Landahl, *Perfect State Transfer in Quantum Spin Networks*, Phys. Rev. Lett. **92** (2004) 187902.
- [3] M. Christandl, N. Datta, T. C. Dorlas, A. Ekert, A. Kay, and A. J. Landahl, *Perfect transfer of arbitrary states in quantum spin networks*, Phys. Rev. A, **71** (2005) 032312.
- [4] A. Bernasconi, C. Godsil, and S. Severini, *Quantum networks on cubelike graphs*, Phys. Rev. A **78** (2008), 052320.
- [5] R.J. Angeles-Canul, R.M. Norton, M.C. Opperman, C.C. Paribello, M.C. Russell, and C. Tamonk, *Quantum perfect state transfer on weighted join graphs*, Int. J. Quantum Inf. **7** (2009), 1429-1445.
- [6] R.J. Angeles-Canul, R.M. Norton, M.C. Opperman, C.C. Paribello, M.C. Russell, and C. Tamonk, *Perfect state transfer, integral circulants and join of graphs*, Quant. Inform. Comput. **10** (2010) 325-342.
- [7] S. Kirkland and S. Severini, *Spin-system dynamics and fault detection in threshold networks*, Phys. Rev. A, **83** (2011), 012310.
- [8] M. Bašić, *Characterization of quantum circulant networks having perfect state transfer*, Quantum Inf. Process. **12** (2013), 345-364.
- [9] C. Godsil, S. Kirkland, S. Severini, and J. Smith, *Number-theoretic nature of communication in quantum spin systems* Phys. Rev. Lett. **109** (2012), 050502.
- [10] A. Kay, *Perfect state transfer: beyond nearest-neighbor couplings*, Phys. Rev. A **73** (2006) 032306.
- [11] S. Kirkland, *Sensitivity analysis of perfect state transfer in quantum spin networks*, Linear Algebra Appl. **472** (2015), 1–30.
- [12] G. De Chiara, D. Rossini, S. Montangero, and R. Fazio, *From perfect to fractal transmission in spin chains*, Phys. Rev. A **72** (2005) 012323.
- [13] R. Ronke, T. P. Spiller, and I. D’Amico, *Effect of perturbations on information transfer in spin chains*, Phys. Rev. A **83** (2011) 012325.
- [14] A. Zwick, G. A. Álvarez, J. Stolze, and O. Osenda, *Robustness of spin-coupling distributions for perfect quantum state transfer*, Phys. Rev. A **84** (2011) 022311.
- [15] C. K. Burrell and T. J. Osborne, *Bounds on the Speed of Information Propagation in Disordered Quantum Spin Chains*, Phys. Rev Lett. **99** (2007) 167201.
- [16] A. Kay, *Quantum Error Correction for Noisy Quantum Wires*, arxiv:1507.06139 (2015)
- [17] G. Golub and C. Van Loan, *Matrix Computations*, 4th edition, Johns Hopkins Press, Baltimore, 2013.
- [18] R. Bhatia, *Matrix Analysis*, Springer, New York, 1997.
- [19] L. Mirsky, *Inequalities for normal and Hermitian matrices*, Duke Math. J. **24** (1957), 591–599.