

2016

Plant characteristics associated with widespread variation in eelgrass wasting disease

ML Groner
Virginia Institute of Marine Science

CA Burge

CJS Kim

E Rees

KL Van Alstyne

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Groner, ML; Burge, CA; Kim, CJS; Rees, E; and Van Alstyne, KL, Plant characteristics associated with widespread variation in eelgrass wasting disease (2016). *Diseases Of Aquatic Organisms*, 118(2), 159-168.
10.3354/dao02962

This Article is brought to you for free and open access by the Virginia Institute of Marine Science at W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Plant characteristics associated with widespread variation in eelgrass wasting disease

Maya L. Groner^{1,7,*}, Colleen A. Burge², Catherine J. S. Kim³, Erin Rees¹,
Kathryn L. Van Alstyne⁴, Sylvia Yang⁴, Sandy Wyllie-Echeverria^{5,6}, C. Drew Harvell³

¹Centre for Veterinary and Epidemiological Research, Department of Health Management, Atlantic Veterinary College, University of Prince Edward Island, Charlottetown, PE C1A 4P3, Canada

²Institute of Marine and Environmental Technology, University of Maryland Baltimore County, Baltimore, MD 21202, USA

³Department of Ecology & Evolutionary Biology, Cornell University, Ithaca, NY 14850, USA

⁴Shannon Point Marine Center, Western Washington University, Anacortes, WA 98221, USA

⁵Friday Harbor Laboratories, University of Washington, Friday Harbor, WA 98250, USA

⁶Center for Marine and Environmental Studies, University of the Virgin Islands, St. Thomas, VI, USA

⁷Present address: Dept. of Fisheries, Virginia Institute of Marine Science, 1208 Greate Road, Gloucester Point, VA 23062, USA

ABSTRACT: Seagrasses are ecosystem engineers of essential marine habitat. Their populations are rapidly declining worldwide. One potential cause of seagrass population declines is wasting disease, which is caused by opportunistic pathogens in the genus *Labyrinthula*. While infection with these pathogens is common in seagrasses, theory suggests that disease only occurs when environmental stressors cause immunosuppression of the host. Recent evidence suggests that host factors may also contribute to disease caused by opportunistic pathogens. In order to quantify patterns of disease, identify risk factors, and investigate responses to infection, we surveyed shoot density, shoot length, epiphyte load, production of plant defenses (phenols), and wasting disease prevalence in eelgrass *Zostera marina* across 11 sites in the central Salish Sea (Washington state, USA), a region where both wasting disease and eelgrass declines have been documented. Wasting disease was diagnosed by the presence of necrotic lesions, and *Labyrinthula* cells were identified with histology. Disease prevalence among sites varied from 6 to 79%. The probability of a shoot being diseased was higher in longer shoots, in patches of higher shoot density, and in shoots with higher levels of biofouling from epiphytes. Phenolic concentration was higher in diseased leaves. We hypothesize that this results from the induction of phenols during infection. Additional research is needed to evaluate whether phenols are an adaptive defense against *Labyrinthula* infection. The high site-level variation in disease prevalence emphasizes the potential for wasting disease to be causing some of the observed decline in eelgrass beds.

KEY WORDS: Opportunistic pathogens · Seagrass declines · Phenols · Plant defenses · *Labyrinthula* · Coastal ecosystems · Restoration · *Zostera marina*

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Recent data suggest that infectious diseases are increasing in both incidence and severity in the ocean (Harvell et al. 2004, Ward & Lafferty 2004). The effects of marine diseases can be especially pro-

nounced when these diseases negatively impact ecosystem engineers such as reef-building corals (Aronson & Precht 2001), sea urchins (Feehan & Scheibling 2014), oysters (Mann et al. 2009), abalone (reviewed by Burge et al. 2014), and seagrasses (Cottam 1933, Short et al. 1987). Ecosystem engineers provide habi-

tat and ecosystem services, and understanding the factors that lead to increased prevalence and severity of disease in these species is critical for conservation of marine biodiversity and ecosystem function in the face of our changing oceans.

Wasting disease is one of many causes of global seagrass declines and threatens the numerous ecosystem functions provided by seagrass (Orth et al. 2006, Sullivan et al. 2013). Caused by intracellular infection of the host mesophyll by opportunistic protists of the genus *Labyrinthula*, wasting disease affects several species of seagrass (Muehlstein et al. 1988, 1991, Sullivan et al. 2013). Disease outbreaks have the potential to cause rapid, dramatic population declines, which can compromise the ecosystem services provided by these plants. In the early 1930s, an outbreak along the Atlantic coasts of North America and Europe resulted in up to 90% mortality of eelgrass *Zostera marina* populations (Cottam 1933, Short et al. 1987). Smaller, more isolated epidemics have been documented in *Z. marina* populations along the Atlantic and Pacific coasts of North America since the 1980s (Short et al. 1987, Groner et al. 2014). However, aside from extreme outbreaks, the impact of chronic infections on seagrass declines is relatively unknown. Loss of seagrass habitats compromises habitat for economically valuable invertebrates, fish, and birds, and affects the ecosystem services provided by these systems, including the stabilization of coastal sediments, biological filtration of terrestrial-derived nutrients, carbon sequestration, and local buffering of ocean pH (Barbier et al. 2011, Fourqurean et al. 2012, Plummer et al. 2013).

Targeted monitoring of diseases and their associated risk factors is critical for early detection and mitigation of marine diseases (e.g. Groner et al. 2015). For many opportunistic pathogens, environmental risk factors cause disease by compromising physiological host defenses (Burge et al. 2013). Recent studies have identified salinity, temperature, and turbidity as factors contributing to wasting disease (McKone & Tanner 2009, Bull et al. 2012, Groner et al. 2014). A potentially overlooked facilitator of disease from opportunistic pathogens is host population structure. Defenses against pathogens often change during development and may be compromised by stressors such as density, reduced light penetration, for example as caused by growth of epiphytes on leaves, or an individual's microbial community (Kazan & Manners 2009). Indeed, recent studies have shown that leaf age and plant size are important determinants of disease in eelgrass (Groner et al. 2014).

Defenses of eelgrass against *Labyrinthula* are not well understood; however, phenolic compounds may play a role (Buchsbbaum et al. 1990, Vergeer et al. 1995, Vergeer & Develi 1997, Sneed 2005, Trevathan-Tackett et al. 2015). Phenolic compounds produced by *Z. marina* have been correlated with resistance to wasting disease (Buchsbbaum et al. 1990) and are capable of inhibiting growth of *Labyrinthula in vitro* (Vergeer et al. 1995). However, it is not known whether phenols produced *in vivo* are capable of inhibiting infections.

Since the early 1990s, a number of eelgrass beds in the Salish Sea, located in the northeast Pacific Ocean, have experienced periodic mortalities and are declining from unknown causes (Wyllie-Echeverria et al. 2003, 2010, Dowty et al. 2010, Washington State Department of Natural Resources 2015). Hypothesized causes for these declines include wasting disease and sulfide toxicity (Dooley et al. 2013, Groner et al. 2014). Wasting disease has long been observed in this system (Muehlstein et al. 1991, Muehlstein 1992), and pathogenic strains of *L. zosterae* have been isolated from diseased leaves (Groner et al. 2014).

In this study, our primary objectives were to identify eelgrass beds with high disease prevalence and to quantify risk factors for disease that could occur at the host level. We surveyed 11 sites throughout the San Juan Archipelago and Puget Sound (i.e. the central Salish Sea). We focused on the role of 3 factors in altering disease prevalence: shoot length, shoot density, and biofouling of leaves by epiphytes. We hypothesized that increases in each of these factors would lead to increased disease prevalence. A second objective of our study was to quantify the relationship between phenolic compounds and disease. We hypothesized that, across sites, diseased leaves would have greater phenolic concentration than healthy leaves.

MATERIALS AND METHODS

Field survey

We sampled 11 sites in the San Juan Archipelago and Northern Puget Sound regions of the Salish Sea in Washington state, USA, between 19 and 24 July 2013 (Fig. 1). We focused on this region because declines in the distribution and density of *Zostera marina* were documented at several sites in the early 2000s (Wyllie-Echeverria et al. 2003, 2010, Ferrier & Berry 2010). Some of these sites show evidence of population decline, while others have remained stable (Fig. 1, Table 1).




Fig. 1. Prevalence of eelgrass wasting disease (as analyzed on the second longest leaf in surveyed shoots of *Zostera marina*) among study sites: (1) Beach Haven, (2) False Bay, (3) Fisherman Bay, (4) Indian Cove, (5) Mosquito Bay, (6) North Cove, (7) Padilla Bay, (8) Picnic Cove, (9) Shallow Bay, (10) Ship Harbor, (11) Shoal Bay

At each site, data were collected along 3 transects (each 50 m long) laid 2 m apart and parallel to the shore in the intertidal region, with the middle transect at approximately -1 m Mean Lower Low Water. Each 50 m transect was divided into 10 m sections ($n = 5$), and in each section, the second oldest leaf was collected from 10 randomly selected shoots. We surveyed the second oldest intact leaf in a shoot to standardize our measurements. We picked this leaf to be conservative, since the older leaves were slightly senescent and much more likely to have lesions, while the youngest leaves may not have been present long enough to present lesions even if the shoot was diseased. In order to determine site-level differences in population density, we also counted the number of shoots in 3 quadrats (0.12 m^2 each) per

transect (at 0, 25, and 50 m). The mean density for each transect was used in the analyses.

Collected leaves were placed in coolers and taken to the Seagrass Lab, Friday Harbor Laboratories, WA, USA, for further measurement. For each leaf, we measured length and width, and described disease status (healthy or diseased) based on visual characteristics. Leaves were diagnosed with wasting disease if they had lesions with irregular, dark, necrotic centers surrounded by a black border (e.g. Burdick et al. 1993, Groner et al. 2014). For 2 of the leaves collected in each 10 m section, we measured the dry weight of all attached epiphytes. We scraped off all epiphytes from both sides of the leaves onto weigh paper and dried these samples in a drying oven until the mass remained constant (minimum of 24 h). We

Table 1. Survey site locations and known information on population trends and summaries (means \pm SD) of eelgrass *Zostera marina* shoot density and length, epiphyte biomass, and wasting disease prevalence at each site. ID numbers as in Fig. 1; (–) sites where population status has not been assessed

ID	Site	Latitude (°N)	Longitude (°W)	Population trend(s)	Shoot density (m ⁻²)	Shoot length (cm)	Epiphytes (mg cm ⁻² of shoot)	Prevalence (%)
1	Beach Haven	48° 41.460'	122° 57.120'	Stable ^a	297 \pm 129	84.6 \pm 22.9	1.17 \pm 1.12	6 \pm 4
2	False Bay	48° 28.975'	123° 04.451'	Declining ^{a,b}	136 \pm 42	57.4 \pm 15.0	3.37 \pm 1.85	47 \pm 17
3	Fisherman Bay	48° 31.572'	122° 55.088'	Stable ^a	213 \pm 29	59.6 \pm 15.8	0.39 \pm 0.46	54 \pm 4
4	Indian Cove	48° 33.773'	122° 56.078'	Stable ^a	336 \pm 106	81.0 \pm 17.8	0.52 \pm 0.28	65 \pm 12
5	Mosquito Pass	48° 35.346'	123° 10.208'	Declining ^a	214 \pm 4	84.9 \pm 19.3	3.70 \pm 2.19	57 \pm 2
6	North Cove	48° 42.287'	123° 03.221'	Declining ^b	542 \pm 31	63.8 \pm 12.0	0.70 \pm 0.49	22 \pm 8
7	Padilla Bay	48° 29.690'	122° 29.244'	–	358 \pm 13	58.7 \pm 10.7	1.10 \pm 1.49	79 \pm 13
8	Picnic Cove	48° 33.942'	122° 55.448'	Declining ^a , Stable ^b	75 \pm 43	66.9 \pm 13.7	0.61 \pm 0.72	44 \pm 2
9	Shallow Bay	48° 45.613'	122° 54.892'	Declining ^a	19 \pm 19	75.4 \pm 17.1	1.46 \pm 1.16	9 \pm 15
10	Ship Harbor	48° 30.307'	122° 40.202'	–	158 \pm 102	73.4 \pm 16.1	4.91 \pm 2.64	43 \pm 2
11	Shoal Bay	48° 33.215'	122° 52.499'	–	164 \pm 4	70.3 \pm 11.9	1.49 \pm 3.38	42 \pm 23

^aPopulation status estimated from shoot density data collected by Wyllie-Echeverria et al. (2010, or unpubl.)
^bPopulation status estimated from data collected by Washington State Department of Natural Resources (2015)

then divided the dry weight of the epiphytes by the surface area of the leaf they were collected from to calculate the surface area-standardized epiphyte load. In order to understand the relationship between disease and phenolic content in these populations, we quantified phenolic concentration in 3 visually healthy and 3 visually diseased leaves that were collected along each transect. The processing of leaves described above typically occurred within 12 h of collection. Plants were kept in bags on ice to prevent further development of disease between collection and processing.

Phenolic measurements

Eelgrass samples frozen at -80°C were transported to the Shannon Point Marine Center in Anacortes, WA, where they were lyophilized and ground to a fine powder with a SPEX mixer/mill. Samples of approximately 10 mg were weighed and extracted in 80% methanol overnight in darkness at -80°C . The extracts were spun down with a centrifuge, and a 50 μl aliquot of each extract was diluted with 950 μl of ANSI Type I water. Three replicate 100 μl aliquots of each diluted sample were dispensed into the cells of a 96-well microplate. Forty μl of 40% Folin & Ciocalteu's phenol reagent (Sigma F9252) were then dispensed into each cell of the microplate. The plates were incubated at 50°C with shaking for 5 min, then 100 μl of 2 N sodium carbonate were added to each well. The absorbance at 765 nm was read after

30 min of incubation. Caffeic acid was initially used as a standard for these assays to generate phenolic measurements that were caffeic acid equivalents, because a native standard was not available at the time the analyses were run. A native standard was produced at a later time and was standardized to caffeic acid (see Fig. S1 in Supplement 1, available at www.int-res.com/articles/suppl/d118p159_supp/). Because of the strong linear relationship ($R^2 = 0.994$, $p < 0.001$) between absorbances obtained with the native standard and absorbances obtained with caffeic acid, we then used this regression to convert the concentrations of phenolic compounds obtained using caffeic acid to concentrations of phenolic compounds based on the native standard. Although the native standard was likely to contain some non-phenolic metabolites, it was more representative of the phenolic compounds in *Z. marina* than the caffeic acid standard.

To obtain the native standard, 10 g of lyophilized, ground *Z. marina* collected from Ship Harbor in July 2014 were extracted once in 1 l of 70% acetone containing 5.5 mM ascorbic acid, and twice in 1 l of 80% methanol containing 5.5 mM ascorbic acid. The extracts were combined and the solvents were evaporated under vacuum with a rotary evaporator until only 200 ml of aqueous extract remained. The extract was partitioned 3 times with hexanes to remove non-polar metabolites. The remaining aqueous extract was mixed with 50 ml of Sephadex LH-20 (Sigma GE17-0090) and placed into a Kontes 4.8 cm diameter chromatography column. Non-phenolic metabolites were removed by washing with 3 column vol-

umes of water. The phenolic compounds were then eluted with 3 column volumes of methanol. Water (100 ml) was added to the methanol fraction, and the methanol was evaporated under vacuum. The remaining aqueous extract was partitioned with 1:1 ethyl acetate:hexanes to remove any remaining non-polar metabolites. Any remaining solvents in the aqueous fraction were then evaporated under vacuum. The fraction was frozen to -80°C and lyophilized to produce brownish-yellow colored flakes.

Histology

We subsampled leaves collected in Indian Cove and False Bay (Fig. 1) to qualitatively confirm visual disease diagnostics with histological analysis. Six healthy and 12 diseased leaves from Indian Cove and 10 healthy and 10 diseased leaves from False Bay were used for this analysis. Assessments of 'healthy' or 'diseased' were based on visual characteristics as described above. For each diseased leaf, we sampled an area including the leading edge of the lesion and adjacent, visually healthy, tissue. Samples were fixed for 24 h in 4% seawater-buffered formalin, and then stored in 70% ethanol. At Cornell University's Diagnostic Laboratory, leaves were embedded in paraffin, sectioned ($5\ \mu\text{m}$), and stained with hematoxylin and eosin (e.g. Groner et al. 2014). Stained sections were randomized and viewed under a bright field Olympus BH-2 microscope with the Olympus DP-20 camera system. Sections were examined blindly (without demarcation of site or disease status) by a single viewer for the presence/absence of *Labyrinthula* spp. cells or other parasites within the tissues. The number of *Labyrinthula* spp. cells was counted at $40\times$ magnification for three $0.5\ \text{mm}$ fields of view on each slide. *Labyrinthula* spp. cells stained purple in contrast to pink plant tissue and were located intracellularly within the mesophyll (e.g. Renn 1936, Porter 1972, Groner et al. 2014). Active *Labyrinthula* spp. cells could be identified by their spindle shape ($\sim 15\ \mu\text{m}$ length), and resting *Labyrinthula* spp. cells could be identified by their slightly smaller round shape. In some cases, extracellular mucus was also observed.

Statistical analyses

We used mixed effects logistic regression to understand the additive effects of shoot density, shoot-standardized epiphyte biomass, and shoot length on the disease status of each individual. We ran all pos-

sible additive combinations of these predictors in separate models and evaluated their fit using Akaike's information criterion (AIC). We included site as a random intercept in all models.

We used a linear mixed effect model to quantify the effects of shoot length and disease status (e.g. diseased or healthy) on the phenolic concentration (% of dry biomass) of leaves. We ran models with each of these predictor variables separately, both together additively, and both together with an interaction. We then used model selection, minimizing AIC values, to select the best fit model. We included site as a random intercept in all models. The model met the assumptions of normality and homoscedasticity.

All analyses were run in R (R Core Team 2014, v. 3.1.2) using the 'lme4' package (Bates et al. 2014). R-scripts and datasets for these analyses are available in Supplements 2 and 3 at www.int-res.com/articles/suppl/d118p159_supp/, respectively.

RESULTS

Field data

In total, 1645 samples were collected from the 11 sites. All sites were diseased; prevalence among sites varied from 6% at Beach Haven to 79% at Padilla Bay (Fig. 1). The geometric mean of disease prevalence across sites was 34%.

The best fitting model included shoot length, shoot density, and epiphyte biomass. The next best fitting model included only shoot length and epiphyte biomass (ΔAIC from best model = 1.3). No other models were strong candidates (ΔAIC from next best model = 6.4). Longer shoots, high shoot density, and high epiphyte biomass on shoots increased the probability of disease (Fig. 2, Table 1). For every 1 cm of shoot length, the odds of being diseased increased by 1.8% ($z = -4.61$, $p < 0.00001$). For every additional shoot in a $100\ \text{cm}^2$ area, the odds of being diseased increased by 19.9% ($z = 1.84$, $p = 0.067$). For every additional 1 mg of epiphytes per cm^2 of shoot, odds of disease increased by 21.9% ($z = 2.88$, $p = 0.003$).

Phenolics

We processed 175 samples for phenolic concentrations. The best model included disease status (healthy or diseased) as a fixed effect and site as a random intercept (ΔAIC from next best model = 7.5). When diseased and healthy shoots were combined,




Fig. 2. Effects of eelgrass *Zostera marina* shoot length, shoot density, and biofouling by epiphytes on the predicted probability that a leaf will have wasting disease. Predicted probabilities were calculated using a logistic regression of shoot length, shoot density, and biofouling by epiphytes on the disease status of a shoot. Vertical lines indicate 25 and 75 % quantiles. For each simulation, 2 variables were held constant, while the variable of interest was allowed to vary. Constants were: shoot length = 70 mm, shoot density = 200 shoots m^{-2} , epiphyte biomass = 120 mg epiphytes per 1 mm shoot

the phenolic concentration (mean \pm SD) varied across sites from $4.9 \pm 0.5\%$ of biomass in False Bay to $14.7 \pm 0.9\%$ of biomass in Padilla Bay. Across sites, the mean phenolic concentration was $7.5 \pm 0.6\%$ of biomass in healthy leaves and $8.5 \pm 0.8\%$ of biomass in diseased leaves. After accounting for random site effects, diseased leaves had an increase in phenolic concentration of 0.76 (in units of % of biomass, $t = 2.33$, $p = 0.021$; Fig. 3). This corresponds to a 10% increase in phenolic concentration in diseased plants relative to plants with no lesions. Shoot length did not improve the model fit and was excluded from the final model.

Histology

Labyrinthula sp. was observed in samples of both healthy and diseased tissue at all sites sampled. Cell counts per field of view were (mean \pm SD): healthy: 7.83 ± 2.16 (Indian Cove) and 5.1 ± 2.8 (False Bay); and diseased: 4.54 ± 2.30 (Indian Cove) and 14.7 ± 3.29 (False Bay). Due to the qualitative nature of histological observations, these data were not analyzed statistically. No other potential infectious agents were noted.

DISCUSSION

Globally, seagrasses are declining at a rate of $7\% \text{ yr}^{-1}$, and identification of factors contributing to these declines is critical for conservation of seagrasses, the habitats that they create, and the ecosystems services that they provide (Waycott et al. 2009). While in a few instances, recent seagrass declines have been associated with disease (Short & Wyllie-Echeverria 1996), the relation between population decline and disease is more frequently unknown. The prevalence of eelgrass wasting disease varied widely across our study region, from 6 to 79%. This result is similar to findings in the Wadden Sea, where disease prevalence varied from 11 to 80%, (Bockelmann et al. 2013). Our study indicates that eelgrass population structure (i.e. shoot length and density) and fouling are correlated

with wasting disease in the Salish Sea. This emphasizes the potential for wasting disease to be contributing to observed population declines. Consistent with previous studies (Buchsbaum et al. 1990, Vergeer et al. 1995, Vergeer & Develi 1997), we show a potential role for phenols as a defense against disease.

Shoot density and shoot length were positive predictors of disease. Correlations between wasting disease and both shoot density and shoot length have been found in previous field studies. For example, disease was positively correlated with density in European populations of eelgrass (Bull et al. 2012),




Fig. 3. Phenolic contents (as percent of dry mass equivalents) of healthy (light bars) and diseased (dark bars) eelgrass *Zostera marina* leaves from 11 sites. The data are means \pm 1 SD. The standard curves used to calibrate the analyses were derived from absorbances obtained from Folin-Ciocalteu analyses using known amounts of extracted phenolic compounds from *Z. marina* collected from Ship Harbor (see Fig. S1 in Supplement 1 at www.int-res.com/articles/suppl/d118p159_supp/)

and a field survey of 3 sites in the San Juan Archipelago found that longer shoots had higher levels of infection (Groner et al. 2014). Experimental infections also were more successful on older shoots compared to younger shoots on a single ramet (Groner et al. 2014). While trade-offs between shoot length and shoot density are common (Olesen & Sand-Jensen 1994a, Yang et al. 2013), our model suggests that disease risk may be higher in areas with increased biomass (e.g. beds with short shoots or low density sites with long shoots). It is possible that such populations are at or near their carrying capacity and are experiencing negative density-dependent growth rates as a result of competition or have attained high enough biomass to allow for spread of disease.

The contribution of epiphytes to wasting disease has not been observed previously. One hypothesis for this effect is that the epiphytes cause a reduction in immune defenses by blocking carbon uptake and light, thereby reducing photosynthesis and allocation of resources towards immune defenses (Sand-Jensen 1977). Another hypothesis is that increased nutrients drive the changes in epiphytes, wasting disease, and phenolics. Increased biofouling by epiphytes is frequently indicative of increased nutrient loading (Borum 1985, Williams & Ruckelshaus 1993), and higher nitrogen levels have been associated with reduced production of phenols and increased wasting disease in experimental treatments (Buchsbaum et al. 1990). Further research is needed to investigate the mechanism behind this relationship.

While disease prevalence was high at several sites, it is unclear what role this may play at the population level. We hypothesize that wasting disease has 2 important effects on eelgrass beds: (1) to lower overall plant fitness, potentially contributing to population declines; and (2) to increase the decay rate of senescing leaves, remineralizing nutrients for availability in the food web (Raghukumar 2002, Raghukumar & Damare 2011). During warm seasons, leaves are turned over rapidly in eelgrass shoots, and it is likely that the sampled leaves were weeks away from senescing (Olesen & Sand-Jensen 1994b). Nonetheless, fitness costs may be incurred from wasting disease at high-prevalence sites in the Salish Sea. While wasting disease has previously caused rapid population declines (Atlantic coastlines in the

1930s), more work will be needed to pinpoint how much wasting is contributing to the current population declines in the Salish Sea. The high prevalence of wasting disease and extent of recent eelgrass population declines warrants further investigation into these topics.

Our study supports previous field observations of increased phenolic content in diseased seagrasses (Vergeer & Develi 1997). This positive association is consistent with some previous laboratory and mesocosm studies (e.g. Buchsbaum et al. 1990, Vergeer et al. 1995, Steele et al. 2005), but not others (McKone & Tanner 2009). Indeed, the wide variation in the association of disease status on phenolic content found across the 11 sites suggests that, in addition to disease, other factors such as nutrient availability, herbivory, and genetics may be influencing the production of phenols.

Phenolic compounds in seagrasses, including *Zostera marina*, are a complex mixture of metabolites that include condensed tannins (proanthocyanidin polymers), phenolic acids, flavonoids, and lignins (Van Alstyne & Padilla in press). In the Salish Sea, the phenolic acids produced by *Z. marina* include ferulic (4-hydroxy-3-methoxycinnamic), vanillic (4-hydroxy-3-methoxybenzoic), p-hydroxybenzoic (4-hydroxybenzoic), caffeic (3,4-dihydroxycinnamic), gallic (3,4,5-trihydroxybenzoic), protocatechuic (3,4-dihydroxybenzoic), and gentisic (2,5-dihydroxybenzoic) acid (Quackenbush et al. 1986, Ferrat et al. 2012). No single assay can measure the concentra-

tions of these structurally diverse compounds. Folin assays, such as the Folin-Ciocalteu assay that was used in this study, provide a measurement of the reducing activity of extracts containing phenolic compounds. This can be used as a proxy for the concentration of the compounds in the plants. The assay measures the ability of hydroxylated aromatic compounds in the extracts to reduce phosphomolybdic and phosphotungstic acid reagents, and reducing ability is thought to be correlated with the biological activity of the compounds (Appel et al. 2001). Because the ability of phenolic compounds to reduce these reagents is dependent on the compounds' structures, changes in 'phenolic contents' of plants, as measured with Folin assays, could result from changes in the amounts of compounds present in the extracts or in their structural types. These assays provide a relatively rapid and inexpensive 'first look' at changes in plant biochemistry that correlate with differences in the plants' environment and disease state, but to understand the specific mechanisms by which plants are responding to pathogens and environmental factors, more detailed analyses of specific compounds (e.g. phenolic acids) or more closely related groups of compounds (e.g. condensed tannins) are needed.

Phenols are associated with disease resistance in many plants (Nicholson & Hammerschmidt 1992); however, it is unclear whether the association found here is indicative of an adaptive response to infection. Increased production of caffeic acid in response to *Labyrinthula* spp. infection has been found in *Z. marina*, and caffeic acid can inhibit growth of *Labyrinthula* spp. *in vitro* (Buchsbaum et al. 1990, Vergeer et al. 1995, Vergeer & Develi 1997). In contrast, a potentially non-adaptive pseudo-induction of phenolics in another seagrass species, turtlegrass *Thalassia testudinum*, is hypothesized to occur when shoots cannot move photosynthates down the leaf past the site of infection (Steele & Valentine 2012, Trevathan-Tackett et al. 2015). This causes carbohydrates to accumulate above the sites of the *Labyrinthula* spp. infections, where they are then used to synthesize phenolic compounds (Steele et al. 2005). Finally, *Labyrinthula* may inhibit production of phenols that are capable of inhibiting growth of *in vitro* strains (e.g. Sneed 2005). Numerous other factors influence phenolic concentrations in plants. Phenol concentration decreases with blade age (Ravn et al. 1994) and is negatively correlated with environmental factors, including turbidity (Sneed 2005), temperature (Vergeer et al. 1995), ocean acidification (Arnold et al. 2012), low salinity (Sneed 2005), and

heavy metal contamination (Ferrat et al. 2012). Mechanistic studies that quantify the locations and types of phenols produced over the course of an infection, and how those phenols affect the growth of *Labyrinthula* under various environmental conditions, are necessary to characterize the roles of these compounds.

The presence of *Labyrinthula* spp. cells in both diseased and healthy tissue is consistent with opportunistic pathogens, whereby the presence of pathogens is not necessarily indicative of disease (reviewed by Burge et al. 2013). The wide range of cell densities in the histology samples could reflect a heterogeneous distribution of *Labyrinthula* spp. in the hosts. Indeed, this is the case with sea fan corals infected with the labyrinthulomycete *Aplanochytrium* (Burge et al. 2012), and indicates that while histology can be an effective qualitative indicator of pathogen presence and infection of host tissues, it is not an accurate method to measure pathogen load, disease state, or strain. This highlights the importance of quantitative PCR methods for quantifying pathogen load in *L. zosterae* infections. Paired measurements of pathogen load and wasting disease status are necessary to understand whether a specific pathogen load and/or strain is associated with disease status.

We have highlighted the influence of host factors on eelgrass wasting disease prevalence across sites in the central region of the Salish Sea. Many questions remain about what other biotic or abiotic factors contribute to or are necessary for disease expression in Pacific Northwest eelgrass meadows. Other potential risk factors have not been fully evaluated in Pacific eelgrass meadows, including temperature, light, salinity, nutrients, and hydrogen sulfide toxicity (Holmer & Bondgaard 2001, Dooley et al. 2013, Yang et al. 2013, Kaldy 2014). Moreover, while our survey provides data on wasting disease prevalence across 11 sites, it is a single snapshot in time. Longitudinal studies to track disease progression and eelgrass population status over seasonal and multi-year fluctuations are critical for clarifying the relationship between disease and declining eelgrass meadows. Finally, the role of strain variation in the incidence and severity of wasting disease is not well understood, although preliminary evidence suggests that this variation may be substantial (Groner et al. 2014). Nonetheless, our study highlights the potential for wasting disease to cause eelgrass population declines in the Salish Sea and provides impetus and direction for further research. More broadly, the important role of host demography in the epidemiology of eelgrass wasting disease suggests that demo-

graphy may be an important area for research in other marine or terrestrial organisms that are infected with pathogenic labyrinthulomycetes.

Acknowledgements. Aissa Yazzie, Holly Moxley-Williams, Diane Ayala, Anna Russell, Nathan Greene, Katherine Boehrer, and Katie Kirk assisted in the field and the laboratory. Thanks also to William Chalmers and 3 anonymous reviewers for insightful comments. This work was funded by the National Science Foundation Ecology and Evolution of Infectious Diseases grant number OCE-1215977 awarded to C.D.H., NSF IOS 1118944 to K.L.V.A., and a Patricia L. Dudley Scholarship awarded to M.L.G. M.L.G. also acknowledges partial support from the Centre for Excellence Research Chair in Aquatic Epidemiology at the Atlantic Veterinary College in Prince Edward Island.

LITERATURE CITED

- Appel HM, Govonor HL, D'Ascenzo M, Siska E, Schultz JC (2001) Limitations of Folin assays of foliar phenolics in ecological studies. *J Chem Ecol* 27:761–778
- Arnold T, Mealey C, Leahey H, Miller AW, Hall-Spencer JM, Milazzo M, Maers K (2012) Ocean acidification and the loss of phenolic substances in marine plants. *PLoS ONE* 7:e35107
- Aronson RB, Precht WF (2001) White-band disease and the changing face of Caribbean coral reefs. In: Porter JW (ed) *The ecology and etiology of newly emerging marine diseases*. Springer, Dordrecht, p 25–38
- Barbier EB, Hacker SD, Kennedy C, Koch EW, Stier AC, Silliman BR (2011) The value of estuarine and coastal ecosystem services. *Ecol Monogr* 81:169–193
- Bates D, Maechler M, Bolker B, Walker S (2014) lme4: Linear mixed-effects models using Eigen and S4. *J Stat Softw* 67:1–48
- Bockelmann AC, Tams V, Ploog J, Schubert PR, Reusch TB (2013) Quantitative PCR reveals strong spatial and temporal variation of the wasting disease pathogen, *Labyrinthula zosterae* in northern European eelgrass (*Zostera marina*) beds. *PLoS ONE* 8:e62169
- Borum J (1985) Development of epiphytic communities on eelgrass (*Zostera marina*) along a nutrient gradient in a Danish estuary. *Mar Biol* 87:211–218
- Buchsbaum RN, Short FT, Cheney DP (1990) Phenolic-nitrogen interactions in eelgrass, *Zostera marina* L.: possible implications for disease resistance. *Aquat Bot* 37:291–297
- Bull JC, Kenyon EJ, Cook KJ (2012) Wasting disease regulates long-term population dynamics in a threatened seagrass. *Oecologia* 169:135–142
- Burdick DM, Short FT, Wolf J (1993) An index to assess and monitor the progression of wasting disease in eelgrass *Zostera marina*. *Mar Ecol Prog Ser* 94:83–90
- Burge CA, Douglas N, Conti-Jerpe, I, Weil E, Roberts S, Friedman CS, Harvell CD (2012) Friend or foe: the association of Labyrinthulomycetes with the Caribbean sea fan *Gorgonia ventalina*. *Dis Aquat Org* 101:1–12
- Burge CA, Kim CJS, Lyles JM, Harvell CD (2013) Special issue oceans and humans health: the ecology of marine opportunists. *Microb Ecol* 65:869–879
- Burge CA, Eakin CM, Friedman CS, Froelich B and others (2014) Climate change influences on marine infectious diseases: implications for management and society. *Annu Rev Mar Sci* 6:249–277
- Cottam CE (1933) Disappearance of eelgrass along the Atlantic coast. *Plant Dis Rep* 17:46–53
- Dooley FS, Wyllie-Echeverria S, Ward PD, Roth M (2013) Tolerance and response of *Zostera marina* seedlings to hydrogen sulfide. *Aquat Bot* 105:7–10
- Dowty P, Berry H, Gaeckle J (2010) Developing indicators and targets for eelgrass in Puget Sound. Nearshore Habitat Program, Washington State Department of Natural Resources, Olympia, WA
- Feehan CJ, Scheibling RE (2014) Effects of sea urchin disease on coastal marine ecosystems. *Mar Biol* 161:1467–1485
- Ferrat L, Wyllie-Echeverria S, Cates RG, Pergent-Martini C and others (2012) *Posidonia oceanica* and *Zostera marina* as potential biomarkers of heavy metal contamination in estuarine and coastal systems. In: Voudouris (ed) *Ecological water quality—water treatment and reuse*. Intech, Rijeka, p 123–140
- Ferrier L, Berry H (2010) Eelgrass (*Zostera marina* L.) abundance and depth distribution along selected San Juan Archipelago shallow embayments. Nearshore Habitat Program, Washington State Department of Natural Resources, Olympia, WA
- Fourqurean JW, Duarte CM, Kennedy H, Marbà N and others (2012) Seagrass ecosystems as a globally significant carbon stock. *Nat Geosci* 5:505–509
- Groner ML, Burge CA, Couch CS, Kim CJS and others (2014) Host demography influences the prevalence and severity of eelgrass wasting disease. *Dis Aquat Org* 108:165–175
- Groner M, Breyta R, Dobson A, Friedman CS and others (2015) Emergency response for marine diseases. *Science* 347:1210
- Harvell D, Aronson R, Baron N, Connell J and others (2004) The rising tide of ocean diseases: unsolved problems and research priorities. *Front Ecol Environ* 2:375–382
- Holmer M, Bondgaard EJ (2001) Photosynthetic and growth response of eelgrass to low oxygen and high sulfide concentrations during hypoxic events. *Aquat Bot* 70:29–38
- Kaldy JE (2014) Effect of temperature and nutrient manipulations on eelgrass *Zostera marina* L. from the Pacific Northwest, USA. *J Exp Mar Biol Ecol* 453:108–115
- Kazan K, Manners JM (2009) Linking development to defense: auxin in plant–pathogen interactions. *Trends Plant Sci* 14:373–382
- Mann R, Southworth M, Harding JM, Wesson JA (2009) Population studies of the native eastern oyster, *Crassostrea virginica*, (Gmelin, 1791) in the James River, Virginia, USA. *J Shellfish Res* 28:193–220
- McKone KL, Tanner CE (2009) Role of salinity in the susceptibility of eelgrass *Zostera marina* to the wasting disease pathogen *Labyrinthula zosterae*. *Mar Ecol Prog Ser* 377:123–130
- Muehlstein LK (1992) The host-pathogen interaction in the wasting disease of eelgrass, *Zostera marina*. *Can J Bot* 70:2081–2088
- Muehlstein LK, Porter D, Short FT (1988) *Labyrinthula* sp., a marine slime mold producing the symptoms of wasting disease in eelgrass, *Zostera marina*. *Mar Biol* 99:465–472
- Muehlstein LK, Porter D, Short FT (1991) *Labyrinthula zosterae* sp. nov., the causative agent of wasting disease of eelgrass, *Zostera marina*. *Mycologia* 83:180–191

- Nicholson RL, Hammerschmidt R (1992) Phenolic compounds and their role in disease resistance. *Annu Rev Phytopathol* 30:369–389
- Olesen B, Sand-Jensen K (1994a) Biomass-density patterns in the temperate seagrass *Zostera marina*. *Mar Ecol Prog Ser* 109:283–291
- Olesen B, Sand-Jensen K (1994b) Demography of shallow eelgrass (*Zostera marina*) populations—shoot dynamics and biomass development. *J Ecol* 82:379–390
- Orth RJ, Carruthers TJ, Dennison WC, Duarte CM and others (2006) A global crisis for seagrass ecosystems. *BioScience* 56:987–996
- Plummer ML, Harvey CJ, Anderson LE, Guerry AD, Ruckelshaus MH (2013) The role of eelgrass in marine community interactions and ecosystem services: results from ecosystem-scale food web models. *Ecosystems* 16: 237–251
- Porter D (1972) Cell division in the marine slime mold, *Labyrinthula* sp., and the role of the bothrosome in extracellular membrane production. *Protoplasma* 74:427–448
- Quackenbush RC, Bunn D, Lingren W (1986) HPLC determination of phenolic acids in the water-soluble extract of *Zostera marina* L. (eelgrass). *Aquat Bot* 24:83–89
- R Core Team (2014) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna. www.r-project.org
- Raghukumar S (2002) Ecology of the marine protists, the Labyrinthulomycetes (Thraustochytrids and Labyrinthulids). *Eur J Protistol* 38:127–145
- Raghukumar S, Damare VS (2011) Increasing evidence for the important role of Labyrinthulomycetes in marine ecosystems. *Bot Mar* 54:3–11
- Ravn H, Pedersen MF, Borum J, Andary C, Anthoni U, Christophersen C, Nielsen PH (1994) Seasonal variation and distribution of two phenolic compounds, rosmarinic acid and caffeic acid, in leaves and roots-rhizomes of eelgrass (*Zostera marina* L.). *Ophelia* 40:51–61
- Renn CE (1936) The wasting disease of *Zostera marina*. I. A phytological investigation of the diseased plant. *Biol Bull (Woods Hole)* 70:148–158
- Sand-Jensen K (1977) Effect of epiphytes on eelgrass photosynthesis. *Aquat Bot* 3:55–63
- Short FT, Wyllie-Echeverria S (1996) Natural and human-induced disturbance of seagrasses. *Environ Conserv* 23: 17–27
- Short FT, Muehlstein LK, Porter D (1987) Eelgrass wasting disease: cause and recurrence of a marine epidemic. *Biol Bull (Woods Hole)* 173:557–562
- Sneed JM (2005) The effects of *Labyrinthula* sp. infection, salinity, and light on the production of phenolic compounds in *Thalassia testudinum*. MSc thesis, University of South Florida, Tampa, FL
- Steele L, Valentine JF (2012) Idiosyncratic responses of seagrass phenolic production following sea urchin grazing. *Mar Ecol Prog Ser* 466:81–92
- Steele L, Caldwell M, Boettcher A, Arnold T (2005) Seagrass-pathogen interactions: ‘pseudo-induction’ of turtlegrass phenolics near wasting disease lesions. *Mar Ecol Prog Ser* 303:123–131
- Sullivan BK, Sherman TD, Damare VS, Lilje O, Gleason FH (2013) Potential roles of *Labyrinthula* spp. in global seagrass population declines. *Fungal Ecol* 6:328–338
- Trevathan-Tackett SM, Lane AL, Bishop N, Ross C (2015) Metabolites derived from the tropical seagrass *Thalassia testudinum* are bioactive against pathogenic *Labyrinthula* sp. *Aquat Bot* 122:1–8
- Van Alstyne KL, Padilla DK (in press) The chemical ecology of seagrasses. In: Puglisi-Weening MP, Becerro MA, Paul VJ (eds) *Chemical ecology: the ecological impacts of marine natural products*. CRC Press, Boca Raton, FL
- Vergeer LH, Develi A (1997) Phenolic acids in healthy and infected leaves of *Zostera marina* and their growth-limiting properties towards *Labyrinthula zosterae*. *Aquat Bot* 58:65–72
- Vergeer LHT, Aarts TL, De Groot JD (1995) The ‘wasting disease’ and the effect of abiotic factors (light intensity, temperature, salinity) and infection with *Labyrinthula zosterae* on the phenolic content of *Zostera marina* shoots. *Aquat Bot* 52:35–44
- Ward JR, Lafferty KD (2004) The elusive baseline of marine disease: Are diseases in ocean ecosystems increasing? *PLoS Biol* 2:e120
- Washington State Department of Natural Resources (2015) Puget Sound submerged vegetation monitoring program 2010–2013 report. Available at: www.dnr.wa.gov/Publications/aqr_nrsh_svmp_report_2013.pdf
- Waycott M, Duarte CM, Carruthers TJ, Orth RJ and others (2009) Accelerating loss of seagrasses across the globe threatens coastal ecosystems. *Proc Natl Acad Sci USA* 106:12377–12381
- Williams SL, Ruckelshaus MH (1993) Effects of nitrogen availability and herbivory on eelgrass (*Zostera marina*) and epiphytes. *Ecology* 74:904–918
- Wyllie-Echeverria S, Mumford TE, Gaydos JK, Buffum S (2003) *Z. marina* declines in San Juan County, WA. Proceedings of the Wescott Bay Taskforce mini-workshop. Available at www.seadocsociety.org/publications (accessed 20 August 2013)
- Wyllie-Echeverria S, Talbot SL, Rearick JR (2010) Genetic structure and diversity of *Zostera marina* (eelgrass) in the San Juan Archipelago, Washington, USA. *Estuaries Coasts* 33:811–827
- Yang S, Wheat EE, Horwith MJ, Ruesink JL (2013) Relative impacts of natural stressors on life history traits underlying resilience of intertidal eelgrass (*Zostera marina* L.). *Estuaries Coasts* 36:1006–1013

Editorial responsibility: Rebecca Gast,
Woods Hole, Massachusetts, USA

Submitted: June 25, 2015; Accepted: December 21, 2015
Proofs received from author(s): February 4, 2016