

2016

Fungal denitrification: *Bipolaris sorokiniana* exclusively denitrifies inorganic nitrogen in the presence and absence of oxygen

R Phillips

G Grelet

A McMillan

BK Song

Virginia Institute of Marine Science

B Weir

See next page for additional authors

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Phillips, R; Grelet, G; McMillan, A; Song, BK; Weir, B; and Et al., "Fungal denitrification: *Bipolaris sorokiniana* exclusively denitrifies inorganic nitrogen in the presence and absence of oxygen" (2016). *VIMS Articles*. 815.

<https://scholarworks.wm.edu/vimsarticles/815>

This Article is brought to you for free and open access by the Virginia Institute of Marine Science at W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Authors

R Phillips, G Grelet, A McMillan, BK Song, B Weir, and Et al.

RESEARCH LETTER – Environmental Microbiology

Fungal denitrification: *Bipolaris sorokiniana* exclusively denitrifies inorganic nitrogen in the presence and absence of oxygen

Rebecca Phillips^{1,*}, Gwen Grelet¹, Andrew McMillan¹, Bongkeun Song², Bevan Weir¹, Thilak Palmada¹ and Craig Tobias³¹Landcare Research, Gerald Street, Lincoln, Canterbury 7608, New Zealand, ²Department of Biological Sciences, Virginia Institute of Marine Science, Gloucester Point, VA 23062, USA and ³Department of Marine Sciences, University of Connecticut, Groton, CT 06269, USA

*Corresponding author: Landcare Research, Gerald Street, Lincoln, Canterbury 7608, New Zealand. Tel: +64-3-3219873;

E-mail: phillipsr@landcareresearch.co.nz

One sentence summary: Inorganic nitrogen addition alters fungal denitrification and respiration in the presence and absence of oxygen.

Editor: Matthias Brock

¹Rebecca Phillips, <http://orcid.org/0000-0003-3881-9065>

ABSTRACT

Fungi may play an important role in the production of the greenhouse gas nitrous oxide (N₂O). *Bipolaris sorokiniana* is a ubiquitous saprobe found in soils worldwide, yet denitrification by this fungal strain has not previously been reported. We aimed to test if *B. sorokiniana* would produce N₂O and CO₂ in the presence of organic and inorganic forms of nitrogen (N) under microaerobic and anaerobic conditions. Nitrogen source (organic-N, inorganic-N, no-N control) significantly affected N₂O and CO₂ production both in the presence and absence of oxygen, which contrasts with bacterial denitrification. Inorganic N addition increased denitrification of N₂O (from 0 to 0.3 µg N₂O-N h⁻¹ g⁻¹ biomass) and reduced respiration of CO₂ (from 0.1 to 0.02 mg CO₂ h⁻¹ g⁻¹ biomass). Isotope analyses indicated that nitrite, rather than ammonium or glutamine, was transformed to N₂O. Results suggest the source of N may play a larger role in fungal N₂O production than oxygen status.

Keywords: nitrite; glutamine; nitrous oxide; carbon dioxide; organic nitrogen; co-denitrification

INTRODUCTION

Fungal denitrification may contribute more to soil emissions of the greenhouse gas nitrous oxide (N₂O) than bacterial denitrification (Laughlin and Stevens 2002; Chen, Mothapo and Shi 2015), but physicochemical factors potentially altering fungal nitrogen (N) cycling require investigation. While many soil bacteria contain the N₂O reductase gene (*nosZ*) (required for N₂O to N₂ conversion), this gene is not found in fungi. Nonetheless, N₂ production by soil fungi has been reported (Shoun et al. 2012; Long

et al. 2013). Some fungi, particularly for genera in the *Hypocreales* order (such as *Fusarium*), are capable of producing N₂O (Maeda et al. 2015) under anaerobic (Zumft 1997; Morozkina and Kurakov 2007; Shoun et al. 2012) or microaerobic conditions (Zhou et al. 2001; Morozkina and Kurakov 2007; Takaya 2009). Two pathways of fungal denitrification have been reported: (a) classical denitrification of nitrate or nitrite when oxygen is limited and insufficient to support aerobic respiration and (b) hybrid formation of N₂O, where two N sources are combined (Spott, Russow

and Stange 2011). It is not clear if induction of anaerobiosis or changes in N source (Takaya 2002; Wei et al. 2014) cause a shift from one pathway to another. Hybrid N_2O and/or N_2 formation (also referred to as co-denitrification) has been observed in the presence of inorganic and organic N (Su, Takaya and Shoun 2004; Spott, Russow and Stange 2011; Long et al. 2013) but may also occur in the presence of two inorganic forms of N (Spott, Russow and Stange 2011). Data are lacking that indicate how inorganic and organic sources of N affect fungal denitrification and co-denitrification to N_2O pre- and postinduction of anaerobiosis. Addition of two N sources (one enriched with ^{15}N) would allow us to determine if N_2O formation could be attributed to denitrification or co-denitrification.

While a number of fungal strains have been tested for denitrification potential (Maeda et al. 2015), many strains, such as *Bipolaris sorokiniana*, have not been tested. *Bipolaris sorokiniana* is the asexual form (anamorph) of the fungus *Cochliobolus sativus*, a common plant pathogen in the order Pleosporales with a wide range of plant hosts. It is particularly common on cereals in the Poaceae family and is widely known as the causal agent for the common root rot. It can also survive as thick-walled conidia or as saprotrophic mycelium in soil or crop debris. Geographic distribution of *B. sorokiniana* is worldwide, particularly near mid-latitudes, in grasslands and agricultural fields (Kumar et al. 2002). It is not known if this soil fungus will denitrify inorganic or organic forms of N and if aerobic respiration and denitrification co-occur under controlled conditions. While a large body of research has reported bacterial N_2O and CO_2 production under microaerobic and anaerobic conditions (Butterbach-Bahl et al. 2013), few reports are available indicating if rates of fungal CO_2 and N_2O production vary with inorganic and/or organic N. It is unclear if fungi incubated with O_2 will respond to induction of anaerobiosis in a manner similar to denitrifying bacteria, with increased rates of N_2O production occurring within hours of induction (Firestone and Tiedje 1979; Smith and Tiedje 1979). The goal of this research was to test how N source (organic or inorganic) influences production of N_2O and CO_2 by *B. sorokiniana* pre and post anaerobiosis. Results add to the emerging body of knowledge regarding controls on fungal denitrification and co-denitrification processes.

MATERIALS AND METHODS

We used pure culture of *B. sorokiniana* (Sacc.) Shoemaker [teleomorph: *C. sativus* (S. Ito & Kurib.) Drechsler ex Dastur] to test how organic N [glutamine ($\text{C}_5\text{H}_{10}\text{N}_2\text{O}_3$)] and inorganic N [ammonium sulfate ($(\text{NH}_4)_2\text{SO}_4$) and sodium nitrite (NaNO_2)] sources might influence fungal production of N_2O and CO_2 under sterile conditions. We controlled O_2 status with air-tight laboratory incubation vessels to evaluate rates of N_2O and CO_2 production pre and post anaerobiosis on the same samples (Firestone and Tiedje 1979; Smith and Tiedje 1979; Zhou et al. 2001; Butterbach-Bahl et al. 2013). We obtained isolate ICMP 6809 from the ICMP culture collection (<http://www.landcareresearch.co.nz/resources/collections/icmp>) in its anamorphic state. It was grown under aerobic conditions in the same media previously used by Rohe et al. (2014a,b) and Shoun et al. (1992), containing 1% glucose, 0.2% peptone, 0.02% $\text{MgSO}_4 \cdot 7 \text{H}_2\text{O}$, 2 ppm $\text{CaCl}_2 \cdot 6 \text{H}_2\text{O}$, 2 ppm $\text{FeSO}_4 \cdot 7 \text{H}_2\text{O}$ and 0.01 mol potassium phosphate (pH 7.4). This media, which contained N as peptone, was used only for fungal growth. Another media, where the N source was omitted, was used for the incubations and referred to here as no-N media. After 7 days,

cultures were washed, drained and stored in a reduced volume of the no-N media. Using the no-N media, four solutions were prepared for fungal inoculation: (a) no-N media (b) 0.5 mmols N as $\text{C}_5\text{H}_{10}\text{N}_2\text{O}_3$, (c) 0.25 mmols N as $\text{Na}^{15}\text{NO}_2$ (99.5 atom%; Cambridge Isotope Laboratory, Andover, MA) and 0.25 mmols N as $(\text{NH}_4)_2\text{SO}_4$, or (d) 0.25 mmols N as $\text{Na}^{15}\text{NO}_2$ and 0.25 mmols N as $\text{C}_5\text{H}_{10}\text{N}_2\text{O}_3$. The $\text{Na}^{15}\text{NO}_2$ was used to determine if these fungi would preferentially use NO_2 to form N_2O (thus forming $^{46}\text{N}_2\text{O}$) or if other sources of N would be utilized.

Approximately 10 ml of fungal biomass were transferred to 125 ml serum bottles, and four replicate bottles were inoculated with 1 ml of each sterile solution and 1 ml of no-N media. Additional bottles containing media solutions without fungi were also prepared. All bottles were sealed and the headspace evacuated and flushed with ultrapure helium, and then injected with O_2 to achieve 0.4% O_2 headspace before setting up on a robotic gas chromatograph (GC) fitted with electron capture and thermal conductivity detectors (Phillips et al. 2014; McMillan et al. 2015). Bottles were placed on the GC and measured every 6 hours for N_2O , CO_2 and O_2 at 19°C. Following this 48-h incubation experiment, bottles were then evacuated and flushed three times to create an anaerobic headspace for the second experiment, where the only difference between experiments was headspace O_2 concentration. Sterility was maintained and conditions remained constant during both incubations, including pH (ranged from 6.2 to 6.9). Data were normalized to the corresponding dry weight of fungal biomass in each bottle. The N isotopes for N_2O gas in the headspace of each sample at the end of the incubations were measured on a continuous-flow isotope ratio mass spectrometer (Thermo Finnigan Delta V; Thermo Scientific, Waltham, MA) in line with an automated gas bench interface (Thermo Gas Bench II) to determine if $^{45}\text{N}_2\text{O}$ or $^{46}\text{N}_2\text{O}$ were present. Precision of the isotopic analysis was <1 atom%. Data were analysed for effects of N source treatment on N_2O and CO_2 production rates for microaerobic and anaerobic experiments separately with a generalized linear model and means were compared with Tukey's test.

RESULTS AND DISCUSSION

Table 1 and Fig. 1 show effects of treatment on CO_2 and N_2O production for the microaerobic incubations were similar to effects of treatment for the anaerobic incubations. Fungi inoculated with only organic-N ($\text{C}_5\text{H}_{10}\text{N}_2\text{O}_3$) or no-N media only linearly produced CO_2 (1–2 mg $\text{CO}_2 \text{ h}^{-1} \text{ g}^{-1}$) but not N_2O . Fungi inoculated with inorganic-N ($\text{Na}^{15}\text{NO}_2$ or $(\text{NH}_4)_2\text{SO}_4$) or inorganic-N plus organic-N ($\text{Na}^{15}\text{NO}_2$ and $\text{C}_5\text{H}_{10}\text{N}_2\text{O}_3$) linearly produced N_2O (0.2–0.3 $\mu\text{g} \text{ N}_2\text{O-N h}^{-1} \text{ g}^{-1}$) but an order of magnitude less CO_2 . The mass $^{46}\text{N}_2\text{O}$ comprised 94.7%–97.3% of all N_2O in the headspace, yielding an equivalent ^{15}N of 98 to 99 (± 0.3) atom%. This enrichment was effectively equivalent to enrichment of the $^{15}\text{NO}_2$ used in the experiment. The lack of headspace $^{45}\text{N}_2\text{O}$ indicated hybrid formation of N_2O by *B. sorokiniana* did not occur under any of these conditions, and NO_2^- was the only N form used to form N_2O . Headspace O_2 (Fig. 2) over the aerobic time course indicated rapid declines for the no-N media and organic-N treatments but not the inorganic-N treatments. Minimal rates of CO_2 respiration (Table 1) in the presence of inorganic-N were consistent with lack of O_2 utilization by *B. sorokiniana* (Fig. 2). We also observed chemodenitrification (6%–8% of total biologically produced N_2O) for media amended with inorganic-N only (van Cleemput 1998; Kampschreur et al. 2011; Jones et al. 2015).

Table 1. Rates of N_2O and CO_2 produced during (1) aerobic and (2) anaerobic incubation experiments by treatment per gram of fungal biomass. Average rates followed by different letters for each experiment represent significant differences between treatments. The same letters indicate no differences between treatments.

Experiment	Treatment	Mean		Mean	
		$ng\ N_2O-N\ h^{-1}\ g^{-1}\ biomass$	Std. Dev.	$\mu g\ CO_2\ h^{-1}\ g^{-1}\ biomass$	Std. Dev.
Anaerobic	No N	0.51 ^a	0.26	896.45 ^a	206.08
	Organic N	0.29 ^a	0.11	728.46 ^a	148.66
	Inorganic N	311.30 ^b	18.98	6.17 ^b	1.34
	Inorganic N + organic N	211.79 ^c	14.44	7.87 ^b	1.48
Aerobic	No N	0.57 ^a	0.23	1179.89 ^a	262.62
	Organic N	0.32 ^a	0.23	988.19 ^a	222.95
	Inorganic N	354.55 ^b	26.00	18.26 ^b	1.48
	Inorganic N + organic N	306.33 ^c	28.61	31.52 ^b	4.03




Figure 1. Average (std. dev; $n = 4$) N_2O and CO_2 measured in the headspace per g of fungal biomass by treatment during aerobic (a, b) and anaerobic (c, d) incubations.

Evidence of chemodenitrification, where N_2O was produced under sterile conditions, may be due to reactions between NO_2^- and reduced metals in the media (van Cleemput 1998; Kampschreur et al. 2011). We chose the media for this experiment because it is commonly used for fungal denitrification investigations (Shoun et al. 1992; Rohe et al. 2014a,b). Kampschreur et al. (2011) used a chemostat to demonstrate that NO_2^- to NO is an equilibrium-based reaction, where emissions of NO and N_2O were coupled to iron oxidation under anoxic conditions. A recent review by Medinets et al. (2015) argued that NO_2^- , a precursor of NO , is central to processes associated with chemodenitrifica-

tion. Additional research is needed to determine the reactants involved in this specific media and the prevalence of chemodenitrification in an oxygenated headspace.

Evidence of biological denitrification coupled with reduced CO_2 respiration in the presence of inorganic-N may be due to *B. sorokiniana* preferentially denitrifying as a defence against accumulation of NO_2^- (Geets, Boon and Verstraete 2006; Clark et al. 2012). Fungi are capable of converting NO_2^- to NO , the precursor to soil N_2O emissions (Russow, Stange and Nueue 2009), through expression of the NO_2^- reductase gene (*nirK*), and NO is a regulator of O_2 consumption in eukaryotes (Thomas et al.




Figure 2. Average (std. dev; $n = 4$) O_2 measured in the headspace by treatment during aerobic incubation.

2001). We suspect that some NO_2^- was converted to NO (both abiotic and biotic), which inhibited mitochondrial respiration (Thomas et al. 2001; Medinets et al. 2015) and therefore CO_2 production, as shown in Fig. 1 and Table 1. One way of discerning the importance of NO_2^- reduction to NO and N_2O by *B. sorokiniana* would be to identify *nirK* gene expression in conjunction with NO measurements. If *B. sorokiniana* does not reduce NO_2^- to NO, then chemical transformation of NO_2^- may be the source of NO (van Cleemput 1998; Kampschreur et al. 2011). In either event, there is a need to identify chemical versus biological sources of NO, the precursor to N_2O (Russow, Stange and Nueue 2009).

Denitrification in soils and sediments is referred to as a keystone ecosystem service with positive water quality implications. However, when N_2O is the final end product of denitrification, there are negative impact on stratospheric ozone and radiative forcing in the troposphere (Erisman et al. 2013). Here, *B. sorokiniana*, a ubiquitous soil fungus, demonstrated the capacity to transform dissolved NO_2^- to gaseous N_2O at the expense of aerobic respiration, a more energetically favourable pathway. Denitrification and aerobic respiration for *B. sorokiniana* were closely linked via NO_2^- . Additional investigations into alternative respiration pathways and effects of inorganic N on C assimilation are warranted, particularly with respect to O_2 status. Here, we focused on how inorganic and organic N sources influence pathways to N_2O production before and after onset of anaerobiosis (Firestone and Tiedje 1979; Smith and Tiedje 1979; Zhou et al. 2001). Future work should be designed with sample in parallel to test how O_2 and N interact to affect fungal denitrification phenotype and genotype. Our results suggest that fungal denitrification may not be driven by anaerobiosis but instead by form of N; however, additional strains need to be tested to determine if this is widespread among soil fungi or unique to *B. sorokiniana*.

ACKNOWLEDGEMENTS

The authors are grateful to Veronica Rolleston, Trish McLennan and Sujatha Senananake for contributing their technical expertise and laboratory facilities. We also thank Mikki Eken for editorial review and Fiona Carswell, Sue Scheele, Chris Phillips, and David Whitehead at Landcare Research for their continued support.

FUNDING

This work was funded by a US Department of Agriculture grant [2014-67019021614]; New Zealand's Ministry of Business, Innovation and Employment; and New Zealand Ecosystems and Global Change Fund.

Conflict of interest. None declared.

REFERENCES

- Butterbach-Bahl K, Baggs EM, Dannenmann M et al. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philos T Roy Soc B* 2013;**368**, DOI: 10.1098/rstb.2013.0122.
- Chen H, Mothapo NV, Shi W. Fungal and bacterial N_2O production regulated by soil amendments of simple and complex substrates. *Soil Biol Biochem* 2015;**84**:116–26.
- Clark IM, Buchkina N, Jhureea D et al. Impacts of nitrogen application rates on the activity and diversity of denitrifying bacteria in the Broadbalk Wheat Experiment. *Philos T Roy Soc B* 2012;**367**:1235–44.
- Erisman JW, Galloway JN, Seitzinger S et al. Consequences of human modification of the global nitrogen cycle. *Philos T Roy Soc B* 2013;**368**, DOI: 10.1098/rstb.2013.0116.
- Firestone MK, Tiedje JM. Temporal change in nitrous oxide and nitrogen from denitrification following onset of anaerobiosis. *Appl Environ Microb* 1979;**38**:673–9.
- Geets J, Boon N, Verstraete W. Strategies of aerobic ammonia-oxidizing bacteria for coping with nutrient and oxygen fluctuations. *FEMS Microb Ecol* 2006;**58**:1–13.
- Jones LC, Peters B, Lezama Pacheco JS et al. Stable isotopes and iron oxide mineral products as markers of chemodenitrification. *Environ Sci Technol* 2015;**49**:3444–52.
- Kampschreur MJ, Kleerebezem R, de Vet WWJM et al. Reduced iron induced nitric oxide and nitrous oxide emission. *Water Res* 2011;**45**:5945–52.
- Kumar J, Schafer P, Huckelhoven R et al. *Bipolaris sorokiniana*, a cereal pathogen of global concern: cytological and molecular approaches towards better control. *Mol Plant Pathol* 2002;**3**:185–95.
- Laughlin RJ, Stevens RJ. Evidence for fungal dominance of denitrification and codenitrification in a grassland soil. *Soil Sci Soc Am J* 2002;**66**:1540–8.
- Long A, Heitman J, Tobias C et al. Co-occurring anammox, denitrification, and codenitrification in agricultural soils. *Appl Environ Microb* 2013;**79**:168–76.
- McMillan A, Pal P, Phillips RL et al. Can pH amendments in grazed pastures help reduce N_2O emissions?—the effects of liming and urine addition on the completion of denitrification in fluvial and volcanic soils. *Soil Biol Biochem* 2015;**93**:90–104.
- Maeda K, Spor A, Edel-Hermann V et al. N_2O production, a widespread trait in fungi. *Sci Rep* 2015;**5**:9697.
- Medinets S, Skiba U, Rennenberg H et al. A review of soil NO transformation: associated processes and possible physiological significance on organisms. *Soil Biol Biochem* 2015;**80**:92–117.
- Morozkina EV, Kurakov AV. Dissimilatory nitrate reduction in fungi under conditions of hypoxia and anoxia: a review. *Appl Biochem Micro+* 2007;**43**:544–9.
- Phillips RL, McMillan AMT, Palmada T et al. Temperature effects on N_2O and N_2 production in a New Zealand pasture soil. *New Zeal J Agr Res* 2014;**58**:89–95.
- Rohe L, Anderson T-H, Braker G et al. Dual isotope and isotopomer signatures of nitrous oxide from fungal

- denitrification—a pure culture study. *Rapid Commun Mass Sp* 2014a;**28**:1893–903.
- Rohe L, Anderson T-H, Braker G et al. Fungal oxygen exchange between denitrification intermediates and water. *Rapid Commun Mass Sp* 2014b;**28**:377–84.
- Russow R, Stange CF, Nueue HU. Role of nitrite and nitric oxide in the processes of nitrification and denitrification in soil: results from ^{15}N tracer experiments. *Soil Biol Biochem* 2009;**41**:785–95.
- Shoun H, Fushinobu S, Jiang L et al. Fungal denitrification and nitric oxide reductase cytochrome P450nor. *Philos T Roy Soc B* 2012;**367**:1186–94.
- Shoun H, Kim D-H, Uchiyama H et al. Denitrification by fungi. *FEMS Microbiol Lett* 1992;**94**:277–81.
- Smith MS, Tiedje JM. Phases of denitrification following oxygen depletion in soil. *Soil Biol Biochem* 1979;**11**: 261–67.
- Spott O, Russow R, Stange CF. Formation of hybrid N_2O and hybrid N_2 due to codenitrification: first review of a barely considered process of microbially mediated N-nitrosation. *Soil Biol Biochem* 2011;**43**:1995–2011.
- Su F, Takaya N, Shoun H. Nitrous oxide-forming co-denitrification catalyzed by cytochrome P450nor. *Biosci Biotech Bioch* 2004;**68**:473–5.
- Takaya N. Dissimilatory nitrate reduction metabolisms and their control in fungi. *J Biosci Bioeng* 2002;**94**:506–10.
- Takaya N. Response to hypoxia, reduction of electron acceptors, and subsequent survival by filamentous fungi. *Biosci Biotech Bioch* 2009;**73**:1–8.
- Thomas DD, Liu X, Kantrow SP et al. The biological lifetime of nitric oxide: implications for the perivascular dynamics of NO and O_2 . *P Natl Acad Sci USA* 2001;**98**:355–60.
- van Cleemput O. Subsoils: chemo- and biological denitrification, N_2O and N_2 emissions. *Nutr Cycl Agroecosys* 1998;**52**:187–94.
- Wei W, Isobe K, Shiratori Y et al. N_2O emission from cropland field soil through fungal denitrification after surface applications of organic fertilizer. *Soil Biol Biochem* 2014;**69**:157–67.
- Zhou Z, Takaya N, Sakairi MAC et al. Oxygen requirement for denitrification by the fungus *Fusarium oxysporum*. *Arch Microbiol* 2001;**175**:19–25.
- Zumft WG. Cell biology and molecular basis of denitrification. *Microbiol Mol Biol R* 1997;**61**:533–616.