

2015

Relative Importance of Nutrient Load and Wind on Regulating Interannual Summer Hypoxia in the Chesapeake Bay (vol 38, pg 1048, 2015)

P Wang

H Wang

Virginia Institute of Marine Science

L Linker

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Wang, P; Wang, H; and Linker, L, "Relative Importance of Nutrient Load and Wind on Regulating Interannual Summer Hypoxia in the Chesapeake Bay (vol 38, pg 1048, 2015)" (2015). *VIMS Articles*. 847.
<https://scholarworks.wm.edu/vimsarticles/847>

This Article is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Erratum to: Relative Importance of Nutrient Load and Wind on Regulating Interannual Summer Hypoxia in the Chesapeake Bay

Ping Wang · Harry Wang · Lewis Linker

Published online: 5 September 2014
© Coastal and Estuarine Research Federation 2014

Erratum to: Estuaries and Coasts
DOI 10.1007/s12237-014-9867-5

The corrected Fig. 6 is as follows:

The online version of the original article can be found at <http://dx.doi.org/10.1007/s12237-014-9867-5>.

P. Wang (✉) · H. Wang
Virginia Institute of Marine Science, PO Box 1346, Gloucester,
VA 23062, USA
e-mail: pwang@chesapeakebay.net

L. Linker
USEPA Chesapeake Bay Program Office, 410 Severn Avenue,
Annapolis, MD 21403, USA




Fig. 6 Tree diagrams for anoxic volume and its influencing variables. **a** For summer anoxic volume, using 1985–2012 data ($n=28$), **b** for summer anoxic volume, using 1991–2005 data ($n=15$), **c** for July anoxic volume,

using 1985–2012 data ($n=28$), and **d** for July anoxic volume, using 1991–2005 data ($n=15$)