

2011

## The role of nitrogen on the growth and colony development of *Phaeocystis globosa* (Prymnesiophyceae)

XD Wang

Y Wang

Walker O. Smith Jr.

*Virginia Institute of Marine Science*

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Aquaculture and Fisheries Commons](#)

---

### Recommended Citation

Wang, XD; Wang, Y; and Smith, Walker O. Jr., "The role of nitrogen on the growth and colony development of *Phaeocystis globosa* (Prymnesiophyceae)" (2011). *VIMS Articles*. 935.

<https://scholarworks.wm.edu/vimsarticles/935>

This Article is brought to you for free and open access by the Virginia Institute of Marine Science at W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).



## The role of nitrogen on the growth and colony development of *Phaeocystis globosa* (Prymnesiophyceae)

Xiaodong Wang , Yan Wang & Walker O. Smith Jr

To cite this article: Xiaodong Wang , Yan Wang & Walker O. Smith Jr (2011) The role of nitrogen on the growth and colony development of *Phaeocystis globosa* (Prymnesiophyceae), European Journal of Phycology, 46:3, 305-314, DOI: [10.1080/09670262.2011.602430](https://doi.org/10.1080/09670262.2011.602430)

To link to this article: <https://doi.org/10.1080/09670262.2011.602430>



Published online: 26 Aug 2011.



Submit your article to this journal [↗](#)



Article views: 338



Citing articles: 5 View citing articles [↗](#)

# The role of nitrogen on the growth and colony development of *Phaeocystis globosa* (Prymnesiophyceae)

XIAODONG WANG<sup>1</sup>, YAN WANG<sup>1</sup> AND WALKER O. SMITH JR<sup>2</sup>

<sup>1</sup>Research Center for Harmful Algae and Aquatic Environment, Jinan University, Guangzhou 510632, China

<sup>2</sup>Virginia Institute of Marine Science, College of William & Mary, Gloucester Point, Virginia 23062, USA

(Received 14 December 2010; revised 14 May 2011; accepted 8 June 2011)

The effects of nitrate, ammonium and urea on the growth and colony formation of three strains of *Phaeocystis globosa* were investigated. Although ammonium and urea supported growth, nitrate was the favoured nitrogen source for the growth of solitary cells for all three strains. *Phaeocystis globosa* CCMP 1528 and 629 formed colonies in all cultures where nitrate was the sole nitrogen source, but only a few colonies were observed in ammonium and urea treatments. Ammonium and urea were far less effective in supporting growth, biomass generation and colony formation in all three strains. Once colonies developed, colonial cells accounted for at least 15% of the total cells when grown with nitrate; colonial chlorophyll also contributed up to 60% to the total chlorophyll. The growth rates of colonial cells when using nitrate were greater than solitary cells. Changes in colony size, colonial cell abundance and total *P. globosa* abundance as affected by the nitrogen source may influence the carbon flux within the pelagic food web.

**Key words:** carbon flux, colony formation, growth, Haptophyta, nitrogen, *Phaeocystis*, solitary cells

## Introduction

The genus *Phaeocystis* (Prymnesiophyceae) is globally distributed, has a distribution that ranges from tropical to polar oceans, and plays a significant role in global carbon and sulphur cycles as well as regional food webs (Lancelot *et al.*, 1998; Schoemann *et al.*, 2005). It has an unusual heteromorphic life cycle that includes gelatinous colonies and solitary cells. Individual cells are generally 3–10 µm in diameter, whereas colonies, with a few to thousands of cells embedded in a mucilaginous matrix, can occasionally be up to 3 cm in diameter (Rousseau *et al.*, 1994; Chen *et al.*, 2002). Among the six species that have been identified, *P. globosa*, *P. pouchetii* and *P. antarctica* have been reported to form massive blooms in colonial form (Medlin & Zingone, 2007). The formation of colonies contributes to the success of *Phaeocystis*, as it has been suggested that it may provide protection against grazers, viruses and bacteria (Hamm *et al.*, 1999; Jakobsen & Tang, 2002; Brussaard *et al.*, 2005). There is also evidence that the colonial matrix can supply energy and nutrients when light and nutrients become limiting (Lancelot & Mathot, 1985; Schoemann *et al.*, 2001).

*Phaeocystis* is regarded as a harmful algal genus in coastal waters (Lancelot *et al.*, 1998) because of negative influences on ecosystem structure, function, fisheries and tourism (Lancelot *et al.*, 1987; Peperzak & Poelman, 2008). Dense blooms have been observed off the coasts of the North Sea (Lancelot *et al.*, 1987), North Atlantic (Gieskes *et al.*, 2007), southeast China (Qi *et al.*, 2004), Vietnam (Tang *et al.*, 2004) and Norway (Larsen *et al.*, 2004). Clones isolated from Chinese and Norwegian waters have been shown to have haemolytic properties (Shen *et al.*, 2004; van Rijssel *et al.*, 2007). All of the locations listed above are considered to be eutrophic, due to riverine and anthropogenic nutrient enrichment (Lancelot *et al.*, 1987; Zhang *et al.*, 2000; Cadée & Hegeman, 2002; Thanh *et al.*, 2004). Increased N:Si and P:Si ratios have been proposed as a mechanism for the increased number and duration of blooms of non-siliceous phytoplankton (Jacobsen *et al.*, 1995), and the development of *Phaeocystis* blooms has been associated with increasing nutrient loads, especially NO<sub>3</sub> and PO<sub>4</sub>, and subsequent changes in the N:P ratio (Riegman *et al.*, 1992).

*Phaeocystis* colony development is controlled mainly by NO<sub>3</sub> supply (Lancelot *et al.*, 2007). Sharp decreases in N:P ratios as a result of high consumption of nitrate have been observed in two

Correspondence to: Xiaodong Wang. E-mail: pouchetii@gmail.com

*P. pouchetii* blooms (Bätje & Michaelis, 1986). Nitrate removal by *P. antarctica* in the Ross Sea has been observed repeatedly (Smith *et al.*, 1998; Arrigo *et al.*, 2000; Cochlan & Bronk 2001), suggesting that *P. antarctica* prefers nitrate as a nitrogen source, or at least can efficiently remove it. Interestingly, Cochlan & Bronk (2003) found that the dependence of colonial *P. antarctica* on nitrate decreased from spring to summer, and Mathot *et al.* (2000) noted a marked increase in the relative contribution of solitary cells after the primary bloom. This may imply that colonies derive their nitrogen from nitrate, whereas solitary cells may depend more heavily on ammonium or reduced nitrogen (which increases in concentration after the primary bloom). In the North Sea the distribution of nitrate showed a gradual decrease with distance from land and was correlated with a decline in the *P. pouchetii* biomass (Veldhuis *et al.*, 1986). *Phaeocystis* dominates the spring bloom in Belgian coastal waters, where nitrate is the major nitrogen source (Tungaraza *et al.*, 2003). Riegman *et al.* (1992) provided experimental evidence that *Phaeocystis* blooms may be restricted to those N-limited environments where colony formation is stimulated by nitrate and inhibited by ammonium. Model studies also demonstrated a significant correlation between *Phaeocystis* colony biomass and winter NO<sub>3</sub> in the North Sea (Lancelot *et al.*, 2005).

Nitrate has long been considered as the critical nitrogenous substrate for phytoplankton; however, studies have also suggested that reduced nitrogen, such as ammonium and urea, can also play a key role in phytoplankton growth (Eppley *et al.*, 1971; Lomas *et al.*, 2002; Berman & Bronk, 2003; Glibert & Burkholder, 2006; Solomon *et al.*, 2010). Elevated ammonium concentrations inhibit nitrate uptake in diatoms as well as in *P. pouchetii* (MacIsaac & Dugdale, 1969; Muggli & Smith, 1993; L'Helguen *et al.*, 2008). The field work of Tungaraza *et al.* (2003) confirmed that nitrate uptake decreased rapidly due to elevated concentrations of NH<sub>4</sub>, and that *Phaeocystis* increased its NH<sub>4</sub> uptake when more NH<sub>4</sub> was available. A 65-fold higher urea uptake rate (relative to other N substrates) by *P. pouchetii* was also observed by Sanderson *et al.* (2008), but utilization of reduced and organic forms by *Phaeocystis* has received relatively less attention compared to uptake of NO<sub>3</sub> and NH<sub>4</sub> (Veldhuis & Admiraal, 1987; Schoemann *et al.*, 2005; Sanderson *et al.*, 2008). Indeed, it is generally assumed that small cells (5 µm diameter and less) utilize reduced nitrogen (NH<sub>4</sub>), whereas net plankton utilize nitrate (Probyn & Painting, 1985; Stolte & Riegman, 1995, 1996). Because of its unusual life cycle, it remains unclear how the different nitrogen sources affect the formation of

*Phaeocystis* colonies and support the growth of solitary and colonial cells. *Phaeocystis globosa* is the most widespread of all known colony-forming *Phaeocystis* species, and forms extensive blooms that have deleterious impacts on coastal ecosystems. Understanding the species' use of nitrogen may help in predicting its appearance and concentrations in coastal environments.

## Materials and methods

### Stock maintenance

Three strains of *Phaeocystis globosa* isolated from two geographical areas were used. *P. globosa* CCMP 627 and CCMP 629 were originally isolated from the North Atlantic, whereas *P. globosa* CCMP 1528 was isolated from the South Pacific. The three strains were maintained in f/2 medium in coastal seawater (Guillard & Ryther, 1962) at 20°C, with a salinity of 32 PSU under an irradiance of 50 µmol photons m<sup>-2</sup> s<sup>-1</sup> on a 12:12 h light:dark cycle. The stocks were maintained in exponential growth by regular dilutions with fresh media. The cultures were non-axenic, and no attempt was made to reduce bacterial growth.

### Experimental cultures


Experiments were conducted in a walk-in growth chamber in the same conditions as for stock maintenance. Prior to the experiments, solitary cells of the three strains of *P. globosa* were isolated by passing the culture through a 10-µm nylon sieve twice under gravity (Tang, 2003), and experiments were initiated with an initial solitary cell density of 10<sup>4</sup> cells ml<sup>-1</sup>. Triplicate one-litre batch cultures of each strain were grown in f/2 medium with a salinity of 32 PSU, where NH<sub>4</sub><sup>+</sup>, NO<sub>3</sub> or urea were added as the nitrogen source with an initial N concentration of 8.82 × 10<sup>-4</sup> M. All cultures were bubbled with filtered air to maintain a uniform distribution (Wang *et al.*, 2010).

### Microscopy and chlorophyll determinations

Samples for biomass assessment were collected every other day. Aliquots of known volume were filtered through Whatman GF/F filters to obtain total chlorophyll *a*, and through 20 µm polycarbonate membranes (Poretics) to obtain the >20 µm fraction, which was assumed to be representative of colonial chlorophyll (Tang *et al.*, 2008; Wang *et al.*, 2010). All chlorophyll *a* samples were extracted in darkness at 0°C for 24 h in 7 ml of 90% acetone; fluorescence of extracted chlorophyll *a* was determined on a TD-700 fluorometer (Turner Designs) before and after acidification (Parsons *et al.*, 1984). Samples for microscopical enumeration of solitary cell and colony abundances were preserved in Lugol's solution (final concentration 2%). Solitary cell concentrations were measured using 1-ml Sedgwick-Rafter chambers. Colony concentration, colony size and cells per colony were measured in 24 multi-plates using a Nikon inverted microscope

**Table 1.** Summary of the growth responses of three clones of *Phaeocystis globosa* as a function of the nitrogen source provided.

<i>P. globosa</i> clone Nitrogen source	CCMP 1528		CCMP 629		CCMP 627	
	Solitary cells	Colonies	Solitary cells	Colonies	Solitary cells	Colonies
Nitrate	++	+++	++	+++	+++	0
Ammonium	+	0	+	0	++	0
Urea	+	0	+	0	++	0

**Fig. 1.** *Phaeocystis globosa* solitary cell abundance (mean  $\pm$  standard deviation;  $n=3$ ) in nitrate (open circles), ammonium (solid circles) and urea (open triangles) cultures.

with a calibrated micro-ruler (Tang, 2003; Wang *et al.*, 2010). Colonies were counted within 24 h of collection to limit colony disruption in the preserved state. Net growth rates (divisions  $d^{-1}$ ) of triplicate samples of each strain growing on each substrate were determined from the slope of a regression between  $\ln(N)$  (cells  $ml^{-1}$ ) vs time (days).

#### Statistical analysis

SigmaStat (v. 3.50 SPSS) was used for statistical analyses. Statistical comparisons of the effects of nitrogen source on cell abundance with time were made by 2-way RM ANOVA. Comparisons of changes in colony concentrations, colony size, colonial cell abundance, partitioning of cell and chlorophyll with time, as well as comparison of growth rates between colonial and solitary cells, were conducted by 1-way RM ANOVA and *t*-tests. Prior to ANOVA, percentage data were normalized by an arcsine

transformation (Zar, 1984). Linear regressions were fit to  $\log \text{ cells colony}^{-1}$  vs.  $\log \text{ colony diameter}$ , and comparison of the slopes of regressions between nitrogen was done by 1-way analysis of covariance (ANCOVA). The significance level for all statistical tests was set *a priori* at a critical *P* value of 0.05.

## Results


### Solitary cell abundances

The maximum solitary cell abundance of *P. globosa* CCMP 627 cultures grown on nitrate was significantly greater than those grown using ammonium or urea ( $P < 0.05$ ), but there was no significant difference in maximum abundance using urea and ammonium as the N source ( $P > 0.05$ ; Fig. 1). The response of *P. globosa* CCMP 1528 was similar to that of CCMP 627, indicating that nitrate was the favoured nitrogen source for the growth of solitary cells in both strains. However, for *P. globosa* CCMP 629 the solitary cell abundance grown on nitrate was higher only on days 12 and 14 ( $P < 0.05$ ) when compared with cultures using ammonium and urea, and the absolute difference was far less than observed with CCMP 627. Regardless of nitrogen source, however, the solitary cell abundance of *P. globosa* CCMP 627 was significantly higher than for the other two strains. For example, even when grown on urea, the maximum mean solitary cell abundance was  $2.85 \pm 1.10 \times 10^6 \text{ cells ml}^{-1}$  (mean  $\pm$  standard deviation), four orders of magnitude higher than in other strains. All three strains showed similar response to urea and ammonium, in that reduced nitrogen was far less effective in supporting growth and biomass generation (Table 1).

### Colony abundances

Unlike in nature, colony development did not occur in our *P. globosa* CCMP 627 culture, regardless of the nitrogen source. *Phaeocystis globosa* CCMP 1528 and CCMP 629 formed colonies in all experiments when nitrate was provided as the nitrogen source, whereas only a few colonies were observed when ammonium and urea were provided as nitrogen sources (Table 1 and Fig. 2).

Furthermore, even these few colonies disappeared soon after being formed (Fig. 2). With nitrate, colony concentrations of CCMP 1528 increased from 95 to 2309 colonies  $\text{ml}^{-1}$  by day 16; colony abundance then decreased slightly toward the end of the experiment. The highest colony numbers for CCMP 629 were also observed on day 16.




**Fig. 2.** Colony abundances (mean  $\pm$  standard deviation;  $n=3$ ) of *Phaeocystis globosa* grown with nitrate (open circles), ammonium (solid circles) and urea (open triangles) as a nitrogen source. Colony formation of CCMP 1528 and 629 within non-nitrate treatments were only observed at the beginning of the experiment.

However, in CCMP 629 colony abundance was significantly less than in CCMP 1528 ( $P < 0.01$ ).

#### Partitioning of cells and chlorophyll

Nitrogen source strongly influenced the partitioning of cells between solitary and colonial forms. CCMP 627 existed exclusively as solitary cells in all nitrogen treatments (Table 1). At the beginning of the experiments (that is, on day 4),  $< 1\%$  of the total cells were in colonial form within non-nitrate treatments for CCMP 1528 and CCMP 629, whereas colonial cells accounted for at least 15% of the total cells when grown with nitrate (Fig. 3). This percentage increased to 50% for both CCMP 1528 and CCMP 629, and then decreased again to *c.* 15% of the total cells at the end of the experiments.

Size-fractionated chlorophyll also differed significantly among nitrogen treatments. Because there were relatively few colonial cells in non-nitrate treatments, the  $> 20 \mu\text{m}$  chlorophyll fraction was not detectable for CCMP 1528 and CCMP 629 (confirming the use of this fraction of chlorophyll as a proxy for colonies). With nitrate as the nitrogen source, the percentage of colonial chlorophyll remained unchanged over time ( $P > 0.05$ ), contributing up to 60% to the total chlorophyll for both CCMP 1528 and CCMP 629 (Fig. 3). In addition, nitrogen source did not influence the chlorophyll *a* per cell in colonies of both CCMP 1528 and CCMP 629 ( $P > 0.05$ , data not shown).



**Fig. 3.** The percentage of colonial cells of *Phaeocystis globosa* relative to the number of total cells (top) and the percentage of chlorophyll in the  $> 20 \mu\text{m}$  fraction relative to total chlorophyll *a* (bottom) during supply of nitrate (open circles), ammonium (solid circles) and urea (open triangles). Due to the relatively small number of colonial cells within non-nitrate treatments, the  $> 20 \mu\text{m}$  chlorophyll fraction was not detectable for CCMP 1528 and CCMP 629.

*Colony size and colonial cell*

With nitrate as the nitrogen source, mean colony size increased from 12.5 to 58.4  $\mu\text{m}$  by day 10 for CCMP 1528, and then decreased slightly but significantly towards the end of the experiment (Fig. 4). Similarly, changes in colony diameter also occurred in the CCMP 629 culture. The colony diameters of CCMP 1528 and 629 were less than 40  $\mu\text{m}$  in the ammonium and urea treatments when colonies developed, significantly less than those in the nitrate-replete cultures after the same period of growth.

The number of cells per colony of CCMP 1528 and 629 as a function of nitrogen were similar (Fig. 4). There were significant increases in colonial cell concentration over time for both strains within nitrate treatments ( $P < 0.01$ ), and the highest mean cell densities were 58.4 and 157 cells colony<sup>-1</sup> for CCMP 1528 and CCMP 629, respectively. However, when grown with ammonium or urea, there were less than 10 cells colony<sup>-1</sup>.

In all cases where colonies developed, there was a significant linear log-log relationship between the number of cells per colony and colony diameter ( $P < 0.01$ ) with a slope  $< 2$ , indicating that the number of cells per unit of colony surface area decreased as the colonies increased in size (Fig. 5). For CCMP 1528, the slope of the regression was significantly smaller when ammonium was the N source when compared to the nitrate treatment ( $P < 0.01$ ). Nitrogen source, however, did not significantly affect cell distribution within colonies of CCMP 629 ( $P > 0.05$ ). Additionally,


within nitrate treatments there was no difference in the slope of the regression lines between CCMP 1528 and 629 ( $P > 0.05$ ).

*Growth rates of solitary and colonial cells*


In the cultures supplied with nitrate, the maximum growth rate of solitary cells for CCMP 1528 was  $0.28 \pm 0.02 \text{ d}^{-1}$ , which was significantly less than that of colonial cells ( $0.47 \pm 0.04 \text{ d}^{-1}$ ;  $P < 0.001$ ). The difference was even greater for CCMP 629; the maximum growth rate of colonial cells was twice as rapid as that of solitary cells ( $0.64 \pm 0.18$  vs  $0.33 \pm 0.07 \text{ d}^{-1}$ ; Table 1).

**Discussion***Nitrogen limitation in the medium*

The objective of this study was to understand how different nitrogen sources influence the growth and biomass generation of *Phaeocystis globosa*, especially in view of the increased nutrient enrichment occurring in many coastal waters. Conversely, studying the nitrogen uptake dynamics of *P. globosa* was secondary to the objective of investigating the role of nitrogen source in colony development, and nitrogen data were not collected during the time-course measurements. However, estimates from rates of particulate matter generation suggest that N-limitation is unlikely. For example, using the ratio of chlorophyll *a*:PON determined by Wang *et al.* (2010), 2.19  $\mu\text{mol N}$  is removed for every 1  $\mu\text{g}$  chlorophyll *a* generated.



**Fig. 4.** Colony diameter and cell abundance (cells per colony) in cultures of *Phaeocystis globosa* using nitrate (open circles), ammonium (solid circles) and urea (open triangle). Colony formation of CCMP 1528 and 629 within non-nitrate treatments was observed only at the beginning of the experiment.



**Fig. 5.** Relationship between cell abundance per colony and colony diameter of *Phaeocystis globosa* CCMP 1528 (top) and 629 (bottom) when grown using different nitrogen sources (left, ammonium; right nitrate). Solid lines are linear regressions, and dotted lines are the 95% confidence intervals. N=number of colonies measured.

Therefore, in f/2 medium with an initial N concentration of 882  $\mu\text{M}$ , we would expect that 402  $\mu\text{g l}^{-1}$  chlorophyll *a* would be generated. Only in the cultures of CCMP 627 growing with nitrate did we observe chlorophyll concentrations of 400  $\mu\text{g l}^{-1}$ , suggesting that nitrogen limitation occurred infrequently. Therefore, we attribute the different response in growth and colony formation to the form of nitrogen supplied, rather than of N-limitation *per se*.

#### Nitrate uptake and colony formation

Our studies suggest strongly that nitrate favours the growth of both solitary cells and colonies of *P. globosa*. Furthermore, reduced forms of nitrogen (ammonium, urea) failed to support extensive growth in this species. Our results are consistent with previous studies that have reported *Phaeocystis* blooms in environments where nitrate was the major N source (Riegman *et al.*, 1992; Smith, 1993; Arrigo *et al.*, 1999; Lancelot *et al.*, 2007). Nitrate is the dominant form of nitrogen in the North Sea (Lancelot *et al.*, 2007), contributing more than 50% of total nitrogen. This region has been dominated by massive *Phaeocystis* blooms each spring, which are tightly coupled to nitrate loading in both space and time (Lancelot *et al.*, 2007). *Phaeocystis antarctica* blooms in the relatively deeply mixed (and nitrate replete) waters of the Ross Sea, where the  $\text{NO}_3:\text{PO}_4$  removal ratio for *P. antarctica* blooms was markedly higher than the Redfield N:P ratio of 16 (Arrigo *et al.*, 2000,

2002). Results from both of these locations suggest that *Phaeocystis* can grow well and reach high abundance using nitrate as a nitrogen source.

The investigations by Hai *et al.* (2010) presented alternative evidence that high ammonium concentrations (4.9  $\mu\text{mol l}^{-1}$ ) favour the development of *P. globosa* blooms in the upwelling waters of the south-central coast of Vietnam. The *Phaeocystis* blooms are induced and supported by offshore upwelling that brings nutrients from the deep ocean to the surface (Tang *et al.*, 2004). The upwelled waters are also in close contact with sediments, potentially providing a significant ammonium source. Hai *et al.* (2010) found that nitrate concentrations were still high (2.1  $\mu\text{mol l}^{-1}$ ) and above saturation levels during the *Phaeocystis* bloom, but that ammonium was even more elevated. It is possible that nitrate simulated colony development close to the upwelling centre and was reduced in concentration during the shoreward advection of the water; then, following significant ammonium regeneration in shallow water, nitrate uptake was inhibited by high concentrations of ammonium (Tungaraza *et al.*, 2003). In addition, the observed differences may be due to the different strains studied. In Vietnam coastal waters, the diameters of colonies of *P. globosa* reached up to 1 cm, and thousands of cells were embedded within the colony matrix (W.O.S., unpublished observations). These giant colonies may represent an entirely different biological response to the variable nitrogen sources than was observed in our cultures.

*Differences in uptake strategies between solitary and colonial cells*

Comparatively few colonies developed when ammonium or urea were provided as N sources, whereas solitary cell abundances increased with time for all three strains, irrespective of nitrogen source. Furthermore, the final biomass of CCMP 627 grown in urea was greater than that grown in nitrate, strongly suggesting that the growth of *P. globosa* solitary cells was largely independent of nitrogen source. *Phaeocystis* can remove ammonium when ammonium concentrations exceed certain concentrations (Smith, 1993; Cochlan & Bronk, 2003; Tungaraza *et al.*, 2003). Unfortunately, none of these studies differentiated between solitary and colonial cells and the nitrogen source of each form, but given *P. globosa*'s failure to form colonies in cultures supplied with reduced nitrogen, we would suggest that ammonium probably was removed by solitary and not colonial cells.

*Nitrogen and colonial architecture*

The slope of the linear log–log relationship between colony cell number and colony diameter that we observed is comparable to previous results in Rousseau *et al.* (1990) and Jakobsen & Tang (2002). In their investigations nitrate was also used as a nitrogen source. The significant difference between the slopes of the regression lines between nitrate and ammonium treatments in our experiments suggests that the cellular density in colonies (the number of cells per unit colony surface) was reduced by ammonium. It is generally considered that ammonium takes less energy to assimilate than nitrate (Syrett, 1981; Thompson *et al.*, 1989), and it is possible that the 'excess' energy of NH<sub>4</sub>-grown cells may have been used to produce more carbon-rich mucus. However, we did not find increased numbers of colonies in ammonium-based cultures; indeed, colonial growth using NH<sub>4</sub> was extremely limited. Given the large increased partitioning of carbon into the mucus, it is possible that the colonies rapidly became structurally unstable and could not maintain colonial integrity. Our data cannot explain the differences in colony architecture, and further work is necessary to provide the underlying physiological or mechanical explanation.

Generally growth of nano-phytoplankton in the ocean is thought to be based largely on ammonium, with nitrate being relatively more important for net phytoplankton (Malone, 1980; Probyn & Painting, 1985). In our study solitary cells with diameters of a few micrometres were able to remove either nitrate or ammonium, whereas

colonies utilized nitrate. This may partly explain why colonies were present only at the beginning of experiments in ammonium-based cultures. Given that individual cells are embedded in the mucoid matrix, it is possible that ammonium diffusion through the mucus was too slow to support growth (perhaps due to the negative charge of the ion). However, Ploug *et al.* (1999) suggested that the mucus was highly permeable to the inorganic substances they tested, which would argue against the possible ammonium limitation via diffusion. Similarly, in the ammonium-replete cultures reported by Riegman *et al.* (1992), cells of *P. globosa* were solitary, whereas they were colonial when supplied nitrate. Although urea concentrations decreased rapidly during both diatom and *Phaeocystis* blooms (Tungaraza *et al.*, 2003), there was no direct evidence that urea was directly used as an N substrate by *Phaeocystis* (Sanderson *et al.*, 2008, Bradley *et al.*, 2010). Only solitary cells of *P. globosa* removed urea in our cultures, strongly suggesting that solitary cells were responsible for the decrease in urea concentration during blooms of *Phaeocystis*.

*Nitrate load within marine systems dominated by Phaeocystis*

Generally colony formation is assumed to represent a potential energy loss, which inevitably results in reduced growth rates of the colonial form (Jakobsen & Tang, 2002). However, colonial and solitary cells have been shown in other studies to have similar growth rates (Hamm, 2000; Jakobsen & Tang, 2002), suggesting that colonial mucus production does not represent a significant energy drain for *P. globosa* nor result in a reduced growth rate of colonial cells. In our cultures, the growth rates of colonial cells were even higher than those of solitary cells, similar to the finding of Shields & Smith (2009) and Veldhuis *et al.* (2005). Because the different morphotypes of *Phaeocystis* have very different trophic roles (e.g. Hamm *et al.*, 2001; Smith *et al.*, 2007), nitrate inputs that induce changes in altered partitioning between solitary and colonial forms will inevitably affect many of the ecological processes in *Phaeocystis*-dominated systems. Colony formation may protect colonial cells against viral infection, bacterial degradation and zooplankton grazing; thus, colony formation and larger size can be considered to be a defence mechanism (Hamm, 2000; Tang, 2003; Nejtgaard *et al.*, 2007). Increases in nitrate concentrations would enhance magnitude and composition of *P. globosa* blooms and result in decreased grazing pressure and, combined with the presumed higher sinking rates of colonies, increased inputs to the benthos (Hamm &

Rousseau, 2003; Reigstad & Wassmann, 2007) and decreased POC retention within the pelagic food web (Hamm, 2000).

Finally, it should be noted that the variations we found among the three strains were substantial, both with respect to colony formation and nitrogen assimilation. While this is not completely unexpected, given that clonal variations have been previously observed (van Rijssel *et al.*, 2000; Long *et al.*, 2007; Mills *et al.*, 2010), the magnitude of these variations was surprising large (e.g. the differences in final standing stocks that were independent of the nitrogen source). While the clones appear morphologically similar, it is possible that significant genetic differences exist among clones that result in substantial ecological and geographic separation. Similar findings for oceanic prochlorophytes have been observed (Johnson *et al.*, 2006) and we know of no reason why similar niche differentiation should not occur in haptophytes. Further detailed molecular analysis and linkage to ecological responses in this critical species would allow a greater understanding of the local responses to nitrogen inputs.

### Acknowledgements

We thank Dr Kam Tang for technical assistance and constructive comments. The authors are also grateful to three anonymous reviewers whose invaluable comments and advice greatly improved the final form of this manuscript. This study was supported by the National Basic Research Program of China 2011CB403603 (X.W.), 2010CB428702 (Y.W.) and U.S. National Science Foundation OCE-0850910 (W.O.S.).

### References

- ARRIGO, K.R., DiTULLIO, G.R., DUNBAR, R.B., ROBINSON, D.H., VAN WOERT, M., WORTHEN, D.L. & LIZOTTE, M.P. (2000). Phytoplankton taxonomic variability in nutrient utilization and primary production in the Ross Sea. *J. Geophys. Res.*, **105**: 8827–8845.
- ARRIGO, K.R., DUNBAR, R.B., LIZOTTE, M. & ROBINSON, D.H. (2002). Taxon specific differences in C/P and N/P drawdown for phytoplankton in the Ross Sea, Antarctica. *Geophys. Res. Lett.*, **29**: 1938.
- ARRIGO, K.R., ROBINSON, D.H., WORTHEN, D.L., DUNBAR, R.B., DiTULLIO, G.R., VAN WOERT, M. & LIZOTTE, M.P. (1999). Phytoplankton community structure and the drawdown of nutrients and CO<sub>2</sub> in the Southern Ocean. *Science*, **283**: 365–367.
- BÄTJE, M. & MICHAELIS, H. (1986). *Phaeocystis pouchetii* blooms in the east Frisian coastal waters (German Bight, North Sea). *Mar. Biol.*, **93**: 21–27.
- BERMAN, T. & BRONK, D.A. (2003). Dissolved organic nitrogen: a dynamic participant in aquatic ecosystems. *Aquat. Microb. Ecol.*, **31**: 279–305.
- BRADLEY, P.B., SANDERSON, M.P., NEJSTGAARD, J.C., SAZHIN, A.F., FRISCHER, M.E., KILLBERG-THORESON, L.M., VERITY, P.G., CAMPBELL, L. & BRONK, D.A. (2010). Nitrogen uptake by phytoplankton and bacteria during an induced *Phaeocystis pouchetii* bloom, measured using size fractionation and flow cytometric sorting. *Aquat. Microb. Ecol.*, **61**: 89–104.
- BRUSSAARD, C.P.D., KUIPERS, B. & VELDHUIS, M.J.W. (2005). A mesocosm study of *Phaeocystis globosa* population dynamics I. Regulatory role of viruses in bloom control. *Harmful Algae*, **4**: 859–874.
- CADÉE, G.C. & HEGEMAN, J. (2002). Phytoplankton in the Marsdiep at the end of the 20th century: 30 years monitoring biomass, primary production, and *Phaeocystis* blooms. *J. Sea Res.*, **48**: 97–110.
- CHEN, Y.Q., WANG, N., ZHANG, P., ZHOU, H. & QU, L.H. (2002). Molecular evidence identifies bloom-forming *Phaeocystis* (Prymnesiophyta) from coastal waters of southeast China as *Phaeocystis globosa*. *Biochem. Syst. Ecol.*, **30**: 15–22.
- COCHLAN, W.P. & BRONK, D.A. (2001). Nitrogen uptake kinetics in the Ross Sea, Antarctica. *Deep Sea Res.*, **48**: 4127–4153.
- COCHLAN, W.P. & BRONK, D.A. (2003). Effects of ammonium on nitrate utilization in the Ross Sea, Antarctica: implications for f-ratio estimates. In *Biogeochemistry of the Ross Sea* (DiTullio, G.R. & Dunbar, R.B., editors), 159–178. American Geophysical Union, Washington, DC.
- EPPLEY, R.W., CARLUCCI, A.F., HOLM-HANSEN, O., KIEFER, D., MCCARTHY, J.J., VENRICK, E. & WILLIAMS, P.M. (1971). Phytoplankton growth and composition in shipboard cultures supplied with nitrate, ammonium, or urea as the nitrogen source. *Limnol. Oceanogr.*, **16**: 741–751.
- GIESKES, W.W.C., LETERME, S.C., PELETIER, H., EDWARDS, M. & REID, P.C. (2007). *Phaeocystis* colony distribution in the North Atlantic Ocean since 1948, and interpretation of long-term changes in the *Phaeocystis* hotspot in the North Sea. *Biogeochemistry*, **83**: 49–60.
- GLIBERT, P.M. & BURKHOLDER, J.M. (2006). The complex relationships between increasing fertilization of the earth, coastal eutrophication and proliferation of harmful algal blooms. In *Ecology of Harmful Algae* (Granéli, E. & Turner, J., editors), 341–354. Springer, New York, NY.
- GUILLARD, R.R.L. & RYTHER, J.J. (1962). Studies of marine planktonic diatoms. *Can. J. Microbiol.*, **8**: 229–239.
- HAI, D.N., LAM, N.N. & DIPPNER, J.W. (2010). Development of *Phaeocystis globosa* blooms in the upwelling waters of the South Central coast of Viet Nam. *J. Mar. Syst.*, **83**: 253–261.
- HAMM, C.E. (2000). Architecture, ecology and biogeochemistry of *Phaeocystis* colonies. *J. Sea Res.*, **43**: 307–315.
- HAMM, C.E. & ROUSSEAU, V. (2003). Composition, assimilation and degradation of *Phaeocystis globosa*-derived fatty acids in the North Sea. *J. Sea Res.*, **50**: 271–283.
- HAMM, C.E., REIGSTAD, M., WEXELS-RISER, C., MÜLEBACH, A. & WASSMANN, P. (2001). On the trophic fate of *Phaeocystis pouchetii*. VII. Sterols and fatty acids reveal sedimentation of *Phaeocystis*-derived organic matter via krill fecal strings. *Mar. Ecol. Progr. Ser.*, **209**: 55–69.
- HAMM, C.E., SIMSON, D.A., MERKEL, R. & SMETACEK, V. (1999). Colonies of *Phaeocystis globosa* are protected by a thin but tough skin. *Mar. Ecol. Progr. Ser.*, **187**: 101–111.
- JACOBSEN, A., EGGE, J.K. & HEIMDAL, B. (1995). Effects of increased concentration on nitrate and phosphate during a spring-bloom experiment in mesocosm. *J. Exp. Mar. Biol. Ecol.*, **187**: 239–251.
- JAKOBSEN, H.H. & TANG, K.W. (2002). Effects of protozoan grazing on colony formation in *Phaeocystis globosa* (Prymnesiophyceae) and the potential costs and benefits. *Aquat. Microb. Ecol.*, **27**: 261–273.
- JOHNSON, Z.I., ZINSER, E.R., COE, A., McNULTY, N.P., WOODWARD, E.M.S. & CHISHOLM, S.W. (2006). Niche partitioning among *Prochlorococcus* ecotypes along ocean-scale environmental gradients. *Science*, **311**: 1737–1740.
- L'HELGUEN, S., MAGUER, J.F. & CARADEC, J. (2008). Inhibition kinetics of nitrate uptake by ammonium in size-fractionated oceanic phytoplankton communities: implications for new production and f-ratio estimates. *J. Plankton Res.*, **30**: 1179–1188.

- LANCELOT, C. & MATHOT, S. (1985). Biochemical fractionation of primary production by phytoplankton in Belgian coastal waters during short- and long-term incubations with  $^{14}\text{C}$ -bicarbonate. II. *Phaeocystis pouchetii* colonial population. *Mar. Biol.*, **86**: 227–232.
- LANCELOT, C., BILLEN, G., SOURNIA, A., WEISSE, T., COLIJN, F., VELDHIJS, M., DAVIES, A. & WASSMAN, P. (1987). *Phaeocystis* blooms and nutrient enrichment in the continental coastal zones of the North Sea. *Ambio*, **16**: 38–46.
- LANCELOT, C., KELLER, M.D., ROUSSEAU, V., SMITH, W.O. JR & MATHOT, S. (1998). Autoecology of the marine haptophyte *Phaeocystis* sp. In *Physiological Ecology of Harmful Algal Blooms* (Anderson, D.M., Cembella, A.D. & Hallegraeff, G.M., editors), 209–224. Springer Verlag, Berlin.
- LANCELOT, C., SPITZ, Y., GYPENS, N., RUDDICK, K., BECQUEVORT, S., ROUSSEAU, V., LACROIX, G. & BILLEN, G. (2005). Modelling diatom–*Phaeocystis* blooms and nutrient cycles in the Southern Bight of the North Sea: the MIRO model. *Mar. Ecol. Progr. Ser.*, **289**: 63–78.
- LANCELOT, C., GYPENS, N., BILLEN, G., GARNIER, J. & ROUBEIX, V. (2007). Testing an integrated river-ocean mathematical tool for linking marine eutrophication to land use: the *Phaeocystis* dominated Belgian coastal zone (Southern North Sea) over the past 50 years. *J. Mar. Syst.*, **64**: 216–228.
- LARSEN, A., FLATEN, G.A.F., SANDAA, R.A., CASTBERG, T., THYRHAUG, R., ERGA, S.R., JACQUET, S. & BRATBAK, G. (2004). Spring phytoplankton bloom dynamics in Norwegian coastal waters: microbial community succession and diversity. *Limnol. Oceanogr.*, **49**: 180–190.
- LOMAS, M.W., TRICE, T.M., GLIBERT, P.M., BRONK, D.A. & MCCARTHY, J.J. (2002). Temporal and spatial dynamics of urea uptake and regeneration rates and concentrations in Chesapeake Bay. *Estuaries*, **25**: 469–482.
- LONG, J.D., FRISCHER, M.E. & ROBERTSON, C.Y. (2007). A *Phaeocystis globosa* bloom associated with upwelling in the subtropical South Atlantic Bight. *J. Plankton Res.*, **29**: 769–774.
- MACISAAC, J.J. & DUGDALE, R.C. (1969). The kinetics of nitrate and ammonia uptake by natural populations of marine phytoplankton. *Deep Sea Res.*, **16**: 45–57.
- MALONE, T.C. (1980). Size-fractionated primary productivity of marine phytoplankton. In *Primary Productivity in the Sea* (Falkowski, P.G., editor), 301–319. Plenum Press, New York.
- MATHOT, S., SMITH, W.O. JR, CARLSON, C.A. & GARRISON, D.L. (2000). Estimate of *Phaeocystis* sp. carbon biomass: methodological problems related to the mucilaginous nature of the colonial matrix. *J. Phycol.*, **36**: 1049–1056.
- MEDLIN, L., ZINGONE, A. (2007) A taxonomic review of the genus *Phaeocystis*. *Biogeochemistry*, **83**: 3–18.
- MILLS, M.M., KROPUENSKA, L.R., VAN DIJKEN, G.L., ALDERKAMP, A.C., BERG, G.M., DALE, H., ROBINSON, D.H., WELSCHMEYER, N.A. & ARRIGO, K.R. (2010). Photophysiology in two southern ocean phytoplankton taxa: photosynthesis of *Phaeocystis antarctica* (Prymnesiophyceae) and *Fragilariopsis cylindrus* (Bacillariophyceae) under simulated mixed-layer irradiance. *Limnol. Oceanogr.*, **46**: 1114–1127.
- MUGGLI, D.L. & SMITH, W.O. JR (1993). Regulation of nitrate and ammonium uptake in the Greenland Sea. *Mar. Biol.*, **115**: 199–208.
- NEJSTGAARD, J.C., TANG, K.W., STEINKE, M., DUTZ, J., KOSKI, M., ANTAJAN, E. & LONG, J.E. (2007). Zooplankton grazing on *Phaeocystis*: a quantitative review and future challenges. *Biogeochemistry*, **83**: 147–172.
- PARSONS, T.R., MAITA, Y. & LALLI, C. (1984). *A Manual of Chemical and Biological Methods for Seawater Analysis*. Oxford: Pergamon Press.
- PEPERZAK, L. & POELMAN, M. (2008). Mass mussel mortality in the Netherlands after a bloom of *Phaeocystis globosa* (Prymnesiophyceae). *J. Sea Res.*, **60**: 220–222.
- PLoug, H., STOLTE, W. & JØRGENSEN, B.B. (1999). Diffusive boundary layers of the colony-forming plankton alga *Phaeocystis* sp.: implications for nutrient uptake and cellular growth. *Limnol. Oceanogr.*, **44**: 1959–1967.
- PROBYN, T.A. & PAINTING, S.J. (1985). Nitrogen uptake by size-fractionated phytoplankton populations in Antarctic surface waters. *Limnol. Oceanogr.*, **30**: 1327–1332.
- QI, Y., CHEN, J., WANG, Z., XU, N., WANG, Y., SHEN, P., LU, S. & HODGKISS, I.J. (2004). Some observations on harmful algal bloom (HAB) events along the coast of Guangdong, southern China in 1998. *Hydrobiologia*, **512**: 209–214.
- REIGSTAD, M. & WASSMANN, P. (2007). Does *Phaeocystis* spp. contribute significantly to vertical export of biogenic matter?. *Biogeochemistry*, **83**: 217–234.
- RIEGMAN, R., NOORDELOOS, A.M. & CADÉE, G.C. (1992). *Phaeocystis* blooms and eutrophication of the continental coastal zones of the North Sea. *Mar. Biol.*, **112**: 479–484.
- ROUSSEAU, V., MATHOT, S. & LANCELOT, C. (1990). Calculating carbon biomass of *Phaeocystis* sp. from microscopic observations. *Mar. Biol.*, **107**: 305–314.
- ROUSSEAU, V., VAULOT, D., CASOTTI, R., CARIU, V., LENZ, J., GUNKEL, J. & BAUMANN, M. (1994). The life cycle of *Phaeocystis* (Prymnesiophyceae): evidence and hypotheses. *J. Mar. Syst.*, **5**: 23–39.
- SANDERSON, M.P., BRONK, D.A., NEJSTGAARD, J.C., VERITY, P.G., SAZHIN, A.F. & FRISCHER, M.E. (2008). Phytoplankton and bacterial uptake of inorganic and organic nitrogen during an induced bloom of *Phaeocystis pouchetii*. *Aquat. Microb. Ecol.*, **51**: 153–168.
- SCHOEMANN, V., WOLLAST, R., CHOU, L. & LANCELOT, C. (2001). Effects of photosynthesis on the accumulation of Mn and Fe by *Phaeocystis* colonies. *Limnol. Oceanogr.*, **46**: 1065–1076.
- SCHOEMANN, V., BECQUEVORT, S., STEFELS, J., ROUSSEAU, V. & LANCELOT, C. (2005). *Phaeocystis* blooms in the global ocean and their controlling mechanisms: a review. *J. Sea Res.*, **53**: 43–66.
- SHEN, P., VAN RIJSEL, M., WANG, Y., LU, S.H., CHEN, J.F. & QI, Y. (2004). Toxic *Phaeocystis globosa* strains from China grow at remarkably high temperatures. In *Harmful Algae, 2002* (Steidinger, K.A., Landsberg, J.H., Tomas, C.R. & Vargo, G.A., editors), 396–398. Florida Fish and Wildlife Conservation Commission, Florida Institute of Oceanography and Intergovernmental Oceanographic Commission of UNESCO, St. Petersburg.
- SHIELDS, A.R. & SMITH, W.O. JR (2009). Size-fractionated photosynthesis/irradiance relationships during *Phaeocystis antarctica*-dominated blooms in the Ross Sea, Antarctica. *J. Plankton Res.*, **31**: 709–712.
- SMITH, W.O. JR (1993). Nitrogen uptake and new production in the Greenland Sea: the spring *Phaeocystis* bloom. *J. Geophys. Res.*, **98**: 4681–4688.
- SMITH, W.O. JR, CARLSON, C.A., DUCKLOW, H.W. & HANSELL, D.A. (1998). Growth dynamics of *Phaeocystis antarctica*-dominated plankton assemblages from the Ross Sea. *Mar. Ecol. Progr. Ser.*, **168**: 229–244.
- SMITH, W.O. JR, AINLEY, D.G. & CATTANEO-VIETTI, R. (2007). Trophic interactions within the Ross Sea continental shelf ecosystem. *Phil. Trans. R. Soc., B*, **362**: 95–111.
- SOLOMON, C.M., COLLIER, J.L., BERG, G.M. & GLIBERT, P.M. (2010). Role of urea in microbial metabolism in aquatic systems: a biochemical and molecular review. *Aquat. Microb. Ecol.*, **59**: 67–88.
- STOLTE, W. & RIEGMAN, R. (1995). The effect of phytoplankton cell size on transient state nitrate and ammonium uptake kinetics. *Microbiology*, **141**: 1221–1229.
- STOLTE, W. & RIEGMAN, R. (1996). A model approach for size-selective competition of marine phytoplankton for fluctuating nitrate and ammonium. *J. Phycol.*, **32**: 732–740.
- SYRETT, P.J. (1981) Nitrogen metabolism of microalgae. In *Physiological Bases of Phytoplankton Ecology* (Platt, T., editor), 182–210. Canadian Bulletin of Fisheries and Aquatic Sciences, vol. 210. Ottawa.
- TANG, D.L., KAWAMURA, H., HAI, D.-N. & TAKAHASHI, W. (2004). Remote sensing oceanography of a harmful algal bloom (HAB)

- off the coast of southeastern Vietnam. *J. Geophys. Res.*, **109**: C03014, doi: 10.1029/2003JC002045.
- TANG, K.W. (2003). Grazing and colony size development in *Phaeocystis globosa* (Prymnesiophyceae): the role of a chemical signal. *J. Plankton Res.*, **25**: 831–842.
- TANG, K.W., SMITH, W.O. JR, ELLIOTT, D.T., SHIELDS, A.R. (2008) Colony size of *Phaeocystis antarctica* (Prymnesiophyceae) as influenced by zooplankton grazers. *J. Phycol.*, **44**: 1372–1378.
- THANH, T.D., SAITO, Y., HUY, D.V., LAP, N.V., OANH, T.T.K. & TATEISHI, M. (2004). Regimes of human and climate impacts on coastal changes in Vietnam. *Reg. Environ. Change*, **4**: 49–62.
- THOMPSON, P.A., LEVASSEUR, M.E. & HARRISON, P.J. (1989). Light-limited growth on ammonium vs. nitrate: what is the advantage for marine phytoplankton?. *Limnol. Oceanogr.*, **34**: 1014–1024.
- TUNGARAZA, C., ROUSSEAU, V., BRION, N., LANCELOT, C., GICHUKI, J., BAUYENS, W. & GOEYENS, L. (2003). Contrasting nitrogen uptake by diatom and *Phaeocystis*-dominated phytoplankton assemblages in the North Sea. *J. Exp. Mar. Biol. Ecol.*, **292**: 19–41.
- VAN RIJSEL, M., ALDERKAMP, A.C., NEJSTGAARD, J.C., SAZHIN, A.F. & VERITY, P.G. (2007). Haemolytic activity of living *Phaeocystis pouchetii* during mesocosm blooms. *Biogeochemistry*, **83**: 189–200.
- VAN RIJSEL, M., JANSE, I., NOORDKAMP, D.J.B. & GIESKES, W.W.C. (2000). An inventory of factors that affect polysaccharide production by *Phaeocystis globosa*. *J. Sea Res.*, **43**: 297–306.
- WANG, X., TANG, K.W., WANG, Y. & SMITH, W.O. JR (2010). Temperature effects on growth, colony development and carbon partitioning in three *Phaeocystis* species. *Aquat. Biol.*, **9**: 239–249.
- VELDHUIS, M.J.W. & ADMIRAAL, W. (1987). Influence of phosphate depletion on the growth and colony formation of *Phaeocystis pouchetii*. *Mar. Biol.*, **95**: 47–54.
- VELDHUIS, M.J.W., BRUSSAARD, C.P.D. & NOORDELOOS, A.A.M. (2005). Living in a *Phaeocystis* colony: a way to be a successful algal species. *Harmful Algae*, **4**: 841–858.
- VELDHUIS, M.J.W., COLIJN, F. & VENEKAMP, L.A.H. (1986). The spring bloom of *Phaeocystis pouchetii* (Haptophyceae) in Dutch coastal waters. *Neth. J. Sea Res.*, **20**: 37–48.
- ZAR, J.H. (1984). *Biostatistical Analysis* 2nd ed. Upper Saddle River, NJ: Prentice-Hall.
- ZHANG, J. (2000). Evidence of trace metal limited photosynthesis in eutrophic estuarine and coastal waters. *Limnol. Oceanogr.*, **45**: 1871–1878.