

2015

Dark chiral symmetry breaking and the origin of the electroweak scale

Christopher D. Carone
College of William and Mary


Raymundo Ramos
College of William and Mary

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Carone, C. D., & Ramos, R. (2015). Dark chiral symmetry breaking and the origin of the electroweak scale. *Physics Letters B*, 746, 424-429.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



Dark chiral symmetry breaking and the origin of the electroweak scale



Christopher D. Carone*, Raymundo Ramos

High Energy Theory Group, Department of Physics, College of William and Mary, Williamsburg, VA 23187-8795, United States

ARTICLE INFO

Article history:

Received 15 March 2015

Received in revised form 14 May 2015

Accepted 17 May 2015

Available online 21 May 2015

Editor: B. Grinstein

ABSTRACT

We study a classically scale-invariant model in which strong dynamics in a dark sector sets the scale of electroweak symmetry breaking. Our model is distinct from others of this type that have appeared in the recent literature. We show that the Higgs sector of the model is phenomenologically viable and that the spectrum of dark sector states includes a partially composite dark matter candidate.

© 2015 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>). Funded by SCOAP³.

1. Introduction

The Lagrangian of the standard model has precisely one dimensionful parameter, the squared mass of the Higgs doublet field. This mass sets the scale of electroweak symmetry breaking, which is communicated to the standard model fermions via their Yukawa couplings. The origin and stability of the hierarchy between the electroweak scale and the Planck scale have motivated many of the leading proposals for physics beyond the standard model. In this Letter, we study the phenomenology of a specific model in which the Higgs mass squared arises as a result of strong dynamics in a dark sector. Other models of this type have been discussed in the recent literature [1,2]; we explain how our model differs from those proposals below.

It is well known that the Yukawa coupling between a scalar ϕ and fermions can lead to a linear term in the scalar potential if the fermions condense. Such a term alters the potential so that the scalar develops a vacuum expectation value (vev). If the scalar squared mass term is absent, then the scale of the scalar vev is set entirely by that of the strong dynamics that produced the condensate. If these fields carry electroweak quantum numbers, then electroweak symmetry will be spontaneously broken. A simple model based on this idea was proposed by Carone and Georgi in Ref. [3]. In this Letter, we consider a similar theory in which the scalar and fermions in question do not carry electroweak charges. The vev of ϕ does not break electroweak symmetry, but provides an origin for the Higgs squared mass via the Higgs portal coupling $\lambda_p \phi^\dagger \phi H^\dagger H$. As long as λ_p has the appropriate sign, electroweak

symmetry breaking is triggered at a scale set by the strong dynamics of the dark sector.

The choice of a classically scale-invariant scalar potential can be justified by various arguments. We place them in two categories:

1. *The model is tuned.* Dimensionful parameters might not assume natural values as a consequence of the probability distribution over the string landscape, which is poorly understood. If one takes this point of view, it is not unreasonable to consider extensions of the standard model that are designed to address its deficiencies (for example, extensions that provide for viable dark matter physics) that appear tuned but are parametrically simple and can be easily tested in experiment. Our model is of this type and could easily be ruled out (or supported) by upcoming dark matter searches.

2. *The model is not tuned.* If there are no physical mass scales between the weak and Planck scales, then the only possible source of a Higgs quadratic divergences is from the cut off of the theory. Although field theoretic completions to low-energy effective theories lead generically to quadratic divergences proportional to the square of the cutoff [4], this may not be the case for quantum gravitational physics at the Planck scale [5]. As argued in Ref. [6], a spacetime description itself may break down at this scale and one's intuition based on quantum field theories may be flawed. If one takes this point of view, it is not unreasonable to assume that a Higgs mass generated via dimensional transmutation in the infrared is only multiplicatively renormalized [7] and to explore the phenomenological consequences. A significant number of recent papers have adopted this perspective [1,2,8,9].

The model we propose has a dark sector $SU(2)_L \times SU(2)_R$ chiral symmetry that is spontaneously broken by a fermion condensate triggered by strong dynamics. An $SU(2)_D$ subgroup of the global symmetry is gauged, and the dark fermions have Yukawa couplings to a scalar that is a doublet under this gauge symmetry. The

* Corresponding author.

E-mail addresses: cdcaro@wm.edu (C.D. Carone), raramos@email.wm.edu (R. Ramos).

dark sector would be an electroweak neutral clone of the model in Ref. [3], except that a U(1) gauge factor is replaced by a discrete subgroup to avoid a massless dark photon. The presence of an $SU(2)_D$ -doublet scalar immediately distinguishes the model from most related ones in the literature which employ a dark singlet to communicate dark sector strong dynamics through the Higgs portal [1]. We note that the model of Ref. [2] has the same dark sector global chiral symmetry as ours, but does not gauge any subgroup. This leads to a different particle spectrum and phenomenology. We also utilize a non-linear chiral Lagrangian approach, familiar from the study of technicolor and QCD, which provides a convenient framework for the systematic description of dark sector phenomenology at low energies.

Our paper is organized as follows. In the next section we define the model. In Section 3, we consider phenomenological constraints. In Section 4, we study the relic density and direct detection of the dark matter candidate in the model, which is a partially composite dark sector state. In Section 5, we present our conclusions.

2. The model

The gauge group of the model is $G_{\text{SM}} \times SU(N) \times SU(2)_D$. The first factor refers to the standard model gauge group, while the second is responsible for confinement in the dark sector. The G_{SM} singlet fields (which we will call the dark sector, henceforth) are: a complex $SU(2)_D$ -doublet scalar ϕ , a left-handed $SU(2)_D$ -doublet fermion $\Upsilon_L \equiv (p_L, m_L)^T$ and two right-handed singlet fermions p_R and m_R . The fermions transform in the fundamental representation of the $SU(N)$ group. The field content is analogous to that of the technicolor model in Ref. [3] with $SU(2)_W$ replaced by $SU(2)_D$ and $U(1)_Y$ replaced by a Z_3 factor. As we will see below, the latter choice is the simplest way to preserve a convenient analogy between the two theories while also eliminating an unwanted massless gauge field. The dark sector has a global $SU(2)_L \times SU(2)_R$ chiral symmetry that is spontaneously broken when the dark fermions condense

$$\langle \bar{p} p + \bar{m} m \rangle \approx 4\pi f^3, \quad (2.1)$$

where f is the dark pion decay constant. We refer to the unbroken $SU(2)$ subgroup of the global symmetry as dark isospin. Spontaneous chiral symmetry breaking results in an isotriplet of dark pions

$$\Pi = \sum_{a=1}^3 \pi^a \frac{\sigma^a}{2}, \quad (2.2)$$

where σ^a are the Pauli matrices. As in the chiral Lagrangian approach of Ref. [3], we adopt a nonlinear representation

$$\Sigma = \exp(2i\Pi/f), \quad (2.3)$$

which transforms under the global chiral symmetry as $\Sigma \rightarrow L\Sigma R^\dagger$, where L and R are the transformation matrices for $SU(2)_L$ and $SU(2)_R$, respectively. It will be convenient to define the following four-by-four matrix field

$$\Phi \equiv (i\sigma^2 \phi^* | \phi), \quad (2.4)$$

and the nonlinear field redefinition

$$\Phi = \frac{\sigma + f'}{\sqrt{2}} \Sigma' \quad (2.5)$$

with $\Sigma' = \exp(2i\Pi'/f')$. The kinetic terms for Φ and Σ are

$$\begin{aligned} \mathcal{L}_{KE} &= \frac{1}{2} \text{tr} \left(D_\mu \Phi^\dagger D^\mu \Phi \right) + \frac{f^2}{4} \text{tr} \left(D_\mu \Sigma^\dagger D^\mu \Sigma \right) \\ &= \frac{1}{2} \partial_\mu \sigma \partial^\mu \sigma + \frac{f^2}{4} \text{tr} \left(D_\mu \Sigma^\dagger D^\mu \Sigma \right) \\ &\quad + \frac{(\sigma + f')^2}{4} \text{tr} \left(D_\mu \Sigma'^\dagger D^\mu \Sigma' \right). \end{aligned} \quad (2.6)$$

Here $D_\mu = \partial_\mu - ig_D A_\mu^a \frac{\sigma^a}{2}$, where A_μ^a is the $SU(2)_D$ gauge field. Study of the terms quadratic in the fields allows one to identify an unphysical linear combination of fields Π_u that becomes the longitudinal component of A_μ^a , and an orthogonal state π_p that is physical:

$$\pi_u = \frac{f\Pi + f'\Pi'}{\sqrt{f^2 + f'^2}}, \quad (2.7)$$

$$\pi_p = \frac{-f'\Pi + f\Pi'}{\sqrt{f^2 + f'^2}}. \quad (2.8)$$

The π_p multiplet will later be identified as the dark matter candidate in the theory.

Explicit breaking of the chiral symmetry originates from the Yukawa couplings. Assuming that the fields transform under the Z_3 symmetry as

$$\Upsilon_L \rightarrow \Upsilon_L, \quad \phi \rightarrow \omega \phi, \quad p_R \rightarrow \omega p_R, \quad m_R \rightarrow \omega^2 m_R, \quad (2.9)$$

where $\omega^3 = 1$, we find that the Yukawa couplings are given as in Ref. [3] by

$$-\mathcal{L}_y = y_+ \bar{\Upsilon}_L \tilde{\phi} p_R + y_- \bar{\Upsilon}_L \phi m_R + \text{h.c.} \quad (2.10)$$

Defining $\Upsilon_R \equiv (p_R, m_R)$ and the matrix $Y \equiv \text{diag}(y_+, y_-)$ this may be re-expressed as

$$-\mathcal{L}_y = \bar{\Upsilon}_L \Phi Y \Upsilon_R + \text{h.c.}, \quad (2.11)$$

which implies that we may treat (ΦY) as a chiral-symmetry-breaking spurion with the transformation property

$$(\Phi Y) \rightarrow L(\Phi Y)R^\dagger. \quad (2.12)$$

The lowest order term in the chiral Lagrangian that involves (ΦY) is

$$\mathcal{L} = c_1 4\pi f^3 \text{tr}(\Phi Y \Sigma^\dagger) + \text{h.c.} \quad (2.13)$$

where c_1 is expected to be of order unity by naive dimensional analysis [10]. This term determines the physical dark pion mass

$$m_\pi^2 = 2c_1 \sqrt{2} \frac{4\pi f}{f'} (f^2 + f'^2) y, \quad (2.14)$$

where $y \equiv (y_+ + y_-)/2$, as well as a linear term in the scalar potential

$$V_y(\sigma) = -8\sqrt{2}\pi c_1 f^3 y \sigma. \quad (2.15)$$

This term sets the scale of the dark scalar vev, which determines the induced mass term for the standard model Higgs doublet H via a coupling in the potential $V = V_0 + V_y$, where V_0 represents the scale-invariant terms:

$$V_0(\phi, H) = \frac{\lambda}{2} (H^\dagger H)^2 - \lambda_p (H^\dagger H) (\phi^\dagger \phi) + \frac{\lambda_\phi}{2} (\phi^\dagger \phi)^2. \quad (2.16)$$

In the ultraviolet (UV), before the dark fermions have condensed, vacuum stability of Eq. (2.16) requires that

$$\lambda > 0 \quad \text{and} \quad \lambda \lambda_\phi > \lambda_p^2. \quad (2.17)$$

Noting that $\phi^\dagger\phi = \text{tr}(\Phi^\dagger\Phi)/2 = (\sigma + f')^2/2$ and working in unitary gauge where $H = [0, (v+h)/\sqrt{2}]^T$, the potential may be re-expressed as

$$V(h, \sigma) = \frac{\lambda}{8}(v+h)^4 - \frac{\lambda_p}{4}(v+h)^2(\sigma + f')^2 + \frac{\lambda_\phi}{8}(\sigma + f')^4 - 8\sqrt{2}\pi c_1 f^3 y \sigma, \quad (2.18)$$

after the dark fermions have condensed. Minimization of Eq. (2.18) leads to the following expressions for the vevs v and f' :

$$v^3 = 2 \left(\frac{\lambda_p}{\lambda} \right)^{3/2} \left(\lambda_\phi - \frac{\lambda_p^2}{\lambda} \right)^{-1} 8\sqrt{2}\pi c_1 f^3 y, \quad (2.19)$$

$$f'^3 = 2 \left(\lambda_\phi - \frac{\lambda_p^2}{\lambda} \right)^{-1} 8\sqrt{2}\pi c_1 f^3 y. \quad (2.20)$$

Of course, we fix $v = 246$ GeV to obtain the correct electroweak gauge boson masses. The mass squared matrix in the (h, σ) basis is given by

$$M^2 = \begin{pmatrix} \lambda & -\sqrt{\lambda\lambda_p} \\ -\sqrt{\lambda\lambda_p} & \frac{1}{2} \left(\frac{3\lambda_\phi\lambda}{\lambda_p} - \lambda_p \right) \end{pmatrix} v^2, \quad (2.21)$$

which is positive definite for positive couplings with $\lambda\lambda_\phi > \lambda_p^2$.

One of the eigenvalues of this matrix corresponds to the squared mass of the Higgs scalar observed at the LHC, $m_{h_0}^2 = (125.09 \text{ GeV})^2$ [11]. We call the remaining mass eigenstate field η below, and define the mixing angle θ by

$$\begin{pmatrix} \cos\theta & -\sin\theta \\ \sin\theta & \cos\theta \end{pmatrix} \begin{pmatrix} h_0 \\ \eta \end{pmatrix} = \begin{pmatrix} h \\ \sigma \end{pmatrix}. \quad (2.22)$$

The value of the angle θ is given by $\tan 2\theta = 2M_{12}^2/(M_{11}^2 - M_{22}^2)$ where M_{jk}^2 are elements of the matrix in Eq. (2.21).

With the Higgs sector of the theory now defined, we proceed in the next section to study its phenomenology. The parameters that define the Higgs sector are y_+ , y_- , c_1 , λ , λ_p , λ_ϕ , f , f' and v . We set the order-one coupling $c_1 = 1$ for definiteness, and fix values for the Yukawa couplings assuming, for simplicity, that $y_+ = y_-$. The remaining six parameters are constrained by $v = 246$ GeV, $m_{h_0} = 125.09$ GeV, and the two minimization conditions given in Eqs. (2.19), (2.20). This leaves two degrees of freedom. We choose the free parameters to be f and λ_p and map out the constraints on the model on the f - λ_p plane. This choice lends itself to easy physical interpretation since f parameterizes the scale of the dark sector strong dynamics, while λ_p indicates how strongly the dark sector couples to the visible one.

3. Phenomenological constraints

We determine whether a given point on the f - λ_p plane is allowed by imposing the following constraints:

1. *Absence of Landau poles below the Planck scale.* The presence of such a Landau pole would suggest the onset of new physics at an intermediate scale, contradicting our initial assumptions. Since a coupling that blows up will become non-perturbative first, we eliminate the possibility of Landau poles by imposing perturbativity constraints on the running couplings. For this purpose, we use the one-loop renormalization group equations (RGEs), which we provide in Appendix A. For the couplings λ , λ_p and λ_ϕ , one-loop corrections become equal in size to tree-level diagrams when, for example, $\lambda \approx 16\pi^2$; to avoid the complete breakdown of the perturbative expansion, we set a generous upper limit on each of

these couplings to be one-third of this value, $16\pi^2/3$, evaluated at all scales between m_Z and the reduced Planck mass M_* . By similar reasoning, we place upper limits on the gauge and Yukawa couplings of $4\pi/\sqrt{3}$. For our numerical results, we choose a perturbative value of the $SU(2)_D$ gauge coupling (35% of $4\pi/\sqrt{3}$ in the example we present) that is large enough to assure that the isotriplet gauge multiplet is heavier than the physical pions π_p ; this will be required for our dark matter solutions, as discussed in the next section. We take the $SU(N)$ gauge coupling to be at our perturbativity limit, $4\pi/\sqrt{3}$, at m_Z and choose $N = 4$. Since the $SU(N)$ gauge coupling is asymptotically free in our theory, it remains perturbative for all scales higher than m_Z (where we evaluate the RGEs), but it blows up quickly below m_Z , consistent with our assumption of strong dynamics in the infrared.

2. *Vacuum stability.* The presence of the non-vanishing Higgs portal coupling requires that vacuum stability be studied in the context of a two-Higgs doublet model. In two-Higgs-doublet models, one can assure that the scalar potential remains bounded from below by taking the stability conditions derived from the tree-level potential and testing whether they continue to hold for values of the couplings evaluated at higher-renormalization scales, up to the Planck scale. The justification for this approach can be found in Ref. [12]. We implement this by evaluating Eq. (2.17) using the one-loop renormalization group equations provided in the appendix. The scale at which a vacuum instability first arises depends on the free parameters of the model. A given point in the f - λ_p plane satisfies the stability criterion if we find numerically that no violation of the stability conditions arises before the Planck scale. For the dark-matter allowed points described later, the Higgs quartic coupling at the weak scale is larger than its standard model value, which contributes to the model's vacuum stability.

3. *Sufficiently standard-model-like Higgs boson.* Standard model Higgs boson couplings are altered in this model by a factor of $\cos^2\theta$, which can be no smaller than 0.7 without spoiling global fits to Higgs data [13]. The η couplings to the visible sector are like those of the Higgs but suppressed by $\sin^2\theta$; non-observation of the η in heavy Higgs search data from the LHC is assured for any η mass within the range experimentally studied, 145–710 GeV, provided that $\sin^2\theta \lesssim 0.1$ [14]. For simplicity, we require that each point in the f - λ_p plane satisfy $\sin^2\theta < 0.1$. Our final set of allowed points in parameter space discussed in Section 4 will correspond to η masses in the range 178–203 GeV, falling within the LHC range. Note that we do not consider potentially tighter mixing angle bounds on very light η from LEP2 since we will see later that this region of parameter space is excluded by our fourth constraint.

4. *Approximate chiral symmetry.* Our effective chiral Lagrangian is valid provided that sources of explicit chiral symmetry breaking are small compared to the chiral-symmetry-breaking scale $\Lambda_\chi \equiv 4\pi f$. We reject points in which the dark fermion masses m_\pm exceed one-third Λ_χ , or equivalently

$$\frac{1}{\sqrt{2}} y_\pm f' < \frac{4}{3} \pi f. \quad (3.1)$$

This assures that our initial assumption of an approximate $SU(2)_L \times SU(2)_R$ global symmetry remains valid.

We show results for a particular choice of y in Fig. 1. We have chosen to study values of f near or below the scale where the $SU(N)$ gauge coupling becomes strong. The shaded regions satisfy the first three of the constraints discussed in this section. The upper branch of points corresponds to an η heavier than the SM Higgs boson, while the lower branch corresponds to the opposite. The points which also satisfy our fourth constraint lie above the solid black line. We find that viable dark matter solutions exist only for $0.23 < y < 0.52$; we have picked an intermediate value of y as a




Fig. 1. Regions of the parameter space consistent with perturbativity and stability constraints, as well as $\sin^2 \theta < 0.1$. Points above the solid black line are consistent with approximate dark sector chiral symmetry. Two branches of points correspond to $m_\eta > m_{h_0}$ (upper branch) and $m_\eta < m_{h_0}$ (lower branch). The triangular points in the upper branch are consistent with current dark matter constraints.

representative choice. The dark matter results included in this figure will be discussed in the following section.

4. Dark matter

The dark sector of the model includes stable dark pions and baryons, provided that the pions are lighter than the baryons and the $SU(2)_D$ gauge multiplet. In the case we consider, where $y_+ = y_-$, the stabilizing symmetry is the residual dark $SU(2)$ isospin, which is non-anomalous and unbroken by higher-dimension operators (which are absent by the assumed scale invariance of the theory). If y_+ and y_- are unequal, then only the lightest of the dark pion triplet would be stable; for simplicity, we consider the degenerate case here. The dark baryons are separately stable due to a conserved dark baryon number. However, estimating the dark baryon–antibaryon annihilation cross section by scaling the analogous quantity measured experimentally in QCD, we find that dark baryon contribution to the relic density is orders of magnitude smaller than that of the π_p for the parameter choices of relevance to our analysis.¹

The Higgs sector mixing angle θ is generally small, and we can estimate the annihilation cross section by the contributions that are lowest order in $\sin \theta$: this selects $\pi_p^a \pi_p^a \rightarrow \eta \eta$, where $\pi_p = \sum_{a=1}^3 \pi_p^a \sigma^a / 2$, with π_p defined in Eq. (2.8). The $\pi_p \pi_p \eta$ and $\pi_p \pi_p h_0$ vertices originate from Eq. (2.13):

$$\mathcal{L} \supset -\frac{m_\pi^2}{2f'} (\eta \cos \theta + h_0 \sin \theta) \pi_p^a \pi_p^a. \quad (4.1)$$

The first term contributes to the annihilation process of interest via t - and u -channel pion exchange diagrams. Working in the non-relativistic limit, we find the thermally averaged annihilation cross section times velocity

¹ We will see in our figures that the relevant π_p masses are comparable to the scale $\Lambda_\chi = 4\pi f$, which we expect to be of order the dark baryon masses; however, in the effective chiral Lagrangian, the baryon mass terms involve additional unknown parameters that we may choose to assure that the dark baryons are heavier than the π_p . We check directly that the $SU(2)_D$ gauge multiplet is also heavier.




Fig. 2. Dark pion–nucleon elastic scattering cross section for the points within the dark-matter-preferred band of Fig. 1. The current bounds from LUX [18] and XENON100 [19] are also shown.

$$\langle \sigma_{\text{ann}} v \rangle = \frac{1}{16\pi} \frac{m_\pi^6}{f'^4} \left(1 - \frac{m_\pi^2}{m_\eta^2} \right)^{1/2} \left[\frac{\cos^2 \theta}{m_\eta^2 - 2m_\pi^2} \right]^2, \quad (4.2)$$

with m_π^2 given by Eq. (2.14). Using this, we calculate the freeze-out temperature T_F and the dark matter relic density by standard methods [15]. Defining $x = m_\pi/T$ and taking into account the dark sector spectrum in evaluating $g_*(x)$, the number of relativistic degrees of freedom at the temperature T , we find freeze-out temperatures near $x_F \approx 26$. The relic density is given by

$$\Omega_D h^2 \approx 3 \cdot \frac{(1.07 \times 10^9 \text{ GeV}^{-1}) x_F}{\sqrt{g_*(x_F)} M_{\text{pl}} (\sigma_{\text{ann}} v)_F}, \quad (4.3)$$

which we require to reproduce the WMAP result 0.1138 ± 0.0045 [16] within two standard deviations. In Fig. 1, the region consistent with π_p dark matter is the band of triangular points in the upper branch of otherwise allowed points. For our choice of $g_D \approx 2.54$, the $SU(2)_D$ gauge bosons are heavier than the π_p for each triangular point shown. We do not display results for other choices of y in the range $0.23 < y < 0.52$ which are similar qualitatively to the plot in Fig. 1. The main effect of increasing y over this range is to enlarge the upper branch of points while moving the solid black exclusion line upwards until it is roughly contiguous with the band preferred by dark matter considerations when $y = 0.52$.

Finally, we compare the direct detection predictions of the model with current experimental bounds. The π_p –nucleon spin-independent elastic scattering cross section is determined by t -channel h_0 and η exchange diagrams following from the vertices in Eq. (4.1). We find

$$\begin{aligned} \sigma_{\text{SI}}(\pi_p N \rightarrow \pi_p N) &= \frac{f_N^2}{16\pi} \frac{m_\pi^2 m_N^2}{v^2 f'^2} \sin^2 2\theta \frac{(m_\eta^2 - m_{h_0}^2)^2}{m_\eta^4 m_{h_0}^4} \left(\frac{m_N m_\pi}{m_N + m_\pi} \right)^2, \end{aligned} \quad (4.4)$$

where f_N parameterizes the Higgs–nucleon coupling and m_N is the nucleon mass. The value of $f_N = 0.35$ is used [17]. Results corresponding to the dark-matter-preferred band in Fig. 1 are shown in Fig. 2, which includes the current LUX [18] and XENON100 [19] bounds for comparison. All the points shown are currently allowed by direct search constraints, though they are in a region not far from the current bounds. This suggests that future results from

the LUX experiment may begin to substantially restrict the preferred dark matter parameter space of the model.

5. Conclusions

We have studied a classically scale-invariant model that provides an origin for the electroweak scale via dark sector strong dynamics. The dark sector has a structure similar to the bosonic technicolor model proposed in Ref. [3]: a fermion condensate is responsible for the instability that leads to a scalar doublet acquiring a vev. In the model of Ref. [3], the fermion condensate and the scalar vev each contribute to the breaking of electroweak symmetries. Here, the analogous fields are electroweak singlets; the scalar vev breaks a dark SU(2) gauge group and induces a mass term for the standard model Higgs doublet field via couplings in the Higgs potential. We found regions in the parameter space of the model where all the couplings can be run up to the Planck scale while remaining perturbative, where the scalar potential satisfies vacuum stability constraints, and where the Higgs boson is sufficiently standard-model-like to be consistent with existing collider data. In addition, we showed that the partially composite dark isotriplet bosons in the model can provide a viable dark matter candidate, providing the desired relic density while evading current direct detection bounds. In addition, the model predicts that the dark matter-nucleon elastic scattering cross section lies just beyond the current LUX bounds. Hence, the model may be ruled out, or given experimental support, as the LUX data set is enlarged.

Acknowledgements

This work was supported by the NSF under Grant PHY-1068008.

Appendix A. RGEs

The RGEs used in our analysis are as follows:

$$16\pi^2 \frac{d\lambda_\phi}{dt} = 4N\lambda_\phi (y_-^2 + y_+^2) - 4N(y_-^4 + y_+^4) - 9g_D^2\lambda_\phi + \frac{9}{4}g_D^4 + 4\lambda_p^2 + 12\lambda_\phi^2, \quad (\text{A.1})$$

$$16\pi^2 \frac{d\lambda}{dt} = \frac{9}{4} \left(\frac{2}{5}g_1^2g_2^2 + \frac{3}{25}g_1^4 + g_2^4 \right) - \lambda \left(\frac{9}{5}g_1^2 + 9g_2^2 \right) + 12\lambda y_t^2 + 12\lambda^2 + 4\lambda_p^2 - 12y_t^4 \quad (\text{A.2})$$

$$16\pi^2 \frac{d\lambda_p}{dt} = \left[2N(y_-^2 + y_+^2) + \frac{9}{2} \left(-\frac{1}{5}g_1^2 - g_2^2 - g_D^2 \right) + 6\lambda - 4\lambda_p + 6\lambda_\phi + 6y_t^2 \right] \lambda_p, \quad (\text{A.3})$$

$$16\pi^2 \frac{dy_t}{dt} = \left[-\frac{17}{20}g_1^2 - \frac{9}{4}g_2^2 - 8g_3^2 + \frac{9}{2}y_t^2 \right] y_t, \quad (\text{A.4})$$

$$16\pi^2 \frac{dy_-}{dt} = \left[\left(N + \frac{3}{2} \right) y_-^2 + \left(N - \frac{3}{2} \right) y_+^2 - \frac{9}{4}g_D^2 - \frac{3(N^2 - 1)}{N}g_N^2 \right] y_-, \quad (\text{A.5})$$

$$16\pi^2 \frac{dy_+}{dt} = \left[\left(N - \frac{3}{2} \right) y_-^2 + \left(N + \frac{3}{2} \right) y_+^2 - \frac{9}{4}g_D^2 - \frac{3(N^2 - 1)}{N}g_N^2 \right] y_+, \quad (\text{A.6})$$

$$16\pi^2 \frac{dg_D}{dt} = \left[\frac{N}{3} - \frac{43}{6} \right] g_D^3, \quad (\text{A.7})$$

$$16\pi^2 \frac{dg_N}{dt} = \left[\frac{4}{3} - \frac{11}{3}N \right] g_N^3, \quad (\text{A.8})$$

$$16\pi^2 \frac{dg_i}{dt} = b_i g_i^3. \quad (\text{A.9})$$

Here $t = \ln(\mu/m_Z)$, where μ is the renormalization scale, $b_i = \left(\frac{41}{10}, -\frac{19}{6}, -7 \right)$, the SU(5) normalization for the hypercharge was used and g_N is the SU(N) gauge coupling.

References

- [1] M. Heikinheimo, C. Spethmann, J. High Energy Phys. 1412 (2014) 084, arXiv:1410.4842 [hep-ph]; J. Kubo, K.S. Lim, M. Lindner, J. High Energy Phys. 1409 (2014) 016, arXiv:1405.1052 [hep-ph]; O. Antipin, M. Redi, A. Strumia, J. High Energy Phys. 1501 (2015) 157, arXiv:1410.1817 [hep-ph]; J. Kubo, K.S. Lim, M. Lindner, Phys. Rev. Lett. 113 (2014) 091604, arXiv:1403.4262 [hep-ph]; M. Holthausen, J. Kubo, K.S. Lim, M. Lindner, J. High Energy Phys. 1312 (2013) 076, arXiv:1310.4423 [hep-ph]; M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann, K. Tuominen, Mod. Phys. Lett. A 29 (2014) 1450077, arXiv:1304.7006 [hep-ph]; T. Hur, P. Ko, Phys. Rev. Lett. 106 (2011) 141802, arXiv:1103.2571 [hep-ph].
- [2] T. Hur, D.W. Jung, P. Ko, J.Y. Lee, Phys. Lett. B 696 (2011) 262, arXiv:0709.1218 [hep-ph].
- [3] C.D. Carone, H. Georgi, Phys. Rev. D 49 (1994) 1427, arXiv:hep-ph/9308205.
- [4] G. Marques Tavares, M. Schmaltz, W. Skiba, Phys. Rev. D 89 (1) (2014) 015009, arXiv:1308.0025 [hep-ph].
- [5] S. Dubovsky, V. Gorbenko, M. Mirbabayi, J. High Energy Phys. 1309 (2013) 045, arXiv:1305.6939 [hep-th].
- [6] W. Altmannshofer, W.A. Bardeen, M. Bauer, M. Carena, J.D. Lykken, J. High Energy Phys. 1501 (2015) 032, arXiv:1408.3429 [hep-ph].
- [7] W.A. Bardeen, FERMILAB-CONF-95-391-T, C95-08-27.3.
- [8] R. Hempfling, Phys. Lett. B 379 (1996) 153, arXiv:hep-ph/9604278; W.-F. Chang, J.N. Ng, J.M.S. Wu, Phys. Rev. D 75 (2007) 115016, arXiv:hep-ph/0701254 [hep-ph]; R. Foot, A. Kobakhidze, K.L. McDonald, R.R. Volkas, Phys. Rev. D 76 (2007) 075014, arXiv:0706.1829 [hep-ph]; R. Foot, A. Kobakhidze, K.L. McDonald, R.R. Volkas, Phys. Rev. D 77 (2008) 035006, arXiv:0709.2750 [hep-ph]; T. Hambye, M.H.G. Tytgat, Phys. Lett. B 659 (2008) 651, arXiv:0707.0633 [hep-ph]; S. Iso, N. Okada, Y. Orikasa, Phys. Lett. B 676 (2009) 81, arXiv:0902.4050 [hep-ph]; M. Holthausen, M. Lindner, M.A. Schmidt, Phys. Rev. D 82 (2010) 055002, arXiv:0911.0710 [hep-ph]; R. Foot, A. Kobakhidze, R.R. Volkas, Phys. Rev. D 82 (2010) 035005, arXiv:1006.0131 [hep-ph]; L. Alexander-Nunneley, A. Pilaftsis, J. High Energy Phys. 1009 (2010) 021, arXiv:1006.5916 [hep-ph]; K.A. Meissner, H. Nicolai, Phys. Lett. B 648 (2007) 312, arXiv:hep-th/0612165; Phys. Lett. B 660 (2008) 260, arXiv:0710.2840 [hep-th]; S. Iso, Y. Orikasa, PTEP, Proces. Teh. Energ. Poljopr. 2013 (2013) 023B08, arXiv:1210.2848 [hep-ph]; C. Englert, J. Jaeckel, V.V. Khoze, M. Spannowsky, J. High Energy Phys. 1304 (2013) 060, arXiv:1301.4224 [hep-ph].
- [9] J. Guo, Z. Kang, P. Ko, Y. Orikasa, arXiv:1502.00508 [hep-ph]; H. Okada, Y. Orikasa, arXiv:1412.3616 [hep-ph]; S. Benic, B. Radovcic, J. High Energy Phys. 1501 (2015) 143, arXiv:1409.5776 [hep-ph]; G.M. Pelaggi, Nucl. Phys. B 893 (2015) 443, arXiv:1406.4104 [hep-ph]; M. Lindner, S. Schmidt, J. Smirnov, J. High Energy Phys. 1410 (2014) 177, arXiv:1405.6204 [hep-ph]; K. Kannike, A. Racioppi, M. Raidal, J. High Energy Phys. 1406 (2014) 154, arXiv:1405.3987 [hep-ph]; A. Farzinnia, J. Ren, Phys. Rev. D 90 (1) (2014) 015019, arXiv:1405.0498 [hep-ph]; C. Tamarit, Phys. Rev. D 90 (5) (2014) 055024, arXiv:1404.7673 [hep-ph]; K. Allison, C.T. Hill, G.G. Ross, Phys. Lett. B 738 (2014) 191, arXiv:1404.6268 [hep-ph]; H. Davoudiasl, I.M. Lewis, Phys. Rev. D 90 (3) (2014) 033003, arXiv:1404.6260 [hep-ph]; V.V. Khoze, C. McCabe, G. Ro, J. High Energy Phys. 1408 (2014) 026, arXiv:1403.4953 [hep-ph]; S. Benic, B. Radovcic, Phys. Lett. B 732 (2014) 91, arXiv:1401.8183 [hep-ph];

- M. Hashimoto, S. Iso, Y. Orikasa, Phys. Rev. D 89 (5) (2014) 056010, arXiv:1401.5944 [hep-ph];
J. Guo, Z. Kang, arXiv:1401.5609 [hep-ph];
C.T. Hill, Phys. Rev. D 89 (7) (2014) 073003, arXiv:1401.4185 [hep-ph];
S. Abel, A. Mariotti, Phys. Rev. D 89 (12) (2014) 125018, arXiv:1312.5335 [hep-ph];
M. Hashimoto, S. Iso, Y. Orikasa, Phys. Rev. D 89 (1) (2014) 016019, arXiv:1310.4304 [hep-ph];
T.G. Steele, Z.W. Wang, D. Contreras, R.B. Mann, Phys. Rev. Lett. 112 (17) (2014) 171602, arXiv:1310.1960 [hep-ph];
E. Gabrielli, M. Heikinheimo, K. Kannike, A. Racioppi, M. Raidal, C. Spethmann, Phys. Rev. D 89 (1) (2014) 015017, arXiv:1309.6632 [hep-ph];
V.V. Khoze, J. High Energy Phys. 1311 (2013) 215, arXiv:1308.6338 [hep-ph];
A. Farzinnia, H.J. He, J. Ren, Phys. Lett. B 727 (2013) 141, arXiv:1308.0295 [hep-ph];
T. Hambye, A. Strumia, Phys. Rev. D 88 (2013) 055022, arXiv:1306.2329 [hep-ph];
C.D. Carone, R. Ramos, Phys. Rev. D 88 (2013) 055020, arXiv:1307.8428 [hep-ph];
P. Humbert, M. Lindner, J. Smirnov, arXiv:1503.03066 [hep-ph];
K. Endo, Y. Sumino, arXiv:1503.02819 [hep-ph].
- [10] A. Manohar, H. Georgi, Nucl. Phys. B 234 (1984) 189;
H. Georgi, L. Randall, Nucl. Phys. B 276 (1986) 241.
[11] G. Aad, et al., ATLAS and CMS Collaborations, arXiv:1503.07589 [hep-ex].
[12] E. Gildener, S. Weinberg, Phys. Rev. D 13 (1976) 3333;
M. Sher, Phys. Rep. 179 (1989) 273;
S. Nie, M. Sher, Phys. Lett. B 449 (1999) 89, arXiv:hep-ph/9811234.
[13] J.R. Espinosa, C. Grojean, M. Muhlleitner, M. Trott, J. High Energy Phys. 1212 (2012) 045, arXiv:1207.1717 [hep-ph].
[14] S. Chatrchyan, et al., CMS Collaboration, Eur. Phys. J. C 73 (2013) 2469, arXiv:1304.0213 [hep-ex].
[15] E.W. Kolb, M.S. Turner, Front. Phys. 69 (1990) 1.
[16] C.L. Bennett, et al., WMAP Collaboration, Astrophys. J. Suppl. Ser. 208 (2013) 20, arXiv:1212.5225 [astro-ph.CO].
[17] J.R. Ellis, A. Ferstl, K.A. Olive, Phys. Lett. B 481 (2000) 304, arXiv:hep-ph/0001005.
[18] D.S. Akerib, et al., LUX Collaboration, Phys. Rev. Lett. 112 (2014) 091303, arXiv:1310.8214 [astro-ph.CO].
[19] E. Aprile, et al., XENON100 Collaboration, Phys. Rev. Lett. 109 (2012) 181301, arXiv:1207.5988 [astro-ph.CO].