

2007

Susceptibility of salt marshes to nutrient enrichment and predator removal

LA Deegan

JL Bowen

D Drake

JW Fleeger

CT Friedrichs

Virginia Institute of Marine Science

See next page for additional authors

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Deegan, LA; Bowen, JL; Drake, D; Fleeger, JW; Friedrichs, CT; and Et al., "Susceptibility of salt marshes to nutrient enrichment and predator removal" (2007). *VIMS Articles*. 1008.

<https://scholarworks.wm.edu/vimsarticles/1008>

This Article is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Authors

LA Deegan, JL Bowen, D Drake, JW Fleegeer, CT Friedrichs, and Et al.

SUSCEPTIBILITY OF SALT MARSHES TO NUTRIENT ENRICHMENT AND PREDATOR REMOVAL

LINDA A. DEEGAN,^{1,7} JENNIFER L. BOWEN,¹ DEANNE DRAKE,¹ JOHN W. FLEEGER,² CARL T. FRIEDRICH,³
KARI A. GALVÁN,² JOHN E. HOBBIÉ,¹ CHARLES HOPKINSON,¹ D. SAMUEL JOHNSON,² J. MICHAEL JOHNSON,¹
LYNSEY E. LEMAY,³ ERIN MILLER,⁴ BRUCE J. PETERSON,¹ CHRISTIAN PICARD,¹ SALLIE SHELDON,⁴
MICHAEL SUTHERLAND,⁵ JOSEPH VALLINO,¹ AND R. SCOTT WARREN⁶

¹The Ecosystems Center, Marine Biological Laboratory, Woods Hole, Massachusetts 02543 USA

²Department of Biological Sciences, Louisiana State University, Baton Rouge, Louisiana 70803 USA

³Virginia Institute of Marine Science, Gloucester Point, Virginia 23062 USA

⁴Middlebury College, Middlebury, Vermont 05753 USA

⁵University of Massachusetts, Amherst, Massachusetts 01003 USA

⁶Department of Botany, Connecticut College, New London, Connecticut 06320 USA

Abstract. Salt marsh ecosystems have been considered not susceptible to nitrogen overloading because early studies suggested that salt marshes adsorbed excess nutrients in plant growth. However, the possible effect of nutrient loading on species composition, and the combined effects of nutrients and altered species composition on structure and function, was largely ignored. Failure to understand interactions between nutrient loading and species composition may lead to severe underestimates of the impacts of stresses. We altered whole salt marsh ecosystems (~60 000 m²/treatment) by addition of nutrients in flooding waters and by reduction of a key predatory fish, the mummichog. We added nutrients (N and P; 15-fold increase over ambient conditions) directly to the flooding tide to mimic the way anthropogenic nutrients are delivered to marsh ecosystems. Despite the high concentrations (70 mmol N/L) achieved in the water column, our annual N loadings (15–60 g N·m⁻²·yr⁻¹) were an order of magnitude less than most plot-level fertilization experiments, yet we detected responses at several trophic levels. Preliminary calculations suggest that 30–40% of the added N was removed by the marsh during each tidal cycle. Creek bank *Spartina alterniflora* and high marsh *S. patens* production increased, but not stunted high marsh *S. alterniflora*. Microbial production increased in the fertilized creek bank *S. alterniflora* habitat where benthic microalgae also increased. We found top-down control of benthic microalgae by killifish, but only under nutrient addition and in the opposite direction (increase) than that predicted by a fish–invertebrate–microalgae trophic cascade. Surprisingly, infauna declined in abundance during the first season of fertilization and with fish removal. Our results demonstrate ecological effects of both nutrient addition and mummichog reduction at the whole-system level, including evidence for synergistic interactions.

Key words: bottom-up; eutrophication; *Fundulus heteroclitus*; multiple stressors; nutrient loading; salt marsh; *Spartina alterniflora*; *Spartina patens*; species change; top-down.

INTRODUCTION

The sustainability of coastal ecosystems in the face of widespread environmental change is an issue of pressing concern throughout the world (Emeis et al. 2001). Coastal ecosystems form a dynamic interface between terrestrial and oceanic systems and are one of the most productive ecosystems in the world. Coastal systems probably serve more human uses than any other ecosystem, and they have always been valued for their rich bounty of fish and shellfish. Coastal areas are also the sites of the nation's and the world's most intense

commercial activity and population growth; worldwide, ~75% of the human population now lives in coastal regions (Emeis et al. 2001).

Since the mid-1970s, nutrient enrichment of coastal and estuarine waters has become the premier issue for both scientists and managers (National Research Council 2000). Our understanding of coastal eutrophication has been developed principally through monitoring of estuaries, with a focus on pelagic or subtidal habitats (National Research Council 2000, Cloern 2001). Because estuarine systems are usually nitrogen limited, NO₃⁻ is the most common nutrient responsible for cultural nutrient enrichment (Cloern 2001). Increased nitrogen delivery to pelagic habitats of estuaries produces the classic response of ecosystems to stress (altered primary producers and nutrient cycles and loss of secondary producer species and production; Nixon 1995, Rapport and Whitford 1999, Deegan et al. 2002).

Manuscript received 15 March 2006; revised 19 October 2006; accepted 15 November 2006; final version received 12 December 2006. Corresponding Editor: A. Townsend. For reprints of this Special Issue, see footnote 1, p. S1.

⁷ E-mail: Ldeegan@mbl.edu

Saltmarsh ecosystems have been thought of as not susceptible to nitrogen overloading, because early studies found added nitrogen increased marsh grass production (primarily *Spartina* spp., cordgrass) and concluded that salt marshes can adsorb excess nutrients in plants and saltmarsh plant-derived organic matter as peat (Verhoeven et al. 2006). Detritus from *Spartina* is important in food webs (Deegan et al. 2000) and in creating peat that forms the physical structure of the marsh platform (Friedrichs and Perry 2001). However, the accumulation of peat and inputs of sediments, and the losses of peat through decomposition and sediment through erosion, may be altered under high nutrient regimes and threaten the long-term stability of marsh systems.

Nitrogen addition may lead to either net gain or loss of the marsh depending on the balance between increased marsh plant production and increased decomposition. Absolute change in marsh surface elevation is determined by marsh plant species composition, production, and allocation to above- and belowground biomass, microbial decomposition, sedimentation, erosion, and compaction (Friedrichs and Perry 2001). Levine et al. (1998) suggested that competitive dynamics among plants might be affected by nutrient enrichment, potentially altering relative abundance patterns and favoring species with less belowground storage and thus lowering rates of peat formation. When combined with the observation that nutrient additions may also stimulate microbial respiration and decomposition (Morris and Bradley 1999), the net effect on the salt marsh under conditions of chronic nitrogen loading is a critical unknown.

Although most research treats nutrient enrichment as a standalone stress, it never occurs in isolation from other perturbations. The effect of nutrient loading on species composition (both plants and animals), and the resultant structure and function of wetlands, has been largely ignored when considering their ability to adsorb nutrients (Verhoeven et al. 2006). Recent studies suggest the response of estuaries to stress may depend on animal species composition (Silliman et al. 2005). Animal species composition may alter the balance between marsh gain and loss, as animals may increase or decrease primary production, decomposition, or N recycling (Pennings and Bertness 2001).

Failure to understand interactions between nutrient loading and change in species composition may lead to underestimating the impacts of these stresses. The "bottom up or top down" theory originated from the observation that nutrient availability (bottom up) sets the quantity of primary productivity, while other studies have shown that species composition (top down), particularly of top consumers, has a marked and cascading effect on ecosystems, including controlling species composition and nutrient cycling (Matson and Price 1992, Pace et al. 1999). Most examples of trophic cascades are in aquatic ecosystems with fairly simple,

algal-grazing pelagic food webs (Strong 1992). The rarity of trophic cascades in terrestrial systems has been attributed to the importance of detrital food webs (Polis 1999). Detritus-based aquatic ecosystems, such as salt marshes, bogs, and swamps, have classically been considered bottom-up or physically controlled ecosystems.

Recent experiments, however, suggest that salt marshes may exhibit top-down control at several trophic levels (Silliman and Zeiman 2001, Silliman and Bertness 2002, Quiñones-Rivera and Fleeger 2005). One abundant, ubiquitous predator, a small (<10 cm total length) killifish (*Fundulus heteroclitus*, mummichog) has been suggested to control benthic algae through a trophic cascade, because they prey on the invertebrates that graze on the benthic algae (Kneib 1997, Sarda et al. 1998). In late summer, killifish are capable of consuming 3–10 times the creek meiofauna production, and meiofauna in the absence of predators appear capable of grazing >60% of the microalgal community per day (Carman et al. 1997). Strong top-down control by grazers is considered a moderating influence on the negative effects of elevated nutrients on algae (Worm et al. 2000). Small-scale nutrient additions and predator community exclusion experiments have demonstrated bottom-up and top-down control of macroinfauna in mudflats associated with saltmarsh creeks (Posey et al. 1999, Posey et al. 2002). Together, these observations suggest mummichogs are at the top of a trophic cascade that controls benthic algae (Sarda et al. 1998).

Mummichogs are also omnivorous and ingest algae, bulk detritus, and the attached microbial community (D'Avanzo and Valiela 1990). As a result, marsh decomposition rates may be limited by top-down controls through trophic pathways or by release from competition with algae for nutrients.

Whole-ecosystem experiments have shown that responses to stress are often not predictable from studies of the individual components (Schindler 1998). Developing the information needed to predict the interacting impacts of nutrient loading and species composition change requires experiments with realistic alterations carried out at scales of space and time that include the complexities of real ecosystems. Whole-ecosystem manipulation experiments have been used effectively in other ecosystems (Bormann and Likens 1979, Carpenter et al. 1995), but they are rare in coastal research. The scale of experiments in salt marshes have traditionally been less than a few square meters. Our understanding of the response of saltmarsh plants to nutrient enrichment is from small (<10 m²), plot-level additions, where uniform levels of dry inorganic fertilizer (20 to >1000 g N·m⁻²·yr⁻¹) are sprinkled on the marsh surface at low tide. Dry-fertilizer additions were usually made every two weeks or monthly, and the duration of elevated nutrient levels after these additions was usually not determined. Tidal water is the primary vector for N delivery to coastal marshes, suggesting that dry fertilizer

addition to the marsh surface may not be the best basis for determining if *Spartina* production responds to nutrient enrichment of tidal waters. Similarly, our understanding of top-down controls in salt marshes also relies on small (1–4 m²) exclusion experiments that use cages to isolate communities from top consumers. While the design of these cage experiments has improved, there are some remaining drawbacks. For example, it is impossible to selectively exclude single species using cages, and recruitment or size-selective movement into or out of the cages may obscure interpretations. In addition, while these small-scale experiments provide insight into controls on isolated ecosystem processes, they do not allow for interaction among different parts of the ecosystem that may buffer or alter the impacts, nor are they appropriate for determining the effects of populations of larger more motile animals on whole ecosystems or the effects of ecosystem changes on populations. For example, interactions may be caused when a motile species alters its distribution among the habitats available to it because of an experimental treatment. Small-scale experiments generally do not allow such events to happen. Complex feedbacks among physical and biological processes can alter accumulation rates and affect marsh elevation relative to sea level rise, making extrapolation of small plot-level experiments to whole-marsh ecosystems problematic.

We are conducting an ecosystem-scale, multiyear field experiment including both nutrient and biotic manipulations to coastal saltmarsh ecosystems. We are testing, for the first time at the ecosystem level, the hypothesis that nutrient enrichment and species composition change have interactive effects across multiple levels of biological organization and a range of biogeochemical processes. We altered whole saltmarsh creek watersheds (~60 000 m² of salt marsh) by addition of nutrients (15-fold above ambient levels) in flooding waters, and by a 60% reduction of a key fish species, the mummichog. Small marsh creek watersheds provide an ideal experimental setting, because they have the spatial complexity, species composition, and processes characteristic of the larger salt marsh ecosystem, which are often hundreds of thousands of square meters. Manipulating entire saltmarsh creeksheds allowed us to examine effects on large motile animals and the interactive effects of motile species changes on ecosystem processes without cage artifacts. Because our manipulations were done on whole-marsh ecosystems, we are able to evaluate the integrated and interactive effects on all habitats (e.g., water column, tidal creeks, and marsh) and on populations. These experiments are similar in many respects to the small watershed experiments carried out in forested catchments.

Our nutrient enrichment is novel compared to past studies in two important ways. We added nutrients (N and P) directly to the flooding tidal creek waters to mimic the way in which anthropogenic nutrients reach

marsh ecosystems. All previous experimental salt marsh nutrient enrichment studies used a dose–response design with spatially uniform dry-fertilizer loading on small plots (<10 m²). Nutrients carried in water will interact and reach parts of the ecosystem differently than dry fertilizer. Our enrichment method also creates a spatial gradient of nutrient loading across the landscape that is proportional to the frequency and depth of inundation in the marsh. Spatial gradients in loading within an ecosystem are typical in real-world situations in many terrestrial and aquatic ecosystems. Because of our enrichment method, at any location in the ecosystem, nutrient load will be a function of the nutrient concentration in the water, the frequency and depth of tidal flooding, and the reduction of nutrients from the flooding waters by other parts of the ecosystem. Uniform loading misses important aspects of the spatial complexity of ecosystem exposure and response.

This work is organized around two questions that are central to understanding the long-term fate of coastal marshes:

- 1) Does chronic nutrient enrichment via flooding water increase primary production more than it stimulates microbial decomposition?
- 2) Do top-down controls change the response of the saltmarsh ecosystem to nutrient enrichment?

Here we present findings on the first two years of these experiments including the following: (1) water chemistry, (2) standing stocks and species composition of benthic microalgae, (3) microbial production, (4) species composition and ecophysiology of macrophytes, (5) invertebrates, and (6) nekton. Because even highly eutrophic waters result in nutrient loading that is an order of magnitude less than most plot-level experiments, we expected little stimulation of saltmarsh vascular plant growth. However, moderate levels of nutrient enrichment in the water column were expected to increase benthic algal biomass and to stimulate bacterial activity and detrital decomposition throughout the ecosystem because of direct uptake of nitrogen from the water column and availability of more high-quality organic matter from increased algal production. We predicted nutrient enrichment would increase invertebrate production because of an increase of high-quality microalgal and microbial production at the base of the food web. Finally, we predicted that fish reduction would reduce predation on benthic invertebrates, resulting in increased abundance of benthic invertebrates that would graze down the benthic algae.

SITE DESCRIPTION

The Plum Island Sound estuary in New England, USA, where we conducted our experiments is a classic saltmarsh estuary that is currently unaffected by nutrient loading (Fig. 1). Tall *Spartina alterniflora* (~200 cm in height; smooth cordgrass) is found in pure stands in low marsh and along creek banks that receive daily tidal inundation, while *Spartina patens* (20–60 cm




FIG. 1. Aerial photograph of the Rowley River region salt marshes. Light areas are *Spartina patens* marsh, and the darker areas are *Spartina alterniflora* marsh. Mottled dark areas seen in the northwest and southwest corners of the map are upland forest. The areas outlined in white are the experimental marshes: SW, Sweeney Creek; WE, West Creek; CL, Clubhead Creek; and NE, Nelson Creek. The base photograph is from (<http://maps.massgis.state.ma.us/MassGISColorOrthos/viewer.htm>); April 2001.

in height; saltmeadow cordgrass) is most abundant in higher elevation areas that are well drained and flood less frequently (see Plate 1). *S. alterniflora* also occurs in a short form (20–60 cm in height) in the high marsh, often in pure stands in areas that are poorly drained. Most of the total marsh area (~80%) is high marsh that floods on spring tides. Mean tide is 2.6 m; mean spring tides are 3 m. During a typical growing season, *S. alterniflora* in Plum Island is under water ~35% of the time, while *S. patens* and short form *S. alterniflora* are inundated ~12% of the time. As is typical of most New England marshes, historically the marsh platform was ditched for saltmarsh haying and mosquito control. Ditch edges show the same vegetation zonation as creek banks. Currently, the all marshes are periodically sprayed with *Bacillus thuringiensis israelensis* (BTI) for mosquito control.

Because characteristics of tidal marsh systems vary along a fresh to salt water gradient, we anticipated that the experimental areas might have to be paired to account for differences associated with position in the estuary. After initial exploratory work, we chose four marsh creek systems (Sweeney, West, Clubhead, and Nelson) with similar geomorphology along the Rowley River in the central portion of Plum Island Sound. Our experimental areas were at the landward end of the creeks, where each creek has two roughly equal-sized branches (700 linear m of branch creek channel and 60 000 m² of marsh per treatment). Prior to experimental manipulation (1998–2002), we collected baseline temperature, salinity, nutrient, and benthic microalgae characteristics on each creek system (four creeks, four years, approximately eight times per year). Additional baseline data on total suspended sediments, marsh plant

TABLE 1. Saltmarsh characteristics (means \pm SE) prior to experimental manipulations (1998–2003).

Characteristic	<i>P</i>	Creek pair 1		Creek pair 2		Plum Island Sound
		Sweeney	West	Clubhead	Nelson	
Marsh creek area (m ² \times 10 ⁴)		12.5	14.8	11.4	12.1	
Water column						
Salinity (psu)	0.001	20.8 ^a \pm 0.8	24.0 ^{a,b} \pm 0.8	27.0 ^b \pm 0.8	25.2 ^b \pm 0.9	29.1 ^b \pm 0.9
Temperature (°C)	0.009	18.7 ^a \pm 0.8	19.4 ^a \pm 0.6	20.0 ^a \pm 0.7	19.0 ^a \pm 0.9	14.1 ^b \pm 1.1
Nitrate (μ m)	0.023	4.3 ^a \pm 0.3	3.5 ^a \pm 0.4	3.2 ^a \pm 0.3	2.7 ^a \pm 0.3	0.7 ^b \pm 0.2
Ammonium (μ m)	0.001	11.7 ^a \pm 1.3	10.0 ^a \pm 1.6	13.8 ^a \pm 1.8	14.9 ^a \pm 2.1	0.8 ^b \pm 0.2
Phosphate (μ m)	0.001	1.2 ^a \pm 0.1	0.9 ^a \pm 0.1	1.1 ^a \pm 0.1	1.3 ^a \pm 0.1	0.4 ^b \pm 0.03
Phytoplankton (mg chl <i>a</i> /L)	0.001	8.4 ^a \pm 0.9	8.8 ^a \pm 1.6	7.4 ^a \pm 0.8	9.0 ^a \pm 1.1	2.5 ^b \pm 0.4
TSS (mg/L)		15 \pm 0.8	18 \pm 1.0	19 \pm 1.3	14 \pm 1.6	
Benthic algae (mg chl <i>a</i> /m ²)						
Mudflat	0.001	62.7 ^b \pm 3.7	93.6 ^a \pm 7.7	134.7 ^c \pm 6.7	110 ^{a,c} \pm 11.1	
<i>S. alterniflora</i>	0.250	197.5 ^a \pm 33.9	250.8 ^a \pm 23.9	215.2 ^a \pm 6.8	249.0 ^a \pm 27.4	
<i>S. patens</i>	0.001	94.6 ^a \pm 19.4	48.6 ^a \pm 6.5	55.1 ^a \pm 4.3	135.9 ^b \pm 27.6	
Vascular plant community (percent cover)						
Tall <i>S. alterniflora</i>	0.738	5.5 \pm 1.1	6.2 \pm 1.1	7.1 \pm 1.2	7.1 \pm 1.1	
Short <i>S. alterniflora</i>	0.001	16.8 ^a \pm 1.5	18.5 ^a \pm 1.6	24.9 ^b \pm 1.9	30.6 ^b \pm 1.8	
<i>S. patens</i>	0.078	41.0 \pm 1.7	42.3 \pm 1.9	43.6 \pm 2.0	37.1 \pm 1.7	
<i>D. spicata</i>	0.001	25.9 ^a \pm 1.5	18.0 ^b \pm 1.4	7.8 ^c \pm 1.1	8.0 ^c \pm 1.1	
No. plant species/m ²	0.001	2.7 ^a \pm 0.1	2.5 ^a \pm 0.1	2.1 ^b \pm 0.0	2.1 ^b \pm 0.1	

Notes: Water column and benthic algae data are from 1998–2002. Total suspended solids (TSS) and vascular plant community (percent cover) are from 2003, prior to experimental treatments. Values of *P* are for ANOVA main effect for creek on water column and benthic algae. Differences between creek branches were always *P* > 0.05. Different superscript letters indicate differences among creeks within a row (Tukey, *P* < 0.05). Error degrees of freedom *N* (see Appendix A): salinity, 240; temperature, 159; nitrate, 230; ammonium, 161; phosphate, 250; phytoplankton, 247; mudflat habitat benthic algae, 247; *Spartina alterniflora* habitat benthic algae, 99; *Spartina patens* habitat benthic algae, 102; total inorganic suspended sediments (TSS), 109; and vascular plants, 23 for each plant species and number of species.

species composition, and distribution was collected in 2003.

The four marsh creek watersheds were very similar, but differences in salinity and vascular plant community composition supported pairing the two sets of experimental and reference creeks. Water column temperature, nutrients (NO₃⁻, NH₄⁺, PO₄³⁻), total suspended sediments, and phytoplankton did not differ among the four creeks, or in branches within creeks (*P* > 0.05 for all constituents except ammonium; see Appendix A), however water column parameters for all creeks were different from Plum Island Sound water (Table 1). Creeks had warmer temperatures, lower salinities, higher nutrients, and higher phytoplankton biomass than did the Sound. Sweeney and West are located further upstream along the tidal Rowley River, closer to sources of freshwater, and had slightly lower salinities than did Clubhead and Nelson.

Marsh vegetation followed the classic distribution of saltmarsh plant communities: tall *Spartina alterniflora* was found along the creek banks in a band that varied in width (1–3 m); short *S. alterniflora*, *S. patens*, and *Distichlis spicata* (saltgrass) were found on the high-marsh platform (see Plate 1). Similarity analysis of the plant community (ANOSIM, Primer-E, Plymouth, UK) found no difference between Sweeney and West (*P* = 0.195) and between Clubhead and Nelson (*P* = 0.152). Analysis of all other creek pairings found that the plant communities were significantly different (for all other

pairings, *P* < 0.004). Total cover of tall *S. alterniflora* and of high-marsh *S. patens* did not differ among creeks. Sweeney and West had less short *S. alterniflora* than did Clubhead and Nelson, and much higher cover of *D. spicata*. *Atriplex patula* (spear saltbush) contributed little to total cover and had higher cover in Sweeney and West compared to Clubhead and Nelson. Sweeney and West also had consistently less plant litter and standing dead than did Clubhead and Nelson. We also mapped the boundaries of individual plant communities using high precision GPS (\pm 2 cm horizontal; maps not shown) to provide a baseline from which to determine long-term shifts in plant community boundaries.

Baseline benthic microalgal biomass differed among habitats (Table 1), and was highest in the intertidal *S. alterniflora* habitat (220 mg chl *a*/m²), intermediate in intertidal mudflat habitat (100 mg chl *a*/m²), and lowest in high-marsh *S. patens* (60 μ g chl *a*/m²; see Plate 1). West had higher mudflat and *S. alterniflora* habitat microalgae biomass compared to Sweeney, while Clubhead and Nelson were very similar to each other. Algal biomass in *S. patens* was highest in Nelson and lowest in Sweeney.

Creek pairings were made on the basis of plant community structure and position in the estuary. Creek pair 1 was Sweeney and West. This pair was closest to the freshwater end of the estuary, had slightly lower salinity and very similar marsh plant communities. Creek pair 2 was Clubhead and Nelson. This pair was

more seaward, with higher salinities and marsh plant communities similar to each other and different than creek pair 1.

EXPERIMENTAL DESIGN

We manipulated the marsh ecosystem in two ways: (1) nutrients were added to one creek of each pair (to ~15-fold above ambient reference concentrations); and (2) fish were reduced in one branch of each creek (~60% decline in mummichog) (Fig. 2). This design resulted in four, ~60 000 m² marsh treatments within each pair: ambient nutrients–ambient mummichog, ambient nutrients–low mummichog, high nutrients–ambient mummichog, and high nutrients–low mummichog. We will refer to the ambient-nutrient creeks as “nutrient reference” and the nutrient-enriched creeks as “nutrient enrichment.” The fish treatments will be called ambient (“fish”) or reduction (“low fish”); these are located in either a reference or nutrient-enrichment creek.

We used a before-and-after control (BACI) type of experimental design in which pairing of experimental units accounts for variability that would contribute to error in a completely randomized design (Underwood 1994). Replication of ecosystem-scale experiments is difficult, because it is often hard to find similar ecosystems (Carpenter et al. 1995, Schindler 1998); the matched-pair approach helps ameliorate this difficulty (Stewart-Oaten and Bence 2001). The use of large experimental units (in our case ~60 000 m²/treatment) is both labor-intensive and expensive (fertilizer alone costs ~\$20 000/yr for each creek). These types of constraints generally necessitate low replication of individual treatments in ecosystem-scale experiments. While subject to the limitations of pseudoreplication (Hurlbert 1984) and small sample size, large-scale manipulative studies provide a realistic environment for examining effects and processes that occur at an ecosystem-scale. Our results include the effects of spatial variation and complexity, interactions between all of the species in the system, gradients across large areas, and large habitat patches.

The experiment was implemented in phases. In 2003 (year 0), standardized procedures and methods were developed and extensive baseline measurements were made in all creeks and branches. In 2004, we began nutrient and fish reduction manipulations in creek pair 1 (year 1). In 2005, manipulations continued in creek pair 1 (year 2) and began in creek pair 2 (year 1). Nutrients were added to Sweeney Creek (pair 1) and Clubhead Creek (pair 2).

Nutrient addition

We implemented an enrichment of 70 $\mu\text{mol NO}_3^-/\text{L}$ and 4 $\mu\text{mol PO}_4^{3-}/\text{L}$ (both 15-fold increases over background concentrations of $<5 \mu\text{mol NO}_3^-/\text{L}$ and $\sim 1 \mu\text{mol PO}_4^{3-}/\text{L}$; Table 1). These concentrations are typical in estuarine systems designated as moderate to highly eutrophic, based on the response of the pelagic




FIG. 2. Diagram of design of creek pairs and experimental treatments. Stars indicate locations of nutrient addition to flooding water. Dotted lines indicate locations of fish barriers; minnow traps were placed behind this barrier with ~25-m spacings.

and subtidal system (National Oceanic and Atmospheric Administration 1999, Environmental Protection Agency [EPA] 2002). We added the NO_3^- and PO_4^{3-} in ~15:1 molar ratio as anthropogenic increases in NO_3^- in coastal waters is usually accompanied by increased PO_4^{3-} (EPA 2002). We added nutrients by pumping a concentrated solution of NO_3^- and PO_4^{3-} to the flooding water of every tide during the growing season (mid-May–October; ~150 d). The pump rate was adjusted every 10 min throughout each incoming tide (based on a hydrologic model) to maintain constant N and P concentrations in incoming waters until a water depth of 3.2 m in the channel (equivalent to ~15 cm of water on the high-marsh platform) was reached. The cessation of nutrient addition at water levels >3.2 m affected approximately the last half hour of the incoming tide on $<20\%$ of the tides between May and October. The solution was distributed through the water column by vertical and horizontal spreader bars and traveled ~100 m allowing in-channel mixing to occur before reaching the experimental area boundary.

Fish reduction

Mummichogs in this estuary have been observed to winter in deep water and move up into marsh systems to spawn in May and June, although some may winter in pools on the marsh platform (Raposa 2003). We excluded adult mummichogs from entering one branch of each creek by using block nets (installed at the branch mouth and small channel boundaries prior to the spring migration) and by constantly deployed minnow traps that fished throughout the season (selective for mummichogs; Layman and Smith 2001). Previous tagging work in these marshes found mummichogs had a home range of ~300 linear meters of creek and little movement between creek branches (Sweeney et al.

1998). Minnow-trapped mummichogs were released >1 km from the treatment area; all other species were returned to the creek. The combination of block netting and minnow trapping reduced mummichog abundance by ~60% (analyses presented in *Results: Fish reduction*).

Sampling locations

Three benchmark sample areas, or "strata," were established on each branch of each creek in 2003. Each stratum began in the creek channel, was 10 m wide, and extended onto the marsh platform for 50 m. Strata were designed to represent the whole treatment area, and to include potential effects of distance from the nutrient addition point. The strata were spaced along each branch ~100 m apart, and each included all major habitat types: intertidal mudflat (MF) in the creek, filamentous microalgae (FA) at the top of the creek bank, tall-form *S. alterniflora* (TSA) at the top the creek bank, *S. patens* (SP) on the marsh platform, and short-form *S. alterniflora* (SSA) on the platform. During 2003–2005, much of the routine/repeated sample collection, such as benthic chlorophyll and invertebrates, sediments, and plant production, was conducted within the strata; replicates from each habitat type was collected from within each stratum. Specific collection points were haphazardly located within habitats. Additional samples of many types were frequently collected outside of the established strata.

METHODS

Nutrients and suspended sediments

We measured baseline water quality characteristics (NO_3^- , NH_4^+ , PO_4^{3-} , TSS, salinity, etc.) for four growing seasons prior to manipulations (May–September, 1998–2002). We collected water samples during mid-ebb once or twice per month from each creek branch and from ~25 m seaward of the confluence of each pair of creek branches. Water samples were also collected two to four times each year from three locations in the Plum Island Sound, New England, USA, to characterize the water that floods the creeks.

In 2003 (premanipulation), 2004, and 2005, we conducted intensive, bimonthly, tidal cycle sampling (hourly sampling over a full, 12-h tidal cycle). Each month (May–September), one spring tide cycle and one neap tide cycle were sampled at each creek branch and confluence location. Tidal cycle sampling allows calculation of nutrient delivery, retention, and export (budgets) for whole tidal cycles.

One-liter water samples were collected in acid washed, Nalgene bottles using an autosampler (SIGMA, Loveland, Colorado, USA). Samples were refrigerated and filtered within 24 h of collection (precombusted 47 mm glass fiber filter [GFF]) and frozen until analysis. Chlorophyll *a* (chl *a*) was determined on the filter (foil wrapped; frozen; 90% acetone extraction and fluorometer analysis, Holm-Hansen et al. 1965). Concentrations of NO_3^- were determined colorimetrically on a flow

injection analyzer (Lachat Quik Chem 8000, Lachat Zellweger Instruments, Milwaukee, Wisconsin, USA) using the cadmium–copper reduction method (Wood et al. 1967), concentrations of PO_4^{3-} were determined colorimetrically using the ascorbic acid method (Parsons et al. 1984), and concentrations of NH_4^+ were determined colorimetrically using the phenolphthorite method modified for small sample size (Solorzano 1969). Both PO_4^{3-} and NH_4^+ samples were run on a Cary Spectrophotometer (Varian, Palo Alto, California, USA).

Water on the marsh platform at high tide was sampled during 2005. Acid washed, narrow-mouthed bottles were attached to stakes and placed throughout the marsh at low tide. The flooding tide filled the bottles, which were retrieved after the tide subsided. These samples were treated as described and analyzed for NO_3^- .

In 2003, total suspended solids samples were collected weekly at slack high tide within each branch of the four creeks. One-liter samples were filtered through pre-weighed, ashed filters (47 mm, 0.7 μm GFF). Filters were then ashed and weighed to determine the amount of inorganic material in suspension.

Vascular marsh plants

To characterize the marsh community prior to manipulations (July 2003), plant species occurrence and visual estimates of percent cover by species were recorded in contiguous 1-m² plots ($N = 50$) along the length of each strata for each creek ($N = 3$ strata per treatment branch; $N = 6$ per creek). Plant frequency (percentage of plots in which a species occurred) and mean percent cover for each species were determined for each strata ($N = 6$ for each species per creek).

Shoot growth of tall *S. alterniflora*, short *S. alterniflora*, and *S. patens* was determined in creek pair 1 (years 1 and 2). Every two weeks (mid-May–August) haphazardly selected shoots ($N = 25$) of each type in each stratum were cut at the peat surface and refrigerated within two hours. Each shoot was individually washed to remove sediment, measured for length, and dried (80°C for ≥ 24 h). Mean shoot length (cm), mass (g), and length-specific leaf mass (mass of an individual shoot divided by the total length; g/cm) were determined for each shoot. Values were normally distributed for all three parameters. Nitrogen and carbon content of tall *S. alterniflora*, short *S. alterniflora*, and *S. patens* were determined (June, July, and August, 2004; $N = 6$ per plant type per creek; PerkinElmer 2400 Series II CHNS/O analyzer, PerkinElmer Life and Analytical Sciences, Wellesley, Massachusetts, USA). Stem density (shoots per unit area) was counted in four plots haphazardly located within 5 m of the centerline of the sample strata in August, and means were calculated for each plant type for each stratum ($N = 6$ per plant type per creek). Because of known differences in stem density between species and growth forms, different size quadrats were used for each species: tall *S. alterniflora*,

square 0.25 m²; short *S. alterniflora*, square 0.0625 m²; *S. patens*, circle 0.008 m².

Microbial production

Surface sediment samples for bacterial production were taken using a core ($N = 10$; 1.5 cm diameter, <10 mm depth) monthly during the growing season in tall *S. alterniflora* and *S. patens*. Replicates were homogenized and then subsampled ($N = 5$) for measurement of bacterial production by uptake of tritiated leucine in sediment slurries (Buesing and Gessner 2003).

Benthic microalgae

Benthic microalgal biomass samples were collected monthly (June–September; 2003, 2004, and 2005; creek pair 2 was not sampled in 2004) in mudflat, tall *S. alterniflora*, and *S. patens* habitats. Macroalgae were rare in these systems. Filamentous (and associated epiphytic) algae occur in specific habitats (i.e., the creek wall near the top and associated with the stems of short-form *S. alterniflora*), and their responses will be discussed in future reports. In September 2005, supplemental samples spaced along the entire creek branch length were taken in each habitat. Cores were taken in each habitat in each sampling area (two per habitat per area; 2.7 cm diameter, 2 cm depth), frozen immediately, and total chlorophyll *a* (mg chl *a*/cm²) determined (Lorenzen 1967; acetone extraction and spectrophotometric analysis). Data were log-transformed.

Benthic microalgal species composition and numerical response were measured monthly on glass slides (five slides per collection; one collection per stratum per month) located parallel to stream flow in the middle of the creek channel and submerged for 2–3 days. Slides were preserved in 70% ethanol, and the species composition and density were determined by microscopic examination.

Benthic invertebrates

Macroinfauna and meiofauna collections were taken coincident with monthly benthic microalgal samples. Macroinfauna samples taken 10 wk after nutrient addition and fish reduction treatments began (2–3 August 2004) in tall *S. alterniflora* in creek pair 1 were analyzed. Duplicate cores (6.6 cm diameter push core; to 5 cm depth) were collected at low tide and fixed with 10% formalin and Rose Bengal. After a minimum of two days, samples were sieved (500- μ m sieve) and macroinfauna were enumerated and identified to the lowest possible taxon. Permanent meiofauna (e.g., nematodes) were not enumerated. Because annelids constituted 98% of all macroinfauna and not all species have been identified, we report only total annelid abundance. Data met the criteria for normal distribution.

To examine the effects of fish reduction on motile saltmarsh epifauna, passive collectors (litterbags) were placed in the tall *S. alterniflora* habitat of reference creeks in July 2005. The bags (13.5 \times 29 cm, with 5-mm

delta weave mesh) were filled with \sim 20 g of dried, standing, dead, *S. alterniflora* leaves (Fell et al. 1998). After two weeks, litterbags were collected and contents preserved in 50% ethanol and Rose Bengal solution. After at least two days, litter was rinsed over a 0.5-mm sieve, and all animals were collected using forceps and a hand lens. Animals were identified to lowest possible taxon and enumerated. Here we report on the most abundant species collected, the amphipod *Uholorchestia spartinophila*, which comprised about 75% of the animals collected.

Nekton abundance, growth, and diet

Nekton were collected on the marsh surface each month (September–October 2003; June–September 2004 and 2005) during nighttime spring tides in each stratum ($N = 3$ per treatment branch). Flume nets (dyed green 3-mm delta mesh) were 3 m wide at the marsh–creek interface and extended perpendicularly 10 m onto the marsh surface (30 m² of marsh surface). Nets were flush with the marsh surface as the tide rose; at slack high tide, the sides were lifted, and the front and back nets attached to enclose the area. The front net collected fish as they moved off of the marsh surface with the falling tide. All nekton collected were identified to species, and \leq 300 individuals of each species measured (total length (TL) \pm 1 mm) and weighed (mass \pm 0.01 g) individually; if we collected $>$ 300 individuals per species, remaining individuals were weighed as an aggregate by species. Abundance and biomass were square-root ($n + 1$) transformed. Mean size (TL \pm 1 mm) of young-of-the-year *Palaemonetes pugio* (daggerblade grass shrimp) was estimated in August and September using length frequency (NORMSEP procedure; Food and Agriculture Organization of the United Nations, 2000, FISAT II, Rome, Italy). Growth was estimated as the change in mean size of the smallest size group. Length frequency was square-root ($n + 1$) transformed before ANOVA analysis. Mummichog diet was analyzed on fish ($N = 20$ per treatment per period) captured in flume nets or minnow traps in summer of 2004 and 2006. Gut fullness was estimated and items identified to lowest practical taxon under a dissecting microscope (Hyslop 1980).

Data analysis

A paired BACI-type ANOVA (Nutrient, Fish reduction, Before/After) analysis with Type III sums of squares was used to analyze the effectiveness of the fish reduction treatment, and the response of benthic microalgal biomass and grass shrimp abundance to nutrient addition and fish reduction treatments. We present computed *P* values, but infer significance when $P < 0.1$ because, in a complex ecosystem experiment, background variability is generally high and replication low, and we wanted to identify hypotheses with support in the data (Oksanen 2001, Hobbs and Hilborn 2006). Evidence from several lines of inquiry was used to further examine support for the responses. Because of

TABLE 2. Incoming and outgoing nutrient concentrations (means \pm SE) during the major tidal water flux periods for all creeks in the premanipulation year (year 0) and nutrient addition creeks in treatment years (years 1 and 2).

Nutrient	Baseline		Nutrient addition					
	All creeks		Creek pair 1				Creek pair 2	
	Year 0		Year 1		Year 2		Year 1	
	In	Out	In	Out	In	Out	In	Out
NO ₃ ⁻	2.6 \pm 0.3	1.3 \pm 0.3	70.7 \pm 38.3	58.4 \pm 38.9	94.9 \pm 18.4	59.2 \pm 17.1	116.3 \pm 20.8	98.2 \pm 22.4
NH ₄ ⁺	9.1 \pm 3.1	4.7 \pm 3.4	9.4 \pm 1.3	4.4 \pm 0.6	3.7 \pm 1.0	6.5 \pm 1.9
PO ₄ ³⁻	0.9 \pm 0.05	1.0 \pm 0.08	5.1 \pm 2.6	4.2 \pm 2.5

Notes: All nutrient concentrations given in $\mu\text{mol/L}$. Missing data for samples that have not yet been analyzed are indicated by ellipses (...).

known differences between creek pairs, and because the pairs were manipulated for different lengths of time, creek pairs were analyzed separately. Replicates of response variables were averaged by sample strata within a treatment ($N = 3$ per treatment). In exploratory analysis, we found no spatial pattern associated with strata for any variable, therefore they were considered independent estimates of the treatment response. Monthly data were nested within treatment Year (0, 1, 2) and Before (0)/After (1, 2). Analysis was performed on the mean response variables (appropriately transformed to meet the assumption of normality), not differences, so that we could examine the interactive effects of nutrient addition and fish reduction. A significant interaction term (Nutrient \times Fish \times Month(Year, Before/After) indicated a difference among treatments over time. If a significant interaction occurred, data were examined visually and Tukey or t tests used to determine differences (Stewart-Oaten and Bence 2001, Parker and Wiens 2005). A two-way ANOVA (Nutrient, Fish reduction) was used to analyze benthic microalgal count and macroinfaunal abundance. An ANOVA (Nutrient, Fish reduction, Month) was used to analyze shrimp body size data. All ANOVAs were calculated using SuperANOVA (1996, Abacus Concepts, Berkeley, California, USA). We expected that fish reduction would have no effect on vascular plants, as the overall nutrient regime (enriched or reference) would not be altered by fish abundance. This was confirmed by initial analyses, and therefore vegetation data were pooled between fish treatments within a creek (nutrient reference or enrichment) and analyzed using the individual measurements. Because the plant response metrics were calculated from the same samples (height- and length-specific leaf mass, $N = 125$ for each species in each creek), we used a Bonferroni adjustment of significance from 0.1 to 0.05. Tukey or t tests were used to examine differences among treatments. Plant community data was analyzed using the ANOSIM procedure in Primer (Primer 5 for Windows Version 5.2.9, 2002, Primer-E, Plymouth, UK). Microbial production was analyzed by plotting nutrient-enriched against reference values and examin-

ing the distribution above and below the one-to-one line with Wilcoxon signed rank test.

RESULTS

Experimental manipulations

Nutrient addition.—In both creek pairs, we achieved our target NO₃⁻ enrichment of ~ 15 -fold increase over reference conditions. Nutrient addition increased the annual mean concentration of NO₃⁻ in the water flooding in creek pair 1 by a factor of ~ 14 -fold (year 1) to ~ 19 -fold (year 2) and ~ 23 -fold in creek pair 2 (Table 2). The nutrient addition increased PO₄³⁻ concentrations in creek pair 1 by a factor of approximately five. We added ~ 3500 kg of N (as NaNO₃) and 250 kg of P (as NaH₂PO₄) each year to each of the nutrient-enriched creeks (creek pair 1, 3251 kg N in year 1, 3546 kg N in year 2, 231 kg P in year 1, 252 kg P in year 2; creek pair 2, 4592 kg N in year 1, 327 kg P in year 1). Mean flooding water nutrient concentrations in year 0 (prior to nutrient addition) in the nutrient-enrichment and reference creeks of both pairs (NO₃⁻, 2.6 $\mu\text{mol/L}$; NH₄⁺, 9.1 $\mu\text{mol/L}$; PO₄³⁻, 0.9 $\mu\text{mol/L}$) were similar to the 1998–2002 concentrations (Table 1). In year 0, as is typical of salt marshes, ebbing waters generally had lower NO₃⁻ concentrations (1.3 $\mu\text{mol/L}$), while NH₄⁺ was highly variable through the tidal cycle (< 1 –24 $\mu\text{mol/L}$), with high concentrations generally occurring at low tide. The concentration of PO₄³⁻ remained at ~ 1 $\mu\text{mol/L}$ through the tidal cycle.

In the first year of nutrient addition to creek pair 1, our addition resulted in an annual mean of 70 $\mu\text{mol NO}_3^-/\text{L}$ and 5.1 $\mu\text{mol PO}_4^{3-}/\text{L}$ for the experimental period. However, examination of specific tidal cycles early in the growing season demonstrated lower than target concentrations during spring tides and higher than target concentrations during neap tides. Consequently, we adjusted the nutrient addition rate in the following year to achieve more uniform nutrient concentrations over a variety of tidal regimes.

The consistent decrease in NO₃⁻ concentration on ebbing tides in both nutrient-enriched creeks suggests uptake of the added NO₃⁻ by the marsh ecosystem. Significantly lower concentrations of NO₃⁻ in ebbing water compared to flooding water in creek pair 1 (year 1,




FIG. 3. Concentration of NO_3^- ($\mu\text{mol/L}$) on the marsh platform of the nutrient addition creeks of each creek pair at high tide, July 2005. Creek pair 1 is Sweeney, and creek pair 2 is Clubhead. The four-pointed star indicates the point of nutrient addition at each study site, and the dotted line indicates the location of the fish barrier. The base aerial photograph is from (<http://maps.massgis.state.ma.us/MassGISColorOrthos/viewer.htm>); April 2001.

58 vs. 70 $\mu\text{mol NO}_3^-/\text{L}$; year 2, 60 vs. 95 $\mu\text{mol NO}_3^-/\text{L}$) and creek pair 2 (98 vs. 116 $\mu\text{mol NO}_3^-/\text{L}$) suggest uptake by the marsh. Concentrations of NH_4^+ did not differ substantially between flooding and ebbing tides.

Despite our initial expectation that the marsh far from the addition point might receive water with lower nutrient concentration due to uptake as the water moved across the marsh, nutrient concentrations at high tide were high across the marsh platform (Fig. 3). The mean NO_3^- concentration on the nutrient-enriched

marsh was $\sim 80 \mu\text{mol NO}_3^-/\text{L}$ (range, 49–105 $\mu\text{mol NO}_3^-/\text{L}$) for creek pair 1. The NO_3^- concentration was $\sim 50 \mu\text{mol NO}_3^-/\text{L}$ (range, 2–78 $\mu\text{mol NO}_3^-/\text{L}$) on the nutrient-enriched marsh of creek pair 2. On highest tides, some dilution of nutrient concentration by water from other creeks outside the study areas was evident on the southern border of creek pair 2. These edges were several hundred meters away from the designated sampling areas and were not sampled for any response variables.




FIG. 4. Abundance (no. fish/30 m²) and biomass (g/m²) of mummichog (*Fundulus heteroclitus*) in experimental creeks. Nutrient reference creeks are indicated by squares; nutrient enrichment creeks are circles. Open symbols are reduction in mummichog density treatments (low fish); solid symbols are ambient mummichog density treatments. Along the x-axis, year 0 corresponds to premanipulation year, years 1 and 2 correspond to first and second years of manipulation, and numbers above year are months (e.g., 6 = June); small triangles indicate when manipulations began.

Fish reduction.—Reduction of mummichog (*Fundulus heteroclitus*) was effective, with a decrease in abundance of ~60% between the ambient-fish (Fish) and low-fish (Low fish) treatments (Fig. 4; see Appendix B). The overall mean was 65 mummichogs/30 m² in ambient-fish treatments compared to 25 mummichogs/30 m² in low-fish treatments. Mummichog was the most abundant fish species captured (19% of total nekton abundance and 92% of fish abundance; 41% of total nekton biomass and 89% of fish biomass over the three years). Mummichog abundance and biomass were near zero in June and increased rapidly until peak abundance in September and October. Mummichogs arrive in June to spawn and leave the marsh shortly after October to winter over in deeper water. Mummichog ranged in size from 15 to 110 mm and exhibited size classes associated with spawning and recruitment events. Our sampling methodology was most effective on larger mummichogs, suggesting that young-of-the-year and thus total mummichog abundances in all treatments may be underestimated. Fish abundance differed between fish treatments (Before/After × Fish treatment: creek pair 1, $F_{1,77} = 18.3$, $P = 0.0001$; creek pair 2, $F_{1,73} = 2.7$, $P = 0.10$), but there was no additional effect of nutrient addition on mummichog abundance for either creek pair (nutrient addition: creek pair 1: $F_{1,77} = 1.3$, $P = 0.25$;

creek pair 2, $F_{1,73} = 0.06$, $P = 0.8$). In creek pair 1 prior to manipulations (year 0), fish abundance in the reference creek branch randomly chosen for fish reduction was significantly higher (seasonal mean, 133 individuals/30 m²) than the paired ambient-fish density branch (43 individuals/30 m², t test, $P < 0.001$); our removal reduced abundance to below the paired ambient density (low fish 17 vs. fish 46 individuals/30 m²; t test, $P < 0.05$). In the nutrient-enriched creek of creek pair 2, initial fish abundance was slightly, but not significantly, lower in the branch designated for fish reduction (46 individuals/30 m²) compared to the ambient density branch (90 individuals/30 m² with high variation); removal lowered abundance in the reduction branch significantly below the previous density in that branch (22 vs. 46 individuals/30 m²) and well below ambient density in the experimental year (22 vs. 86 individuals/30 m²; t test, $P < 0.05$). Reduction was most effective for the larger size individuals (>40 mm TL), as smaller fish were not retained by the minnow traps or excluded by the block net. Thus, the effect of removing fish made a larger difference in reducing the total biomass compared to the number of individuals of mummichog. The overall seasonal average was ~30 g/30 m² in low-fish vs. ~100g/30 m² in ambient-fish treatment (Fig. 4).




FIG. 5. Growth and percentage nitrogen content (mean \pm SE) of tall *Spartina alterniflora* (TSA), short-form *S. alterniflora* (SSA), and *S. patens* (SP) of creek pair 1 after one and two years of nutrient addition. (a and b) Mid-August shoot height (cm), (c and d) mid-August length-specific leaf mass (g dry mass/cm shoot length), and (e and f) nitrogen content (percentage of dry mass) of above- and belowground biomass in year 1 (SP belowground not measured). Significantly different pairs are indicated by an asterisk (*t* test, $P < 0.05$, $N = 125$ for each creek for plant growth, and $N = 6$ for each creek for %N).

Ecosystem response

Vascular marsh plants.—Both *S. alterniflora* and *S. patens* growth increased with the nutrient addition, but stem density did not change (Fig. 5, Table 3). Nutrient enrichment increased tall *S. alterniflora* stem length by 20% in year 1 and increased length-specific leaf mass (g/cm of stem) by 25% in both years compared to the nutrient reference creek in creek pair 1 (Fig. 5a–d). Mean short-form *S. alterniflora* length-specific leaf mass was greater in both years, but the difference was

significant only in the first year of nutrient enrichment. *S. patens* length-specific leaf mass was indistinguishable between nutrient-enriched and reference creeks in either year of nutrient enrichment (Fig. 5c, d). In the first year of nutrient enrichment, mean height of *S. alterniflora* (tall and short) and *S. patens* began to diverge between the enriched and reference creeks in July, and by mid-August the height differences were significant. In the second year of nutrient enrichment, by mid-August *S. patens* was again significantly taller in the nutrient-

TABLE 3. Stem density (no. stems/m²; means \pm SE) in 2005 for tall *Spartina alterniflora*, short *Spartina alterniflora*, and marsh platform *Spartina patens*.

Marsh plant	Creek pair 1		Creek pair 2	
	Reference	Nutrient addition	Reference	Nutrient addition
Tall <i>Spartina alterniflora</i>	263 ^b \pm 15.9	206 ^a \pm 12.4	197 ^a \pm 8.9	173 ^a \pm 8.7
Short <i>Spartina alterniflora</i>	1336 ^a \pm 60	1336 ^a \pm 118	1358 ^a \pm 71	1532 ^a \pm 87
<i>Spartina patens</i>	6938 ^b \pm 2356	6263 ^{a,b} \pm 1246	5025 ^a \pm 1314	5575 ^a \pm 1767

Notes: Data for creek pair 1 are year-2 nutrient addition; data for creek pair 2 are year-1 nutrient addition. Different letters indicate differences among creeks within a row (Tukey, $P < 0.05$). $N = 24$ for each species for each creek.




FIG. 6. Benthic microbial production in tall *Spartina alterniflora* and *S. patens* habitats in creek pair 1 ambient-fish treatment after two years of nutrient enrichment. A significant difference from the one-to-one line was found for *S. alterniflora* (Wilcoxon signed-rank test, $Z = 3.5$, $P < 0.01$), but not for *S. patens* (Wilcoxon signed-rank test, $Z = 0.72$, $P > 0.10$).

enriched creek, however tall *S. alterniflora* height did not differ between creeks and short *S. alterniflora* was taller in the reference.

For both tall and short *S. alterniflora*, nitrogen content in aboveground shoots was higher in plants from the nutrient enrichment creek compared to the reference creek, while nitrogen content of *S. patens* was identical (Fig. 5e). For belowground biomass, tall *S. alterniflora* in the nutrient enrichment creek had a higher percentage N than the reference, but the short *S. alterniflora* did not differ in its percentage N. Over the growing season, nitrogen content of shoot tissue declined by ~35% in tall *S. alterniflora*, 42% in short *S. alterniflora*, and 55% in *S. patens*. Nitrogen content in belowground roots and rhizomes was higher in the nutrient enrichment creek for tall *S. alterniflora*, but short *S. alterniflora* had higher N content in the reference creek.

After two years of nutrient enrichment, we found no differences in stem density associated with nutrient enrichment for all plant species (Table 3). West had higher tall *S. alterniflora* stem density compared to the other three creeks, while short *S. alterniflora* stem density did not differ among creeks. Sweeney and West had higher *S. patens* stem densities than Clubhead and Nelson, but only West was significantly different.

Microbial production.—The bacterial production assay suggests that the productivity of microbes was

enhanced by nutrient enrichment in some marsh habitats, but was unaffected in others. In the tall *S. alterniflora* zone, rates of production were significantly higher in nutrient-enriched areas compared paired reference areas, but bacterial production in the *S. patens* zone was unaffected by nutrient enrichment (Fig. 6). These results suggest that the microbial response to nutrient enrichment is via some indirect mechanism that has differing effects in each marsh habitat.

Benthic microalgae.—Several lines of evidence demonstrate that benthic algae had a cumulative response to the nutrient addition and that the response to nutrient enrichment was greatest in the fish reduction treatment (Figs. 7 and 8). By the end of the second year of nutrient enrichment, under the ambient-fish treatment, benthic microalgal biomass was 60% higher than initial baseline and the reference creek for both tall-form *S. alterniflora* (TSA) and mudflat habitats (interaction term: mudflat, $F_{1,25} = 1.5$, $P < 0.09$; TSA, $F_{1,25} = 2.3$, $P < 0.004$; see Appendix B). Nutrient enrichment had a strong positive effect on benthic microalgal biomass in the fish reduction treatment in the second year (Fig. 7; t test, $P < 0.05$). Nutrient enrichment also altered the seasonal pattern and supported a high benthic microalgal biomass in mudflat and tall *S. alterniflora* habitats throughout the summer, while in reference creeks and pretreatment years microalgal biomass was lowest in midsummer. The pattern (1998–2002) of higher benthic microalgal biomass in tall *S. alterniflora* and mudflat habitats in the reference creek of creek pair 1 was also observed in the year immediately prior to manipulations (year 0). In the first year of nutrient enrichment of creek pair 1, microalgal biomass in the tall *S. alterniflora* habitat increased over baseline in the nutrient-enriched creek and was approximately equal to the mean for the reference creek (1998–2003).

The benthic microalgal response was spatially variable. Benthic microalgal biomass in *S. patens* habitat did not differ among any treatment or years ($F_{1,25} = 1.2$, $P = 0.25$). Creek pair 2 had no detectable change in benthic microalgal biomass associated with any treatment in any habitat (mudflat, $F_{1,11} = 0.5$, $P = 0.9$; TSA, $F_{1,11} = 1.18$, $P = 0.3$; *S. patens*, $F_{1,11} = 0.6$, $P = 0.8$; see Appendix B). Fish reduction had no discernable effect on microalgal biomass in the reference creeks (Fig. 7).

Microalgal numerical abundance (creek pair 1) followed the same pattern as mudflat microalgal biomass, with a significant increase in the nutrient-enriched creek by the end of year 2, although no differences were apparent among treatments in the first year. Again, the greatest response to nutrient enrichment was observed in the low-fish treatment (Fig. 8a).

Algal species richness increased in response to both the nutrient enrichment and the fish reduction (Fig. 8b). Similar to other measures of microalgal response, species richness increased more in response to the nutrient enrichment in the fish reduction treatment. Species composition differed with nutrient addition,




FIG. 7. Biomass of benthic microalgae (mg chl a/m^2) in mudflat, *Spartina alterniflora*, and *S. patens* habitats. Nutrient reference creeks are indicated by squares; nutrient enrichment creeks are circles. Open symbols are reduction in mummichog density treatments (low fish); solid symbols are ambient mummichog density treatments. Asterisks indicate significantly different ($P < 0.05$) creek pairs (ANOVA, Appendix B).

with more diatom species and no cyanobacteria species found in the nutrient-enriched creek.

Benthic invertebrates.—Infaunal annelid abundance declined with both nutrient enrichment and fish reduction. Total abundance ranged within 10 000–40 000 individuals/ m^2 (Fig. 9a), with a 50% reduction in abundance in the nutrient-enriched creek ($F_{1,20} = 19.13$, $P < 0.001$), and within a nutrient treatment slightly lower abundances in the low-fish treatment ($F_{1,20} = 3.02$, $P = 0.098$); effects were independent ($F_{1,20} = 0.00$, $P = 0.95$). Abundance of the amphipod (*U. spartinophila*) increased approximately threefold (Fig. 9b) with reduction of mummichog (t value = 4.70, $P < 0.0001$) in reference creeks.

Nekton community.—Daggerblade grass shrimp (*Palaeomonetes pugio*; 79% of abundance, 42% of biomass) and mummichog were the dominant nekton species (combined 98% of total abundance) out of 11 species. The other species were typical of salt marshes: *Crangon septemspinosus* (sand shrimp), *Carcinus maenas* (green crab), Xanthidae sp. (crabs), *Anguilla rostrata* (American eel), Clupeidae sp. (herring), *Menidia menidia* (Atlantic silversides), *Apeltes quadracus* (fourspine stick-

leback), *Pungitius pungitius* (ninespine stickleback), and *Syngathus fuscus* (northern pipefish). These species were not abundant or captured frequently enough to examine the effects of the experimental treatments.

Grass shrimp responded to the nutrient addition with increased growth, but abundance did not differ among treatments (Fig. 10). Grass shrimp abundance increased over the season, with peak abundance in the fall. The mean abundance of *P. pugio* varied within 50–1000 individuals/30 m^2 (sample range 0–1319 individuals/30 m^2). Grass shrimp abundance was variable among years and treatments and did not appear to be affected by any treatment in either creek pair (see Appendix B). Although we found a significant interaction term (creek pair 1: $F_{1,77} = 2.3$, 0.003) in creek pair 1, this did not appear to be attributable to any treatment effect (Fig. 10). Creek pair 2 did not have a significant interaction ($F_{1,73} = 0.89$, $P = 0.6$).

Growth of grass shrimp in nutrient addition creeks was 50% higher (~6 mm/month) than growth achieved in the reference creeks (~4 mm/month; Table 4). Shrimp grew between August (overall mean = 21 mm TL) and September (overall mean = 27 mm TL), with an overall




FIG. 8. (a) Cell counts (mean \pm SE) and (b) species richness (no. species) of benthic algae in creek pair 1. (a) Nutrient reference creeks are indicated by squares; nutrient enrichment creeks are circles. Open symbols are reduction in mummichog density treatments (low fish); solid symbols are ambient mummichog density treatments. Triangles indicate the start of manipulations. (b) "F" indicates filamentous species, and "S" indicates single-cell species. Data are from year 2.

effect of larger size in the nutrient-enriched marshes (nutrient: creek pair 1, $F_{1,3275} = 75.25$, $P < 0.001$; creek pair 2, $F_{1,1677} = 22.01$, $P < 0.0001$; see Appendix C); fish reduction had a small negative effect (creek pair 1, $F_{1,3275} = 43.89$, $P < 0.0001$; creek pair 2, $F_{1,1667} = 5.1$, $P = 0.023$) in both reference and nutrient addition creeks.

Mummichog diet was a mixture of detritus, algae, aquatic invertebrates (amphipods, polychaete worms, shrimp, isopods, copepods, and snails), and terrestrial insects (spiders, flies, and grasshoppers), as has been found in other saltmarshes, with an ontogenetic shift from largely carnivorous at small sizes to a more omnivorous, plant-based diet at larger sizes (Smith et al. 2000, Wainright et al. 2000, Fell et al. 2003). All fish examined had food in their guts; gut fullness was usually $>60\%$. In 2006, 20% of fish >40 mm TL in the nutrient-enriched, fish reduction treatment had algae as the most abundant food item in their stomachs. Detritus was ranked the most abundant item in all other fish.

DISCUSSION

Our results demonstrate ecological effects of both nutrient addition and predator reduction at the whole-system level, including evidence for synergistic interac-

tions. Our nutrient-loading rates were approximately 10 times less than previous dry-fertilizer plot-level experiments, yet we detected responses at several trophic levels within one to two years, suggesting that application of nutrients via daily water flooding is critical to understanding the impacts of coastal N enrichment. The results also suggest that eutrophication may have cumulative effects that are not apparent in a single year of nutrient enrichment and interactive effects with species composition. Observed changes in primary producers and decomposers suggest a long-term change in N-processing capacity in response to chronic N-enrichment.

Nutrient loading

Our annual N loadings were low compared to most plot-level experiments, despite the high concentrations we achieved in the water column. The simplest approach to estimating the loading rate in our experiment is to divide the total mass of N added by the area of the marsh, resulting in a watershed mean loading of $30 \text{ g N}\cdot\text{m}^{-2}\cdot\text{yr}^{-1}$. This approach does not account for the spatial gradient of nutrient delivery set up by the differential flooding by the tide and marsh elevation. We can estimate habitat-specific loading rates by using a marsh-flooding model (L. Harris, J. Vallino, and C. T. Friedrichs, unpublished data) that estimates the duration and depth of flooding over the experimental addition period of 150 d and measured nutrient concentrations.




FIG. 9. Abundance of (a) infauna (total annelids; no. individuals/m²) in creek pair 1 after one year of treatments (August 2004), and (b) epifaunal (no. *Uhlorchestia*) in ambient-fish and low-fish treatments in reference creeks (2005).




FIG. 10. Abundance of grass shrimp (*Palaemonetes pugio*; no. individuals/30 m²) in experimental creeks. Nutrient reference creeks are indicated by squares; nutrient enrichment creeks are circles. Open symbols are reduction in mummichog density treatments (low fish); solid symbols are ambient mummichog density treatments. Triangles on x-axes indicate the start of manipulations.

The intertidal tall *S. alterniflora* habitat experiences a tidally averaged 20 cm of water twice daily, resulting in an annual loading of ~ 60 g N·m⁻²·yr⁻¹. In the less frequently flooded *S. patens* marsh, the annual loading was ~ 15 g N·m⁻²·yr⁻¹. Nitrogen loading in plot-level experiments was generally 100–500 g N·m⁻²·yr⁻¹ (Mendelssohn 1979, Valiela 1983, Dai and Wiegert 1997, Boyer and Zedler 1998, Tyler et al. 2003).

We increased the average loading rate approximately a factor of 10 above background for this estuary;

ambient watershed loading to the whole Plum Island Sound estuary is ~ 3 g N·m⁻²·yr⁻¹ (Williams et al. 2004). Another perspective on our loading rate is that the amount of N that we added was roughly equivalent to the nutrient loading from the addition of 1000 houses in the upland watershed of each of the nutrient-enriched marsh creeks (Cape Cod Commission 1992).

The consistent decrease in NO₃⁻ concentration in ebbing water in both creek pairs suggests retention of some of the added NO₃⁻ by the marsh ecosystem. Of course, the entire nutrient content of the water is not removed, because at high tides water depths are too deep and residence time too short to allow for complete assimilation by marsh processes. A rough calculation, based on the incoming and outgoing nutrient concentrations and a net water balance, suggests that 30–40% of the added nitrogen did not leave the marsh in the ebbing water. Hydrodynamic modeling and δ¹⁵N tracer addition experiments are currently being used to obtain a more precise estimate of the retention and fate of the added nutrients.

Question 1: response of vascular plants and microbial decomposers

The dominant primary producers, *Spartina* spp. and benthic microalgae, generally respond positively to N fertilization (Mendelssohn and Morris 2000, Sullivan and Currin 2000). The combined results of previous studies suggests the dose–response of *Spartina* production was nonlinear, with little increase with low N additions (20 g N·m⁻²·yr⁻¹; Valiela 1983), steep increases at medium amounts (100–500 g N·m⁻²·yr⁻¹), and then no further increases at higher N loading rates (Mendelssohn 1979).

Contrary to our expectation that, because of the relatively low loading, the plant response would not be detectable, *S. alterniflora* and perhaps *S. patens* production has increased. The first two years of nutrient enrichment in creek pair 1 increased the length-specific biomass in tall, creek bank *S. alterniflora*, resulting in increased aboveground plant production. As in most plot-level nutrient-loading experiments (e.g., Valiela 1983), we found no difference in stem density for any plant species or growth form. Increased production

TABLE 4. Young-of-the-year grass shrimp (*Palaemonetes pugio*) size (mean total length ± SE) and growth rate (TL) between August and September 2005.

Sample	Reference		Nutrient addition	
	Fish (mm)	Low fish (mm)	Fish (mm)	Low fish (mm)
Creek pair 1				
August	21.69 ± 0.03	21.55 ± 0.02	23.02 ± 0.03	21.02 ± 0.01
September	26.97 ± 0.01	24.52 ± 0.01	27.93 ± 0.01	24.71 ± 0.01
Growth	5.28	2.97	4.91	6.39
Creek pair 2				
August	21.85 ± 0.02	21.74 ± 0.03	21.53 ± 0.01	21.25 ± 0.01
September	26.16 ± 0.01	26.07 ± 0.01	28.53 ± 0.01	27.76 ± 0.01
Growth	4.31	4.33	7.00	6.01



PLATE 1. Early season (June 2004) sampling for benthic algae and invertebrates on the marsh platform near the head of the West Creek of the Plum Island Sound estuary in New England, USA (David Johnson on the left and Kari Galván on the right standing in tall *Spartina alterniflora* habitat). Intertidal creek mudflat habitat can be seen at the bottom of the photo. The expanse of the lighter grass in the background is high marsh platform *S. patens* habitat. Photo credit: L. Deegan.

resulted from greater biomass per unit length of shoot or longer shoots, or in some cases both. We observed a response in year 1 of creek pair 1, but not in year 1 of creek pair 2. In the first year of some studies, researchers found little response (Valiela 1983), while others found differences with nutrient enrichment (Tyler et al. 2003).

The higher aboveground plant nitrogen content in nutrient-enriched plants supports the suggestion that both growth forms of *S. alterniflora* responded to our low nutrient loading. Nitrogen content was consistently greater in both tall and short forms of *S. alterniflora* in the nutrient-enriched compared to the reference creek, as has been found in many other nutrient enrichment studies (Tyler et al. 2003). Interestingly, despite taller plants in both years, nitrogen content of *S. patens* was identical in the two creeks. Nitrogen content in roots and rhizomes was less clear-cut, with a tendency toward greater nitrogen in tall *S. alterniflora* in the nutrient-enriched creek compared to the reference, and no difference in short *S. alterniflora* from the high marsh. This result is similar to Boyer et al. (2000) who found that <2% of N added was incorporated into root tissue and that there was only a slight trend toward increased belowground biomass with N nutrient enrichment in a created *Spartina foliosa* marsh.

Spartina spp. exhibits low uptake rates of NO_3^- from overlying water (Wright et al. 1996) consistent with *Spartina* obtaining its nutrients from belowground. The

probable mechanism for increased growth of *S. alterniflora* near creek edges is tidal pumping (Mendelssohn and Morris 2000), which flushed nutrient-rich water through the root zone at the creek edge. The ditching of the marsh increased water inundation and nutrient transport onto the high marsh, potentially increasing the *S. alterniflora* edge effect. The response of *S. patens* was probably limited by the infrequent flooding of the high marsh with the enriched water and because the nutrients do not penetrate deeply into the already water-saturated soil of the high marsh during the short period of time (3–4 h) that the marsh is flooded each tide.

Microbial production.—Nutrient enrichment was expected to stimulate bacterial activity and decomposition throughout the ecosystem because of direct uptake of N from the water column and availability of more high-quality organic matter from increased algal production (Howarth and Hobbie 1982). We expected that in sparsely vegetated saltmarsh habitats, such as creek bank *S. alterniflora*, where there is sufficient light penetration for benthic microalgae to proliferate (Pinckney and Zingmark 1993), rates of bacterial production would increase as an indirect result of the increase in highly labile carbon from benthic microalgae. Conversely in more densely vegetated habitats, such as *S. patens* high marsh, we anticipated that, because of the large store of carbon in marsh peat, bacteria would be

nitrogen limited and bacterial production would be directly enhanced by new nitrogen inputs.

We found a microbial production increase only in the tall *S. alterniflora* habitat, not in the *S. patens* high-marsh habitat. If bacteria in these salt marsh sediments were responding directly to the increased nitrogen input, we expected to see stimulation of bacterial production in both creek bank *S. alterniflora* and *S. patens* habitats. The lack of a response to the nutrient addition in the *S. patens* high-marsh habitat was surprising, given the large store of carbon and the importance of microbial decomposition in salt marshes (Howes et al. 1984, Buchan et al. 2003). Nitrate additions in laboratory experiments have stimulated aerobic saltmarsh detritus decomposition (Howarth and Hobbie 1982, Morris and Bradley 1999) and anoxic decomposition via NO_3^- reduction (Zumft 1991).

Because we see microbial production stimulation only in the nutrient-enriched creek bank *S. alterniflora* habitat where benthic microalgae also increased, this suggests that bacteria are responding primarily to increases in high-quality organic matter from benthic microalgae, not to increases in dissolved nitrogen. Although saltmarsh macrophytes dominate the total primary productivity in marsh ecosystems, stable isotopic evidence indicates that it may not be a preferred carbon source for marsh sediment bacteria even in the presence of elevated inorganic nitrogen. Stable carbon isotope ratios of bacterial specific polar lipid-derived fatty acids from the Waarde Marsh, The Netherlands, indicated that the dominant carbon source for marsh sediment bacteria was algal (Boschker et al. 1999). Similarly, in a $\delta^{15}\text{N}$ tracer addition experiment in the Rowley River, Massachusetts, USA, the $\delta^{15}\text{N}$ value of the bacterial-specific biomarker diaminopimelic acid indicated that the bacteria were more closely linked to benthic microalgae than to bulk-sediment detritus (Tobias et al. 2003). These findings support the hypothesis that bacterial production is indirectly stimulated through increased production of benthic microalgae. We are examining this hypothesis through a $\delta^{15}\text{N}$ tracer addition experiment in the nutrient-enriched and reference marshes and with litterbag experiments.

Question 2: top-down controls

We found evidence for top-down control on benthic microalgae and on marsh epifauna, but not on benthic infauna. Top-down effects on benthic microalgae were seen only under nutrient enrichment, suggesting that killifish influence microalgae via direct effects (e.g., herbivory). The response of the benthic microalgae is suggestive of a synergistic effect of nutrients and top-down control, and perhaps a self-shading limitation (Hillebrand 2005). Benthic algal biomass and productivity typically increased in response to N additions, but these increases were often limited by marsh grass shading and grazing (Pinckney and Zingmark 1993, Sullivan and Currin 2000). The lack of response in

microalgal biomass in nutrient-enriched *S. patens* was consistent with previous studies that found light-limited benthic algal production under the dense *S. patens* canopy; we found mean light under the *S. patens* canopy was <3% of ambient (S. Sheldon, unpublished data).

In the *S. alterniflora* habitat, where light is often not limiting, the simplest microalgal-response model assumes a linear increase in biomass with nutrient addition and no consumer control. This model predicts that benthic microalgae would increase from 150 (background) to 2000 mg chl *a*/m² with our 15-fold increase in nutrient enrichment, if there were no other limiting factors. While we saw nearly a doubling of microalgal biomass in year 2 with nutrient enrichment and fish reduction (to 250 mg chl *a*/m²), the increase was clearly not directly proportional to added nutrients. Species composition also changed, with more diatom species and few cyanobacteria species found in the nutrient-enriched creek, suggesting potential increases in the palatability of algae and loss of N fixation. Typical values for benthic microalgal biomass in the *S. alterniflora* habitat are ~100 mg chl *a*/m² (Sullivan and Currin 2000), suggesting that the initial high microalgal biomass might have limited the scope for response before self-shading became limiting (Hillebrand 2005). The highest microalgal biomass measured in any salt marsh habitat was 800 mg chl *a*/m² in a cyanobacteria mat in a high-marsh salt panne (Sullivan and Currin 2000). Our nutrient addition decreased the abundance of cyanobacteria, thus development of a dense cyanobacteria mat is not likely to occur. It is doubtful that we will ever measure 2000 mg chl *a*/m² unless the algal community changes from domination by diatoms to filamentous algae or macroalgae.

Grazers also clearly play a role in the response of benthic microalgae to nutrient additions (Sullivan and Currin 2000). We found a top-down control on benthic microalgae, but only under conditions of nutrient addition. This is consistent with other studies that found the largest increases in benthic microalgae in response to increased nitrogen occurred when grazers were removed (Sullivan and Currin 2000). Killifish is considered a secondary consumer, with a diet of small invertebrates, benthic algae, and detritus (Deegan et al. 2000, Currin et al. 2003). Contrary to our expectations, benthic microalgae increased with mummichog reduction, suggesting either that mummichog exert top-down control by direct grazing, or that the trophic cascade operated in ways we do not understand. Based on previous work, we expected that mummichog removal would result in a trophic cascade in which infauna increase and benthic microalgal biomass would decrease in response to increased infaunal grazing. Instead, our results suggest that, under conditions of high algal productivity, killifish may drop in trophic level from carnivore to herbivorous grazer and become an important grazer in a nutrient-enriched environment.

Counter-intuitively, our results show that small infauna abundance decreased slightly with mummichog reduction. We did not find the predicted increase in small infauna expected if release from predation by mummichog controlled their abundance. Our methods of lower mummichog abundance were most effective on fish >40 mm, and larger mummichogs probably prey more heavily on larger epifauna (e.g., amphipods and grass shrimp) and algae than on small infauna (Curran et al. 2003, Fell et al. 2003), perhaps explaining the absence of direct top-down control on infauna. An increase in the importance of algae in the diet of large mummichogs remaining in the fish reduction treatment supports the potential importance of direct grazing; this idea is also being examined by stable-isotope analysis. Additionally, *Spartina* stems may inhibit foraging by *F. heteroclitus* (Walters et al. 1996, Carson and Merchant 2005), thereby precluding strong direct effects in grass habitats.

Our experimental reduction of mummichogs may have affected an intermediate predator of infauna, the grass shrimp, and perhaps initiated an indirect interaction. *P. pugio* graze epiphytic algae and forage for macroinfauna (Fleeger et al. 1999). Smaller mummichogs compete with grass shrimp for infaunal food, and larger mummichog prey on grass shrimp (Kneib and Stiven 1982, Cross and Stiven 1999, Weis et al. 2000). Mummichog removal was thus predicted to increase grass shrimp growth and abundance and thereby increase grazing pressure on microalgae and infauna. Although we found no effect on abundance of grass shrimp with mummichog reduction, fish reduction may have allowed grass shrimp to modify their behavior. Carson and Merchant (2005) found that grass shrimp reduce swimming behavior in the absence of large mummichogs, resulting in increased benthic contact time that may facilitate grass shrimp feeding activities. Thus, observed declines in infauna in areas where mummichog abundance was experimentally reduced could be mediated by increased grass shrimp predation. We are currently examining infaunal responses in other marsh habitats and quantifying species composition to determine if responses to mummichog reduction were habitat or species specific (Posey et al. [1999] found species-specific responses).

An important prey species for large mummichog is the intertidal amphipod (*Uholchestia spartinophila*), a large epifaunal amphipod (Covi and Kneib 1995; K. Galván, unpublished manuscript). Amphipod abundance in *S. alterniflora* habitats more than doubled with mummichog reduction (Fig. 9), suggesting release from predation. *U. spartinophila* is widely distributed through the vegetated intertidal marsh, unlike other talitrid amphipods that are restricted to supralittoral zones (Covi and Kneib 1995), and our results demonstrate that their abundances may be regulated by mummichog predation. This may have consequences for long-term gains or losses in marsh elevation, as these amphipods are detritivores that can accelerate the decomposition of *S.*

alterniflora litter by stimulating microbial activity (Lopez et al. 1977, Bousfield and Heard 1986). These amphipods also feed on benthic microalgae (K. A. Galván, unpublished data), and it is possible that they are important grazers in this system. Additional studies are planned to examine treatment effects on amphipod activity patterns.

Unexpectedly, infauna responded to nutrient addition with reduced abundance in the first year of nutrient addition. We are currently examining annelid biomass to determine if there was a per capita increase in biomass with nutrient addition, as found by Posey et al. (2006). This first year of nutrient addition did not bring about large increases in benthic microalgae, perhaps explaining why infauna did not respond in a bottom-up cascade. Furthermore, the increased body size of grass shrimp in nutrient-enriched creeks suggests they exerted increased predation pressure on infauna. Collections from the second year of nutrient addition are currently being processed.

Overall, we found that top-down effects may modify the responses of benthic microalgae to nutrient enrichment, i.e., a synergistic effect on benthic microalgae biomass occurred when nutrients were added and when mummichog was reduced in saltmarsh creeks, but that light limitation prevented similar responses in the high marsh. Although nutrient addition and predator reduction effects noted to date are best expressed as independent on benthic infauna, additional collections across the marsh landscape are being processed. Most tests for interactions among environmental factors are based on small-scale exclusion (Posey et al. 1999) or mesocosm studies (Breitburg et al. 1999); we anticipate that our whole-ecosystem study will provide a more realistic picture of how nutrient addition and predator reduction interact throughout a salt marsh.

CONCLUSIONS

Human activity is changing local and global environments at unprecedented rates. Some of the most significant changes involve global warming, sea level rise, widespread nutrient enrichment, and species changes. All of these affect saltmarsh ecosystems. Understanding the responses of saltmarsh ecosystems to stressors, as well as the mechanisms and processes that underlie these responses, is fundamental to the pressing regulatory and cultural decisions that many communities now face. Our project fills a unique role in provision of an ecosystem-scale, multiyear perspective for management decisions, such as installation of sewage treatment plants or regulating housing density, that will affect coastal communities and ecosystems for generations.

The initial results and analyses presented here demonstrate ecological effects of both nutrient addition and fish reduction at the whole-system level. Evidence of synergistic interactions between these stressors is also emerging, especially at higher trophic levels. Given the

cumulative effects we have observed, it is far too early to make long-term predictions about the relative stimulation of plant production vs. microbial decomposition. Although we have identified some early effects of these stressors, we know that saltmarsh ecosystems will respond to stress over time periods much longer than two years; some effects may be cumulative, and some, such as plant community change, may take many years to develop. This highlights the importance and potential benefit of continuing the manipulations for several additional years. Related, ongoing studies within the project include sediment dynamics, hydrodynamics, benthic microalgal physiological responses, phytoplankton community and nutrient cycling, and isotopic tracer studies examining food webs and biogeochemistry; these will add substantially to our understanding of the effects of multiple anthropogenic stressors on saltmarsh ecosystems. Our continuing work examines response thresholds and time lags, the mechanisms regulating saltmarsh responses, cumulative responses over longer periods of disturbance, and, we hope eventually, recovery.

ACKNOWLEDGMENTS

The National Science Foundation (grant DEB 0213767, OCE 9726921, and OCE 0423565) supported this work. Additional funding was provided by a National Science Foundation postdoctoral fellowship in Microbial Biology (DBI-0400819), the NOAA Coastal Intern grant (NA04NOS4780182), the Office of Environmental Education of Louisiana, Middlebury College, and Connecticut College. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the funding agencies. We thank Bonnie Keeler for her role in the initial design and implementation of the experiment and Lora Harris for thoughtful comments on the manuscript. Finally, the authors thank the many interns and undergraduates for their many hours of work and contributions to this project. It would not have been as fun without them. Note also that, after the first author, names are in alphabetical order.

LITERATURE CITED

- Bormann, F., and E. Likens. 1979. Pattern and process in a forested ecosystem. Springer-Verlag, New York, New York, USA.
- Boschker, H. T. S., J. F. C. de Brouwer, and T. E. Cappenberg. 1999. The contribution of macrophyte-derived organic matter to microbial biomass in salt-marsh sediments: stable carbon isotope analysis of microbial biomarkers. *Limnology and Oceanography* 44:309–319.
- Bousfield, E. L., and R. W. Heard. 1986. Systematics, distributional ecology, and some host–parasite relationships of *Uhlorchestia uhleri* (Shoemaker) and *U. spartinophila*, new species (Crustacea: Amphipoda), endemic to salt marshes of the Atlantic coast of North America. *Journal of Crustacean Biology* 6(2):264–274.
- Boyer, K. E., J. C. Calloway, and J. B. Zedler. 2000. Evaluating the progress of restored cordgrass (*Spartina foliosa*) marshes: belowground biomass and tissue nitrogen. *Estuaries* 23:711–721.
- Boyer, K., and J. Zedler. 1998. Effects of nitrogen additions on the vertical structure of a constructed cordgrass marsh. *Ecological Applications* 8:692–705.
- Breitburg, D. L., J. G. Sanders, C. C. Gilmour, C. A. Hatfield, R. W. Osman, G. F. Riedel, S. B. Seitzinger, and K. G. Sellner. 1999. Variability in responses to nutrients and trace elements, and transmission of stressor effects through an estuarine food web. *Limnology and Oceanography* 44:837–863.
- Buchan, A., S. Y. Newell, M. Butler, E. J. Biers, J. T. Hollibaugh, and M. A. Moran. 2003. Dynamics of bacterial and fungal communities on decaying salt marsh grass. *Applied and Environmental Microbiology* 69:6676–6687.
- Buesing, N., and M. O. Gessner. 2003. Incorporation of radiolabeled leucine into protein to estimate bacterial production in plant litter, sediment, epiphytic biofilms, and water samples. *Microbial Ecology* 45:291–301.
- Cape Cod Commission. 1992. Nitrogen loading. Technical Bulletin 91-001, Cape Cod Commission Water Resource Office, Barnstable, Massachusetts, USA.
- Carman, K. R., J. W. Fleeger, and S. Pomarico. 1997. Response of a benthic food web to hydrocarbon contamination. *Limnology and Oceanography* 42:561–571.
- Carpenter, S. R., S. W. Chisholm, C. J. Krebs, D. W. Schindler, and R. F. Wright. 1995. Ecosystem experiments. *Science* 26:324–327.
- Carson, M. L., and H. Merchant. 2005. A laboratory study of the behavior of two species of grass shrimp (*Palaemonetes pugio* Holthuis and *Palaemonetes vulgaris* Holthuis) and the killifish (*Fundulus heteroclitus* Linnaeus). *Journal of Experimental Marine Biology and Ecology* 314:187–201.
- Cloern, J. E. 2001. Our evolving conceptual model of the coastal eutrophication problem. *Marine Ecology Progress Series* 210:223–253.
- Covi, M. P., and R. T. Kneib. 1995. Intertidal distribution, population dynamics and production of the amphipod *Uhlorchestia spartinophila* in a Georgia, USA, salt marsh. *Marine Biology* 121:447–455.
- Cross, R. E., and A. E. Stiven. 1999. Size-dependent interactions in salt marsh fish (*Fundulus heteroclitus* Linnaeus) and shrimp (*Palaemonetes pugio* Holthuis). *Journal of Experimental Marine Biology and Ecology* 242:179–199.
- Currin, C. A., S. C. Wainright, K. W. Able, M. P. Weinstein, and C. M. Fuller. 2003. Determination of food web support and trophic position of the Mummichog, *Fundulus heteroclitus*, in New Jersey smooth cordgrass (*Spartina alterniflora*), common reed (*Phragmites australis*), and restored salt marshes. *Estuaries* 26:495–510.
- Dai, T., and R. G. Wiegert. 1997. A field study of photosynthetic capacity and its response to nitrogen fertilization in *Spartina alterniflora*. *Estuarine and Coastal Shelf Science* 45:273–283.
- D'Avanzo, C., and I. Valiela. 1990. Use of detrital foods and assimilation of nitrogen by coastal detritivores. *Estuaries* 13(1):20–24.
- Deegan, L. A., J. E. Hughes, and R. A. Rountree. 2000. Salt marsh ecosystem support of marine transient species. Pages 333–365 in M. P. Weinstein and D. A. Kreeger, editors. Concepts and controversies in tidal marsh ecology. Kluwer Academic, Dordrecht, The Netherlands.
- Deegan, L. A., A. Wright, S. G. Ayvazian, J. T. Finn, H. Golden, R. Rand Merson, and J. Harrison. 2002. Nitrogen loading from upland areas alters seagrass support of higher trophic levels. *Aquatic Conservation: Freshwater and Marine Ecosystems* 12:193–212.
- Emeis, K. C., J. R. Benoit, L. Deegan, A. J. Gilbert, V. Lee, J. M. Glade, M. Meybeck, S. B. Olsen, and B. von Bodungen. 2001. Unifying concepts for integrated coastal management. Pages 341–364 in B. Von Bodungen and R. K. Turner, editors. Science and integrated coastal management. Dahlem University Press, Berlin, Germany.

- Environmental Protection Agency. 2002. Mid-Atlantic integrated assessment (MAIA) estuaries 1997–1998. Summary report. U.S. Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Atlantic Ecology Division, Narragansett, Rhode Island, USA.
- Fell, P. E., R. S. Warren, J. K. Light, R. L. Rawson, Jr., and S. M. Fairley. 2003. Comparison of fish and microinvertebrate use of *Typha angustifolia*, *Phragmites australis*, and treated *Phragmites* marshes along the lower Connecticut River. *Estuaries* 26:534–551.
- Fell, P. E., S. P. Weissbach, D. A. Jones, M. A. Fallon, J. A. Zeppieri, E. K. Faison, K. A. Lennon, K. J. Newberry, and L. K. Reddington. 1998. Does invasion of oligohaline tidal marshes by reed grass, *Phragmites australis*, affect the availability of prey resources for the mummichog, *Fundulus heteroclitus*? *Journal of Experimental Marine Biology and Ecology* 222:59–77.
- Fleeger, J. W., K. R. Carman, S. Webb, N. Hilbun, and M. C. Pace. 1999. Consumption of microalgae by the grass shrimp, *Palaemonetes pugio*. *Journal of Crustacean Biology* 19:324–336.
- Friedrichs, C. T., and J. E. Perry. 2001. A review of tidal salt marsh morphodynamics. *Journal of Coastal Research, Special Issue* 27:7–37.
- Hillebrand, H. 2005. Light regime and consumer control of autotrophic biomass. *Journal of Ecology* 93:758–769.
- Hobbs, N. T., and R. Hilborn. 2006. Alternatives to statistical hypothesis testing in ecology: a guide to self teaching. *Ecological Applications* 16:5–19.
- Holm-Hansen, O., C. Lorenzen, R. Holmes, and J. Strickland. 1965. Fluorometric determination of chlorophyll. *Journal du Conseil Permanent International Pour l'Exploration de la Mer* 30:3–15.
- Howarth, R. W., and J. E. Hobbie. 1982. The regulation of decomposition and heterotrophic microbial activity in salt marsh soils. Pages 183–207 in V. S. Kennedy, editor. *Estuarine comparisons*. Academic Press, New York, New York, USA.
- Howes, B. L., J. W. H. Dacey, and G. M. King. 1984. Carbon flow through oxygen and sulfate reduction pathways in salt marsh sediments. *Limnology and Oceanography* 29:1037–1051.
- Hurlbert, S. H. 1984. Pseudoreplication and the design of ecological field experiments. *Ecological Monographs* 54:187–211.
- Hyslop, E. J. 1980. Stomach contents analysis—a review of methods and their application. *Journal of Fish Biology* 17: 411–429.
- Kneib, R. T. 1997. The role of tidal marshes in the ecology of estuarine nekton. *Oceanography and Marine Biology: An Annual Review* 35:163–220.
- Kneib, R. T., and A. E. Stiven. 1982. Benthic invertebrate responses to size and density manipulations of the common mummichog, *Fundulus heteroclitus*, in an intertidal salt marsh. *Ecology* 63:1518–1532.
- Layman, C. A., and D. E. Smith. 2001. Sampling bias of minnow traps in shallow aquatic habitats on the Eastern Shore of Virginia. *Wetlands* 21:145–154.
- Levine, J., S. Brewer, and M. Bertness. 1998. Nutrients, competition and plant zonation in a New England salt marsh. *Journal of Ecology* 86:285–292.
- Lopez, G. R., J. S. Levington, and L. B. Slobodkin. 1977. The effect of grazing by the detritivore *Orchestia grillus* on *Spartina* litter and its associated microbial community. *Oecologia* 30: 111–127.
- Lorenzen, G. L. 1967. Determination of chlorophyll and phaeopigments; spectrophotometric equations. *Limnology and Oceanography* 12:343–346.
- Matson, P., and M. D. Price. 1992. The relative importance of top-down and bottom-up forces in populations and community ecology. *Ecology* 73:723.
- Mendelsohn, I. A. 1979. The influence of nitrogen level, form and application method on the growth response of *Spartina alterniflora* in North Carolina. *Estuaries* 2:106–112.
- Mendelsohn, I. A., and J. T. Morris. 2000. Eco-physiological controls on the productivity of *Spartina alterniflora* Loisel. Pages 59–80 in M. P. Weinstein and D. A. Kreeger, editors. *Concepts and controversies in tidal marsh ecology*. Kluwer Academic, Dordrecht, The Netherlands.
- Morris, J. M., and P. Bradley. 1999. Effects of nutrient loading on the carbon balance of coastal wetland sediments. *Limnology and Oceanography* 44:699–702.
- National Oceanic and Atmospheric Administration (NOAA). 1999. National estuarine eutrophication assessment: effects of nutrient enrichment in the nation's estuaries. National Ocean Service, NOAA, U.S. Department of Commerce.
- National Research Council. 2000. *Clean coastal waters: understanding and reducing the effects of nutrient pollution*. National Academy Press, Washington, D.C., USA.
- Nixon, S. W. 1995. Coastal marine eutrophication: a definition, social causes and future concerns. *Ophelia* 41:199–219.
- Oksanen, L. 2001. Logic of experiments in ecology: is pseudoreplication a pseudoissue? *Oikos* 94:27–38.
- Pace, M. L., J. J. Cole, S. R. Carpenter, and J. F. Kitchell. 1999. Trophic cascades revealed in diverse ecosystems. *Trends in Ecology and Evolution* 14(12):483–488.
- Parker, K. R., and J. A. Wiens. 2005. Assessing recovery following environmental accidents: environmental variation, ecological assumptions, and strategies. *Ecological Applications* 15:2037–2051.
- Parsons, T. R., Y. Maita, and C. M. Lalli. 1984. *A manual of chemical and biological methods for seawater analysis*. Pergamon Press, New York, New York, USA.
- Pennings, S., and M. D. Bertness. 2001. Salt marsh communities. Pages 289–316 in M. D. Bertness, S. D. Gaines, and M. Hay, editors. *Marine community ecology*. Sinauer Associates, Sunderland, Massachusetts, USA.
- Pinckney, J. L., and R. G. Zingmark. 1993. Modeling the annual production of intertidal benthic microalgae in estuarine ecosystems. *Journal of Phycology* 29:396–407.
- Polis, G. A. 1999. Why are parts of the world green? Multiple factors control productivity and the distribution of biomass. *Oikos* 86:3–15.
- Posey, M. H., T. D. Alphin, and L. Cahoon. 2006. Benthic community responses to nutrient enrichment and predator exclusion: influence of background nutrient concentrations and interactive effects. *Journal of Experimental Marine Biology and Ecology* 30:105–118.
- Posey, M. H., T. D. Alphin, L. Cahoon, D. Lindquist, and M. E. Becker. 1999. Interactive effects of nutrient additions and predation on infaunal communities. *Estuaries* 22:785–792.
- Posey, M. H., T. D. Alphin, L. Cahoon, D. Lindquist, M. A. Mallin, and M. B. Nevers. 2002. Top-down versus bottom-up limitation in benthic infaunal communities: direct and indirect effects. *Estuaries* 25:999–1014.
- Quiñones-Rivera, Z. J., and J. W. Fleeger. 2005. The grazing effects of grass shrimp, *Palaemonetes pugio*, on epiphytic microalgae associated with *Spartina alterniflora*. *Estuaries* 28: 274–285.
- Raposa, K. 2003. Overwintering habitat selection by the mummichog, *Fundulus heteroclitus*, in a Cape Cod (USA) salt marsh. *Wetlands Ecology and Management* 11:175–182.
- Rapport, D. J., and W. G. Whitford. 1999. How ecosystems respond to stress. *Bioscience* 49:193–203.
- Sarda, R., K. Foreman, C. E. Werme, and I. Valiela. 1998. The impact of epifaunal predation on the structure of macro-

- infaunal invertebrate communities of tidal saltmarsh creeks. *Estuarine and Coastal Shelf Science* 46:657–669.
- Schindler, D. W. 1998. Replication versus realism: the need for ecosystem-scale experiments. *Ecosystems* 3:323–334.
- Silliman, B. R., and M. D. Bertness. 2002. A trophic cascade regulates salt marsh primary productivity. *Proceedings of the National Academy of Sciences (USA)* 99:10500–10505.
- Silliman, B. R., J. van de Koppel, M. D. Bertness, L. E. Stanton, and I. A. Mendelsohn. 2005. Drought, snails, and large-scale die-off of southern U.S. salt marshes. *Science* 310:1803–1806.
- Silliman, B. R., and J. C. Zeiman. 2001. Top-down control of *Spartina alterniflora* production by periwinkle grazing in a Virginia salt marsh. *Ecology* 82:2830–2845.
- Smith, K. J., G. L. Taghon, and K. W. Able. 2000. Trophic linkages in marshes: ontogenetic changes in diet for young-of-the-year mummichog *Fundulus heteroclitus*. Pages 221–237 in M. P. Weinstein and D. A. Kreeger, editors. *Concepts and controversies in tidal marsh ecology*. Kluwer Academic, Dordrecht, The Netherlands.
- Solorzano, L. 1969. Determination of ammonium in natural waters by the phenylhypochlorite method. *Limnology and Oceanography* 14:799–801.
- Stewart-Oaten, A., and J. R. Bence. 2001. Temporal and spatial variation in environmental impact assessment. *Ecological Monographs* 71:305–339.
- Strong, D. 1992. Are trophic cascades all wet? Differentiation and donor-control in speciose ecosystems. *Ecology* 73:747–755.
- Sullivan, M. J., and C. A. Currin. 2000. Community structure and functional dynamics of benthic microalgae in saltmarshes. Pages 81–106 in M. P. Weinstein and D. A. Kreeger, editors. *Concepts and controversies in tidal marsh ecology*. Kluwer Academic, Dordrecht, The Netherlands.
- Sweeney, J., L. Deegan, and R. Garritt. 1998. Population size and site fidelity of *Fundulus heteroclitus* in a macrotidal saltmarsh creek. *Biological Bulletin* 195:238–239.
- Tobias, C. R., M. Cieri, B. J. Peterson, L. A. Deegan, J. Vallino, and J. Hughes. 2003. Processing watershed-derived nitrogen in a well-flushed New England estuary. *Limnology and Oceanography* 48(5):1766–1778.
- Tyler, A. C., T. A. Mastrorcola, and K. J. McGlathery. 2003. Nitrogen fixation and nitrogen limitation of primary production along a natural marsh chronosequence. *Oecologia* 136:431–438.
- Underwood, A. 1994. On beyond BACI: sampling designs that might reliably detect environmental disturbances. *Ecological Applications* 4:3–15.
- Valiela, I. 1983. Nitrogen in salt marsh ecosystems. Pages 649–678 in E. J. Carpenter and D. G. Capone, editors. *Nitrogen in the marine environment*. Academic Press, New York, New York, USA.
- Verhoeven, J. T. A., B. Arheimer, C. Yin, and M. M. Hefting. 2006. Regional and global concerns over wetlands and water quality. *Trends in Ecology and Evolution* 21:96–103.
- Wainright, S. C., M. P. Weinstein, K. W. Able, and C. A. Currin. 2000. Relative importance of benthic microalgae, phytoplankton and the detritus of smooth cordgrass *Spartina alterniflora* and the common reed *Phragmites australis* to brackish-marsh food webs. *Marine Ecology Progress Series* 200:77–91.
- Walters, K., E. Jones, and L. Etherington. 1996. Experimental studies of predation on metazoans inhabiting *Spartina alterniflora* stems. *Journal of Experimental Marine Biology and Ecology* 195:251–265.
- Weis, J. S., G. Smith, and C. Santiago-Bass. 2000. Predator/prey interactions: a link between the individual level and both higher and lower level effects of toxicants in aquatic ecosystems. *Journal of Aquatic Ecosystem Stress and Recovery* 7:145–153.
- Williams, M., C. Hopkinson, E. Rastetter, and J. Vallino. 2004. N budgets and aquatic uptake in the Ipswich River Basin, northeastern Massachusetts. *Water Resources Research* 11:1–12.
- Wood, E. D., F. A. J. Armstrong, and F. A. Richards. 1967. Determination of nitrate in sea water by cadmium–copper reduction to nitrite. *Journal of the Marine Biological Association of the United Kingdom* 47:23.
- Worm, B., H. Lotze, and U. Sommers. 2000. Coastal food web structure, carbon storage, and nitrogen retention regulated by consumer pressure and nutrient loading. *Limnology and Oceanography* 45:339–349.
- Wright, A. L., R. W. Weaver, and J. W. Webb. 1996. Concentrations of N and P in floodwater and uptake of ¹⁵N by *Spartina alterniflora* in oil-contaminated mesocosms. *Bioresource Technology* 56:257–264.
- Zumft, W. G. 1991. The denitrifying bacteria. Pages 554–582 in A. Balows, M. Dworkin, H. G. Schlegel, and H. Trueper, editors. *The prokaryotes*. Springer-Verlag, Berlin, Germany.

APPENDIX A

Two-way ANOVA (creek, branch) on routine sampling (1998–2002) prior to manipulations (*Ecological Archives* A017-063-A1).

APPENDIX B

BACI-type ANOVA results for effects of treatments and interactions on benthic algae biomass, nekton abundance, and nekton biomass (*Ecological Archives* A017-063-A2).

APPENDIX C

ANOVA of body size of young-of-the-year grass shrimp (*Palaemonetes pugio*) in 2005 (*Ecological Archives* A017-063-A3).