
Reports

2014

Estimation of juvenile striped bass relative abundance in the Virginia portion of Chesapeake Bay : annual progress report : January 2013-December 2013

Mary C. Fabrizio
Virginia Institute of Marine Science

Troy D. Tuckey
Virginia Institute of Marine Science

Christopher D. Davis
Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Aquaculture and Fisheries Commons](#), and the [Marine Biology Commons](#)

Recommended Citation

Fabrizio, M. C., Tuckey, T. D., & Davis, C. D. (2014) Estimation of juvenile striped bass relative abundance in the Virginia portion of Chesapeake Bay : annual progress report : January 2013-December 2013. Virginia Institute of Marine Science, William & Mary. <http://dx.doi.org/doi:10.21220/m2-8rz1-3k08>

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Estimation of Juvenile Striped Bass Relative Abundance in the Virginia Portion of Chesapeake Bay

ANNUAL PROGRESS REPORT: January 2013-December 2013

**Mary Fabrizio
Troy Tuckey
Christopher Davis**

Department of Fisheries Science
Virginia Institute of Marine Science
College of William and Mary
Gloucester Point, Virginia



U.S. Fish and Wildlife Service Sportfish Restoration Project F87R24
Submitted to Virginia Marine Resources Commission March 2013



TABLE OF CONTENTS

EXECUTIVE SUMMARY	<i>ii</i>
LIST OF TABLES	<i>iii</i>
LIST OF FIGURES	<i>iv</i>
PREFACE	<i>v</i>
INTRODUCTION.....	1
METHODS.....	3
RESULTS AND DISCUSSION	6
CONCLUSION.....	20
ACKNOWLEDGMENTS.....	21
LITERATURE CITED.....	22
TABLES.....	25
FIGURES.....	42

EXECUTIVE SUMMARY

The 2013 Striped Bass juvenile abundance index was 10.94 and was significantly higher than the historic average of 7.04. Abundance indices from individual rivers varied compared with their respective historic averages, and significantly higher recruitment was observed in the York River (particularly the Mattaponi River). Average recruitment occurred in the other rivers. Increased catches of young-of-the-year Striped Bass at upriver and downriver auxiliary sites suggest an expansion of Striped Bass nursery grounds in 2013. The evident increase in occurrence of young-of-the-year Striped Bass throughout the tidal tributaries in Virginia further supports the observation of an above-average recruitment year in 2013.

In contrast with Striped Bass, juvenile White Perch abundance indices in 2013 were considerably lower than the historic average. A strong year class for Atlantic Croaker appears to have occurred in 2013. Average to above-average indices for forage fishes suggest adequate trophic support for populations of commercial and recreational piscivores in Virginia waters.

LIST OF TABLES

Table 1.Catch of young-of-the-year Striped Bass per seine haul in 2013	25
Table 2.Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) summarized by year.....	26
Table 3.Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia using only the 1 st haul summarized by year.....	27
Table 4.Catch of young-of-the-year Striped Bass per seine haul in the primary nursery area in 2013 summarized by drainage and river.....	28
Table 5.Site specific Striped Bass indices and average site salinity during 2013 compared to historic (1967 – 2013) index values with corresponding average salinities.....	29
Table 6.Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia in 2013 summarized by sampling round and month.....	30
Table 7.Water temperature (°C) recorded at seine survey stations in 2013.....	31
Table 8.Catch of young-of-the-year Striped Bass per seine haul in the primary nursery areas of Virginia in 2013 summarized by water temperature.....	32
Table 9.Catch of young-of-the-year Striped Bass per seine haul in the primary nursery areas of Virginia in 2013 summarized by salinity.....	33
Table 10.Salinity (ppt) recorded at seine survey stations in 2013.....	34
Table 11.Dissolved oxygen concentrations (mg/L) at seine survey stations in 2013.....	35
Table 12.Species collected during the 2013 seine survey (index and auxiliary stations).....	36
Table 13.Preliminary catch of Spottail Shiner from select juvenile Striped Bass seine survey stations using only the 1 st haul summarized by year.....	38
Table 14.Preliminary catch of Atlantic Silverside from select juvenile Striped Bass seine survey stations using only the 1 st haul summarized by year.....	39
Table 15.Preliminary catch of Inland Silverside from select juvenile Striped Bass seine survey stations using only the 1 st haul summarized by year.....	40
Table 16.Preliminary catch of Banded Killifish from select juvenile Striped Bass seine survey stations using only the 1 st haul summarized by year.....	41

LIST OF FIGURES

Figure 1. Juvenile Striped Bass seine survey stations.....	42
Figure 2. Scaled geometric mean of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) by year.....	43
Figure 3. Scaled geometric mean of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) by drainage and river.....	44
Figure 4. Catch of young-of-the-year Striped Bass by station in the James River drainage in 2013.....	45
Figure 5. Catch of young-of-the-year Striped Bass by station in the York and Mattaponi rivers in 2013.....	46
Figure 6. Catch of young-of-the-year Striped Bass by station in the York and Pamunkey rivers in 2013.....	47
Figure 7. Catch of young-of-the-year Striped Bass by station in the Rappahannock River in 2013.....	48
Figure 8. Geometric mean of young-of-the-year Spot from select seine survey stations in Virginia tributaries of Chesapeake Bay by year.....	49
Figure 9. Delta-lognormal mean of young-of-the-year White Perch from select seine survey stations by river and year.....	50
Figure 10. Delta-lognormal mean of young-of-the-year White Perch from the James River nursery area by year.....	51
Figure 11. Delta-lognormal mean of young-of-the-year White Perch from the Pamunkey River nursery area by year.....	52
Figure 12. Delta-lognormal mean of young-of-the-year White Perch from the Mattaponi River nursery area by year	53
Figure 13. Delta-lognormal mean of young-of-the-year White Perch from the Rappahannock River nursery area by year	54
Figure 14. Delta-lognormal mean of young-of-the-year Atlantic Croaker from select seine survey stations in Virginia tributaries of Chesapeake Bay by year	55

PREFACE

The primary objective of the Virginia Institute of Marine Science juvenile Striped Bass survey is to monitor the relative annual recruitment success of juvenile Striped Bass in the major Virginia nursery areas of Chesapeake Bay. The U.S. Fish and Wildlife Service initially funded the survey from 1967 to 1973. Beginning in 1980, funds were provided by the National Marine Fisheries Service under the Emergency Striped Bass Study program. Commencing with the 1989 annual survey, the work was jointly supported by Wallop-Breaux funds (Sport Fish Restoration Act), administered through the U.S. Fish and Wildlife Service, and the Virginia Marine Resources Commission. This report summarizes the results of the 2013 sampling period and compares these results with previous years.

INTRODUCTION

Striped Bass (*Morone saxatilis*) is one of the most commercially and recreationally sought-after fish species on the east coast of the United States. Decreases in the commercial harvest of Striped Bass in the 1970s paralleled the steady decline in abundance of Striped Bass along the east coast; Chesapeake Bay stock abundances were particularly depressed. Declines in commercial harvests mirrored declines in juvenile recruitment (Goodyear 1985). Because the tributaries of Chesapeake Bay were identified as primary spawning and nursery areas, fishery managers enacted regulations intended to halt and reverse the decline of Striped Bass in Chesapeake Bay and elsewhere within its native range (ASMFC 2003).

In 1981, the Atlantic States Marine Fisheries Commission (ASMFC) developed the Atlantic Coast Striped Bass Interstate Fisheries Management Plan (FMP), which included recommendations aimed to improve the stock status. The Virginia Marine Resources Commission (VMRC) adopted the plan in March 1982 (Regulation 450-01-0034), but the ASMFC did not have regulatory authority for fisheries management in individual states at that time. As Striped Bass populations continued to decline, Congress passed the Atlantic Striped Bass Conservation Act (PL 98-613) in 1984, which required states to follow and enforce management measures in the FMP or face a moratorium on Striped Bass harvests. Since 1981 the FMP has been amended six times to address changes in the management of the stocks. Amendment VI to the plan, adopted in February 2003, requires "producing states" (i.e., Virginia, Maryland, Delaware and New York) to develop and support programs monitoring Striped Bass recruitment.

In 1967, before the FMP requirement, Virginia began monitoring the annual recruitment of juvenile Striped Bass using funding from the Commercial Fisheries Development Act of 1965 (PL88-309). This monitoring continued until 1973 when funding was discontinued. Monitoring of striped bass recruitment was re-instituted in 1980 using Emergency Striped Bass Study funds (PL 96-118, 16 U.S.C. 767g, the "Chafee Amendment"), and since 1989, the annual survey has been funded by the Wallop-Breaux expansion of the Sport Fish Restoration and Enhancement Act of 1988 (PL 100-448, "the Dingle-Johnson Act"). Funds are administered through the VMRC.

Initially, the Virginia program used a 6 ft. x 100 ft. x 0.25 in. mesh (2 m x 30.5 m x 6.4 mm) bag seine, but comparison hauls with Maryland gear (4 ft. x 100 ft. x 0.25 in. mesh; 1.2m x 30.5m x 6.4mm mesh) showed virtually no statistical differences in catch, and Virginia adopted the "Maryland seine" after 1987 (Colvocoresses 1987). The gear comparison study aimed to standardize methods and promote a bay-wide recruitment estimate (Colvocoresses and Austin 1987). This was never realized due to remaining differences in the methods of estimation of means (MD: arithmetic index; VA: geometric index). A bay-wide index using a geometric mean weighted by spawning area in each river was proposed in 1993 (Austin et al. 1993) but has not been implemented. Recent computations of a bay-wide geometric mean juvenile abundance index (JAI) were found to be correlated with fishery-independent monitoring (Woodward 2009).

Objectives for the 2013 program were to:

1. estimate the relative abundance of the 2013 year class of Striped Bass in the James, York and Rappahannock river systems,

2. quantify environmental conditions at the time of collection, and
3. examine relationships between juvenile Striped Bass abundance and environmental and biological data.

METHODS

Field sampling was conducted during five biweekly periods (rounds) from 8 July through 13 September 2013. During each round, seine hauls were conducted at 18 index stations and 21 auxiliary stations in the James, York and Rappahannock river systems (Figure 1). Auxiliary sites were added to the survey in 1989 to provide better geographic coverage and increase sample sizes within each river system. Such monitoring was desirable in light of increases in stock size during the 1980s and hypothesized expansion of the nursery ground in years of high recruitment.

Collections were made by deploying a 100 ft. (30.5 m) long, 4 ft. (1.2 m) deep, 0.25 in (6.4 mm) mesh minnow seine perpendicular to the shoreline until either the net was fully extended or a depth of approximately 4 ft. (1.2 m) was encountered and then pulling the offshore end down-current and back to the shore. During each round a single haul was made at each auxiliary station and duplicate hauls, with a 30-minute interlude, were made at each index station. Every fish collected during a haul was removed from the net and placed into water-filled buckets. All Striped Bass were measured to the nearest mm fork length (FL), and for all other species, a sub-sample of up to 25 individuals was measured to the nearest mm FL (or total length if appropriate). At index stations, fish collected during the first haul were held in a water-filled bucket until the second haul was completed. All captured fish, except those preserved for life

history studies, were returned to the water at the conclusion of sampling. Sampling time, tidal stage, and weather conditions were recorded at each sampling location. Salinity, water temperature and dissolved oxygen concentrations were measured after the first haul using a YSI water quality sampler.

In this report, comparisons of Striped Bass recruitment indices with prior years are made for the “primary nursery” area only (Colvocoresses 1984), using data collected from months and areas sampled during all years (i.e., index stations). Catch data from auxiliary stations are not included in the calculation of the annual indices. The index of relative abundance for young-of-the-year Striped Bass is calculated as the adjusted overall mean catch per seine haul such that

$$\text{Index} = (\exp(\ln(\text{totnum} + 1)) - 1) \times 2.28$$

where *totnum* is the total number of Striped Bass per seine haul; includes catches from the first and second seine haul at each index station are included in this calculation. Because the frequency distribution of the catch is skewed (Colvocoresses 1984), a logarithmic transformation ($\ln(\text{totnum}+1)$) was applied to the data prior to analysis (Sokal and Rohlf 1981). Mean values are back-transformed and scaled up arithmetically ($\times 2.28$) to allow comparisons with Maryland data. Thus, a “scaled” index refers to an index that is directly comparable with the Maryland index.

In accordance with suggestions made by Rago et al. (1995), the Virginia juvenile Striped Bass index has also been recomputed using only the first haul at each index station. Additionally, the rehabilitation of Chesapeake Bay Striped Bass stocks, and subsequent relaxation of commercial and recreational fisheries regulations in

Chesapeake Bay in 1990 (ASMFC 2003) allows examination of the recruitment of Striped Bass during three distinct periods:

- 1967 – 1973: an early period of monitoring;
- 1980 – 1989: a decade reflecting severe population depression during which temporary fishing moratoria were in place; and,
- 1990 – Present: a period of post-recovery and regulation targeting the development of a sustainable fishery.

An average index value for 1990 – 2013 was calculated using only the first haul at each index site and was compared with the annual index value to provide a benchmark for interpreting recruitment strength during the post-recovery period.

In previous years, the historic average was calculated as the geometric mean across all stations. However, survey effort has not been equal through time. The number of hauls completed annually ranged from 42 (1967) to 180 (post-1988) resulting in an estimate of the historic average that was biased by recruitment years with higher effort (i.e., more recent years). The historic average should instead be calculated as the mean of the annual abundance estimates ($n = 41$, one estimate per year). Equal weight is thus given to years of lower sampling effort (which in this time series happen to be years of low abundance), and years with higher sampling effort (which occurred during the latter part of the time series and represent years of higher abundance).

Throughout this report, mean catch rates are compared using 95% confidence intervals. Reference to “significant” differences between geometric means in this context will be restricted to cases of non-overlapping confidence intervals. Because

standard errors are calculated from transformed (logarithmic) values, confidence intervals for the back-transformed and scaled indices are non-symmetrical.

RESULTS AND DISCUSSION

Juvenile Index of Abundance for Virginia

In 2013, 1,620 young-of-the-year Striped Bass were collected from 180 seine hauls at index stations and 705 individuals were collected from 105 hauls at auxiliary stations (Table 1). Using index station data from both hauls, the estimated Striped Bass recruitment index in 2013 was 10.94 (LCI = 8.97, UCI = 13.25; Table 2), which was significantly higher than the historic average of 7.04 (LCI = 5.56, UCI = 8.81; Figure 2).

Even with a 30-minute interlude between hauls at index stations, second hauls are not independent samples and their use violates a key assumption necessary for making inferences from a sample mean (Rago et al. 1995). Previous reports consistently documented fewer catches in the second haul (e.g. Hewitt et al. 2007, 2008), a result which artificially lowers the geometric mean when data from both hauls are included in the index computation. Thus, the annual and historic indices were recalculated using only the first haul at each index station. In 2013, 877 young-of-the-year Striped Bass were collected in the first haul resulting in an index of 11.85 (LCI = 8.92, UCI = 15.54, Table 3), which is similar to the recomputed first-haul historic index of 8.36 (LCI = 6.57, UCI = 10.52). All annual Striped Bass abundance estimates presented in Table 3 are based on single hauls. The 2013 Virginia-wide index of 11.85 (LCI = 8.92, UCI = 15.54) is similar to the mean index estimated for the post-recovery period (index = 11.9; LCI = 9.35, UCI = 15.01).

Prior to 2011, annual recruitment indices were derived from all collections made during a sampling year. From 1967 to 1973, seine sampling extended into October and occasionally into December (1973). Current sampling concludes in mid-September because after this time, sampling efficiency decreases due to mortality of juveniles, increased avoidance of the sampling gear, and the dispersal of juveniles into deeper waters. Indices calculated from data that include catches after mid-September are therefore biased low. Starting in 2011, recruitment calculations were made using catch data from the currently established sampling season (July through mid-September) to permit uniform comparisons of annual recruitment (Tables 2– 4).

Striped Bass recruitment success in the Virginia portion of Chesapeake Bay is variable among years and among nursery areas within years. Average to above-average recruitment years occurred between 2003 and 2011, and in 2013 (Figure 2). The highest recruitment index observed by the Virginia seine survey occurred in 2011. Since the termination of the Striped Bass fishing moratorium in 1990, strong year classes have been observed approximately every decade (1993, 2003, and 2011). Similarly, single years of low recruitment have occurred approximately every ten years (1991, 1999, and 2002). Poor recruitment in 2012 is therefore consistent with past observations. Individual years of poor recruitment after the moratorium (i.e., since 1990) were followed by years of average or above-average recruitment and that pattern was repeated in 2013 (Figure 2). Under current ASMFC regulations (ASMFC 2010), management action is triggered after three consecutive years of below-average recruitment in producing states.

Continued monitoring of regional recruitment success will be important in identifying management strategies to protect the spawning stock of Chesapeake Bay Striped Bass. Research suggests that a Chesapeake Bay-wide index, computed from Virginia and Maryland data combined, will provide a better estimate of recruitment strength and serve as a better predictor of subsequent adult Striped Bass abundance within the Bay (Woodward 2009; Fabrizio et al. *in review*). This may be particularly appropriate in years when individual state JAIs provide divergent estimates of year-class strength; such differences may arise due to annual changes in the spatial distribution and contribution of nursery areas throughout Chesapeake Bay.

Juvenile Index of Abundance for Individual Watersheds

Recruitment indices observed in the three Virginia watersheds during 2013 were considered average (James and Rappahannock rivers) or above-average (York River) compared with their respective historic averages. The 2013 JAI for the James River drainage was 14.30 (LCI = 10.35, UCI = 19.48), compared with the historic James River index of 8.96 (LCI = 6.82, UCI = 11.61; Table 4). The 2013 JAI value for the York River drainage was 11.23 (LCI = 8.48, UCI = 14.69), compared with the historic York River index of 5.38 (LCI = 4.16, UCI = 6.83). The 2013 JAI value for the Rappahannock River was 7.49 (LCI = 4.63, UCI = 11.54), compared with the historic Rappahannock River index of 7.14 (LCI = 5.36, UCI = 9.32).

Throughout the James River watershed, four of six index sites were characterized by above-average relative abundances compared with their respective historic means (Table 5). Catches at mid-river index stations dominated the catches in the James River

watershed (Machut and Fabrizio 2011, 2012), with 47% of all young-of-the-year Striped Bass occurring in the core nursery zone (C1, C3, and J46). The remaining striped bass occurred about equally at downriver (29%) and upriver (25%) sites.

The 2013 York River JAI (11.23, LCI = 8.48, UCI = 14.69) was significantly greater than the historic average of 5.38 (LCI = 4.16, UCI = 6.83) for this river (Table 4; Figure 3). No index sites are located along the main stem of the York River although three auxiliary stations are sampled; the watershed JAI is compiled from sites located within the two principle York River tributaries, the Mattaponi and Pamunkey rivers. The 2013 Pamunkey River JAI of 11.31 (LCI = 6.76, UCI = 18.16) was not significantly different than the historic index of 6.04 (LCI = 4.33, UCI = 8.19), but the 2013 Mattaponi River index (11.17; LCI = 8.08, UCI = 15.19) was significantly higher than the historic average (4.92; LCI = 3.91, UCI = 6.09). About 49% of York River Striped Bass were collected from the Pamunkey River and 51% from the Mattaponi River (Table 4).

The 2013 Rappahannock River index of 7.49 (LCI = 4.63, UCI = 11.54) was similar to the historic average of 7.14 (LCI = 5.36, UCI = 9.32; Table 4). In 2013, 85% of the total Rappahannock River catch was taken from the two uppermost index sites (R50 and R55; Table 1); these two sites have dominated the catches in this drainage for several years.

Weak year classes in 1991, 1999, 2002, and 2012 were all followed by above-average recruitment in the following year (Figure 2). Large changes in young-of-the-year abundance estimates are typical of estuarine-dependent species. Catches of young-of-the-year Striped Bass from individual watersheds in Virginia during 2013 indicate an average or above-average year class was produced throughout Virginia

waters. After the moratorium was lifted in 1990, individual years of poor recruitment have occurred sporadically and were usually driven by poor recruitment within a single watershed except for 2012, when recruitment was poor in all three watersheds. However, because the 2012 year class followed the strong 2011 year class, regional production of juvenile Striped Bass in Chesapeake Bay and along the Atlantic coast was not of concern. In 2013, recruitment of striped bass returned to above average conditions.

Striped Bass Collections from Auxiliary Stations

Figures 4 – 7 illustrate the spatial distribution of the 2013 year class throughout the nursery area. Note that scaling is not constant across figures. The 1989 addition of auxiliary stations has provided increased spatial coverage for the James, York and Rappahannock drainages and the upriver and downriver auxiliary sites allow delineation of the upper and lower limits of the nursery. These auxiliary stations reveal spatial changes in the nursery areas that occur due to annual changes in river flow. Additionally, in years of low or high juvenile abundance, the nursery area may contract or expand spatially. Inter-annual changes in the occurrence of young-of-the-year Striped Bass at auxiliary sites was particularly evident in 2013: we observed increased catches at upriver and downriver auxiliary stations which we interpret as expansion in the spatial extent of Striped Bass nursery grounds.

Juvenile Striped Bass were captured at all auxiliary stations in the James River during 2013, though catches were lowest at upper- and lower-most stations (Table 1; Figure 4). The greatest number of Striped Bass collected from stations in the James

River was from an upriver auxiliary station (J62, n = 107 Striped Bass; Tables 1 and 5). The nursery grounds also extended downstream in 2013 (to J22 and J12; Table 1).

Striped Bass were collected from all auxiliary sites in the Pamunkey and Mattaponi rivers in 2013 (Tables 1 and 5). Additionally, Striped Bass were collected from all three auxiliary stations within the York River watershed (Table 1). The greatest number of Striped Bass captured in the York River system in 2013 was at Y28, a downriver auxiliary station (Table 5).

We previously suggested that the lack of Striped Bass at upriver auxiliary stations in the York River watershed may have been due to the inability to accurately sample in the dense hydrilla vegetation that occurs at these sites (Machut and Fabrizio 2010). In 2013, we detected juvenile Striped Bass at the York River auxiliary sites (Table 1), but not all fish may have been detected due to low capture efficiencies associated with hauling a seine net through dense aquatic vegetation. Furthermore, recent catches at P55, and to a lesser extent at M52, may have been affected by the altered state of the near-shore zone at these sites. Striped Bass may have been forced into deeper waters by the dense hydrilla beds; alternatively, Striped Bass may preferentially use hydrilla habitats but remain unavailable to the sampling gear. The continued sampling difficulties at these stations suggest a need to examine alternative collection methods within this region to determine the abundance of juvenile Striped Bass in near-shore areas where hydrilla is present.

Striped Bass were collected at all Rappahannock River auxiliary stations during 2013. In recent years, few fish were collected at the lower auxiliary stations in the

Rappahannock River (R12, R21) even though these sites have favorable substrate and no obstructions to compromise seining. The consistent low capture rates at R12 and R21 suggest these sites may have lower value as nursery areas in the Rappahannock River. The same is not true for upstream auxiliary locations. Historic JAI values at auxiliary stations upstream of Tappahannock (near R37) appear comparable to JAIs at index stations R28, R37, and R44 (Table 5).

The evident increase in occurrence of young-of-the-year Striped Bass at downstream and upstream auxiliary sites throughout tidal tributaries in Virginia further supports the observation of an above-average recruitment year in 2013. Direct comparisons between auxiliary and index sites are problematic due to slightly different sampling protocols (index station catches are reported as an average of two hauls, whereas only a single haul is made at auxiliary stations). Past analyses demonstrate that catches are consistently greater in the first of two hauls at a given site. Because only one haul is completed at auxiliary sites, abundance estimates at these sites may be biased high relative to abundances estimated at index sites.

Comparison among Sampling Rounds

Fewer young-of-the-year Striped Bass were collected during each subsequent round in 2013, a pattern that has been observed in previous years (Table 6). Historically, 21% fewer Striped Bass are observed in the second round relative to the first round. In 2013, the reduction was only 15% and the decrease in observed numbers of Striped Bass between rounds 2 and 3 was 22% compared with the historic average of 26% (Table 6).

Environmental Conditions and Potential Relationships to Juvenile Striped Bass

Abundance

In Virginia tidal tributaries, water temperatures follow a well-defined pattern of high temperatures in July followed by declining temperatures towards the end of the sampling season in mid September. In 2013, this pattern was slightly altered: water temperatures peaked during July (rounds 1 and 2), and remained above 25°C throughout August (except for two sites; rounds 3 and 4) and into early September (round 5; Table 7). A more traditional temperature pattern occurred in 2009 and 2010 (Machut and Fabrizio 2010, 2011) when temperatures in September were below 25°C. In September 2013 (round 5), no sites exhibited water temperatures below 25°C compared with about 50% of sites in 2010. This unusual temperature pattern did not seem to affect catches, however. Catch rates in 2013 followed the historic pattern with respect to water temperature: most fish (96%) were captured at temperatures between 25.0 and 34.9°C (Table 8). Water temperatures in tidal tributaries reflect not only long-term, regional climate patterns, but also significant day-to-day and local variation. Shallow shoreline areas are easily affected by local events such as thunderstorms and small-scale spatial and temporal variations associated with time of sampling (e.g., morning versus afternoon, riparian shading, tidal stage). As noted in previous reports, the relationship between declining catches and decreasing temperature is considered to be largely the result of a coincident downward decline in catch rates and water temperatures as the season progresses (beyond early August) rather than any direct effect of water temperature on juvenile fish distribution.

In 2013, average salinity at all stations was lower than that observed historically except for J68 and Y21 (Table 5). As observed in the past, greater catches of young-of-the-year Striped Bass in 2013 were obtained at sites exhibiting low salinities within the primary nursery area (Table 9). No index station exceeded 11.0 ppt salinity, although salinity as high as 22.1 ppt was observed at one auxiliary site in the York River (Table 10). In 2013, the percentage of catch observed in low salinities (0.0 – 4.9 ppt) was similar to that observed historically (97.7% in 2013 vs. 93% all years; Table 9). Conversely, the catch in mid-range salinities (5.0 – 9.9 ppt) was below the historic average (1% in 2013 vs. 6%, all years). Although juvenile Striped Bass were captured at downstream sites with average salinities up to 16 ppt, catches were distinctly lower at such sites than the catches observed in lower salinity areas.

Dissolved oxygen (DO) concentrations were lower in 2013 at many sites compared with the historic average (Table 11). Within the nursery area, approximately half of the measurements (93 of 190 measurements) exhibited DO levels that were more than one standard error (SE) less than the site's historic average (Table 11). Twelve DO levels measured during 2013 (all occurring in the York River system) were particularly depressed. Correlations between DO and juvenile Striped Bass catches are difficult to ascertain. Lower-than-average values occurred inconsistently through time and across sampling sites. DO values more than one SE less than the mean at a given station (shaded values in Table 11) do not necessarily correspond with low catches at that station (Table 1). For example, juvenile Striped Bass were captured at M33 during periods approaching hypoxia and in normoxic conditions. Farther upriver, high seasonal

catches at index stations occurred during periods when DO was more than one SE below the historic average and when DO measures were within one SE of the historic average. Similarly, lower than average DO conditions were prevalent throughout the Rappahannock River in 2013 and catches varied greatly between sites.

Striped Bass recruitment variability may be partially explained by regional climate patterns during winter and spring (Wood 2000). For example, abundance of young Striped Bass has been positively associated with high river flows during the preceding winter (Wingate and Secor 2008). One of the strongest Striped Bass year classes was produced in 2011, which was characterized by above normal precipitation in winter and spring (Machut and Fabrizio 2012). Precipitation in the winter of 2013 (December 2012 - February 2013) was “near normal” in Maryland and “above normal” in Virginia (NCDC 2014). Spring precipitation (March – May 2013) was “below normal” in Maryland but remained “above normal” in Virginia. These regional precipitation conditions resulted in our observation that salinities were below the historic averages at most Virginia seine sites (Table 5). It is still unclear if fine-scale climate patterns or other factors exert effects on variations in recruitment of juvenile Striped Bass.

Additional Abundance Indices Calculated from the Seine Survey

Due to a sampling regime that spans from the euryhaline to the freshwater zone, a variety of species are collected by the juvenile Striped Bass seine survey annually. The five most common species encountered in 2013 were Atlantic Silverside (*Menidia menidia*), Spottail Shiner (*Notropis hudsonius*), White Perch (*Morone americana*), Hogchoker (*Trinectes maculatus*), and Bay Anchovy (*Anchoa mitchilli*; Table 12).

Juvenile Striped Bass were the 7th most common species. In 2013, more than 36,000 individuals comprising 62 species were collected (Table 12).

To fulfill commonwealth compliance requirements to the ASMFC, several annual indices are reported for species of management importance that are presently derived from seine survey collections. These species include American Shad (Watkins et al. 2011) and Atlantic Menhaden (VMRC 2010). Abundance estimates for juvenile American Shad from the seine survey were highly correlated with those from push-net sampling (Wilhite et al. 2003), providing validation for this seine survey-based index. Additionally, requests for an index of abundance or raw catch data occur throughout the year on an *ad hoc* basis.

One of the most common species annually captured by the seine survey, White Perch, supports important recreational and commercial fisheries in Chesapeake Bay (Murdy et al. 1997, NMFS 2012). The general overlap in spawning time and use of nursery grounds by White Perch and Striped Bass suggest that the seine survey may adequately sample juvenile White Perch and that calculation of a recruitment index for this species is appropriate. Colvocoresses (1988) found a strong correlation between a young-of-the-year geometric mean calculated from seine survey sampling and a trawl survey estimate of White Perch fishable stock. However, examination of raw catch data suggests high annual variability in White Perch catches. In years of low abundance (e.g., 1985) the proportion of hauls containing White Perch may be as low as 40%; whereas in years of high abundance (e.g., 2011), White Perch may be found in 95% of hauls. A delta-lognormal index was developed to address this inter-annual variation and to

accommodate data with a high proportion of zero hauls. We used Cox's method (Fletcher 2008) to estimate the mean abundance based on the delta-lognormal distribution, and calculated 95% confidence intervals from 1,000 bootstrap samples as described by Fletcher (2008). This approach remains under development, so we report only the means here.

From July through September 2013, 4,762 young-of-the-year White Perch were collected from 30 sampling stations during five sampling rounds. Because White Perch movement among Virginia tributaries is unlikely (Mulligan and Chapman 1989), we presume each tributary supports a distinct stock and report juvenile abundance for each river separately (Figures 9 - 13). Generally, 2013 river-specific JAIs for White Perch suggest below average recruitment for this species throughout Virginia. Numerically, the highest 2013 JAI values for White Perch were observed in the James, Rappahannock, and Mattaponi rivers. Near complete recruitment failure occurred in the Pamunkey River in 2013, when the index was 17% of the historic average for this river. At present, development of river-specific indices for White Perch is incomplete. Although we feel confident in the estimation of annual mean relative abundance, alternative approaches for estimating confidence intervals should be examined. The White Perch JAI developed by the seine survey compliments the juvenile White Perch index currently reported by the Juvenile Fish Trawl Survey (Tuckey and Fabrizio 2012); however, unlike the index reported by the trawl survey, the seine survey index is based on catches from tidal brackish and freshwater zones.

Atlantic Croaker (*Micropogonias undulatus*) is another commercially and recreationally important fish (Murdy et al. 1997, NMFS 2012) regularly collected by the seine survey. Young-of-the-year Atlantic Croaker are collected at predominately mesohaline sampling sites during July and early August (rounds 1, 2 and 3) before fish are able to avoid the net and reduce their detection probability (Williams and Fabrizio 2011). Murdy et al. (1997) report peak spawning of Atlantic Croaker from August – October; thus, young-of-the-year fish collected during 2013 were spawned during the fall of 2012. Similar to White Perch, Atlantic Croaker raw catch data exhibit high annual variability in the proportion of non-zero hauls. To address this variation and accommodate data with a high proportion of zero hauls we developed a delta-lognormal index for Atlantic Croaker. The delta-lognormal approach remains under development, so only means are currently reported. Alternative approaches for estimating confidence intervals should be examined.

From July through early August 2013, 974 young-of-the-year Atlantic Croaker were collected from 36 sampling locations. Because Atlantic Croaker are coastal shelf spawners with larval migration into Chesapeake Bay, we report a Virginia-wide estimate of juvenile abundance (Figure 14). A strong year class for Atlantic Croaker appears to have occurred during 2013. Periods of strong recruitment from 1991 – 1995, 1997 – 1998, and 2008 – 2009 correspond with patterns observed by the VIMS Juvenile Fish Trawl Survey (Tuckey and Fabrizio 2012).

Spot (*Leiostomus xanthurus*), like Atlantic Croaker, is another commercially and recreationally important fish that is collected by the seine survey and reported as a

Virginia-wide estimate of juvenile abundance (Figure 8). From July through early August 2013, 695 young-of-the-year Spot were collected from 44 sampling locations. An average year class for Spot was observed in 2013. The spatial proximity of sites sampled by the seine and trawl surveys provides a more complete picture of fish distribution in estuarine areas. Near-shore shallow zones are sampled by the seine survey whereas the trawl survey samples adjacent benthic main stem sections. Indices of abundance for common forage species within the tidal near-shore zone have been computed for: Spottail Shiner (*Notropis hudsonius*; Table 13), Atlantic Silverside (*Menidia menidia*; Table 14), Inland Silverside (*Menidia beryllina*; Table 15), and Banded Killifish (*Fundulus diaphanus*; Table 16). The 2013 Spottail Shiner geometric mean of 14.98 (LCI = 11.49, UCI = 19.44) was significantly greater than the historic average (Index = 9.07, LCI = 7.43, UCI = 11.04). The 2013 Atlantic Silverside index of 11.17 (LCI = 6.12, UCI = 19.79) was similar to the historic average of 10.17 (LCI = 8.10, UCI 12.71; Table 14). The 2013 Inland Silverside abundance index of 5.15 (LCI = 3.79, UCI = 6.89) was significantly higher than the historic geometric mean of 1.79 (LCI = 1.41, UCI = 2.23). However, the 2013 Banded Killifish abundance index of 2.45 (LCI = 1.74, UCI = 3.36) was not significantly different from the historic average of 2.02 (LCI = 1.46, UCI = 2.72). Average to above-average indices for these four species in 2013 suggest that a robust population of forage fishes was available for commercially and recreationally important piscivores in Virginia waters.

CONCLUSION

The 2013 juvenile abundance index (JAI) for Striped Bass (10.94) was significantly higher than the historic average (7.04) for Virginia waters. Compared with historic averages, we observed significantly higher recruitment in the York River (particularly in the Mattaponi River), but average recruitment in the other rivers. Continued evaluation of juvenile Striped Bass recruitment is important in predicting recruitment to the commercial and recreational Striped Bass fisheries in the Chesapeake Bay and along the Atlantic coast. A critical characteristic of the long-term annual seine survey conducted in the Chesapeake Bay is the ability to identify years of poor recruitment which, if persistent, serve as an early warning to managers of potential declines in Striped Bass standing stock biomass. In contrast with Striped Bass, the juvenile White Perch abundance indices in 2013 were considerably lower than the historic average for this species. Forage fish abundance index values were average or significantly above average in 2013.

ACKNOWLEDGMENTS

We are indebted to the many landowners who have graciously allowed us access to their waterfront properties. We thank the Mariners' Museum, Jamestown 4-H Camp, June Parker Marina, Chickahominy Riverfront Park, and the United States Army at Fort Eustis for their permission to sample. Additional thanks go to Jordan Point Marina, June Parker Marina and Chickahominy Riverfront Park for permission to use their boat ramps. Summer technicians were Bruce Pfirrmann and Anya Voloshin. We also thank VIMS

students/staff who assisted in field sampling, including Wendy Lowery, Hank Brooks, Aimee Comer, Jennifer Greaney, Lauren Nys, Ryan Schloesser, and Beth Dzula. Funding was provided by a grant from the United States Fish and Wildlife Service Sport Fish Restoration Project (F-87-R-25) through the Virginia Marine Resources Commission to the Virginia Institute of Marine Science.

LITERATURE CITED

- ASMFC (Atlantic States Marine Fisheries Commission). 2003. Amendment #6 to the Interstate Fishery Management Plan for Atlantic Striped Bass. Fisheries Management Report 41. Atlantic States Marine Fisheries Commission, Washington, D.C. 63 p.
- ASMFC (Atlantic States Marine Fisheries Commission). 2010. Addendum II to Amendment #6 to the Interstate Fishery Management Plan: definition of recruitment failure. Atlantic States Marine Fisheries Commission, Washington, D.C. 18 p.
- Austin, H.M., J.A. Colvocoresses and T.A. Mosca III. 1993. Develop a Chesapeake Bay-wide young-of-the-year Striped Bass index. Final Report, CBSAC Cooperative Agreement NA16FUO393-01, 59 p. + 2 app.
- Colvocoresses, J. A. 1984. Striped Bass research, Virginia. Part I: Juvenile Striped Bass seining program. Annual Report 1987-88. Virginia Institute of Marine Science, Gloucester Point, Virginia. 64 p.
- Colvocoresses, J. A. 1987. Intercalibration and refinement of estimates of abundance of Chesapeake Bay juvenile Striped Bass. NOAA Tech. Rept. TRS-SAC-91-010, 28 p.
- Colvocoresses, J. A. 1988. Comparisons among York River White Perch stock abundance measures. NOAA Tech. Rept. TRS-SAC-91-021, 18 p.
- Colvocoresses, J.A. and H.M. Austin. 1987. Development of an index of juvenile Striped Bass abundance for the Chesapeake Bay System: I. An evaluation of present measures and recommendations for future studies. Special Science Report 120. Virginia Institute of Marine Science, Gloucester Point, VA. 108 p.
- Fabrizio, M. C., R. J. Latour, J. R. Woodward, C. F. Bonzek. In Prep. Evaluating relationships among indices from fishery-independent surveys.
- Fletcher, D. 2008. Confidence intervals for the mean of the delta-lognormal distribution. Environmental and Ecological Statistics 15: 175 – 189.
- Goodyear, C. P. 1985. Relationship between reported commercial landings and abundance of young Striped Bass in Chesapeake Bay, Maryland. Transactions of the American Fisheries Society 114: 92 – 96.
- Hewitt, A. H., J.K. Ellis and M.C. Fabrizio. 2007. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2006. Virginia Institute of Marine Science, Gloucester Point, VA. 31 p.

- Hewitt, A. H., L. S. Machut and M.C. Fabrizio. 2008. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2007. Virginia Institute of Marine Science, Gloucester Point, VA. 28 p.
- Machut, L. S., and M.C. Fabrizio. 2010. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2009. Virginia Institute of Marine Science, Gloucester Point, VA. 45 p.
- Machut, L. S., and M.C. Fabrizio. 2011. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2010. Virginia Institute of Marine Science, Gloucester Point, VA. 47 p.
- Machut, L. S., and M.C. Fabrizio. 2012. Estimation of juvenile Striped Bass relative abundance in the Virginia portion of Chesapeake Bay. Annual Report 2011. Virginia Institute of Marine Science, Gloucester Point, VA. 55 p.
- Mulligan, T. J., and R. W. Chapman. 1989. Mitochondrial DNA analysis of Chesapeake Bay White Perch, *Morone americana*. *Copeia* 3: 679 – 688.
- Murdy, E. O., R. S. Birdsong, and J.A. Musick. 1997. Fishes of Chesapeake Bay. Smithsonian Institution Press, Washington, D. C. 324 p.
- NCDC (National Climate Data Center). 2013. <http://www.ncdc.noaa.gov/oa/ncdc.html>. Site accessed February 2014.
- NMFS (National Marine Fisheries Service). 2012. <http://www.st.nmfs.noaa.gov/st1/commercial/landings/annuallandings.html>. Site accessed March 2012.
- Rago, P., D. Stephan, and H. Austin. 1995. ASMFC Special Report 48. Report of the juvenile indices abundance workshop, January 1992, Kent Island, MD. 83 p.
- Sokal, R.R. and F.J. Rohlf. 1981. Biometry. W.H. Freeman and Co., San Francisco, CA. 851 p.
- Tuckey, T.D., and M.C. Fabrizio. 2012. Estimating relative juvenile abundance of ecologically important finfish in the Virginia portion of Chesapeake Bay. Final Report to the Virginia Marine Resources Commission.
- Watkins, B.J. Olney, and R. O'Reilly. 2011. A summary of Virginia's American Shad fisheries in 2010 and results of monitoring and restoration programs: annual compliance report to the Atlantic States Marine Fisheries Commission American Shad Technical Committee, Virginia Institute of Marine Sciences, Gloucester Point, VA. 43 pp.
- Wilhite, M.L., K. L. Maki, J. M. Hoenig, and J. E. Olney. 2003. Towards validation of a juvenile index of abundance for American Shad in the York River, Virginia. Pages 285 - 294 in K. E. Limburg and J. A. Waldman (eds.) Biodiversity Status and

- Conservation of the World's Shads. American Fisheries Society Symposium 35, Bethesda, MD.
- Williams, B. D. and M. C. Fabrizio. 2011. Detectability of estuarine fishes in a beach seine survey of tidal tributaries of lower Chesapeake Bay. *Transactions of the American Fisheries Society* 140: 1340-1350.
- Wingate, R. L., and D. H. Secor. 2008. Effects of winter temperature and flow on a summer-fall nursery fish assemblage in the Chesapeake Bay, Maryland. *Transactions of the American Fisheries Society* 137: 1147 – 1156.
- Wood, R.J. 2000. Synoptic scale climatic forcing of multispecies fish recruitment patterns in Chesapeake Bay. Ph.D. Dissertation. College of William and Mary, Williamsburg, VA.
- Woodward, J.R. 2009. Investigating the relationships between recruitment indices and estimates of adult abundance for Striped Bass, Weakfish, and Atlantic Croaker. Master's thesis. College of William and Mary, Williamsburg, VA.
- VMRC (Virginia Marine Resources Commission). 2010. Atlantic Menhaden compliance report for Virginia: Report to the Atlantic States Marine Fisheries Commission. Fisheries Management Division, Newport News, VA. 16 pp.

Table 1. Catch of young-of-the-year Striped Bass per seine haul in 2013. Two hauls were completed at each index station (bold). Sampling was completed in June (round 1), July (round 2), August (rounds 3 and 4), and September (round 5).

Drainage															Round			
JAMES	Station	J12	J22	J29	J36	J42	C1	C3	J46	J51	J56	J62	J68	J78	Total			
	Round	1	1	4	16/12	29/24	13	54/42	4/5	0/1	30	13/5	38	2	2	295		
		2	0	23	2/1	6/9	0	16/39	12/8	2/4	17	15/15	23	0	1	193		
		3	0	10	0/3	11/6	5	62/50	4/3	17/24	9	1/1	14	0	0	220		
		4	0	2	14/18	1/1	30	6/17	3/2	4/10	7	3/0	14	0	0	132		
		5	0	4	9/4	1/3	14	13/17	2/1	6/20	4	2/4	18	0	1	123		
															James Total	963		
YORK	Station	Y15	Y21	Y28	P36	P42	P45	P50	P55							Total		
	Round	1	2	12	50	70	14/5	22/11	33/31	1						251		
		2	2	2	44	6	4/3	7/0	23/5	11						107		
		3	0	1	29	13	0/0	9/3	22/1	4						82		
		4	0	2	7	12	7/5	12/8	6/0	3						62		
		5	0	0	1	15	3/7	6/17	0/0	0						49		
	Station					M33	M37	M41	M44	M47	M52						Total	
	Round	1					11/2	8	7/5	7/7	7/6	2						62
		2					9/10	0	31/5	5/11	4/3	2						80
		3					23/3	3	8/6	8/3	3/6	1						64
		4					0/3	2	21/6	8/3	15/10	1						69
		5					0/0	14	7/2	1/0	7/3	0						34
															York Total	860		
RAPPAHANNOCK	Station	R12	R21	R28	R37	R41	R44	R50	R55	R60	R65	R69	R76			Total		
	Round	1	4	0	6/6	0/0	20	0/1	12/6	18/47	10	4	5	5	144			
		2	0	2	7/0	0/0	5	0/6	77/40	5/15	0	7	0	1	165			
		3	0	0	1/3	0/0	3	2/1	4/9	2/14	7	8	5	0	59			
		4	0	0	3/9	0/0	3	7/6	8/7	19/19	3	0	0	0	84			
		5	0	0	2/0	0/0	6	0/1	4/5	22/9	0	0	1	0	50			
															Rappahannock Total	502		
															2013 Catch	2,325		

Table 2. Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) summarized by year, where x = total fish, $\text{Index} = (\exp(\ln(x + 1)) - 1) \times 2.28$, SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean ln (x+1)	SD	Index	C.I. (± 2 SE)	N (hauls)
1967	191	1.18	1.00	5.17	3.20-7.86	42
1968	184	1.04	0.92	4.15	2.68-6.06	50
1969	193	0.97	0.94	3.73	2.39-5.46	55
1970	345	1.39	1.11	6.88	4.52-10.06	56
1971	165	0.90	0.90	3.34	2.17-4.81	60
1972	84	0.45	0.59	1.28	0.87-1.75	90
1973	133	0.60	0.82	1.86	1.12-2.76	70
1980	228	0.74	0.90	2.52	1.68-3.53	89
1981	165	0.52	0.69	1.56	1.10-2.09	116
1982	323	0.78	0.97	2.71	1.85-3.74	106
1983	296	0.91	0.83	3.40	2.53-4.42	102
1984	597	1.09	1.06	4.47	3.22-6.02	106
1985	322	0.72	0.86	2.41	1.78-3.14	142
1986	669	1.12	1.04	4.74	3.62-6.06	144
1987	2191	2.07	1.23	15.74	12.40-19.80	144
1988	1348	1.47	1.13	7.64	6.10-9.45	180
1989	1978	1.78	1.12	11.23	9.15-13.70	180
1990	1249	1.44	1.10	7.34	5.89-9.05	180
1991	667	0.97	0.95	3.76	2.96-4.68	180
1992	1769	1.44	1.24	7.35	5.72-9.31	180
1993	2323	2.19	0.98	18.11	15.35-21.30	180
1994	1510	1.72	1.03	10.48	8.66-12.60	180
1995	926	1.22	1.05	5.45	4.33-6.75	180
1996	3759	2.41	1.23	23.00	18.80-28.10	180
1997	1484	1.63	1.10	9.35	7.59-11.40	180
1998	2084	1.92	1.14	13.25	10.80-16.10	180
1999	442	0.80	0.86	2.80	2.19-3.50	180
2000	2741	2.09	1.24	16.18	13.06-19.92	180
2001	2624	1.98	1.27	14.17	11.33-17.60	180
2002	813	1.01	1.09	3.98	3.05-5.08	180
2003	3406	2.40	1.18	22.89	18.84-27.71	180
2004	1928	1.88	1.04	12.70	10.54-15.22	180
2005	1352	1.61	1.05	9.09	7.45-11.02	180
2006	1408	1.69	1.04	10.10	8.31-12.18	180
2007	1999	1.83	1.18	11.96	9.66-14.70	180
2008	1518	1.50	1.17	7.97	6.33-9.93	180
2009	1408	1.55	1.10	8.42	6.80-10.32	180
2010	1721	1.61	1.25	9.07	7.14-11.40	180
2011	4189	2.56	1.19	27.09	22.30-32.80	178
2012	408	0.78	0.83	2.68	2.10-3.33	179
2013	1620	1.76	1.08	10.94	8.97-13.25	180
Overall (1967-2013)	52900	1.41	0.55	7.04	5.56-8.81	41 (years)

Table 3. Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia using only the 1st haul (Rago et al. 1995) summarized by year, Index = $(\exp(\ln(x + 1)) - 1) \times 2.28$, SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean ln (x+1)	SD	Index	C.I. (± 2 SE)	N (hauls)
1967	191	1.18	1.00	5.17	3.20-7.86	42
1968	184	1.04	0.92	4.15	2.68-6.06	50
1969	193	0.97	0.94	3.73	2.39-5.46	55
1970	345	1.39	1.11	6.88	4.52-10.06	56
1971	165	0.90	0.90	3.34	2.17-4.81	60
1972	84	0.45	0.59	1.28	0.87-1.75	90
1973	133	0.60	0.82	1.86	1.12-2.76	70
1980	216	0.82	0.96	2.90	1.85-4.21	72
1981	112	0.64	0.74	2.05	1.28-2.99	58
1982	172	0.86	0.96	3.10	1.86-4.71	54
1983	185	0.99	0.94	3.86	2.44-5.71	51
1984	377	1.27	1.09	5.81	3.72-8.63	53
1985	216	0.94	0.92	3.54	2.4-4.97	71
1986	449	1.35	1.07	6.53	4.56-9.06	72
1987	1314	2.27	1.22	19.77	14.25-27.13	72
1988	820	1.57	1.21	8.66	6.2-11.85	90
1989	1427	2.06	1.18	15.68	11.71-20.77	90
1990	720	1.58	1.12	8.76	6.44-11.7	90
1991	462	1.17	1.05	5.04	3.59-6.85	90
1992	1143	1.65	1.31	9.63	6.76-13.41	90
1993	1241	2.34	0.89	21.36	17.31-26.25	90
1994	969	1.93	1.09	13.37	10.17-17.4	90
1995	559	1.37	1.07	6.71	4.89-8.99	90
1996	2326	2.60	1.27	28.29	21.11-37.69	90
1997	931	1.83	1.14	11.92	8.9-15.76	90
1998	1365	2.12	1.22	16.66	12.35-22.23	90
1999	274	0.92	0.91	3.43	2.43-4.64	90
2000	1528	2.22	1.23	18.70	13.91-24.9	90
2001	1671	2.16	1.32	17.52	12.7-23.89	90
2002	486	1.17	1.13	5.03	3.48-7.01	90
2003	2042	2.50	1.26	25.61	19.09-34.13	90
2004	1129	2.07	1.04	15.75	12.19-20.19	90
2005	835	1.79	1.07	11.42	8.64-14.9	90
2006	767	1.76	1.06	11.02	8.34-14.36	90
2007	1271	2.09	1.21	16.07	11.95-21.39	90
2008	867	1.70	1.11	10.15	7.56-13.42	90
2009	861	1.72	1.11	10.47	7.81-13.83	90
2010	994	1.75	1.26	10.83	7.78-14.82	90
2011	2397	2.70	1.17	31.69	24.29-41.16	90
2012	265	0.92	0.87	3.47	2.50-4.63	90
2013	877	1.82	1.10	11.85	8.92-15.54	90
(1967- 2013)	32563	1.54	0.59	8.36	6.57-10.52	41 (years)
(1990-2013)	25980	1.83	0.49	11.9	9.35-15.01	24 (years)

Table 4. Catch of young-of-the-year Striped Bass per seine haul in the primary nursery area in 2013 summarized by drainage and river.

Drainage River	<u>2013</u>				<u>All Years Combined</u> (1967-2013)			
	Total Fish	Index	C.I. (±2 SE)	N (hauls)	Total Fish	Index	C.I. (±2 SE)	N (years)
JAMES	677	14.30	10.35 – 19.48	60	21161	8.96	6.82 – 11.61	41
James	317	11.13	7.55 – 16.00	40	12974	8.17	6.01 – 10.90	41
Chickahominy	360	23.06	13.31– 38.93	20	8187	11.15	8.38 – 14.64	41
YORK	540	11.23	8.48 – 14.69	70	15411	5.38	4.16 – 6.83	41
Pamunkey	264	11.31	6.76 – 18.16	30	7643	6.04	4.33 – 8.19	41
Mattaponi	276	11.17	8.08 – 15.19	40	7768	4.92	3.91 – 6.09	41
RAPPAHANNOCK	403	7.49	4.63 – 11.54	50	16188	7.14	5.36 – 9.32	41
Overall	1620	11.27	9.18 - 13.75	180	52760	8.02	7.71 – 8.34	41

Table 5. Site specific Striped Bass indices and average site salinity during 2013 compared to historic (1967 – 2013) index values with corresponding average salinities (Avg. Sal., ppt). The York drainage includes Pamunkey and Mattaponi rivers. Index stations are indicated by bold font.

Drainage															
JAMES		Station	J12	J22	J29	J36	J42	C1	C3	J46	J51	J56	J62	J68	J77
1967-2013	Avg. Sal.	14.4	7.8	4.8	2.5	1.6	1.5	1.3	0.6	0.3	0.2	0.2	0.1	0.2	
	Index	1.8	14.5	7.5	12.7	13.3	16.7	7.4	20.0	16.6	6.3	10.0	6.7	2.5	
2013	Avg. Sal.	12.1	4.6	2.2	0.6	0.1	0.2	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
	Index	0.4	18.3	11.7	12.6	37.2	39.3	8.4	12.2	24.5	8.4	45.4	0.6	1.5	
YORK		Station	Y15	Y21	Y28	P36	P42	P45	P50	P55					
1967-2013	Avg. Sal.	16.6	13.9	10.7	4.2	1.7	0.7	0.4	0.3						
	Index	1.2	1.8	5.3	11.5	4.2	8.8	12.4	4.5						
2013	Avg. Sal.	16.0	14.6	9.8	2.9	0.5	0.1	0.1	0.1						
	Index	1.3	4.5	34.6	36.6	7.8	21.0	11.6	5.6						
		Station				M33	M37	M41	M44	M47	M52				
1967-2013	Avg. Sal.				4.6	2.4	1.2	0.4	0.3	0.1					
	Index				5.9	8.0	6.8	5.5	4.8	1.4					
2013	Avg. Sal.				3.5	1.0	0.1	0.0	0.0	0.0					
	Index				7.2	7.7	21.7	11.3	13.0	2.4					
RAPPAHANNOCK		Station	R12	R21	R28	R37	R41	R44	R50	R55	R60	R65	R69	R75	
1967-2013	Avg. Sal.	14.2	12.9	10.0	5.4	3.2	2.0	1.0	0.6	0.2	0.2	0.1	0.1		
	Index	0.5	0.8	2.7	3.2	5.7	7.9	13.0	39.2	5.7	4.0	3.0	2.1		
2013	Avg. Sal.	13.8	12.4	9.8	4.3	1.7	0.8	0.1	0.1	0.0	0.0	0.0	0.0		
	Index	0.9	0.6	6.0	0.0	13.1	3.4	28.9	30.4	5.1	5.1	3.1	1.5		

Table 6. Catch of young-of-the-year Striped Bass in the primary nursery areas of Virginia in 2013 summarized by sampling round and month.

Month (Round)	N (hauls)	Total Fish	<u>2013</u>			<u>All Years Combined (1967-2013)</u>				
			Scaled Mean	C.I. (± 2 SE)	Decrease From Previous Round	N (years)	Total Fish	Scaled Mean	C.I. (± 2 SE)	Decrease From Previous Round
July (1 st)	36	469	17.06	11.07 – 25.75		41	16329	11.17	8.72 – 14.16	
(2 nd)	36	399	13.44	8.59 – 20.46	14.9 %	41	12956	8.30	6.40 – 10.62	20.7%
Aug. (3 rd)	36	313	9.50	5.87 – 14.75	21.6%	41	9556	6.53	5.05 – 8.30	26.2%
(4 th)	36	261	11.02	7.43 – 15.93	16.6%	37	8175	6.13	4.69 – 7.88	14.5%
Sept. (5 th)	36	178	6.20	3.79 – 9.57	31.8%	34	5744	4.95	3.82 – 6.29	29.7%

Table 7. Water temperature (°C) recorded at seine survey stations in 2013. The York drainage includes the Pamunkey and Mattaponi rivers. Index stations are indicated by bold font. Red colors denote temperatures over 30°C; blue colors denote temperatures below 25°C.

Drainage		Station	J12	J22	J29	J36	J42	C1	C3	J46	J51	J56	J62	J68	J78
JAMES	Round	1	29.9	33.0	28.5	27.0	28.8	27.8	28.8	28.8	28.9	27.6	28.7	27.4	27.8
		2	28.0	29.5	29.3	28.0	27.9	28.6	28.7	28.8	29.4	28.6	30.6	31.3	31.6
		3	27.3	25.2	29.2	26.4	28.9	27.7	27.6	28.6	26.9	26.3	27.6	28.4	29.9
		4	27.0	27.6	28.2	23.8	26.0	25.4	25.5	26.9	26.2	29.9	27.5	27.3	26.1
		5	27.1	26.6	28.9	25.3	27.3	25.9	26.0	27.9	25.8	25.5	27.8	28.4	27.4
YORK	Round	1	Y15	Y21	Y28	P36	P42	P45	P50	P55					
		2	28.8	27.8	27.0	27.8	27.9	28.3	20.1	28.3					
		3	26.6	25.6	25.7	28.1	28.4	29.4	29.4	29.7					
		4	26.1	25.0	26.9	27.8	28.3	29.2	29.8	29.5					
		5	26.4	25.7	24.4	25.8	26.4	26.9	27.2	26.6					
	Round	1	M33	M37	M41	M44	M47	M52							
		2	28.2	28.5	28.8	30.5	31.4	30.3							
		3	26.6	27.8	28.0	28.6	29.5	29.5							
		4	27.7	28.0	28.1	28.7	29.1	29.2							
		5	26.3	26.4	27.0	27.4	29.0	27.2							
RAPPAHANNOCK	Round	1	R12	R21	R28	R37	R41	R44	R50	R55	R60	R65	R69	R76	
		2	30.8	29.5	27.3	28.7	30.0	29.8	29.2	29.8	29.8	31.5	28.5	29.6	
		3	31.3	30.9	29.1	30.6	31.2	30.7	30.0	30.5	30.2	30.5	30.5	31.4	
		4	27.9	27.0	25.6	26.3	27.4	26.9	26.3	26.5	26.4	25.9	26.9	ns	
		5	26.6	27.1	26.1	27.1	27.3	27.1	25.0	25.9	25.7	26.3	26.6	27.3	

Table 8. Catch of young-of-the-year Striped Bass per seine haul in the primary nursery areas of Virginia in 2013 summarized by water temperature.

Temp. (°C)	<u>2013</u>				<u>All Years Combined</u> (1967-2013)			
	Total Fish	Scaled Mean	C.I. (± 2 SE)	N (sites)	Total Fish	Scaled Mean	C.I. (± 2 SE)	N (sites)
15.0 - 19.9	N/A			0	54	2.30	0.66 – 4.85	20
20.0 - 24.9	66	16.24	1.39 – 91.15	4	2653	3.97	3.50 – 4.49	645
25.0 - 29.9	1374	10.70	8.72 – 13.04	162	40,022	8.59	8.21 – 8.99	4389
30.0 - 34.9	180	12.57	5.21 – 27.16	14	9656	9.62	8.70 – 10.62	931
Overall	1620	10.94	8.97 – 13.25	180	52,760	8.02	7.71 – 8.34	6050

Table 9. Catch of young-of-the-year Striped Bass per seine haul in the primary nursery areas of Virginia in 2013 summarized by salinity.

Salinity (ppt)	<u>2013</u>				<u>All Years Combined</u> (1967-2013)			
	Total Fish	Scaled Mean	C.I. (± 2 SE)	N (sites)	Total Fish	Scaled Mean	C.I. (± 2 SE)	N (sites)
0.0 - 4.9	1583	11.61	9.48 – 14.12	168	49,042	9.28	8.90 – 9.68	5023
5.0 - 9.9	19	3.89	0.25 – 12.76	6	3294	4.35	3.88 – 4.85	755
10.0 - 14.9	18	4.88	1.54 – 11.16	6	422	2.16	1.75 – 2.61	244
15.0 - 19.9	N/A			0	2	0.12	0.00 – 0.29	28
Overall	1620	10.94	8.97 – 13.25	180	52,760	8.02	7.71 – 8.34	6050

Table 10. Salinity (ppt) recorded at seine survey stations in 2013. The York drainage includes the Pamunkey and Mattaponi rivers. Index stations are indicated by bold font.

Drainage															
JAMES		Station	J12	J22	J29	J36	J42	C1	C3	J46	J51	J56	J62	J68	J78
Round	1	5.7	1.5	0.1	0.0	0.0	0.1	0.0	0.0	0.1	0.1	0.0	0.0	0.0	
	2	13.1	4.5	1.7	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
	3	13.8	5.4	2.4	0.4	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1	
	4	13.0	5.0	3.2	1.1	0.2	0.3	0.3	0.1	0.1	0.1	0.1	0.1	0.1	
	5	14.9	6.8	3.7	1.7	0.4	0.5	0.3	0.1	0.1	0.1	0.2	0.1	0.1	
YORK		Station	Y15	Y21	Y28	P36	P42	P45	P50	P55					
Round	1	13.3	10.2	8.2	0.4	0.0	0.0	0.0	0.0	0.0					
	2	16.7	13.5	11.0	4.5	1.1	0.1	0.1	0.1	0.1					
	3	17.1	22.1	10.6	2.6	0.7	0.1	0.1	0.1	0.1					
	4	15.5	12.7	8.6	4.2	0.3	0.1	0.1	0.1	0.1					
	5	17.3	14.3	10.7	3.0	0.6	0.1	0.1	0.1	0.1					
Round	Station					M33	M37	M41	M44	M47	M52				
	1					1.4	0.0	0.0	0.0	0.0	0.0				
	2					4.8	1.9	0.2	0.0	0.0	0.0				
	3					4.7	1.2	0.1	0.0	0.0	0.0				
	4					2.6	0.6	0.1	0.0	0.0	0.0				
5					4.0	1.3	0.2	0.1	0.1	0.0					
RAPPAHANNOCK		Station	R12	R21	R28	R37	R41	R44	R50	R55	R60	R65	R69	R76	
Round	1	13.6	10.9	8.2	3.5	0.1	0.1	0.0	0.0	0.0	0.0	0.0	0.0		
	2	13.3	11.6	9.1	3.6	1.0	0.2	0.0	0.0	0.0	0.0	0.0	0.0		
	3	13.6	12.6	10.3	4.4	1.7	0.7	0.1	0.1	0.0	0.0	0.0	ns		
	4	14.2	12.9	10.2	4.7	2.1	1.1	0.2	0.1	0.1	0.0	0.0	0.0		
	5	14.3	13.8	11.0	5.5	3.5	1.7	0.3	0.1	0.1	0.1	0.1	0.1		

Table 11. Dissolved oxygen concentrations (mg/L) at seine survey stations in 2013. The York drainage includes the Pamunkey and Mattaponi rivers. Blue shaded values are more than one standard error (SE) less than the mean dissolved oxygen concentrations recorded at that station from 1989 to 2013. Yellow shaded values denote particularly low dissolved oxygen values (also more than 1 SE below the long-term mean). Index stations are indicated by bold font.

Drainage		Station	J12	J22	J29	J36	J42	C1	C3	J46	J51	J56	J62	J68	J78
JAMES	Round	1	6.3	8.3	6.3	5.9	9.2	7.5	6.9	6.7	6.4	6.4	6.1	6.0	6.1
		2	6.7	8.2	5.0	4.8	7.2	6.5	5.8	5.3	5.2	6.3	7.2	5.5	5.1
		3	5.7	7.2	6.8	6.6	7.8	7.2	6.4	6.1	6.7	6.9	10.6	6.2	5.7
		4	6.5	7.5	8.5	6.4	7.7	7.2	5.9	5.3	5.8	7.1	8.2	6.6	7.3
		5	4.5	6.4	6.6	5.5	7.7	6.9	6.0	5.6	6.4	6.3	9.5	6.3	6.8
YORK	Round	1	6.1	5.0	3.5	4.8	5.2	5.1	4.5	5.7					
		2	5.4	5.3	4.5	3.1	4.1	3.4	4.2	7.0					
		3	5.9	5.6	3.9	5.1	5.8	5.4	5.1	5.2					
		4	4.8	4.6	7.6	4.9	5.6	5.3	5.8	5.1					
		5	4.3	6.1	5.2	4.5	5.2	5.5	4.4	4.8					
	Round	1				M33	M37	M41	M44	M47	M52				
		2				3.5	4.0	3.9	3.7	4.6	3.9				
		3				7.5	3.7	3.9	4.4	6.2	5.1				
		4				3.3	3.9	4.3	4.8	5.9	4.8				
		5				3.7	4.6	4.9	6.2	7.3	5.9				
RAPPAHANNOCK	Round	1	R12	R21	R28	R37	R41	R44	R50	R55	R60	R65	R69	R76	
		2	7.2	5.2	4.5	4.8	6.5	6.6	7.7	7.3	6.6	7.4	4.4	4.0	
		3	7.1	5.8	4.7	5.6	6.1	7.2	5.9	6.1	4.2	5.3	5.2	5.6	
		4	6.8	6.7	5.8	6.2	6.7	6.8	5.9	6.3	5.4	6.9	6.8	ns	
		5	6.2	6.4	5.9	5.8	6.7	6.5	5.9	7.0	5.8	6.2	5.9	6.3	
5	6.8	5.9	6.1	6.1	5.9	6.8	6.7	7.0	6.0	4.5	5.5	4.3			

Table 12. Species collected during the 2013 seine survey (index and auxilliary stations).

Scientific Name	Common Name	Total Caught
<i>Menidia menidia</i>	Atlantic Silverside	7,217
<i>Notropis hudsonius</i>	Spottail Shiner	5,224
<i>Morone americana</i>	White Perch	4,762
<i>Trinectes maculatus</i>	Hogchoker	2,806
<i>Anchoa mitchilli</i>	Bay Anchovy	2,594
<i>Menidia beryllina</i>	Inland Silverside	2,416
<i>Morone saxatilis</i>	Striped Bass	2,325
<i>Fundulus heteroclitus</i>	Mummichog	1,083
<i>Micropogonias undulatus</i>	Atlantic Croaker	974
<i>Fundulus diaphanus</i>	Banded Killifish	940
<i>Brevoortia tyrannus</i>	Atlantic Menhaden	823
<i>Leiostomus xanthurus</i>	Spot	695
<i>Dorosoma petenense</i>	Threadfin Shad	655
<i>Dorosoma cepedianum</i>	Gizzard Shad	552
<i>Notropis analostanus</i>	Satinfin Shiner	420
<i>Ictalurus furcatus</i>	Blue Catfish	404
<i>Hybognathus regius</i>	Eastern Silvery Minnow	381
<i>Etheostoma olmstedi</i>	Tessellated Darter	288
<i>Bairdiella chrysoura</i>	Silver Perch	279
<i>Membras martinica</i>	Rough Silverside	221
<i>Fundulus majalis</i>	Striped Killifish	195
<i>Menticirrhus americanus</i>	Southern Kingfish	164
<i>Alosa aestivalis</i>	Blueback Herring	143
<i>Lepomis macrochirus</i>	Bluegill	108
<i>Lepomis gibbosus</i>	Pumpkinseed	80
<i>Anchoa hepsetus</i>	Striped Anchovy	79
<i>Ictalurus punctatus</i>	Channel Catfish	51
<i>Cynoscion regalis</i>	Weakfish	50
<i>Alosa sapidissima</i>	American Shad	48
<i>Perca flavescens</i>	Yellow Perch	33
<i>Micropterus salmoides</i>	Largemouth Bass	28
<i>Mugil cephalus</i>	Striped Mullet	27
<i>Anguilla rostrata</i>	American Eel	26
<i>Ictalurus catus</i>	White Catfish	24
<i>Lepomis auritus</i>	Redbreast Sunfish	21
<i>Alosa pseudoharengus</i>	Alewife	20
<i>Gambusia affinis</i>	Mosquitofish	20

Table 12. (cont'd.)

Scientific Name	Common Name	Total Caught
<i>Lepomis microlophus</i>	Redear Sunfish	14
<i>Micropterus punctulatus</i>	Spotted Bass	13
<i>Moxostoma macrolepidotum</i>	Shorthead Redhorse	12
<i>Syngnathus fuscus</i>	Northern Pipefish	11
<i>Mugil curema</i>	White Mullet	8
<i>Pomatomus saltatrix</i>	Bluefish	7
<i>Enneacanthus gloriosus</i>	Bluespotted Sunfish	7
<i>Paralichthys dentatus</i>	Summer Flounder	7
<i>Strongylura marina</i>	Atlantic Needlefish	5
<i>Cynoscion nebulosus</i>	Spotted Seatrout	4
<i>Micropterus dolomieu</i>	Smallmouth Bass	4
<i>Lepomis cyanellus</i>	Green Sunfish	4
<i>Elops saurus</i>	Ladyfish	4
<i>Cyprinus carpio</i>	Common Carp	3
<i>Notemigonus crysoleucas</i>	Golden Shiner	3
<i>Syngnathus louisianae</i>	Chain Pipefish	3
<i>Alosa mediocris</i>	Hickory Shad	2
<i>Chasmodes bosquianus</i>	Striped Blenny	2
<i>Lepisosteus osseus</i>	Longnose Gar	2
<i>Symphurus plagiusa</i>	Blackcheek Tonguefish	1
<i>Gobiesox strumosus</i>	Skilletfish	1
<i>Lagodon rhomboides</i>	Pinfish	1
<i>Trachinotus falcatus</i>	Permit	1
<i>Noturus gyrinus</i>	Tadpole Madtom	1
<i>Gobiosoma bosci</i>	Naked Goby	1
	Total	36,297

Table 13. Preliminary catch of Spottail Shiner from select juvenile Striped Bass seine survey stations using only the 1st haul (Rago et al. 1995) summarized by year, where x = total fish, Index = $(\exp(\ln(x + 1)) - 1)$, SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean ln (x+1)	SD	Index	C.I. (± 2 SE)	N (hauls)
1989	2940	2.64	1.15	12.99	10.34-16.25	121
1990	2068	2.12	1.30	7.35	5.62-9.54	124
1991	1429	1.87	1.24	5.49	4.17-7.14	119
1992	2357	2.02	1.40	6.50	4.83-8.65	123
1993	1713	1.96	1.27	6.13	4.65-8.01	118
1994	2498	2.29	1.34	8.91	6.77-11.66	120
1995	2216	2.10	1.36	7.16	5.37-9.46	120
1996	2280	2.28	1.27	8.74	6.72-11.29	119
1997	3605	2.17	1.53	7.77	5.67-10.53	125
1998	2092	2.12	1.32	7.36	5.53-9.72	114
1999	1252	1.48	1.30	3.38	2.48-4.52	126
2000	4882	2.73	1.43	14.39	10.92-18.86	125
2001	2848	2.39	1.33	9.92	7.64-12.82	128
2002	1541	1.30	1.40	2.67	1.86-3.70	128
2003	2972	2.42	1.40	10.21	7.76-13.34	129
2004	5113	3.25	1.13	24.72	19.98-30.54	123
2005	3585	2.63	1.40	12.85	9.71-16.91	119
2006	3451	2.47	1.51	10.85	7.96-14.68	117
2007	3823	2.58	1.47	12.22	9.09-16.33	118
2008	2152	1.97	1.46	6.16	4.51-8.31	124
2009	3033	2.21	1.54	8.10	5.89-11.02	122
2010	3983	2.38	1.54	9.79	7.16-13.26	121
2011	6194	3.20	1.41	23.50	17.84-30.85	117
2012	3997	2.40	1.58	10.03	7.27-13.71	121
2013	4502	2.77	1.38	14.98	11.49-19.44	125
Overall (1989-2013)	76,526	2.31	0.45	9.07	7.43-11.04	25 (years)

Table 14. Preliminary catch of Atlantic Silverside from select juvenile Striped Bass seine survey stations using only the 1st haul (Rago et al. 1995) summarized by year, where x = total fish, Index = $(\exp(\ln(x + 1)) - 1)$, SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean $\ln(x+1)$	SD	Index	C.I. (± 2 SE)	N (hauls)
1989	881	1.49	1.57	3.42	1.93-5.68	58
1990	1430	1.47	1.46	3.33	1.97-5.31	60
1991	2532	2.53	1.71	11.51	6.89-18.84	55
1992	5564	2.88	2.08	16.79	9.39-29.45	60
1993	2166	2.21	1.80	8.12	4.71-13.56	59
1994	2174	1.98	1.73	6.26	3.64-10.35	60
1995	2701	2.43	1.81	10.39	6.11-17.26	59
1996	4666	2.50	2.17	11.24	5.96-20.52	59
1997	973	1.83	1.48	5.26	3.24-8.23	58
1998	2182	2.61	1.60	12.64	8.02-19.64	60
1999	6227	3.37	1.50	28.03	18.49-42.23	57
2000	2936	2.83	1.72	15.99	9.81-25.71	58
2001	3487	2.92	1.69	17.48	11.02-27.41	62
2002	4582	3.48	1.53	31.38	20.82-47.04	60
2003	3470	2.16	2.15	7.63	3.95-14.04	60
2004	1473	1.76	1.79	4.78	2.64-8.19	60
2005	1843	2.48	1.50	10.97	7.18-16.52	62
2006	2613	2.56	1.68	11.96	7.52-18.72	64
2007	2021	2.68	1.51	13.61	8.84-20.70	58
2008	3107	2.04	1.78	6.71	3.93-11.06	63
2009	2618	2.76	1.68	14.80	9.35-23.13	63
2010	1347	2.38	1.26	9.78	6.87-13.78	64
2011	2953	2.63	1.80	12.94	7.87-20.92	63
2012	1079	1.86	1.46	5.42	3.43-8.31	62
2013	5745	2.50	2.14	11.17	6.12-19.79	64
Overall (1989-2013)	70,770	2.41	0.51	10.17	8.10-12.71	25 (years)

Table 15. Preliminary catch of Inland Silverside from select juvenile Striped Bass seine survey stations using only the 1st haul (Rago et al. 1995) summarized by year, where x = total fish, Index = $(\exp(\ln(x + 1)) - 1)$, SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean ln (x+1)	SD	Index	C.I. (± 2 SE)	N (hauls)
1989	471	1.15	0.96	2.17	1.63-2.81	107
1990	574	1.09	1.14	1.97	1.39-2.70	110
1991	285	0.86	0.87	1.37	1.00-1.81	105
1992	326	0.67	0.90	0.96	0.65-1.33	110
1993	368	0.76	0.97	1.14	0.77-1.59	106
1994	166	0.53	0.76	0.70	0.46-0.97	106
1995	104	0.44	0.62	0.56	0.38-0.75	107
1996	772	0.82	1.13	1.27	0.83-1.83	107
1997	175	0.54	0.76	0.71	0.48-0.98	110
1998	204	0.69	0.80	0.99	0.70-1.33	104
1999	298	0.72	0.93	1.06	0.73-1.45	113
2000	718	1.06	1.19	1.89	1.31-2.62	113
2001	626	0.96	1.15	1.61	1.10-2.24	115
2002	447	0.78	1.04	1.18	0.80-1.66	114
2003	545	1.21	0.99	2.37	1.80-3.06	113
2004	753	1.23	1.17	2.44	1.75-3.29	113
2005	368	0.93	0.94	1.53	1.11-2.03	110
2006	1161	1.32	1.32	2.73	1.90-3.79	112
2007	807	1.06	1.20	1.88	1.29-2.62	111
2008	658	1.15	1.11	2.14	1.56-2.87	114
2009	1691	1.88	1.29	5.56	4.16-7.35	114
2010	908	1.19	1.30	2.29	1.57-3.21	111
2011	1334	1.32	1.27	2.76	1.95-3.79	110
2012	901	1.45	1.10	3.26	2.49-4.25	113
2013	1608	1.82	1.34	5.15	3.79-6.89	115
Overall (1989-2013)	16,268	1.03	0.37	1.79	1.41-2.23	25 (years)

Table 16. Preliminary catch of Banded Killifish from select juvenile Striped Bass seine survey stations using only the 1st haul (Rago et al. 1995) summarized by year, where x = total fish, Index = $(\exp(\ln(x + 1)) - 1)$, SD = Standard Deviation, and SE = Standard Error.

Year	Total Fish (x)	Mean $\ln(x+1)$	SD	Index	C.I. (± 2 SE)	N (hauls)
1989	231	0.56	0.82	0.75	0.49 - 1.05	106
1990	235	0.65	0.88	0.92	0.63 - 1.28	109
1991	247	0.59	0.93	0.80	0.50 - 1.16	104
1992	153	0.46	0.77	0.59	0.37 - 0.84	108
1993	258	0.59	0.95	0.80	0.49 - 1.17	103
1994	200	0.53	0.84	0.70	0.44 - 1.01	105
1995	287	0.66	1.01	0.93	0.59 - 1.35	105
1996	600	1.14	1.20	2.12	1.46 - 2.94	104
1997	365	0.88	1.00	1.41	0.99 - 1.92	110
1998	304	0.92	0.94	1.52	1.07 - 2.05	95
1999	335	0.79	1.01	1.20	0.81 - 1.68	107
2000	312	0.81	0.95	1.24	0.86 - 1.69	105
2001	374	0.99	0.95	1.68	1.23 - 2.22	108
2002	478	0.82	1.12	1.26	0.83 - 1.80	109
2003	841	1.16	1.24	2.18	1.50 - 3.03	109
2004	1388	1.79	1.31	5.00	3.63 - 6.77	103
2005	721	1.29	1.22	2.64	1.86 - 3.65	100
2006	498	0.93	1.18	1.53	0.99 - 2.21	97
2007	677	1.32	1.18	2.73	1.94 - 3.74	98
2008	1017	1.62	1.19	4.05	3.00 - 5.37	105
2009	1202	1.74	1.29	4.72	3.43 - 6.39	102
2010	1927	2.15	1.37	7.63	5.57 - 10.34	101
2011	1920	1.95	1.95	6.00	4.25-8.32	97
2012	1831	2.11	1.33	7.22	5.32-9.70	102
2013	653	1.24	1.19	2.45	1.74-3.36	105
Overall (1989-2013)	17,054	1.11	0.52	2.02	1.46-2.72	25 (years)




Figure 1. Juvenile Striped Bass seine survey stations. Numeric portion of station designation indicates approximate mile from mouth of river.




Figure 2. Scaled geometric mean of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) by year. Vertical bars are 95% confidence intervals as estimated by ± 2 standard errors of the mean. Horizontal lines indicate the historical geometric mean (solid) and confidence intervals (dotted) for 1967-2013.




Figure 3. Scaled geometric mean of young-of-the-year Striped Bass in the primary nursery areas of Virginia (index stations) by drainage and river.




Figure 4. Catch of young-of-the-year Striped Bass by station in the James River drainage in 2013. Index station catch represents an average of two hauls; auxiliary station (starred) catch represents one haul.




Figure 5. Catch of young-of-the-year Striped Bass by station in the York and Mattaponi rivers in 2013. Index station catch represents an average of two hauls; auxiliary station (starred) catch represents one haul.




Figure 6. Catch of young-of-the-year Striped Bass by station in the York and Pamunkey rivers in 2013. Index station catch represents an average of two hauls; auxiliary station (starred) catch represents one haul.




Figure 7. Catch of young-of-the-year Striped Bass by station in the Rappahannock River in 2013. Index station catch represents an average of two hauls; auxiliary station (starred) catch represents one haul.




Figure 8. Scaled geometric mean of young-of-the-year Spot by year from select seine survey stations in Virginia tributaries of Chesapeake Bay. The vertical bars are 95% confidence intervals as estimated by 2 standard errors of the mean. Horizontal lines indicate the historical geometric mean (solid) and confidence intervals (dotted) for 1967-2013.




Figure 9. Delta-lognormal mean of young-of-the-year White Perch from select seine survey stations by river and year.




Figure 10. Delta-lognormal mean of young-of-the-year White Perch from the James River nursery area by year. The horizontal line indicates the historical mean for 1967-2013.




Figure 11. Delta-lognormal mean of young-of-the-year White Perch from the Pamunkey River nursery area by year. The horizontal line indicates the historical mean for 1967-2013.




Figure 12. Delta-lognormal mean of young-of-the-year White Perch from the Mattaponi River nursery area by year. The horizontal line indicates the historical mean for 1967-2013.




Figure 13. Delta-lognormal mean of young-of-the-year White Perch from the Rappahannock River nursery area by year. The horizontal line indicates the historical mean for 1967-2013.




Figure 14. Delta-lognormal mean of young-of-the-year Atlantic Croaker from select seine survey stations in Virginia tributaries of Chesapeake Bay by year. The horizontal line indicates the historical mean for 1980-2013.