

2010

Modeling the Effects of Genetic Manipulations of Calsequestrin on Local Calcium Release and Depletion in Cardiac Myocytes

Ryan P. Carpenter
William & Mary

Gregory D. Smith
William & Mary, gdsmit@wm.edu

Sandor Gyorke

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Carpenter, R. P., Gyorke, S., & Smith, G. D. (2010). Modeling the Effects of Genetic Manipulations of Calsequestrin on Local Calcium Release and Depletion in Cardiac Myocytes. *Biophysical Journal*, 98(3), 104a.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

situ (an SCR) occurs in a synchronized fashion; however, the mechanism responsible for synchronizing SCR activity in coupled myocardium is unknown. Since others have reported that Ca can diffuse through gap junctions (GJ), we hypothesize that uncoupling cells by blocking GJ will desynchronize, and attenuate SCR activity. Methods: To test this hypothesis high resolution optical mapping of Ca (Indo-1AM) from the anterior surface of the Langendorff perfused guinea pig heart (n=5) was performed in hearts under high Ca conditions ($[Ca^{2+}]_e=5.5mM$), with and without carbonoxolone (CBX, 50 μM) to reduce GJ coupling. Endocardial cryoablation were performed to eliminate Purkinje fibers and cytochalasin-D (7 μM) was administered to remove motion artifact. Fifteen seconds of rapid pacing (350-160 ms cycle length) followed by a pause was used to induce SCR activity. Results: In all preparations, SCR activity was observed across the entire mapping field before and after CBX. With CBX, the amplitude of SCR activity increased (+14.8%, $p < 0.05$) and its time to peak occurred earlier (-11.2%, $p < 0.01$) compared to no CBX. CBX also decreased the range of local SCR time to peaks across the mapping field (-17.2%, $p < 0.05$), suggesting that uncoupling myocytes synchronizes spontaneous calcium release across cells. There was no statistical difference in the occurrence of triggered activity before and during CBX. Conclusions: These results demonstrate that the occurrence of spontaneous calcium release in tissue (an SCR) does not require Ca diffusion through GJs. In fact, spontaneous calcium release in tissue is paradoxically enhanced during GJ inhibition.

546-Pos

Synchronization of Spontaneous Calcium Release Waves Among Myocytes in Intact Heart Determines the Magnitude of Delayed Afterdepolarizations and Triggered Activity

J. Andrew Wasserstrom¹, Yohannes Shiferaw², Satvik Ramakrishna¹, Heetabh Patel¹, Matthew J. O'Toole¹, Amanda Pappas¹, Asrar Khan¹, Rishi Arora¹, Gary L. Aistrup¹.

¹Northwestern University Feinberg School of Medicine, Chicago, IL, USA,

²California State University - Northridge, Northridge, CA, USA.

Intracellular Ca waves occur as the result of spontaneous Ca release (SCR) during Ca overload. Ca waves activate Na-Ca exchange, causing delayed afterdepolarizations (DAD) which can achieve threshold and produce triggered extrasystoles. It is not known how these single cell events depolarize enough myocytes in intact heart to produce a triggered beat. We combined experimental observations with computer simulations to explain how SCR synchronization among myocytes brings a critical tissue mass to threshold. Confocal microscopy was used to measure SCR waves in groups of myocytes in the LV epicardium of rat hearts loaded with fluo-4AM. Contraction was abolished with cytochalasin-D and blebbistatin. Raising extracellular [Ca] and rapid pacing protocols were used to increase sarcoplasmic reticulum (SR) Ca load and induce Ca waves. As Ca load increased, the number of myocytes giving waves increased. Both the wave latency and the variability (SD) of wave latency decreased with increasing Ca load. Similar results were obtained in isolated rat ventricular myocytes, indicating that the reduction in latency interval and variability represent intrinsic properties of SR release in Ca overload. Computer simulations demonstrate that decreasing wave latency and variability determine the rate and magnitude of increased cytoplasmic [Ca] and therefore determine the timing and magnitude of the DAD. The synchrony of SCR waves among myocytes therefore determines the likelihood of achieving threshold and producing a triggered beat. These results demonstrate that intrinsic properties of SR Ca release are responsible for Ca wave synchronization during Ca overload, causing DADs to reach threshold and produce triggered arrhythmias.

547-Pos

Looking at the Trigger for CICR During Rat Cardiac Action Potentials

Cherrie H.T. Kong, Mark B. Cannell.

University of Auckland, Auckland, New Zealand.

It is generally accepted that cardiac CICR is triggered by L-type Ca²⁺ channel activation during the action potential. At the microscopic level, CICR is revealed by calcium sparks whose activation by single L-type Ca²⁺ channels has been demonstrated. In this study, we have tried to visualize the Ca²⁺ influx (that triggers CICR) during normal action potentials and when the L-type Ca²⁺ channel is partially blocked and/or during the application of an L-type Ca²⁺ channel gating modifier FPL64176. Using Fluo-4 and high speed confocal line scanning we have detected the rise in Ca²⁺ that precedes SR Ca²⁺ release. When ~90% of L-type Ca²⁺ channels are blocked with 10 μM nifedipine, L-type Ca²⁺ influx is seen as an increase in fluorescence of ~2 %/ms, which is 8% of the rate of rise of Ca²⁺ associated with Ca²⁺ sparks. This is associated with a latency for Ca²⁺ spark activation of typically 9 ms; assuming a K_d for fluo-4 of 800 and a resting Ca²⁺ of 65nM the trigger in these condi-

tions equates to a current of ~1 nA for a 30 pL cell. An unexpected finding was that FPL64176 did not dramatically decrease the latency for Ca²⁺ spark activation, as might be expected if many short L-type Ca²⁺ channel openings are required to activate CICR. In addition, no 'sparklets' were observed as might be expected if L-type Ca²⁺ channels are located almost exclusively in the junctional space and Ca²⁺ sparks activate with minimal delay.

548-Pos

Modeling the Effects of Genetic Manipulations of Calsequestrin on Local Calcium Release and Depletion in Cardiac Myocytes

Ryan P. Carpenter¹, Sandor Gyorke², Gregory D. Smith¹.

¹The College of William and Mary, Williamsburg, VA, USA, ²The Ohio State University, Columbus, OH, USA.

Cardiac calsequestrin (CASQ2), a Ca buffer localized to the junctional SR (jSR) of cardiac myocytes, is known to bind to the RyR-triadin-junctin complex, participate in the luminal regulation of RyRs, and modulate Ca spark activity. To investigate the functional role of CASQ2 during spontaneous Ca sparks, we constructed a hybrid CTMC-ODE stochastic simulation of a Ca release site model composed of 100 Lee-Keener RyRs [J. Theor. Biol. 253:668-679, 2008] that includes Ca activation, Ca inactivation, CASQ2-RyR binding, and the dynamics of myoplasmic and luminal domain Ca and buffer concentrations. Myoplasmic and network SR [Ca] were determined by balancing the simulated average release flux and reuptake mediated by SERCA. The model reproduces average properties of spontaneous sparks in normal myocytes including spark amplitude, blink nadir, and junctional SR recovery time. Parameter studies were performed to interpret the effects of known arrhythmogenic CASQ2 mutants [Terentyev et al., Biophys. J. 95(4):2037-2048, 2008] on average spark properties. Increases in the total amount of CASQ2 resulted in increased spark amplitudes and increased jSR recovery times (observed in myocytes overexpressing wildtype CASQ2); shallower nadirs were observed in simulations but not experiment. Increasing the K_d of CASQ2 and Ca binding decreases spark amplitude and jSR recovery time (observed in myocytes expressing the dominant negative mutation CASQ2-DEL that suppresses CASQ2-Ca binding); in simulations (but not experiment) release-reuptake balance leads to increased network SR [Ca]. Increasing the K_d of CASQ2 and RyR binding is associated with decreased network SR [Ca], as in myocytes expressing CASQ2-R33Q, a variant with dominant negative effects on interactions of CASQ with RyRs. The implications of these results to mechanism of release regulation by CASQ2 will be discussed.

549-Pos

Loss of Calsequestrin (Casq2) in the Heart Increases Spark Frequency and Alters Spark Properties

Eleonora Savio Galimberti¹, Sabine Huke¹, Sylvain J. Le Marchand², David W. Piston², Bjorn C. Knollmann¹.

¹Vanderbilt University, Department of Medicine, Division of Clinical Pharmacology, Nashville, TN, USA, ²Vanderbilt University, Department of Molecular Physiology & Biophysics, Nashville, TN, USA.

Lack of Casq2 causes spontaneous Ca²⁺ releases from the sarcoplasmic reticulum (SR) and catecholaminergic-polymorphic ventricular tachycardia. We tested the hypothesis that lack of Casq2 alters elementary Ca²⁺ release events (Ca²⁺ sparks) by comparing spark properties of ventricular myocytes isolated from wild-type (WT) and Casq2 null (Casq2^{-/-}) mice. Sparks were recorded in line-scan mode and analyzed with SparkMaster. Spark mass was calculated as amplitude \times 1.206 \times FWHM³. SR Ca²⁺ load was measured by rapid application of caffeine. In intact Casq2^{-/-} myocytes stimulated with 100nM isoproterenol, spark amplitude and spark width (FWHM) increased compared to WT (0.77 \pm 0.019 vs. 0.39 \pm 0.02 $\Delta(F/Fo)$, and 2.4 \pm 0.03 vs. 1.3 \pm 0.05 μm , 1095 and 105 sparks respectively), resulting in larger spark mass (20 \pm 1.4 vs. 2.2 \pm 0.4 $\Delta(F/Fo)$, μm^3). Time-to-peak and spark duration (FDHM) were 2.5-fold longer and spark frequency was 4-fold higher in Casq2^{-/-} myocytes (2.4 \pm 0.2 vs. 0.5 \pm 0.08 sparks \times 100 $\mu m^{-1} \times s^{-1}$). Spark-mediated leak (spark mass \times spark frequency) was much larger (47.6 vs. 1.2 $\Delta(F/Fo)$ μm^3). In saproin-permeabilized myocytes, spark-mediated leak and spark frequency were also higher in Casq2^{-/-} myocytes (249.2 vs. 171.3 $\Delta(F/Fo)$ μm^3 , and 9.5 \pm 1 vs. 4.2 \pm 1 sparks \times 100 $\mu m^{-1} \times s^{-1}$, 587 and 333 sparks for Casq^{-/-} and WT respectively), but the differences between Casq^{-/-} and WT were less pronounced compared to intact myocytes. This may be a consequence of the increased spark-mediated SR Ca²⁺ leak resulting in significantly decreased SR Ca²⁺ load in permeabilized Casq2^{-/-} cells (-25% at baseline and -31% with 50 μM cAMP, n = 10-16 myocytes per group).

Conclusions: Lack of Casq2 in cardiac myocytes increases the spark frequency and the spark-mediated leak. This is still observed in permeabilized cells despite decreased SR Ca²⁺ load, suggesting that this is due to a primary Casq2