

8-19-2002

Are Pfiesteria species toxicogenic? Evidence against production of ichthyotoxins by Pfiesteria shumwayae

JP Berry

KS Reece

Virginia Institute of Marine Science

KS Rein

DG Baden

LW Haas

See next page for additional authors

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Immunology and Infectious Disease Commons](#)

Recommended Citation

Berry, JP; Reece, KS; Rein, KS; Baden, DG; Haas, LW; Ribeiro, WL; Shields, JD; Snyder, RV; Vogelbein, WK; and Gawley, RE, "Are Pfiesteria species toxicogenic? Evidence against production of ichthyotoxins by Pfiesteria shumwayae" (2002). *VIMS Articles*. 1379. <https://scholarworks.wm.edu/vimsarticles/1379>

This Article is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Authors

JP Berry, KS Reece, KS Rein, DG Baden, LW Haas, WL Ribeiro, JD Shields, RV Snyder, WK Vogelbein, and RE Gawley

Are *Pfiesteria* species toxicogenic? Evidence against production of ichthyotoxins by *Pfiesteria shumwayae*

J. P. Berry*, K. S. Reece†, K. S. Rein‡, D. G. Baden§, L. W. Haas†, W. L. Ribeiro†, J. D. Shields†, R. V. Snyder†, W. K. Vogelbein†, and R. E. Gawley*^{¶1}

*Department of Chemistry/National Institute of Environmental Health Sciences, Marine and Freshwater Biomedical Science Center, University of Miami, P.O. Box 249118, Coral Gables, FL 33124; †Virginia Institute of Marine Science, College of William and Mary, Route 1208, Gloucester Point, VA 23062; ‡Department of Chemistry, Florida International University, 11200 SW 8th Street, Miami, FL 33199; and §Center for Marine Science, University of North Carolina, 5600 Marvin K. Moss Lane, Wilmington, NC 28409

Edited by John H. Law, University of Arizona, Tucson, AZ, and approved May 29, 2002 (received for review April 12, 2002)

The estuarine genus *Pfiesteria* has received considerable attention since it was first identified and proposed to be the causative agent of fish kills along the mid-Atlantic coast in 1992. The presumption has been that the mechanism of fish death is by release of one or more toxins by the dinoflagellate. In this report, we challenge the notion that *Pfiesteria* species produce ichthyotoxins. Specifically, we show that (i) simple centrifugation, with and without ultrasonication, is sufficient to “detoxify” water of actively fish-killing cultures of *Pfiesteria shumwayae*, (ii) organic extracts of lyophilized cultures are not toxic to fish, (iii) degenerate primers that amplify *PKS* genes from several polyketide-producing dinoflagellates failed to yield a product with *P. shumwayae* DNA or cDNA, and (iv) degenerate primers for *NRPS* genes failed to amplify any *NRPS* genes but (unexpectedly) yielded a band (among several) that corresponded to known or putative *PKS*s and fatty acid synthases. We conclude that *P. shumwayae* is able to kill fish by means other than releasing a toxin into bulk water. Alternative explanations of the effects attributed to *Pfiesteria* are suggested.

Members of the estuarine dinoflagellate genus *Pfiesteria* have been linked to numerous fish kills in the estuaries of the mid-Atlantic coast of the U.S. in the past decade. Since first being identified and reported as the causative agent of fish kills in 1992 (1), there have been many follow-up investigations. An online literature search reveals over 100 articles with the keyword “*Pfiesteria*” published since 1995. These include reports of ichthyotoxicity and cytotoxicity (1–7) ascribed to the dinoflagellate as well as links between exposure to *Pfiesteria* and human health effects (6, 8–19). An entire issue of the journal *Environmental Health Perspectives* [(2001) *Environ. Health Perspect.*, Suppl. 5, 109, 633–808] was devoted exclusively to the *Pfiesteria* phenomenon and summarizes the state of affairs as of October 2000. *Pfiesteria* has also received considerable attention in the lay press, sometimes in a spectacularly sensationalized fashion [e.g., “The Coming Plague” (20)]. The emotion surrounding the *Pfiesteria* phenomenon, including accusations of data suppression and questionable ethics, has even sparked debate in anthropology journals (21–24).

Many authors either presume or explicitly state that *Pfiesteria* species are toxicogenic (i.e., toxin-producing) dinoflagellates, and that a “small-molecule” toxin is responsible for the observed effects on fish and mammals. This is a reasonable position, because a structurally diverse array of toxins are produced by many dinoflagellates and related microalgae, commonly referred to as harmful algal blooms (25–28). Based on the presumed toxicogenicity, we set out to isolate and characterize the putative toxin using bioassay-guided fractionation. Furthermore, to characterize possible secondary metabolites of *Pfiesteria* structurally, we used molecular genetic techniques to search for evidence of polyketide synthase (*PKS*) and nonribosomal peptide (*NRPS*) genes such as those involved in the biosynthesis of typical secondary metabolites of algae. We report the results of our investigations herein.

Materials and Methods

Dinoflagellate Cultures. A clonal isolate of *Pfiesteria shumwayae* [Center for Culture of Marine Phytoplankton (CCMP) 2089] was maintained in cultures as described below. The identity of *P. shumwayae* was confirmed additionally by PCR using species-specific primers for regions of the ribosomal RNA gene complex (29).

Algal prey. CCMP 2089 was cultured clonally on algal prey, *Rhodomonas salina* (CCMP 1319), in York River water (estuarine water adjusted to 12 psu with distilled water, autoclaved, and 0.22- μ m-filtered) with bacterial growth minimized by the addition of penicillin and streptomycin (final concentrations 2,500 units/liter and 2,500 μ g/liter, respectively).

Tilapia prey. CCMP 2089 was cultured in 38-liter aquaria containing artificial seawater (ASW) and tilapia (*Oreochromis niloticus*) in a Biosafety Level 3 containment facility. Water-quality parameters in tanks including pH, temperature, salinity, ammonia, nitrites, and dissolved oxygen were monitored weekly. Water was changed (\approx 60–75% aquarium volume using ASW) after fish mortalities in tanks ceased and ammonia levels were high (>30 ppm). Cell densities were determined daily during fish mortalities by fixing 50-ml aliquots of culture in Lugol’s iodine, centrifuging at 4,000 \times g for 15 min, decanting, and counting cells with a hemacytometer (Neubauer, Brightline, Horsham, PA).

Cell-Free Fractions. Raw tank water consisting of a culture of *P. shumwayae* that had been killing tilapia (*O. niloticus*, 20–50% daily mortality) for 8 months before the assay was treated as follows.

Supernatant. Tank water was centrifuged (4,500 rpm, 1 h) with and without prior sonication (Brinkman Polytron PT 10/35 at 27,000 rpm) to obtain cell-free aqueous lysates and supernatants, respectively.

Solvent extracts. Raw tank water was lyophilized and extracted sequentially with dichloromethane and methanol (MeOH) to obtain soluble extracts of cellular material. For assays, organic extracts were evaporated to dryness and redissolved in volumes of water equivalent to the original volume of the culture material to achieve concentrations comparable to whole cultures.

Ichthyotoxicity Controls. To ensure that our experiments were done with toxic cultures, ichthyotoxicity of CCMP 2089 was assessed using either adult *Gambusia holbrooki* or larval *Cyprinodon variegatus*. *G. holbrooki* were collected from the wild and maintained in 20-gallon aquaria filled with ground water. They

This paper was submitted directly (Track II) to the PNAS office.

Abbreviations: *PKS*, polyketide synthase; *NRPS*, nonribosomal peptide synthase; CCMP, Center for Culture of Marine Phytoplankton; ASW, artificial seawater; RT, reverse transcription; ITS, internal transcribed spacer.

Data deposition: The sequence reported in this paper has been deposited in the GenBank database (accession no. AF525290).

^{¶1}To whom reprint requests should be addressed. E-mail: rgawley@miami.edu.

were tested in 15-ml volumes of ground water (with or without solvent extracts) in 150-ml beakers (two fish per beaker) supplied with aeration. Mortality was assessed up to 14 days.

To evaluate toxicity of cell-free supernatants and lysates, *C. variegatus* were placed in six-well plates (3–5 fish per well). ASW alone was used as a negative control. To verify ichthyotoxicity of live dinospores, *C. variegatus* were placed in 10 ml of ASW medium containing dinoflagellates (diluted 1:1 in ASW to give a final concentration of 500 dinospores per ml). In both cases, mortality was monitored for 7 days, after which the water levels and water quality deteriorated.

Preparation of DNA and cDNA for PCR. Genomic DNA and cDNA were prepared from samples of clonal cultures of CCMP 2089 grown on algal prey as described above. For genomic DNA, cells from 100-ml samples were collected by vacuum filtration onto 3- μ m Nuclepore (Costar) filters, and DNA was prepared using the DNeasy tissue kit (Qiagen, Valencia, CA). For cDNA, total RNA was isolated from CCMP 2089 obtained from a 125-ml sample of the algal-grown culture and a 125-ml sample of the algal-grown culture that was exposed to *C. variegatus* for 18 h before RNA extraction. For comparison, cells from a dense culture (60 ml) of the algal food, *R. salina*, were collected and processed identically for RNA extraction and cDNA preparation. Cells were collected on 3- μ m, RNase-free Nuclepore (Costar) filters by vacuum filtration. RNA was extracted from cells by denaturation with TRIzol followed by chloroform/isoamyl alcohol (25:1) extraction and overnight precipitation of RNA at -20°C with an equal volume of isopropanol. RNA was subsequently redissolved in nuclease-free water (25 μ l), quantified spectrophotometrically, and shown by formaldehyde gel electrophoresis to be of high quality (30).

The RNA was reverse-transcribed by Moloney murine leukemia virus reverse transcriptase and random hexamers. Aliquots (5 μ l) of RNA were transferred to RNase-free Eppendorf tubes and incubated for 15 min at 65°C with random hexamers (2 μ l \times 50 ng/ μ l) in 17 μ l of nuclease-free water. To each, 3.5 μ l of 5 mM dNTPs, 7 μ l of 5 \times reverse transcription (RT) buffer, and 0.5 μ l of Moloney murine leukemia virus reverse transcriptase were added, and reactions were incubated at 42°C for 2 h. cDNA was stored at -20°C for subsequent PCR.

PCR and RT-PCR Evaluation of PKS and NRPS Genes. The presence and expression of *PKS* and *NRPS* genes were evaluated by PCR of total genomic DNA and total cDNA, respectively, with primers (synthesized by Invitrogen Life Technologies, Carlsbad, CA) specific to these genes (31–34). PCR of *PKS*s and *NRPS*s used the following primer sequences, based on the ketosynthase domain and adenylation domain of *PKS*s and *NRPS*s, respectively: *PKSI*, forward/reverse, 5'-MGIGARGCIYTICARATG-GAYCCICARMG-3'/5'-GGRTCNCIAR YTGIGTIC-CIGTICCRTGIGC-3'; *PKSII*, forward/reverse, 5'-TSGCST-GCTTCGAYGCSATC-3'/5'-TGGAANCCGAABCCGCT-3'; *NRPSI*, forward/reverse, 5'-TWYCGIACIGGIGA-YYKIGKICG-3'/5'-AWIGARKSICCIARRSIMRAARA-A-3'; and *NRPSII*, forward/reverse, 5'-GCNGCNGGNGCI-TAYGTICC-3'/5'-CCNCKDATYTTNACYTG-3'. Reactions (25 μ l) consisted of 50 mM KCl, 10 mM Tris-HCl, 1.5 mM MgCl₂, 200 μ M dNTPs, 1.5 units of *Taq* polymerase and reverse and forward primers (2.4 pM), and 0.5 or 2 μ l of template for genomic DNA and cDNA, respectively. Cycling conditions, based on a touchdown method, were: 94°C for 2 min; 10 cycles of 94°C for 30 s, 65°C (decreasing 1°C per cycle) for 90 s, 72°C for 2 min; 25–35 cycles of 94°C for 30 s, 55°C for 60 s, and 72°C for 2 min; and a final extension at 72°C for 10 min. As a positive control, type I *PKS* sequences were amplified with the *PKSI* primer from the plasmids PL29-*PKS* and

AK112 containing known *PKS* sequences from *Prorocentrum lima* and *Aphidinium klebsii*, respectively.

In addition to specific primers, PCR with primers (Invitrogen Life Technologies) specific to the 18S ribosomal subunit or the internal transcribed spacer (ITS) region of the ribosomal RNA gene complex were used to confirm the presence of eukaryotic, and specifically *P. shumwayae*, DNA. The universal 18S primers, 16s-A/16s-B (5'-GGTTGATCCTGCCAGTAGTCATATGCTG-3'/5'-GATCCTTCCGCAGGTTACCTACGGAAACC-3') were derived from Rowan and Powers (35). The species-specific primers PfbSSU/16s-B and PfbSSU/PfBITSR201L were developed based on comparing the SSU gene, or SSU gene and ITS region sequences, of *P. shumwayae* to sequences for these regions from *Pfiesteria piscicida* and several *Pfiesteria*-like organisms (29). PfbSSU has been described previously (29); the sequence of PfbSSU/ITSR201L is as follows: 5'-CCAGCTTCTGGATTTT-GTCGC-3'/5'-GCGATGAGGAAGAGAAAATGACG-3'. Reactions for universal 18S primers were conducted identically to *PKS*-specific primers. Reactions with species-specific primers required the addition of 0.4 mg/ml nonacetylated BSA as a stabilizer. Conditions for amplification with the PfbSSU/16s-B and PfbSSU/PfBITSR201L primer sets were as follows: 94°C for 4 min; 40 cycles of 94°C for 30 s, 56°C for 30 s, and 72°C for 90 s; and 72°C for 5 min. PCR products were analyzed by gel electrophoresis (30).

Results

Fish-Killing Activity of Dinoflagellate Cultures, Supernatant, Lysate, and Extracts. Relatively high densities ($>1,000$ zoospores per ml) of *P. shumwayae* that had been maintained on a supply of *R. salina* in culture or *O. niloticus* in aquaria were found to cause mortality of larval fish in less than 24 h (Fig. 1). *P. shumwayae* raised on *R. salina* and that had not been previously exposed to fish demonstrated 100% mortality of larval fish in 24–48 h compared with 72 h for aquarium cultures that had been killing *O. niloticus* in aquaria before assay with *C. variegatus*.

However, cell-free supernatants and lysates (prepared by sonication), failed to kill larval *C. variegatus* in ichthyotoxicity assays lasting up to 7 days. Likewise, organic extracts (dichloromethane and MeOH) of lyophilized zoospores, after reconstitution in ASW, were not ichthyotoxic when tested with *G. holbrooki* for 14 days at concentrations equivalent to or greater than the concentrations present in whole culture. Furthermore, no obvious behavioral changes were observed in fish in the cell-free fractions as compared with fish tested in the whole cultures that showed characteristic responses to an irritant, such as hyperactivity, “wavering,” and flick responses of the fins.

PCR and RT-PCR with *PKS*- and *NRPS*-Specific Primers. We used primers specific for type I and II *PKS* and *NRPS* genes (31, 33, 34, 36) to investigate the possible presence and expression of these genes by *P. shumwayae* using PCR and RT-PCR. Amplification of 1.8-kbp and 100- or 400-bp products with universal 18S (35) and *P. shumwayae*-specific primers (29), respectively, confirmed the presence of *P. shumwayae* DNA and cDNA (Figs. 2–4). Neither *PKS*-specific primer set amplified DNA from either total genomic DNA or cDNA isolated from *P. shumwayae* (Figs. 2 and 3). However, amplification of the ketosynthase region (700 bp) of the *PKS* (type I) sequence from plasmids PL29-*PKS* and AK112 confirmed the effectiveness of these primers (Fig. 3).

Neither *NRPS*-specific primer set yielded a product of the expected size (600 or 1,000 bp; see *Supporting Text*, which is published as supporting information on the PNAS web site, www.pnas.org) from either *P. shumwayae* genomic DNA or cDNA. However, both *NRPS* primer sets yielded several bands of various sizes from the genomic PCR and RT-PCR (see Fig. 5, which is published as supporting information on the PNAS web




Fig. 1. Fish-killing activity of *P. shumwayae* grown on algae and tilapia, plus supernatant and lysate, were assayed with *C. variegatus* ($n =$ three per well \times two wells per treatment). Cell-free lysates and supernatants were from raw tank water cultured on tilapia prey and centrifuged with or without ultrasonication. For lysate, no data were collected after 4 days, because the water in the wells evaporated. MeOH and dichloromethane (DCM) extracts of lyophilized raw tank water were tested against *G. holbrooki* ($n =$ two fish per beaker \times two beakers per treatment); no mortality was observed in this case up to 14 days (last 7 days not shown). "Untreated" and "ASW only" controls refer to fish tested in ground water (for *G. holbrooki*) or ASW (for *C. variegatus*), respectively.

site). Several of the bands from NRPSII were cloned, and the inserts were sequenced. Analysis by BLAST (37) revealed (i) no homologies to any genes in the GenBank database (several bands

of various sizes), (ii) homology to succinate dehydrogenases (495-bp band), or (iii) homology to the N-terminal portion of the ketosynthase domain of known and putative PKSs or fatty acid synthases (401-bp band). For details on the cloning, sequencing, and PCR studies with nested primers of this band, see *Supporting Text*.




Fig. 2. Gel (2% agarose) showing a lack of PCR amplification with genomic DNA from *P. shumwayae* with primers specific to ketosynthase domain of type I and II PKSs (PKSI and PKSII, lanes 1 and 2, respectively). Shown is positive amplification of 1.8-kb (lane 3) and 100-bp (lane 4) products, respectively, with "universal" 18S primer (35) and *P. shumwayae*-specific primers (PfB5SU/16s-B) confirm the presence of *P. shumwayae* DNA. Lane 5 is a 100-bp ladder.

Discussion

In general, demonstration of toxicogenicity requires that chemical constituents capable of causing the attributed effect(s) can be isolated from the suspected source when a given toxicosis is observed and subsequently shown to induce the same effect *in vitro*. Specifically therefore, in terms of the production of an exogenous ichthyotoxin by *Pfiesteria*, it must be shown that a compound can be separated from the medium in which fish are dying or dinoflagellate biomass itself and subsequently shown to induce mortality in fish.

The proposed "gold standard" for detecting ichthyotoxic *Pfiesteria* is a standardized fish bioassay (2, 38), although DNA probes have been used also to indicate the presence of *Pfiesteria* but not toxicity (39). In our hands, clonal cultures of *P. shumwayae* grown on either *Rhodomonas* algae or in aquaria with fish (*O. niloticus*) produced rapid mortality (Fig. 1) at rates comparable to those of the allegedly toxicogenic "TOX-A" strain of *P. shumwayae* reported by the Burkholder group (2, 38). Interestingly, and in contrast to previous reports (2), prior exposure to fish is not required since *P. shumwayae* grown on *R. salina* algae produced rapid mortality (Fig. 1).




Fig. 3. Gel (1% agarose) of RT-PCR products showing lack of expression of type I or II *PKS* genes by *P. shumwayae*. No amplification was observed with cDNA from *P. shumwayae* using type I *PKSI* (lane 2) or type II *PKSII* (lane 3) primer. Expression, likewise, is not seen with primers *PKSI* and *PKSII* (lanes 5 and 6, respectively) for the same culture exposed to larval fish (*C. variegatus*) 18 h before RNA extraction or with negative control of the algal food, *R. salina*, only (lanes 8 and 9, respectively). Positive amplification was observed with universal eukaryotic 18S primers (35) for CCMP 2089 (lane 4), CCMP 2089 exposed to *C. variegatus* (lane 7), and *R. salina* (lane 10). Amplification of the type I *PKS* sequences was observed for positive control plasmids PL29 (lane 11) and AK112 (lane 12). Negative controls of no template (lanes 13, 14, and 15, respectively) showed no amplification with *PKSI*, *PKSII*, or 18S primers. Lane 1 is a 1-kb DNA ladder.

If fish were being killed or even stunned by a toxin released by the dinoflagellate, that toxin should be present in the bulk water of the tank in which the fish are dying. However, we find that centrifugation of water from actively killing cultures of *Pfiesteria* renders the medium nontoxic. Conceivably, a toxin might be released by the dinoflagellate after physical contact with a fish. If this were true, then the toxin would have to be stored in the cell. However, lysis of the cells by ultrasonication followed by removal of cellular matter by centrifugation produced a solution that was completely nontoxic. In other words, *removal of cells or cellular solids detoxifies the medium*.

The only way a toxin that is present in the cultures could be removed from solution through centrifugation of cellular matter is by adsorption onto the solid cellular matter. However, methanolic and dichloromethane extracts of the lyophilized cell mass failed to produce any mortality after reconstitution in water. *These observations, by themselves, refute the hypothesis that there is a toxin in the cultures killing the fish*. Conceivably, the toxin could be unstable, such that it decomposes during the centrifugations and sonications described above. This seems unlikely, since it would be difficult to reconcile the presence of a labile toxin with the massive fish kills ascribed to these species. Others

have also tried to find a toxin (2–7, 40), but the only isolate characterized thus far is a phthalate artifact presumed to originate in an ASW preparation (41). Fish in control aquaria without *Pfiesteria* show no mortality; thus, ASW is not toxic.

As far as we are aware, all known ichthyotoxins produced by dinoflagellates are polyketides (27, 28, 42–45), which are biosynthesized by a family of homologous enzymes known as *PKSs* (45). Polyketide toxins include the brevetoxins, the ciguatoxins, maitotoxin, the recently elucidated ichthyotoxin prymnesin (46), pectenotoxin, prorocentrolides, zooxanthellotoxins, okadaic acid, luteophanols, and amphidinols (45).

The presence of a *PKS* gene in *Pfiesteria* would be an indication that a toxin indeed might be produced, but fractionation bioassays had simply missed it. To that end, we tested the *Pfiesteria* genome for the presence of *PKS* genes. Neither *PKS* primer set yielded a product from either genomic DNA (Fig. 2) or total cDNA (Fig. 3). Apparently, the primers lack sufficient homology to *PKS* genes in *Pfiesteria*, although the type I *PKS* primer set amplifies products with significant homologies to known and putative *PKSs* from the DNA of polyketide toxin-producing dinoflagellates such as *Karenia brevis* (brevetoxins), *P. lima* (okadaic acid), *A. klebsii* (amphidinolides), and *Symbio-*




Fig. 4. Gel (2% agarose) of RT-PCR products showing amplification with *P. shumwayae*-specific primer. Amplification of the expected 400-bp sequence was observed with cDNA from *P. shumwayae* (CCMP 2089) as well as the same culture exposed to larval *C. variegatus* 18 h before RNA extraction using the primers PFBUS/PFBITSR201L (lanes 2 and 5), confirming the presence of *P. shumwayae* cDNA. No amplification was seen (lane 8) with the algal food, *R. salina*. Positive amplification was also observed for all positive controls of *P. shumwayae* genomic DNA (lanes 3, 6, and 9). No amplification was observed for any of the negative controls of no template (lanes 4, 7, and 10). Lanes 1 and 11 are 100-bp DNA ladders.

dinium (zooxanthellatoxins) (refs. 31 and 32; further details will be published elsewhere).

Given this homology in PKSs, the production of a 401-bp product having homology to known and putative PKS or fatty acid synthase genes, using the NRPSII primer set, was very surprising! Inspection of the sequence revealed that the forward NRPSII primer primed both the forward and reverse directions. Control experiments confirmed that this amplification product originated in *P. shumwayae* and was not found in genomic or cDNAs of the cultured algal prey, *R. salina*, or ASW that had been incubated with fish in the absence of CCMP 2089 (see *Supporting Text*). In fact, one of the highest scoring sequences in the BLAST analysis of the 401-bp product was a fatty acid synthase homologue (GenBank accession no. AF082993) from *Cryptosporidium parvum* (47). *C. parvum* belongs to the Coccidia, a sister clade to the Dinophyceae within the Alveolata (48), suggesting that this gene does indeed originate from *P. shumwayae* and not from bacteria associated with either the cultures or fish. The function of the AF082993 gene is unclear. Although described as a fatty acid synthase homologue (47), it contains a loading domain found only in other PKSs such as FK506 and rapamycin. Dinoflagellates are prodigious producers of polyunsaturated fatty acids (PUFAs). Recently, type I PKS-like enzymes have been demonstrated to be involved in the production of PUFAs in both marine prokaryotes and lower marine eukaryotes (49). Careful inspection of these enzymes indicates that they are unique among type I PKSs in both their domain organization and their amino acid sequence. Furthermore, amplification of the ketosynthase domain of these PKSs using our PKS1 primer set would be unlikely (see *Supporting Text*). Thus we conclude that *P. shumwayae* does express PKS-encoding genes. However, these genes may be involved in PUFA biosynthesis rather than toxin production. Sequencing of the full-length gene and further characterization will be needed before we can understand the function of the encoded enzyme.

While these experiments do not eliminate polyketides as possible toxins produced by *Pfiesteria*, they strongly suggest that if *Pfiesteria* is capable of producing this class of secondary metabolites, the polyketide is either not excreted into the bulk medium and not released by cell lysis or is nontoxic. Taken together with the lack of ichthyotoxicity in centrifuged medium or solvent extracts, the conclusion is virtually inescapable that *P. shumwayae* does not produce an exogenous ichthyotoxin. Nevertheless, cultures of *P. shumwayae* (CCMP 2089) kill fish at rates that classify these cultures in Burkholder's TOX-A category (1, 2, 40). This paradox can be resolved by invoking other mechanisms.

P. shumwayae zoospores can attach to dermal cells with a peduncle and feed via myzocytosis (29). In tilapia, this causes a loss of mucus coat and scales as well as mild petechial hemorrhage, thereby precipitating epidermal erosion and bacterial colonization (29). Interestingly, lesions such as those ascribed to *Pfiesteria* have been recently shown to be caused by a fungal pathogen without prior exposure to other stressors (50). The fungus is highly pathogenic, particularly to menhaden exhibiting

a prior portal of entry. Although numerous reports have demonstrated mortality and morbidity of tilapia exposed to *Pfiesteria* (1–7, 35), all have used live cultures of the dinoflagellates.

The taxonomy of the *Pfiesteria* genus is becoming increasingly complex and unclear. Currently, two species, *P. piscicida* (51) and *P. shumwayae* (40), are recognized. As originally described (1, 3), *P. piscicida* allegedly had up to 24 life-cycle stages. Using a variety of techniques, Litaker *et al.* have challenged this assertion (52). Using species-specific hybridization probes, no evidence could be found for a chrysophyte-like cyst stage or the amoeba stages originally proposed. Some authors (3, 7, 53, 54) describe a more intricate “complex” of *Pfiesteria*-like organisms, often occurring in association with organisms such as fungi, bacteria, and similar dinoflagellates (53, 55). Moreover, most work on *Pfiesteria* is conducted using nonaxenic cultures that undoubtedly contain innumerable microorganisms unrelated to *Pfiesteria*. Further complicating matters, some authors assert a variety of “functional types” of *Pfiesteria* having different abilities to kill fish and produce a toxin (2, 7). Accordingly, it might be suggested that the lack of toxicogenicity observed here is specific to the species and isolate used; however, we have observed a similar lack of ichthyotoxicity for several isolates of *P. shumwayae*, *P. piscicida*, and other *Pfiesteria*-like organisms (unpublished observations made by the present authors in Miami, Wilmington, NC, and Gloucester Point, VA; see also ref. 29).

It may be that at least some of the effects attributed to *Pfiesteria* are due to other algae. Recently, one raphidophyte species, *Chattonella cf. verruculosa*, recovered from Rehoboth Bay, DE, during a massive fish-kill event in August, 2000 was found to produce the potent ichthyotoxin brevetoxin (56). It is possible that large fish kills are not the result of a single organism but ensue from a combination of ichthyotoxins such as a *Chattonella*-derived brevetoxin, seasonal hypoxia (57), and predation by *Pfiesteria* as well as subsequent or concurrent fungal invasion (50). Parenthetically, it is noteworthy that brevetoxin is a potent neurotoxin (58–64), so some of the neurological problems reported in humans and associated with *Pfiesteria* blooms (65–68) could be the result of brevetoxin insult from a previously unidentified raphidophyte.

In conclusion, we confirm that *P. shumwayae* can kill fish within 24 h. However, we find no evidence of ichthyotoxins from *P. shumwayae* and no evidence of genes encoding for PKSs typical of those found in toxin-producing dinoflagellates. We therefore conclude that *P. shumwayae* is not toxicogenic.

We thank T. Shedd and the U.S. Army Center for Environmental Health Research for the lease of the BSL3 facility at Virginia Institute of Marine Science (VIMS). We also thank C. Squyars, A. Miller, V. Foster, L. Ott, L. Walker, N. Stokes, and K. Hudson for technical assistance at VIMS (VIMS contribution no. 2480). This work was supported by Environmental Protection Agency and National Oceanic and Atmospheric Administration ECOHAB Grants R826655, R828225-01-0, and R826791-01-0 and National Institute of Environmental Health Sciences Grants T32-ES07320 and P30-ES05705. Work at the University of North Carolina was funded in part by a subcontract on P01-ES09563 awarded to the University of Maryland.

- Burkholder, J. M., Noga, E. J., Hobbs, C. H. & Glasgow, H. B., Jr. (1992) *Nature (London)* **358**, 407–410.
- Burkholder, J. M., Glasgow, H. B., Jr. & Deamer-Melia, N. (2001) *Phycologia* **40**, 186–214.
- Burkholder, J. M. & Glasgow, H. B., Jr. (1997) *Limnol. Oceanogr.* **42**, 1052–1075.
- Fairey, E. R., Edmunds, J. S., Deamer-Melia, N. J., Glasgow, H. B., Jr. Johnson, F. M., Moeller, P. D., Burkholder, J. M. & Ramsdell, J. S. (1999) *Environ. Health Perspect.* **107**, 711–714.
- Glasgow, H. B., Jr., Burkholder, J. M., Schmechel, D. E., Tester, P. A. & Rublee, P. A. (1995) *J. Toxicol. Environ. Health* **46**, 501–522.
- Kimm-Brinson, K. L., Moeller, P. D., Barbier, M., Glasgow, H. B., Jr., Burkholder, J. M. & Ramsdell, J. S. (2001) *Environ. Health Perspect.* **109**, 457–462.
- Marshall, H. G., Gordon, A. S., Seaborn, D. W., Dyer, B., Dunstan, W. M. & Seaborn, A. M. (2000) *J. Exp. Mar. Biol. Ecol.* **255**, 51–74.
- Shoemaker, R. C. & Hudnell, H. K. (2001) *Environ. Health Perspect.* **109**, 539–545.
- Hudnell, H. K., House, D., Schmid, J., Koltai, D., Stopford, W., Wilkins, J., Savitz, D. A., Swinker, M. & Music, S. (2001) *J. Toxicol. Environ. Health* **109**, 457–462.
- El-Nabawi, A., Quesenberry, M., Saito, K., Silbergeld, E., Vasta, G. & Eldefrawi, A. (2000) *Toxicol. Appl. Pharmacol.* **169**, 84–93.
- Levin, E. D., Rezvani, A. H., Christopher, N. C., Glasgow, H. B., Jr., Deamer-Melia, N. J., Burkholder, J. M., Moser, V. C. & Jensen, K. (2000) *Neurotoxicol. Teratol.* **22**, 533–540.
- Levin, E. D., Simon, B. B., Schmechel, D. E., Glasgow, H. B., Jr., Deamer-Melia, N. J., Burkholder, J. M., Moser, V. C., Jensen, K. & Harry, G. J. (1999) *Neurotoxicol. Teratol.* **21**, 215–221.

13. Grattan, L. M., Oldach, D., Perl, T. M., Lowitt, M. H., Matuszak, D. L., Dickson, C., Parrot, C., Shoemaker, R. C., Kauffman, C. L., Wasserman, M. P., Hebel, J. R., Charache, P. & Morris, J. G., Jr. (1998) *Lancet* **352**, 532–539.
14. Bever, C. T., Grattan, L. & Morris, J. G., Jr. (1998) *Md. Med. J.* **47**, 120–123.
15. Shoemaker, R. C. (1998) *Md. Med. J.* **47**, 64–66.
16. Matuszak, D. L., Sanders, M., Taylor, J. L. & Wasserman, M. P. (1997) *Md. Med. J.* **46**, 515–520.
17. Rullan, J. V. & Jenkins, S. R. (1998) *Va. Med. Q.* **125**, 78–80.
18. Shoemaker, R. C. (1997) *Md. Med. J.* **46**, 521–523.
19. Levin, E. D., Schmechel, D. E., Burkholder, J. M., Deamer-Melia, N. J., Moser, V. C. & Harry, G. J. (1997) *Environ. Health Perspect.* **105**, 1320–1325.
20. Wise, J. (2001) *MAXIM*, October, 75–184.
21. Griffith, D. (1999) *Hum. Organ.* **58**, 119–127.
22. Burkholder, J. M. & Glasgow, H. B., Jr. (1999) *Hum. Organ.* **58**, 443–454.
23. Lewitus, A. J., Rublee, P. A., Mallin, M. A. & Shumway, S. E. (1999) *Hum. Organ.* **58**, 455–458.
24. Oldach, D. (1999) *Hum. Organ.* **58**, 459–462.
25. Fenical, W., Baden, D., Burg, M., de Goyet, C. d. V., Grimes, D. J., Katz, M., Marcus, N., Pomponi, S., Rhines, P., Tester, P. & Vena, J. (1999) *From Monsoons to Microbes: Understanding the Ocean's Role in Human Health* (Natl. Acad. Press, Washington, DC).
26. Botana, L. M. (2000) *Seafood and Freshwater Toxins: Pharmacology, Physiology, and Detection* (Dekker, New York).
27. Yasumoto, T. & Murata, M. (1993) *Chem. Rev.* **93**, 1897–1909.
28. Shimizu, Y. (1993) *Chem. Rev.* **93**, 1685–1698.
29. Vogelbein, W. K., Shields, J. D., Haas, L. W., Reece, K. S. & Zwerner, D. E. (2001) *Environ. Health Perspect.* **109**, 687–693.
30. Sambrook, J., Fritsch, E. F. & Maniatis, T. (1989) *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Lab. Press, Plainview, NY).
31. Metsä-Ketelä, M., Salo, V., Halo, L., Hautala, A., Hakala, P. M. & Ylihonko, K. (1999) *FEMS Microbiol. Lett.* **180**, 1–6.
32. Brown, D. W., Adams, T. H. & Keller, N. P. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 14873–14877.
33. Marahiel, M. A., Stachelhause, T. & Mootz, H. D. (1997) *Chem. Rev.* **97**, 2651–2673.
34. Neilan, B. A., Dittmann, E., Rouhiainen, L., Bass, R. A., Schaub, V., Sivonen, K. & Börner, T. (1999) *J. Bacteriol.* **181**, 4089–4097.
35. Rowan, R. & Powers, D. A. (1992) *Proc. Natl. Acad. Sci. USA* **89**, 3639–3643.
36. Brown, D. W., Yu, J. H., Kelkar, H. S., Fernandes, M., Nesbitt, T. C., Keller, N. P., Adams, T. H. & Leonard, T. J. (1996) *Proc. Natl. Acad. Sci. USA* **93**, 1418–1422.
37. Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990) *J. Mol. Biol.* **215**, 403–410.
38. Glasgow, H. B., Jr., Burkholder, J. M., Mallin, M. A., Deamer-Melia, N. J. & Reed, R. E. (2001) *Environ. Health Perspect.* **109**, 715–730.
39. Melo, A. C., Moeller, P. D. R., Glasgow, H. B., Jr., Burkholder, J. M. & Ramsdell, J. S. (2001) *Environ. Health Perspect.* **109**, Suppl. 5, 731–737.
40. Glasgow, H. B., Jr., Burkholder, J. M., Morton, S. L. & Springer, J. (2001) *Phycologia* **40**, 234–245.
41. Moeller, P. D. R., Morton, S. L., Mitchell, B. A., Sivertsen, S. K., Fairey, E. R., Mikulski, T. M., Glasgow, H. B., Jr., Deamer-Melia, N. J., Burkholder, J. M. & Ramsdell, J. S. (2001) *Environ. Health Perspect.* **109**, 739–743.
42. Hall, S. & Strichartz, G. (1990) *Marine Toxins: Origin, Structure, and Molecular Pharmacology* (Am. Chem. Soc., Washington, DC).
43. Yasumoto, T., Oshima, Y. & Fukuyo, Y. (1996) *Harmful and Toxic Algal Blooms* (Intergovernmental Oceanographic Commission of UNESCO, New York).
44. Cooksey, K. E. (1998) *Molecular Approaches to the Study of the Ocean* (Chapman & Hall, London).
45. Rein, K. & Borrone, J. (1999) *Comp. Biochem. Physiol.* **124**, 117–131.
46. Igarashi, T., Satake, M. & Yasumoto, T. (1999) *J. Am. Chem. Soc.* **121**, 8499–8511.
47. Zhu, G., Marchewka, M. J., Woods, K. M., Upton, S. J. & Keithly, J. S. (2000) *Mol. Biochem. Parasitol.* **105**, 253–260.
48. Cavalier-Smith, T. (1993) *Microbiol. Rev.* **57**, 953–994.
49. Metz, J. G., Roessler, P., Facciotti, D., Levering, C., Dittrich, F., Lassner, M., Valentine, R., Lardizabal, K., Domergue, F., Yamada, A., Yazawa, K., Knauf, V. & Browse, J. (2001) *Science* **293**, 290–293.
50. Kiryu, Y., Shields, J. D., Vogelbein, W. K., Zwerner, D. E., Kator, H. & Blazer, V. S. (2002) *J. Aquat. Anim. Health* **14**, 11–24.
51. Steidinger, K. A., Burkholder, J. M., Glasgow, H. B., Jr., Ruby, E., Garrett, J., Noga, E. J. & Smith, S. A. (1996) *J. Phycol.* **32**, 157–164.
52. Litaker, R. W., Vandersea, M. W., Kibler, S. R., Madden, V. J., Noga, E. J. & Tester, P. A. (2002) *J. Phycol.* **38**, 442–463.
53. Dykstra, M. J. & Kane, A. S. (2000) *J. Aquat. Anim. Health* **12**, 18–25.
54. Lewitus, A. J., Glasgow, H. B., Jr. & Burkholder, J. M. (1999) *J. Phycol.* **35**, 303–312.
55. Alavi, M., Miller, T., Erlanson, K., Schneider, R. & Belas, R. (2001) *Environ. Microbiol.* **3**, 380–396.
56. Bourdelais, A. J., Tomas, C. R., Naar, J., Kubanek, J. & Baden, D. G. (2002) *Environ. Health Perspect.* **110**, 465–470.
57. Paerl, H. W., Bales, J. D., Ausley, L. W., Buzzelli, C. P., Crowder, L. B., Eby, L. A., Fear, J. M., Go, M., Peierls, B. L., Richardson, T. L. & Ramus, J. S. (2001) *Proc. Natl. Acad. Sci.* **98**, 5655–5660.
58. Baden, D. G., Fleming, L. E. & Bean, J. A. (1995) in *Handbook of Clinical Neurology: Intoxications of the Nervous System, Part III*, ed. de Wolff, F. A. (Elsevier, Amsterdam), Vol. 21, pp. 141–175.
59. Gawley, R. E., Rein, K. S., Kinoshita, M. & Baden, D. G. (1992) *Toxicol.* **30**, 780–785.
60. Gawley, R. E., Rein, K. S., Jeglitsch, G., Adams, D. J., Theodorakis, E. A., Tiebes, J., Nicolaou, K. C. & Baden, D. G. (1995) *Chem. Biol.* **2**, 533–541.
61. Jeglitsch, G., Rein, K. S., Baden, D. G. & Adams, D. J. (1998) *J. Pharmacol. Exp. Ther.* **284**, 516–525.
62. Poli, M. A., Mende, T. J. & Baden, D. G. (1986) *Mol. Pharmacol.* **30**, 129–135.
63. Purkerson, S. L., Baden, D. G. & Fieber, L. A. (1999) *Neurotoxicology* **20**, 909–920.
64. Rein, K. S., Lynn, B., Gawley, R. E. & Baden, D. G. (1994) *J. Org. Chem.* **59**, 2107–2113.
65. Moe, C. L., Turf, E., Oldach, D., Bell, P., Hutton, S., Savitz, D., Koltai, D., Turf, M., Ingrisawang, L., Hart, R., et al. (2001) *Environ. Health Perspect.* **109**, 781–786.
66. Morris, J. G., Jr. (2001) *Environ. Health Perspect.* **109**, 787–790.
67. Shoemaker, R. C. (2001) *Environ. Health Perspect.* **109**, 791–796.
68. Backer, L. C., Niskar, A. S., Rubin, C., Blindauer, K., Christianson, D., Naeher, L. & Rogers, H. S. (2001) *Environ. Health Perspect.* **109**, 797–801.