

12-1986

End-Products Of Anaerobic Chitin Degradation By Salt-Marsh Bacteria As Substrates For Dissimilatory Sulfate Reduction And Methanogenesis

JN Boyer

Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Environmental Microbiology and Microbial Ecology Commons](#), and the [Marine Biology Commons](#)

Recommended Citation

Boyer, JN, End-Products Of Anaerobic Chitin Degradation By Salt-Marsh Bacteria As Substrates For Dissimilatory Sulfate Reduction And Methanogenesis (1986). *Applied and Environmental Microbiology*, 52(6), 1415-1418.

<https://scholarworks.wm.edu/vimsarticles/1384>

This Article is brought to you for free and open access by the Virginia Institute of Marine Science at W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

NOTES

End Products of Anaerobic Chitin Degradation by Salt Marsh Bacteria as Substrates for Dissimilatory Sulfate Reduction and Methanogenesis†

JOSEPH N. BOYER

*Virginia Institute of Marine Science, School of Marine Science, College of William and Mary,
Gloucester Point, Virginia 23062*

Received 23 July 1986/Accepted 8 September 1986

The anaerobic pathway of chitin decomposition by chitinoclastic bacteria was examined with an emphasis on end product coupling to other salt marsh bacteria. Actively growing chitinoclastic bacterial isolates produced primarily acetate, H₂, and CO₂ in broth culture. No sulfate-reducing or methanogenic isolates grew on chitin as sole carbon source or produced any measurable degradation products. Mixed cultures of chitin degraders with sulfate reducers resulted in positive sulfide production. Mixed cultures of chitin-degrading isolates with methanogens resulted in the production of CH₄ with reductions in headspace CO₂ and H₂. The combination of all three metabolic types resulted in the simultaneous production of methane and sulfide, with more methane being produced in mixed cultures containing CO₂-reducing methanogens and acetoclastic sulfate reducers because of less interspecific H₂ competition.

Chitin and cellulose are the major structural polymers in invertebrates and plants, respectively. Both are polysaccharides composed of β {1-4}-linked monomers of *N*-acetyl-D-glucosamine in chitin or glucose in cellulose. The biosynthesis of chitin and cellulose proceeds via parallel enzymatic pathways (13), as does aerobic degradation by bacteria (4, 12). Billions of tons of chitin are produced annually in the oceans (5). It is not known how much of this production is incorporated into anaerobic habitats, but the amount could conceivably be substantial because its particulate nature favors rapid sedimentation and it occurs in many benthic organisms (15).

The occurrence of anaerobic chitin degradation has been largely ignored in the literature. A study by Alshina (1) showed that the amendment of chitin to anoxic sediments stimulated sulfate reduction, a result which led the author to conclude that sulfate-reducing bacteria were responsible for anaerobic chitin cycling. Billy (3) and Timmis et al. (14) have shown that certain clostridia can colonize and use chitin as a carbon source. In contrast, the amount of information concerning anaerobic cellulolysis is immense, mostly from work with the rumen system (9).

Considering the similar polymeric structures of chitin and cellulose, it was hypothesized that under anoxic conditions (i) chitin would be broken down by anaerobic sediment bacteria with an enzyme system similar to that of cellulolytic organisms, (ii) the products of chitin fermentation would be comparable to those of cellulolysis, and (iii) the fermentation products could be coupled via interspecies transfer to both sulfate reduction and methanogenesis.

Sediment samples from the Great Sippewissett Salt Marsh, Mass., were taken with core tubes (5-cm inner diameter). The anoxic portions of the cores (5 to 10 cm below the surface) were extruded into Whirl-Pac bags (American Sci-

entific Products, McGaw Park, Ill.). Air was expressed from the bags before they were sealed. Upon return, the cores were pooled and homogenized by being mixed in a covered beaker under a constant stream of N₂. Samples were streaked onto bottle plates (6) containing prerduced medium selective for the isolation of anaerobic chitinolytic, sulfate-reducing, and methanogenic bacteria. Medium components used for the selective isolation of salt marsh bacteria are listed in Table 1. Chitin degrader medium consisted of a heterotrophic marine agar base with an agar overlay of finely ground reprecipitated chitin (Calbiochem-Behring, San Diego, Calif.) as sole carbon source (8). Sulfate-reducing bacteria were isolated on a minimal salts medium containing sodium sulfate with acetate instead of lactate used as the carbon source (10) under an atmosphere of 80% H₂-20% CO₂. Ferrous iron was incorporated into the medium as a sulfide indicator. Methanogen medium was modified from that of Balch et al. (2) with acetate and CO₂ as carbon sources. Resazurin was incorporated into all media as an E_h indicator.

Plates were inoculated at 30°C for 5 days. Zones of clearing around colonies on chitin agar indicated the presence of extracellular chitinase. Formation of black colonies caused by precipitation of FeS on minimal salts medium was considered positive sulfate reduction. Methane production in the headspace was determined by a Carle gas chromatograph equipped with a thermal conductivity detector with a Porapak column (Supelco, Bellefonte, Pa.). Anaerobic transfers were performed until pure cultures were obtained as confirmed by light microscopy of gram-stained slides.

Axenic cultures of the bacterial types were inoculated into Balch tubes (2) containing chitin coculture medium, a mineral salts broth with reprecipitated chitin as sole carbon source and kept under N₂ (Table 1). The purpose was to ascertain whether sulfate reducers or methanogens could degrade chitin. Growth was defined as the presence of CO₂

† Virginia Institute of Marine Science, contribution 1335.

TABLE 1. Medium composition

Medium component	Amt (per liter of distilled water)				
	Chitin degrader		Sulfate reducer	Methanogen	Chitin coculture
	Base	Overlay			
K ₂ HPO ₄	0.25 g		0.5 g		
NH ₄ Cl			1.0 g		
CaCl ₂ · 2H ₂ O			0.1 g		
Mg ₂ SO ₄ · 6H ₂ O			2.0 g	3.0 g	
NaCl	30.0 g	18.0 g	30.0 g	25.0 g	18.0 g
NaSO ₄			1.0 g		
NaHCO ₃		3.0 g	3.0 g	2.0 g	
Fe(NH ₄) ₂ (SO ₄) ₂ · 7H ₂ O	0.04 g	0.04 g		0.04 g	0.04 g
FeSO ₄			0.1 g		0.1 g
1 N HCl				9.0 g	
Mineral 2 ^a				40.0 ml	
Mineral 3 ^b		500.0 ml			500.0 ml
Trace minerals ^c		10.0 ml	10.0 ml	10.0 ml	10.0 ml
Trace vitamins ^d		10.0 ml	10.0 ml	10.0 ml	10.0 ml
Yeast extract ^e	0.5 g	0.2 g	0.2 g	0.2 g	
Peptone ^f	1.0 g				
Sodium glycerophosphate	0.1 g				
Acetate			2.0 g	2.0 g	
Chitin		4.0 g			4.0 g
Cysteine-sodium sulfate (1:1)	0.5 g	0.5 g		0.5 g	
Ascorbic acid			0.02 g		0.02 g
Sodium thioglycolate			0.02 g		0.02 g
0.1% Resazurin	1.0 ml	1.0 ml	1.0 ml	1.0 ml	1.0 ml
Agar	15.0 g	10.0 g	15.0 g	15.0 g	
N ₂	+ ^g	+			+
80% H ₂ -20% CO ₂			+	+	

^a Grams per liter of distilled water: KH₂PO₄, 6; (NH₄)₂SO₄, 6; NaCl, 12; MgSO₄ · 7H₂O, 2.6; CaCl₂ · 2H₂O, 0.16.

^b Grams per liter of distilled water: KCl, 0.67; MgCl₂ · 2H₂O, 5.5; MgSO₄ · 7H₂O, 6.9; NH₄Cl, 0.5; CaCl₂ · 2H₂O, 0.28; K₂HPO₄, 0.28.

^c Grams per liter of distilled water: nitroloacetic acid, 1.5; MgSO₄ · 7H₂O, 3.0; MnSO₄ · 2H₂O, 0.5; NaCl, 1.0; FeSO₄ · 7H₂O, 0.1; CoSO₄, 0.1; CaCl₂ · 2H₂O, 0.1; ZnSO₄, 0.1; CuSO₄ · 5H₂O, 0.01; AlK(SO₄)₂, 0.01; H₃BO₃, 0.01; Na₂MoO₄ · 2H₂O, 0.01 (pH to 7.0 with KOH).

^d Grams per liter of distilled water: biotin, 0.002; folic acid, 0.002; pyridoxine hydrochloride, 0.01; thiamine hydrochloride, 0.005; riboflavin, 0.005; nicotinic acid, 0.005; DL-calcium pantothenate, 0.005; vitamin B₁₂, 0.0001; *p*-aminobenzoic acid, 0.005; lipoic acid, 0.005.

^e Difco Laboratories, Detroit, Mich.

^f BBL Microbiology Systems, Cockeysville, Md.

^g +, Present.

or CH₄ in the headspace, blackening of the medium, or formation of volatile fatty acids, particularly acetate. Volatile fatty acids were extracted from the medium with ether (7) and separated on a Resoflex column (Burrell Corp., Pittsburgh, Pa.) by a Carle gas chromatograph equipped with a flame ionization detector.

A final experiment was performed to evaluate whether sulfate reducers or methanogens could grow on the fermentative products of chitin degradation. Various combinations of culture types were inoculated into chitin coculture medium, and growth was ascertained as previously described.

A total of 17 anaerobic chitinoclastic colony types were observed (Table 2). Eight types were obligate anaerobes, and nine types were facultative. Five of the obligately anaerobic chitinoclastic isolates were gram-positive motile rods, while the other three were gram-negative motile rods. Of the nine facultative isolates, all were motile and gram negative. One of the three slender rods with gliding motility was pigmented. The other six were motile by flagella. Morphologically, these ranged from coccoid to curved and straight rods. Colonies of coccoid cells were luminescent when grown aerobically. All isolates produced some CO₂, H₂, and various types and amounts of volatile fatty acids.

Five distinct colonies of sulfate reducers were isolated on acetate. All colonies were made up of obligately anaerobic, motile rods that reduced Fe³⁺ to FeS. The four methano-

genic isolates were distinctly different in their morphology, motility, and gram-staining characteristics, but all grew on acetate under H₂-CO₂ and produced methane. None of the sulfate-reducing or methanogenic isolates grew on chitin as sole carbon source or produced any measurable degradation products.

Results of mixed culture experiments with various combinations of chitinoclasts, sulfate reducers, and methanogens were similar. The most metabolically active cocultures (Table 3) are representative of the entire data set. As previously mentioned, the chitinoclastic bacteria degraded chitin and produced CO₂, H₂, and acetate. In some cases, a series of volatile fatty acids was produced, including propionate and butyrate.

Combinations of chitinoclasts with sulfate reducers showed two distinct pathways of sulfate reduction, i.e., an acetoclastic route, by which acetate was oxidized to CO₂ and sulfide was produced, and an H₂-utilizing route, by which sulfate was directly reduced. These results are consistent with the findings of others (11). Because sulfide was only qualitatively determined, the relative magnitude of sulfate reduction by each isolate was unknown. Cocultures of methanogens and chitinoclasts resulted in the production of CH₄, with decreases in H₂ and CO₂. Unfortunately, none of the methanogenic isolates could ferment acetate to CH₄.

Cocultures of chitin degraders with both sulfate reducers and methanogens showed the ability of methanogens to

TABLE 2. Sediment isolates on agar plates

Type and name of isolate	Characteristic of isolate					
	Obligate anaerobe	Gram stain	Motility	Morphology	Pigment	Luminescence
Chitin degrading						
CD1	+	+	+	Rod	—	—
CD2	+	+	+	Rod	—	—
CD3	—	—	+	Curved rod	—	—
CD4	+	+	+	Rod	—	—
CD5	—	—	Gliding	Rod	—	—
CD6	—	—	+	Cocoid	—	+
CD7	+	—	+	Rod	—	—
CD8	+	+	+	Rod	—	—
CD9	+	—	+	Rod	—	—
CD10	—	—	Gliding	Rod	Yellow	—
CD11	—	—	+	Curved rod	—	—
CD12	+	+	+	Rod	—	—
CD13	—	—	+	Rod	—	—
CD14	—	—	+	Cocoid	—	+
CD15	—	—	Gliding	Rod	—	—
CD16	+	—	+	Rod	—	—
CD17	—	—	+	Curved rod	—	—
Sulfate reducing						
SR1	+	—	+	Curved rod	—	NA ^a
SR2	+	—	+	Curved rod	—	NA
SR3	+	+	+	Rod	—	NA
SR4	+	—	+	Rod	—	NA
SR5	+	—	+	Curved rod	—	NA
Methanogenic						
M1	+	+	+	Cocoid	—	NA
M2	+	+	—	Cocoid	—	NA
M3	+	+	+	Rod	—	NA
M4	+	—	+	Rod	—	NA

^a NA, Not applicable.

compete for the limiting resource, hydrogen. An H₂-utilizing methanogen in coculture with an acetoclastic sulfate reducer produced more methane than the same methanogen cultured with an H₂-utilizing sulfate reducer. Because no acetoclastic methanogens were isolated, observations on interspecific competition between acetoclastic sulfate reducers and methanogens were not possible.

It must be emphasized that this study was not designed to taxonomically characterize the bacteria involved in anaerobic chitin decomposition, but to determine which metabolic types are important. Clearly, there exists in anoxic salt marsh sediments a consortium of chitin-fermenting bacteria whose end products are coupled to both sulfate reduction and methanogenesis in pure culture. The relative contribution of these metabolic types to the total flux

of carbon from chitin in anoxic sediments awaits further study.

I thank the teaching staff of the Microbial Ecology course at the Marine Biology Laboratory, Woods Hole, Mass., and especially Ralph S. Wolfe for his invaluable discussions on theory and practice of anaerobic microbiology.

ADDENDUM IN PROOF

After this paper had been accepted, I became aware of a recent publication by Roel Pel and Jan C. Gottschal, *FEMS Microb. Ecol.* **38**:39–49, 1986. This article describes cultivation and characterization of eight strains of obligately anaerobic chitinolytic bacteria and the fermentation products resulting from their growth: acetate, ethanol, formate, hydrogen, carbon dioxide, and ammonium.

LITERATURE CITED

1. **Alshina, V. I.** 1938. Destruction of chitin by sulfate-reducing bacteria and changes in the oxidation-reduction conditions in the reduction of sulfate. *Microbiol. SSSR* **7**:850–859.
2. **Balch, W. E., G. E. Fox, L. J. Magrum, C. R. Woese, and R. S. Wolfe.** 1979. Methanogens: reevaluation of a unique biological group. *Microbiol. Rev.* **43**:260–296.
3. **Billy, C.** 1968. A chitinolytic anaerobic bacterium *Clostridium chitinophilum*. *C. R. Acad. Sci. Ser. D.* **266**:1535–1536.
4. **Burns, R. G.** 1982. Carbon mineralization by mixed cultures, p. 475–543. *In* A. T. Bull and J. H. Slater (ed.), *Microbial interactions and communities*, vol. 1. Academic Press, Inc., New York.
5. **Goodrich, T. D., and R. Y. Morita.** 1977. Incidence and estima-

TABLE 3. Coculture experimental results

Product	Amt produced (mmol/100 mmol of chitin)					
	CD3	CD3 cocultured with:				
		SR4	SR1	M2	M2 + SR4	M2 + SR1
Acetate	47.9	35.2	44.6	51.3	31.4	46.1
H ₂	10.9	9.1	ND ^a	ND	ND	ND
CO ₂	11.7	27.2	ND	ND	12.4	ND
CH ₄	ND	ND	ND	20.6	15.4	2.1
S ²⁻	— ^b	+	+	—	+	+

^a ND, Not detectable.

^b —, Not present; +, present.

- tion of chitinase activity associated with marine fish and other estuarine samples. *Mar. Biol.* **41**:349-353.
6. **Hermann, M., K. M. Noll, and R. S. Wolfe.** 1986. Improved agar bottle plate for isolation of methanogens or other anaerobes in a defined gas atmosphere. *Appl. Environ. Microbiol.* **51**:1124-1126.
 7. **Holdeman, L. V., and W. E. C. Moore (ed).** 1973. The anaerobe laboratory manual, 2nd ed. Virginia Polytechnic Institute and State University, Blacksburg.
 8. **Hsu, S. C., and J. L. Lockwood.** 1975. Powdered chitin agar as a selective medium for enumeration of actinomycetes in water and soil. *Appl. Microbiol.* **29**:422-426.
 9. **Hungate, R. E.** 1985. Anaerobic transformations of organic matter, p. 39-95. *In* E. R. Leadbetter and J. S. Poindexter (ed.), *Bacteria in nature*, vol. 1. Bacterial activities in perspective. Plenum Publishing Corp., New York.
 10. **Lapage, S. P., J. E. Shelton, T. G. Mitchell.** 1970. Media for the maintenance and preservation of bacteria, p. 120. *In* J. R. Norris and D. W. Ribbons (ed.), *Methods in microbiology*, vol. 3A. Academic Press, Inc., New York.
 11. **Nedwell, D. B.** 1982. The cycling of sulphur in marine and freshwater sediments, p. 73-106. *In* D. B. Nedwell and C. M. Brown (ed.), *Sediment microbiology*. Academic Press, Inc., New York.
 12. **Okafor, N.** 1966. Estimation of the decomposition of chitin in soil by the method of carbon dioxide release. *Soil Sci.* **102**:140-142.
 13. **Pahlic, M., and J. R. Stevenson.** 1978. Glucosamine-6-phosphate synthesis in the crayfish epidermis. *Comp. Biochem. Physiol.* **60B**:281-285.
 14. **Timmis, K., G. Hobbs, and R. C. W. Berkely.** 1974. Chitinolytic clostridia isolated from marine mud. *Can. J. Microbiol.* **20**:1284-1285.
 15. **ZoBell, C. E., and S. C. Rittenberg.** 1938. The occurrence and characteristics of chitinoclastic bacteria in the sea. *J. Bacteriol.* **35**:275-287.