

2017

Interactive effects of elevated temperature and CO₂ on nitrate, urea, and dissolved inorganic carbon uptake by a coastal California, USA, microbial community

JL Spackeen
Virginia Institute of Marine Science

RE Sipler
Virginia Institute of Marine Science

K Xu

et al

DA Bronk
Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Marine Biology Commons](#), and the [Oceanography Commons](#)

Recommended Citation

Spackeen, JL; Sipler, RE; Xu, K; al, et; and Bronk, DA, Interactive effects of elevated temperature and CO₂ on nitrate, urea, and dissolved inorganic carbon uptake by a coastal California, USA, microbial community (2017). *Marine Ecology Progress Series*, 577, 49-65.
10.3354/meps12243

This Article is brought to you for free and open access by the Virginia Institute of Marine Science at W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Interactive effects of elevated temperature and CO₂ on nitrate, urea, and dissolved inorganic carbon uptake by a coastal California, USA, microbial community

Jenna L. Spackeen^{1,*}, Rachel E. Sipler¹, Kai Xu², Avery O. Tatters²,
Nathan G. Walworth², Erin M. Bertrand^{3,4,5}, Jeffrey B. McQuaid^{4,5},
David A. Hutchins², Andrew E. Allen^{4,5}, Deborah A. Bronk¹

¹Virginia Institute of Marine Science, College of William & Mary, PO Box 1346, Gloucester Point, VA 23062, USA

²The University of Southern California, Department of Biological Sciences, 3616 Trousdale Parkway, Los Angeles, CA 90089, USA

³Department of Biology, Dalhousie University, Halifax, NS B3H 4R2, Canada

⁴Microbial and Environmental Genomics, J. Craig Venter Institute, La Jolla, CA 92037, USA

⁵Integrative Oceanography Division, Scripps Institution of Oceanography, UC San Diego, La Jolla, CA 92037, USA

ABSTRACT: Average global temperatures and carbon dioxide (CO₂) levels are expected to increase in the coming decades. Implications for ocean ecosystems include shifts in microbial community structure and subsequent modifications to nutrient pathways. Studying how predicted future temperature and CO₂ conditions will impact the biogeochemistry of the ocean is important because of the ocean's role in regulating global climate. We determined how elevated temperature and CO₂ affect uptake rates of nitrate, urea, and dissolved inorganic carbon (DIC) by 2 size classes (0.7–5.0 and >5.0 μm) of a microbial assemblage collected from coastal California, USA. This microbial community was incubated for 10 d using an ecostat continuous culture system that supplied the microorganisms with either nitrate or urea as the dominant nitrogen source. Biomass parameters, nutrient concentrations, and uptake rates were measured throughout the experiment. In all treatments, urea uptake rates were greater than nitrate, and larger microorganisms had higher uptake rates than smaller microorganisms. Uptake rates of urea and DIC within both size fractions were higher at elevated temperature, and uptake rates of nitrate by smaller microorganisms increased with elevated CO₂. These findings suggest that the rate at which nutrients cycle in temperate coastal waters will increase as temperature and CO₂ levels rise and that the effect will vary between nitrogen substrates and different microorganisms.

KEY WORDS: Nitrate · Urea · DIC · Uptake · Temperature · CO₂ · Southern California Bight · Microbial communities

—Resale or republication not permitted without written consent of the publisher—

INTRODUCTION

Global temperatures and atmospheric concentrations of carbon dioxide (CO₂) are increasing at unprecedented rates compared to Earth's geologic past (Beardall et al. 2009), with the rise predicted to continue in the future (Cubasch & Wuebbles 2014). Depending on the model, average global surface temperature is currently predicted to increase by 1.4 to 5.8°C

over the next century (IPCC 2014), and atmospheric CO₂ levels are projected to reach 750 to 1000 ppm by 2100 (Meehl et al. 2007). Considering the magnitude of the predicted change, it is crucial to understand how marine microorganisms will respond to increasing temperature and CO₂ at both physiological and ecological levels, because they form the base of the marine food web and largely control the cycling of nutrients throughout the ocean (Boyd & Hutchins 2012).

*Corresponding author: j.spackeen@gmail.com

§Corrections were made after publication. For details see www.int-res.com/articles/meps2017/580/m580p247.pdf
This version: September 29, 2017

© Inter-Research 2017 · www.int-res.com

Temperature changes will affect the physiology of microorganisms, as well as their global distribution. For example, microorganisms capable of fixing nitrogen typically live in warmer oceans, and their distributions may expand as the temperature of the surface water increases (Boyd & Doney 2002). Additionally, the supply of nutrients may be affected by changes to ocean currents. For example, model studies have found that nitrate (NO_3^-) supply to the coastal California (USA) ecosystem has increased since 1980 (Jacox et al. 2015); in the model, NO_3^- input to the euphotic zone was projected to increase by 80% by the year 2100 due to an increased flux of NO_3^- from deep waters (Rykaczewski & Dunne 2010). Such a large increase in NO_3^- supply has the potential to influence the community composition of microorganisms and their physiology.

In the case of CO_2 , the large increase in atmospheric CO_2 concentrations translates to a substantial change in the availability of dissolved inorganic carbon (DIC) in the ocean. The present-day concentration of dissolved CO_2 in the ocean is $\sim 13 \mu\text{M}$, and as the ocean becomes more CO_2 -saturated, concentrations are expected to more than double, causing the pH of the ocean to drop from 8.07 to 7.77 (Raven et al. 2005, Beardall et al. 2009). We know that microorganisms can have differing levels of sensitivity to CO_2 and ocean acidification (Riebesell 2004, Doney et al. 2009). It is well documented that calcifying phytoplankton, like coccolithophores, are sensitive to ocean acidification because their liths start dissolving with declining pH (Riebesell et al. 2000, Kroeker et al. 2013), although non-calcifying microorganisms may also be affected by ocean acidification (Tortell et al. 2002, 2008, Rost et al. 2008, Wu et al. 2010). How phytoplankton species respond to ocean acidification is likely affected by whether they have carbon-concentrating mechanisms (CCMs), which help them concentrate inorganic carbon into their cells. As the partial pressure of ambient CO_2 increases, it has been suggested that there will be less of a need for CCMs (Tortell et al. 2008) and that microorganisms will be able to allocate the energy and resources required to maintain CCMs toward other physiological processes (Raven 1991). Likewise, microorganisms that obtain CO_2 solely through passive diffusion may also benefit as DIC concentrations increase (Beardall et al. 2009).

If microorganisms are able to take up DIC more effectively, growth rates and the demand for other nutrients, such as nitrogen and phosphorus, will likely increase. The source of these nutrients varies

from system to system. In coastal systems, nutrient supply is influenced by terrigenous sources, including riverine discharge or runoff that contains nutrients from wastewater treatment plants and agricultural production. In the past few decades, the use of fertilizers made with organic forms of nitrogen, primarily urea, has increased more than 100-fold (Glibert et al. 2006). The delivery of excess nutrients to the coastal environment is one factor that has the potential to increase the prevalence of harmful algal blooms (HABs).

In the last few decades, HABs have become a recurring issue off the coast of California (Schnitzer et al. 2007, Lewitus et al. 2012), and within the Southern California Bight region, nutrients supplied from natural sources (upwelling) and anthropogenic sources, namely treated wastewater effluent, can be on the same order of magnitude (Howard et al. 2014). One frequent bloomer in the Southern California Bight is *Pseudo-nitzschia* spp., a genus capable of producing domoic acid, the toxin responsible for amnesic shellfish poisoning (Bates et al. 1989, Bates & Trainer 2006). The response of *Pseudo-nitzschia* spp. to organic forms of nitrogen, such as urea, is particularly important to consider, as urea can sustain blooms of this species when other sources of nitrogen are in low supply (Cochlan et al. 2008), and certain strains produce more domoic acid when grown on urea than when grown on other nitrogen sources (Howard et al. 2007, Auro & Cochlan 2013, Martin-Jézéquel et al. 2015). Additional and different nutrient substrates, coupled with high temperatures, may create ideal conditions for blooms of *Pseudo-nitzschia*. For example, in 2014 and 2015, the North Pacific, including waters off the coast of southern California, was characterized by unusually warm sea surface temperatures that were as much as 3°C higher than average. The anomalously warm water moved into coastal waters, resulting in the largest and most prolonged bloom of *Pseudo-nitzschia* spp. along the US west coast within the past 15 yr (Bond et al. 2015, Du et al. 2016, McCabe et al. 2016).

As sea surface temperatures and CO_2 concentrations continue to rise, it is likely that nutrient regimes will shift, new microbial communities will become established (Sarmiento et al. 2010), and rates of nitrogen and carbon uptake will be altered. In this study, a factorial design was used to assess the individual and combined effects of increases in temperature and CO_2 on the rate of NO_3^- , urea, and DIC uptake by a Southern California Bight microbial community.

MATERIALS AND METHODS

Sample collection

Seawater was collected from Fish Harbor at the Southern California Marine Institute's (SCMI) dock in Terminal Island, California (33° 44' 59" N, 118° 12' 54" W) on 10 May 2012. Seawater was drawn from just below the surface using a manual suction pump and collected in acid washed (10% HCl) cubitainers that were pre-rinsed with seawater. Collected seawater was transported directly to the University of Southern California (USC), where it was immediately dispensed into 2.7 l polycarbonate bottles and placed within 2 ecostat continuous culture systems. Diluent (i.e. seawater amended with either NO₃⁻ or urea) was continually added to the ecostats over the duration of the 10 d experiment. Samples for this study were collected at 48 h intervals to determine the rate of uptake of 2 stable isotopically labeled nitrogen substrates, NO₃⁻ and urea, as well as labeled HCO₃⁻ (referred to as DIC uptake henceforth) by the microbial community. Additional samples were collected at the start and end of the experiment so that changes in nutrient pools could be assessed.

Ecostat design

Ecostats are continuous culture systems that create a controlled experimental setting to culture microorganisms over extended periods of time (Hutchins et al. 2003, Hare et al. 2007, Feng et al. 2009, 2010). Continuous culturing is advantageous because nutrients are delivered at a constant rate, there is quantitative removal of biomass, and the response of microbial communities to changing variables can be examined over extended periods of time (i.e. weeks to months; Hutchins et al. 2003, Pickell et al. 2009). As a result, the microbial community has sufficient time to acclimate to experimental conditions and competitive displacement will occur, meaning that microorganisms better suited to the experimental conditions will thrive, while those less adapted will have a lower abundance or be lost. One difference between nature and ecostats, however, is that loss processes, such as grazing, can be selective for certain species while cells lost from the ecostat are removed indiscriminately.

In this experiment, temperature, CO₂ level, and nutrient inputs were manipulated using 2 ecostat systems, 1 for each temperature level, each of which contained 12 individual bottles. The ecostats were set to

test present-day (19°C and 380 ppm CO₂) and potential future temperature (23°C) and CO₂ (800 ppm) conditions. Control (19°C, 380 ppm CO₂), +CO₂ (19°C, 800 ppm CO₂), +Temp (23°C, 380 ppm CO₂), and Combined (23°C, 800 ppm CO₂) are the terms that will be used to describe the treatments throughout the manuscript.

Each of the treatment bottles in the Control, +CO₂, +Temp, and Combined factorial matrix received either NO₃⁻ or urea as the primary nitrogen source. Seawater used for diluent was collected from Redondo Beach Harbor on 9 May 2012, filtered using an acid-washed, in-line 0.2 µm cartridge filter, and placed in two 50 l reservoirs. The second collection of water that occurred on 10 May 2012 was necessary because biomass was low in the Redondo Beach Harbor water and could not be used as the inoculant. NO₃⁻ was added to 1 reservoir to a final concentration of 15.1 ± 0.6 (SD) µmol N l⁻¹, and urea was added to the other to a final concentration of 11.1 ± 0.4 µmol N l⁻¹ (Table 1). We note that, while the diluent contained these elevated concentrations of NO₃⁻ and urea, the cells within the ecostats were never exposed to concentrations this high, because the diluent was added slowly over time and the cells were taking the substrates up as the diluent was added. Both diluent reservoirs were amended with silicic acid (H₄SiO₄) and phosphate (PO₄³⁻) to a final concentration of ~42 µmol Si l⁻¹ and ~2 µmol P l⁻¹ (Table 1). The concentration of ammonium (NH₄⁺) in both diluent types was ~2 µmol N l⁻¹; we note that this concentration is ~1 µmol N l⁻¹ higher than the site water (Table 1), and may be due to contamination that occurred during the experimental set-up. Diluent from each reservoir was supplied to triplicate incubation bottles for each treatment using Teflon lines and adjustable peristaltic pumps. The Teflon lines entered the incubation bottles through the top and extended to the bottom where the diluent was released. Temperature levels were maintained using thermostatically controlled, recirculating heater/chiller systems set to either 19 or 23°C (Hare et al. 2007, Feng et al. 2009, 2010). The CO₂ concentrations within the +CO₂ and Combined treatments were manipulated using commercially prepared CO₂/air mixtures (Praxair) bubbled into the incubation bottles through Teflon tubes.

The ecostats were positioned on the roof of the Allan Hancock Foundation building so that they would be exposed to ambient light conditions. The incubation bottles were secured to a plexiglass rack designed to automatically flip on its side every 5 min. This motion re-suspends the cells that may have set-

Table 1. Nutrient concentrations of the seawater collected off the Southern California Marine Institute (SCMI, the starting community) dock and in Redondo Harbor (diluent-base), as well as the 2 diluent types (NO_3^- - and urea-based). Nutrient concentrations measured on the final day of the experiment for each treatment (Treatment Tfinals) are displayed below the respective diluent type that the treatment received. Treatment details are provided in the 'Materials and methods'. Concentrations are in units of $\mu\text{mol C, N, P, or Si l}^{-1} \pm 1 \text{ SD}$ ($n = 3$). DIC (DOC): dissolved inorganic (organic) carbon; DON: dissolved organic carbon; DPA: dissolved organic nitrogen; DPA: dissolved organic nitrogen; DPA: dissolved organic nitrogen; DPA: dissolved organic nitrogen; DPA: dissolved organic nitrogen; NM: not measured; BD: below detection

Source	DIC	DOC	NH_4^+	NO_2^-	NO_3^-	DON	Urea	DPA	PO_4^{3-}	H_4SiO_4	NH_4^+	NO_3^-	% Urea	% DON
SCMI	NM	129 ± 4	0.93 ± 0.0	0.04 ± 0.0	0.51 ± 0.0	6.37 ± 0.6	0.84 ± 0.5	0.21 ± 0.0	0.41 ± 0.0	3.0 ± 0.0	11.9	6.5	10.7	81.2
Redondo	NM	115 ± 2	0.85 ± 0.1	0.19 ± 0.0	4.30 ± 0.0	5.81 ± 0.2	1.44 ± 0.0	0.10 ± 0.0	0.83 ± 0.0	10.4 ± 0.0	7.6	38.6	12.9	52.1
NO_3^- diluent	NM	116 ± 1	1.89 ± 0.9	0.19 ± 0.0	15.10 ± 0.6	5.67 ± 2.3	1.22 ± 0.0	0.09 ± 0.0	2.17 ± 0.2	41.9 ± 1.7	8.3	66.0	5.3	24.9
Treatment Tfinals														
Control	2056 ± 100	117 ± 10	0.12 ± 0.0	BD	0.03 ± 0.0	5.56 ± 0.2	0.27 ± 0.1	0.13 ± 0.0	BD	9.0 ± 2.1	2.1	0.5	4.8	97.2
+Temp	2095 ± 67	135 ± 5	0.14 ± 0.0	BD	0.06 ± 0.0	5.76 ± 0.3	0.21 ± 0.0	0.15 ± 0.0	BD	27.2 ± 8.8 ^a	2.3	1.0	3.6	96.3
+CO ₂	2053 ± 52	113 ± 1	0.16 ± 0.0	BD	0.06 ± 0.0	5.43 ± 0.1	0.36 ± 0.0	0.14 ± 0.0	BD	7.4 ± 0.8	2.8	1.1	6.4	96.1
Combined	2188 ± 64	156 ± 11	0.15 ± 0.0	BD	0.05 ± 0.0	6.17 ± 0.3	0.18 ± 0.0	0.16 ± 0.0	BD	20.4 ± 8.2	2.4	0.8	2.9	96.9
Urea diluent	NM	126 ± 10	2.10 ± 0.8	0.2 ± 0.0	4.30 ± 0.1	16.50 ± 2.1	11.10 ± 0.4	0.11 ± 0.0	2.17 ± 0.1	41.5 ± 0.8	9.1	18.6	48.1	71.4
Treatment Tfinals														
Control	2133 ± 42	117 ± 11	0.15 ± 0.0	BD	0.05 ± 0.0	5.90 ± 0.4	0.25 ± 0.1	0.13 ± 0.0	BD	17.4 ± 14.2	2.5	0.8	4.1	96.9
+Temp	2052 ± 49	132 ± 6	0.18 ± 0.1	BD	0.04 ± 0.0	5.40 ± 0.2	0.28 ± 0.0	0.12 ± 0.0	BD	8.6 ± 0.4	3.2	0.7	5.0	95.9
+CO ₂	2156 ± 139	112 ± 7	0.10 ± 0.0	BD	0.04 ± 0.0	5.40 ± 0.2	0.23 ± 0.1	0.13 ± 0.0	BD	8.0 ± 6.1	1.8	0.7	4.2	97.7
Combined	2162 ± 18	150 ± 22	0.21 ± 0.1	BD	BD	6.30 ± 0.2	0.28 ± 0.0	0.12 ± 0.0	BD	17.4 ± 11.8	3.1	0.3	4.2	96.5

^aIndicates $n = 2$ and error is half the range

tled on the bottom, and allows the biomass to be homogeneously removed through the outflow lines, which are situated on the upper shoulder of the incubation bottles. Consistent, quantitative removal of biomass is key to the ecostat design, as it is this mechanism that allows growth rates of the community to equal loss rates, which enables the conditions within the ecostat to become and remain relatively stable over time. We note that although conditions were generally uniform during the experiment, changes were continuously occurring, which is typical of continuous culture experiments looking at natural microbial assemblages (MacIntyre & Cullen 2005, Pickell et al. 2009). The dilution rate was set to 0.3 d^{-1} . This rate was selected based on typical specific growth rates of phytoplankton communities in California coastal waters (Landry et al. 2009, Li et al. 2010). Within the ranges reported in those studies (~ 0.3 – 0.6), we conservatively established our dilution rate so we would not risk washing out microorganisms with slower growth rates.

The SCMI community incubated for 24 h in the ecostat bottles before any diluent (NO_3^- or urea-based media) was added. After dilutions began, the first sampling event occurred 24 h later. These steps were completed in sequence to ensure that cells within the ecostats were growing and biomass was increasing before we began measuring rates.

Nutrient uptake experiments

To determine rates of nitrogen and DIC uptake in the ecostat treatments, an acid-washed syringe was used to collect 250 ml from each bottle. Care was taken to only draw off approximately 10% of the bottle volume to avoid substantial disruptions to the continuous culture nutrient inflow/biomass accumulation balance (Hutchins et al. 2003). Sub-samples were collected to measure concentrations of either NO_3^- or urea, corresponding to the diluent type that the treatment received. The remaining volume was used to measure uptake rates in acid-washed (10% HCl with multiple rinses) 230 ml polycarbonate conical bottles. Uptake experiments were performed using 2 different ^{15}N -labeled substrates: potassium nitrate (K^{15}NO_3 ; 98%) and ^{15}N -urea (98%; Cambridge Isotope Laboratories). To measure DIC uptake, ^{13}C -labeled $\text{NaH}^{13}\text{CO}_3^-$ (99%; Cambridge Isotope Laboratories) was used and added to all treatments (Hama et al. 1983). We note that the use of $\text{NaH}^{13}\text{CO}_3^-$ prevented the use of dual ^{15}N and ^{13}C -labeled urea, and so uptake of urea-C was not measured; generally

uptake of urea-C is very low (Bradley et al. 2010a, Bronk et al. 2014) or below detection (Bradley et al. 2010b). Uptake of NO₃⁻ was measured in the treatments that received NO₃⁻-based diluent, while uptake of urea was measured in the treatments that were supported by urea-based diluent. Concentrations of NO₃⁻ and urea were not known at the start of the ¹⁵N incubations, so additions of 1 μmol N l⁻¹ as ¹⁵NO₃⁻ or ¹⁵N-labeled urea were used. These additions were above tracer level for NO₃⁻ and urea throughout the experiment. The choice to use a larger addition was made to increase the likelihood that substrate concentrations would be similar across all treatments during the ¹⁵N incubations.

The bottles with added tracer were incubated within the ecostats, under the same temperature and light conditions as the larger ecostat experimental bottles. During the incubations, CO₂ was not bubbled into the uptake bottles; however, the uptake bottles were filled to capacity, which minimized head space and CO₂ exchange. Bottles were incubated for approximately 3 to 4 h and then filtered sequentially to collect 2 size fractions (0.7–5.0 and >5.0 μm). The larger size fraction (>5.0 μm) was collected on a Sterlitech silver membrane filter. The filtrate was then passed through a Whatman glass fiber filter (GF/F; nominal pore size of 0.7 μm) to collect the smaller size fraction (0.7–5.0 μm). Uptake experiments were conducted every other day throughout the duration of the experiment for a total of 5 time points over 10 d.

Dissolved and particulate nutrient analyses

Concentrations of NO₃⁻ and urea were measured on the same days that uptake experiments were performed. Additional nutrient samples (NH₄⁺, nitrite [NO₂⁻], PO₄³⁻, H₄SiO₄, dissolved primary amines [DPA], total dissolved nitrogen [TDN], and dissolved organic carbon [DOC]) were collected at the beginning and end of the experiment so that net changes in specific nutrient pools could be assessed. All nutrient samples were filtered through Whatman GF/F filters that had been combusted for 2 h at 450°C. Samples were stored at -20°C until analysis.

Concentrations of NH₄⁺ were measured in triplicate using the phenol-hypochlorite method (detection limit [DL] 0.05 μmol N l⁻¹, Koroleff 1983) on a Shimadzu UV-1601 spectrophotometer. Concentrations of NO₃⁻, NO₂⁻, PO₄³⁻, and H₄SiO₄ were measured in duplicate on a Lachat QuickChem 8500 autoanalyzer (DL 0.03 μmol N l⁻¹, DL 0.03 μmol P

l⁻¹, DL 0.05 μmol Si l⁻¹; Parsons et al. 1984). The manual diacetyl monoxime thiosemicarbazide method was used to analyze the concentration of urea in duplicate (DL 0.025 μmol N l⁻¹; Price & Harrison 1987). DPA samples were measured in triplicate using the *o*-phthalaldehyde method followed by analysis on a Shimadzu RF-1501 spectrofluorometer (DL 0.025 μmol N l⁻¹; Parsons et al. 1984). TDN and DOC were measured on a Shimadzu 5000A TOC-V/TNM (DL 2 μmol N l⁻¹, 5 μmol C l⁻¹; Sharp et al. 2004); deep-sea reference water samples from the University of Miami consensus reference material program were included to ensure analytical accuracy (Hansell 2005). Dissolved organic nitrogen (DON) was calculated by taking the difference between TDN and the sum of the inorganic nitrogen species (NH₄⁺, NO₃⁻, and NO₂⁻); standard deviations for the final DON concentrations were calculated using propagation of error. DIC was measured using coulometry (CM5230, UIC) following King et al. (2011); CO₂SYN software (Lewis & Wallace 1998) was used to calculate pCO₂ (in μatm). A Europa 20/20 isotope ratio mass spectrometer was used to analyze particulate nitrogen (PN) and particulate carbon (PC) concentrations and to measure sample isotopic enrichment of ¹⁵N and ¹³C.

Samples were collected at the start of the experiment to determine the initial community composition. Cells were preserved using an acidified Lugol's solution, identified and enumerated according to Tomas (1997) and the Utermöhl method (Utermöhl 1931), respectively, using an inverted compound light-microscope with an Accu0Scope 3032.

Uptake calculations

Specific (*V*) and absolute (*ρ*) uptake rates of NO₃⁻ and urea were calculated using Eqs. (1) and (2), respectively (Dugdale & Wilkerson 1986):

$$V = \frac{\text{PN at\%xs}}{\text{SP at\%xs} \times \text{Time}} \quad (1)$$

$$\rho = \frac{\text{PN at\%xs}}{\text{SP at\%xs} \times \text{Time}} \times [\text{PN}] \quad (2)$$

where *V* is specific uptake rate (h⁻¹), and *ρ* is absolute uptake rate (μmol N l⁻¹ h⁻¹); PN at%xs is the ¹⁵N atom% enrichment of PN minus 0.366, which is the typical ¹⁵N atom% enrichment of an atmospheric N₂ gas standard. SP at%xs is the initial enrichment of the nitrogen source pool minus ¹⁵N atom% enrichment of the atmospheric standard; in

this study the source pool was either NO_3^- or urea. The ambient concentration for NO_3^- and urea were measured each time an uptake experiment was conducted, and these values were used to calculate the initial enrichment of the nitrogen source pool. [PN] is the concentration of PN at the end of the incubation ($\mu\text{mol N l}^{-1}$). V expresses the physiological response of the community, and ρ normalizes for differences in biomass between treatments. Specific and absolute uptake rates of DIC were also calculated using the same equations; however, the nitrogen parameters were substituted with carbon, and PC at%xs is the ^{13}C atom% enrichment of PC minus 1.08, which is the natural ^{13}C enrichment of phytoplankton (Slawyk et al. 1977, Lefebvre et al. 2012). DIC concentrations were measured at 2 time points during the experiment, and the average DIC concentration for each experimental condition (Control, +Temp, + CO_2 , Combined; $n = 12$) was used in the calculation of uptake rates.

Statistical analyses

To calculate average uptake rates and average stoichiometric uptake ratios for the 4 treatments (Control, +Temp, + CO_2 , and Combined), all replicates within each treatment were combined and averaged ($n = 15$; 3 replicates of each treatment \times 5 time points). While the magnitude of V and ρ varied throughout the experiment, the trends were generally the same across time points. Expressing uptake rates as a combined average of the time points increased variability, but was done to produce greater statistical power.

Data were analyzed using the R statistical program (R Development Core Team 2010). Data were checked for normality and homogeneity of variance, and those that were not normally distributed were log transformed prior to statistical analysis. A 1-way repeated measures analysis of variance (ANOVA) was used to determine if there were significant differences in uptake rates between treatments, nitrogen source (NO_3^- and urea), and size fractions. A post hoc Tukey's test was run in order to locate the uptake means that were significantly different from one another. Means were considered to be significantly different at $p \leq 0.05$.

RESULTS

Source water characterization and nutrient concentrations

Concentrations of NO_3^- , NO_2^- , and urea were higher in Redondo Harbor relative to the SCMI dock. Concentrations of NO_3^- and urea, respectively, were 4.30 and 1.44 $\mu\text{mol N l}^{-1}$ in Redondo Harbor compared to 0.51 and 0.84 $\mu\text{mol N l}^{-1}$ at the SCMI dock. At both locations, DON made up $>50\%$ of the TDN pool, with urea accounting for 25% of the DON pool in Redondo Harbor and 13% of the DON pool at the SCMI dock. The concentration of PO_4^{3-} at Redondo Harbor was double the concentration at the SCMI dock, and the concentration of H_4SiO_4 was more than triple that seen off the SCMI dock (Table 1).

Every other day during the experiment, concentrations of NO_3^- were measured in the treatments that received NO_3^- diluent and urea in the treatments that received urea diluent (Table 2). Concentrations of NO_3^- were $<1 \mu\text{mol N l}^{-1}$ at all time points, with the exception of the + CO_2 treatment on Day 4. Concentrations of urea were consistently higher than concentrations of NO_3^- during the experiment. The 2 treatments exposed to elevated temperatures were more efficient at drawing down the available urea in the ecostats (Table 2).

The full suite of nutrients was measured in the ecostat bottles on the final day of the experiment. Nutrient concentrations were generally similar between treatments (Table 1). The concentration of TDN was 5.5–6.5 $\mu\text{mol N l}^{-1}$ in all treatments, with NH_4^+ comprising the largest portion of the inorganic nitrogen

Table 2. NO_3^- and urea concentrations measured on Days 2, 4, 6, 8, and 10 for the different treatments. NO_3^- concentrations are from the treatments that received NO_3^- diluent, and urea concentrations are from the treatments that received urea diluent. Treatment details are provided in 'Materials and methods'

	Day 2	Day 4	Day 6	Day 8	Day 10
NO_3^- concentrations					
Control	0.75 \pm 0.3	0.22 \pm 0.1	0.09 \pm 0.0	0.09 \pm 0.1	0.03 \pm 0.0
+Temp	0.17 \pm 0.1	0.39 \pm 0.1	0.07 \pm 0.0	0.12 \pm 0.1	0.06 \pm 0.0
+ CO_2	0.25 \pm 0.1	2.74 \pm 3.1 ^a	0.27 \pm 0.1	0.21 \pm 0.1	0.06 \pm 0.0
Combined	0.18 \pm 0.1	0.35 \pm 0.1	0.08 \pm 0.0	0.14 \pm 0.1	0.05 \pm 0.0
Urea concentrations					
Control	5.08 \pm 0.5	4.51 \pm 3.2	3.05 \pm 2.3	0.36 \pm 0.1	0.25 \pm 0.1
+Temp	0.51 \pm 0.1	1.15 \pm 0.9	0.45 \pm 0.1	0.42 \pm 0.0	0.28 \pm 0.0
+ CO_2	4.91 \pm 0.7	4.60 \pm 2.5	1.25 \pm 0.7	0.37 \pm 0.1	0.23 \pm 0.1
Combined	0.43 \pm 0.2	0.52 \pm 0.1	0.42 \pm 0.1	0.44 \pm 0.2	0.28 \pm 0.0

^aThere was an issue with the diluent pump prior to sampling

fraction. NO₃⁻ concentrations were low, measuring approximately 0.05 μmol N l⁻¹ in all treatments, and concentrations of PO₄³⁻ were below the limit of detection (Table 1). Although NO₃⁻ and PO₄³⁻ concentrations were low or below detection on the final day, both were continually supplied in the seawater diluent at a constant rate throughout the experiment. Concentrations of H₄SiO₄ and DOC varied between treatments at the end of the experiment. For both diluent types, DOC concentrations in the +Temp and Combined treatments were approximately 20 μmol C l⁻¹ higher in concentration than the Control and +CO₂ treatments.

Initial community and biomass

Diatoms dominated the initial community that was collected off of the SCMI dock, accounting for 99% of the taxa that were present. Approximately half of the diatom community was comprised of *Pseudo-nitzschia* species, including both *P. multiseriis* and *P. hasleana*, whose relative abundances were 48 and 4%, respectively. Other dominant diatoms included *Leptocylindrus danicus* (39%) and *Chaetoceros* spp. (5%).

PC and PN concentrations of the whole community (>0.7 μm) were more constant in the treatments receiving NO₃⁻ diluent, but more variable and higher in the treatments receiving urea diluent (Fig. 1).

Nitrogen uptake

Results from the community incubated in the ecostats had 2 general trends. Uptake rates of NO₃⁻ (both V_{NO_3} and ρ_{NO_3}) were generally lower than uptake rates of urea (both V_{UREA} and ρ_{UREA}), and uptake rates were higher in the larger size fraction (>5.0 μm) relative to the smaller size fraction (0.7–5.0 μm; Fig. 2). Elevated CO₂ resulted in higher NO₃⁻ uptake rates by smaller microorganisms, and elevated temperature resulted in higher urea uptake rates by both large and small microorganisms. These general trends were

also observed over the course of the experiment (Fig. 3). Uptake rates changed the most within the first 2 to 4 d of the experiment, particularly in the case of urea. After the fourth day, uptakes rates remained relatively stable in most treatments (Fig. 3).

NO₃⁻ uptake rates (V_{NO_3} and ρ_{NO_3}) in the larger size fraction did not significantly differ between any of the treatments (Fig. 2, $p > 0.05$). In the +CO₂ treatment, however, V_{NO_3} and ρ_{NO_3} within the smaller size fraction were significantly higher (~50%) than in the Control (Fig. 2, $p < 0.007$). Although CO₂ was also manipulated in the Combined treatment, no significant increase in NO₃⁻ uptake was observed in that treatment compared to the Control.




Fig. 1. Mean particulate (A) carbon (PC) and (B) nitrogen (PN) concentrations \pm 1 SD ($n = 3$) of NO₃⁻ diluent treatments (shaded markers) and urea diluent treatments (white markers) of the whole community (>0.7 μm) over the course of the experiment. Treatment details are provided in 'Materials and methods'




Fig. 2. Mean (A) specific (V) and (B) absolute (ρ) uptake rates \pm 1 SD ($n = 15$) of NO_3^- (solid bars) and urea (dotted bars) by small (0.7–5.0 μm , black bars) and large (>5.0 μm , white bars) microorganisms in the incubated community. Bars are an average of all treatment replicates across 5 time points. Bars that have different letters are significantly different from one another ($p \leq 0.05$). Treatment details are provided in 'Materials and methods'

Urea uptake rates (V_{UREA} and ρ_{UREA}) showed a distinctly different pattern. For V_{UREA} and ρ_{UREA} in the large size fraction, the +Temp treatment and the Combined treatment were significantly higher than both the Control and the +CO₂ treatment (Fig. 2, $p < 0.01$). In the smaller size fraction, the +Temp and Combined treatments had significantly higher V_{UREA} than the Control and +CO₂ treatment ($p < 0.001$). The same results were seen for ρ_{UREA} ($p < 0.01$), except that the Combined treatment did not have significantly higher uptake rates than the +CO₂ treatment.

DIC uptake

DIC uptake rates in the larger size fraction were significantly greater than uptake rates in the smaller size fraction for both V_{DIC} and ρ_{DIC} . This pattern occurred in all experimental treatments (Fig. 4, $p < 0.001$). Higher DIC uptake rates were observed in the +Temp and Combined treatments, following the same general trends observed for urea. In the smaller size fraction, this trend was significant for both V_{DIC}

and ρ_{DIC} when comparing the +Temp treatment to the Control ($p < 0.006$). The trend was also significant in the smaller size fraction when comparing V_{DIC} and ρ_{DIC} of the Combined treatment grown on NO_3^- to the Control ($p < 0.001$). In the larger size fraction, ρ_{DIC} of the +Temp treatment grown on NO_3^- had significantly higher rates than the Control, and ρ_{DIC} of the Combined treatment grown on urea was significantly higher than the Control ($p < 0.03$). None of the +CO₂ treatments was significantly different than the Controls; however, V_{DIC} and ρ_{DIC} of the larger size fraction in the +CO₂ treatments were significantly lower than the +Temp treatments ($p < 0.02$).

Stoichiometry

Particulate concentrations and C:N ratios were impacted by elevated temperature, although the trends were not statistically significant ($p > 0.05$). Concentrations of PC for the +Temp and Combined treatments were higher than the Control and +CO₂ treatments for both size fractions and both diluent types (Table 3). This trend was also found for PN concentrations in the treatments that received NO_3^- diluent, but not in the treatments that received urea diluent. For all treatments, PC:PN ratios increased in the +Temp and Combined treatments, while the +CO₂ treatment had similar ratios to the Control (Table 3).

Temperature also impacted the stoichiometric uptake ratio of DIC to NO_3^- . In the larger size fraction, the ratio of $\rho_{\text{DIC}}:\rho_{\text{NO}_3}$ for the +Temp treatment was significantly higher than the Control (Fig. 5, $p < 0.05$). In the smaller size fraction, the Combined treatment had a significantly higher $\rho_{\text{DIC}}:\rho_{\text{NO}_3}$ ratio than the Control (Fig. 5, $p < 0.02$). In contrast, the ratios of $\rho_{\text{DIC}}:\rho_{\text{UREA}}$ in all treatments were not significantly different from the Control.

DISCUSSION

The goal of this study was to determine how uptake of NO_3^- , urea, and DIC by Southern California Bight microorganisms will be affected by elevated temperature and CO₂ conditions that will likely develop over the next century. The SCMI microbial community used to set up the ecostats was dominated by diatom species that are commonly found in coastal southern California waters (Cupp 1943, Reid et al. 1970, 1978, Cullen et al. 1982, Venrick 2015), including *Pseudo-nitzschia* spp., *Leptocylindrus danicus*, and *Chaetoceros* spp. Both diluent types used in this experiment




Fig. 3. Mean (A,B) specific (V) and (C,D) absolute (ρ) uptake rates \pm 1 SD ($n = 3$) of NO_3^- (shaded markers) and urea (white markers) by small (0.7–5.0 μm , A and C) and large (>5.0 μm , B and D) microorganisms in the incubated community over the course of the experiment. Panels C and D have a different y-axis scale due to large differences in uptake rates between the small and large size fraction. Treatment details are provided in 'Materials and methods'




Fig. 4. Mean (A) specific (V) and (B) absolute (ρ) uptake rates \pm 1 SD ($n = 15$) of dissolved inorganic carbon (DIC) by microorganisms sustained on NO_3^- -based diluent (solid bars) and urea-based diluent (dotted bars) by small (0.7–5.0 μm , black bars) and large (>5.0 μm , white bars) microorganisms in the incubated community. Bars are an average of all treatment replicates across 5 time points. Bars that have different letters are significantly different from one another ($p \leq 0.05$). Treatment details are provided in 'Materials and methods'

were supplemented with H_4SiO_4 , ensuring that the diatom community would not become H_4SiO_4 limited. Thus, our study demonstrates how a coastal California microbial assemblage primarily composed of diatoms would be affected by climate change parameters and changes to the dominant nitrogen substrate. At the end of the experiment, *L. danicus* remained dominant in all treatments, and the relative abundance of other diatom species varied significantly between treatments (A. O. Tatters et al. unpubl.).

Nutrient uptake in the +Temp treatment

Many studies have documented that a temperature increase of a few degrees increases microbial growth rates and photosynthesis (Eppley 1972, Raven & Geider 1988, Boyd et al. 2013). Studies using both cultured phytoplankton and field communities found that the relationship between temperature and nitrogen uptake varies between species (Goldman 1977, Kristiansen 1983, Glibert et al. 1995, Lomas & Glibert 1999). Of these studies, that of Lomas & Glibert (1999) is the most similar to our study, as it used ¹⁵N incorporation to measure both NO_3^- and urea uptake by diatom-dominated field communities subjected to temperature manipulations that were similar to the temperatures used in our study. In that study, for every 1°C change in temperature, NO_3^- uptake decreased by 2.9% on average. We found that temper-

Table 3. Average particulate carbon (PC) and particulate nitrogen (PN) concentrations and PC:PN ratios for 2 size classes of microorganisms (0.7–5.0 and >5.0 μm) at the Southern California Marine Institute (SCMI, the starting community) dock and in the different treatments (treatment details are provided in 'Materials and methods'). PC and PN concentrations are in units of $\mu\text{mol C or N l}^{-1} \pm 1 \text{ SD}$ ($n = 15$; 3 reps $\times 5$ time points)

Source	0.7–5.0 μm			>5.0 μm		
	PC	PN	PC:PN	PC	PN	PC:PN
SCMI	29.0 \pm 1.1	3.38 \pm 0.2	8.60 \pm 0.6	36.6 \pm 1.6	3.97 \pm 0.1	9.18 \pm 0.2
NO₃⁻ diluent						
Control	35.6 \pm 6.1	3.72 \pm 1.1	10.2 \pm 2.3	153.6 \pm 50	11.0 \pm 2.3	14.2 \pm 4.7
+Temp	39.9 \pm 4.8	3.81 \pm 0.9	10.8 \pm 1.7	208.2 \pm 50	13.5 \pm 3.1	15.8 \pm 3.5
+CO ₂	37.4 \pm 7.6	3.96 \pm 0.8	9.6 \pm 1.4	132.5 \pm 46	10.2 \pm 2.4	13.3 \pm 5.2
Combined	45.4 \pm 9.4	4.09 \pm 1.1	11.3 \pm 1.5	210.2 \pm 73	13.9 \pm 5.0	15.6 \pm 3.2
Urea diluent						
Control	28.7 \pm 7.3	3.32 \pm 1.1	9.0 \pm 1.6	181.0 \pm 100	13.7 \pm 4.2	12.5 \pm 3.8
+Temp	32.6 \pm 7.2	3.23 \pm 1.0	10.3 \pm 1.3	205.3 \pm 51	14.7 \pm 2.1	13.9 \pm 2.7
+CO ₂	29.6 \pm 6.9	3.54 \pm 1.2	8.8 \pm 1.9	196.3 \pm 106	15.5 \pm 4.7	12.0 \pm 3.7
Combined	35.9 \pm 9.6	3.43 \pm 1.2	10.7 \pm 1.6	260.6 \pm 78	19.0 \pm 4.7	13.9 \pm 3.9




Fig. 5. Stoichiometric ratios of p_{DIC} to p_{NO_3} (solid bars) and p_{DIC} to p_{UREA} (dotted bars) $\pm 1 \text{ SD}$ ($n = 15$) for small (0.7–5.0 μm , black bars) and large (>5.0 μm , white bars) microorganisms in the incubated community, where p is the absolute uptake rate and DIC is dissolved inorganic carbon. Bars that have different letters are significantly different from one another ($p \leq 0.05$). Treatment details are provided in 'Materials and methods'

ature did not impact NO_3^- uptake, but consistent with Lomas & Glibert (1999), temperature did significantly increase uptake of urea (Fig. 2).

There are a few possible explanations for why temperature did not significantly affect NO_3^- uptake in our study. Seawater was collected from the surface close to shore, so microorganisms in this region may have already been accustomed to temperature fluctuations of several degrees. The amplitude of sea surface seasonal variation off the coast of Southern California can reach 3.1°C (Nezlin et al. 2004). Thus, our temperature increase of 4°C is close to the realm of nat-

ural seasonal variation. It is also possible that the ambient (19°C) and elevated (23°C) temperatures within this study were above the optimal physiological range for diatom NO_3^- uptake. In the diatom *Skeletonema costatum* that typically inhabits seawater that is 12–15°C, the activity of nitrate reductase (NR) is unstable above ~16°C (Gao et al. 2000). Likewise, Berges et al. (2002) showed that ^{15}N uptake and NR activity of the temperate diatom *Thalassiosira pseudonana* declines between 17 and 25°C. The optimal temperature range for growth of *L. danicus*, a dominant phytoplankton species in our study, is between 15 and 20°C (Verity 1982), so it is possible that NO_3^- uptake physi-

ology falls in this range as well.

In the case of urea, our data show a positive relationship between temperature and urea uptake. The majority of studies investigating temperature effects on urea uptake measured the activity and expression of enzymes, like urease, involved in urea metabolism rather than directly quantifying urea uptake. These studies showed that urease activity (Fan et al. 2003) and urea uptake (Lomas & Glibert 1999) are positively correlated with temperature. In our study, raising the temperature by 4°C increased uptake rates of urea (V_{UREA}) by 80% for small microorganisms and 54% for large microorganisms. The substantial increase in urea uptake rates suggests that as sea surface temperatures continue to rise, organisms capable of using urea may have an advantage, especially in temperate coastal systems that receive wastewater effluent or agricultural runoff, both of which contain urea (Bronk et al. 2010, Howard et al. 2014). Urea is also the most common form of nitrogen used in fertilizers for agricultural production (Glibert et al. 2006, Mérigout et al. 2008), and California is the top agricultural center in the US (California Department of Food and Agriculture 2015). With respect to the Southern California Bight, ~60% of the total nitrogen in riverine runoff is comprised of organic forms, including urea, and riverine runoff has been identified as the primary source of urea to this system (Howard et al. 2014). Urea is a small component, however, of the total nitrogen load delivered to the Southern California Bight when both natural and anthropogenic sources are included.

Similar to urea, there was also an increase in DIC uptake with increasing temperature (Fig. 4). This trend occurred regardless of the nitrogen substrate added to the diluent. The general model is that increased temperature elevates carbon fixation and respiration up to a certain threshold that is species-specific and contingent upon the organism's geographic location and inherent ability to acclimate (Eppley 1972, Goldman 1977, Verity 1982, Thomas et al. 2012, Boyd et al. 2013). In the past decade, more work has been done on communities rather than cultures, and the relationship between temperature and carbon consumption is proving to be more complex. A few studies even show that elevated temperature decreases carbon drawdown (Wohlers et al. 2009, Kim et al. 2011), which is contrary to our results. Not only did DIC uptake rates increase as a function of temperature, but concentrations of PC and DOC also increased in the +Temp treatments (Tables 1 & 3). Our findings resemble the results reported by Taucher et al. (2012), where a Baltic Sea microbial community was grown for a month at a range of temperatures resulting in a net increase of particulate and dissolved carbon. The authors hypothesized that increased temperatures may lead to carbon overconsumption and a decoupling between C:N ratios in both the particulate and dissolved form, leading to a build up of particulate and dissolved carbon and an increase in C:N ratios. This hypothesis is also supported by our study, where the PC:PN ratios (Table 3) and the ratio of absolute DIC uptake to absolute NO₃⁻ uptake were both elevated in the +Temp treatment (Fig. 5). The average ratio of $\rho_{\text{DIC}}:\rho_{\text{NO}_3^-}$ was significantly higher in the +Temp treatment compared to the Control for the larger size fraction. Higher DIC to NO₃⁻ uptake ratios at elevated temperature may be linked to the presence of diatoms. In a culture experiment, the diatom *Thalassiosira weissflogii* had higher DIC to NO₃⁻ uptake ratios when temperature was elevated by 5°C (Taucher et al. 2015). In this same study, however, the opposite pattern occurred for the diatom *Dactyliosolen fragilissimus*. This finding warrants further investigation because of the implications for shaping ocean carbon biogeochemistry and C:N stoichiometry in the future.

Nutrient uptake in the +CO₂ treatment

The physiological effects that CO₂ has on the uptake of nutrients are complex. Elevated CO₂ can significantly impact enzyme activity and subsequent metabolic processes such as photosynthesis (Mercado et al. 1999, Zou et al. 2011), calcification (Doney et al.

2009 and references therein), and nutrient uptake and assimilation (Magnusson et al. 1996, Zou 2005, Hofmann et al. 2013). Uptake of nitrogen is intrinsically linked to carbon utilization, and the mechanism by which microorganisms use carbon varies between species. Some microorganisms contain CCMs, and those that do not, rely on passive diffusion to acquire CO₂ (Raven 1991, Kaplan & Reinhold 1999, Giordano et al. 2005). Cells that actively use CCMs to acquire CO₂/and or HCO₃⁻ are rendered less 'leaky' when surrounded by seawater with higher CO₂ concentrations, causing the regulatory enzymes that operate the CCM to be down-regulated resulting in more energy that can be allocated to support physiological processes such as growth and nutrient assimilation (Burkhardt et al. 2001, Rost et al. 2003). Production and growth of species that rely on passive diffusion should be enhanced by higher ambient CO₂ concentrations. The metabolic processes of microorganisms without CCMs are more likely to be enhanced when CO₂ is increased than those microorganisms with CCMs (Giordano et al. 2005). Under elevated CO₂ conditions, community shifts can occur depending on the effectiveness of the mechanism that microorganisms use to acquire inorganic carbon (Tortell et al. 2002, 2008, Hare et al. 2007, Feng et al. 2009, 2010), translating to differences in uptake rates of other nutrients. In our study, elevated CO₂ concentrations influenced the uptake of NO₃⁻, while urea uptake was unaffected (Fig. 2).

Although few studies have examined how NO₃⁻ uptake by microalgae is affected by CO₂ concentration, the response appears to differ between microbial groups. We found this to be true in our study, given that uptake rates of NO₃⁻ by smaller microorganisms were significantly higher at elevated CO₂, while uptake rates of NO₃⁻ by larger microorganisms increased, but not significantly. Consistent with our results for smaller microorganisms, Beardall & Koss (unpubl. data described by Beardall et al. 2009) found that uptake of nitrogen was enhanced by elevated CO₂ concentrations in species of phytoplankton; however, the nitrogen substrate and phytoplankton group(s) that they used were not identified. Other mesocosm studies, however, have shown that uptake of NO₃⁻ is unaffected by varying levels of CO₂ in communities where diatoms were present (Riebesell et al. 2007) and absent (Engel et al. 2005).

Like urea, the uptake of DIC in both nitrogen diluent types and both size fractions was unaffected under high CO₂, contrary to several studies (Hutchins et al. 2007, Trimborn et al. 2009). The coastal California ecosystem is an upwelling system charac-

terized by high levels of DIC and low pH (Hauri et al. 2009) with a high degree of temporal variability (Carpone & Hutchins 2013, Leinweber & Gruber 2013) relative to other oceanic regions, making it likely that the microorganisms in our study have already optimized for high CO₂ conditions. Diatoms in particular may be less susceptible to high CO₂ conditions than other microalgae groups because of their efficient CCMs (Tortell et al. 2002, Fabry et al. 2008).

Nutrient uptake in the Combined treatment

We did not detect a significant additive effect of temperature combined with CO₂ on the uptake of NO₃⁻, urea, or DIC. Even though V_{UREA} and ρ_{UREA} were significantly higher than the Control in the Combined treatment, this was likely a function of temperature, as the combined effect was not significantly greater than the impact of temperature alone, and urea uptake was unaffected by elevated CO₂. Our results for DIC uptake in the Combined treatment were also likely driven by temperature. It is interesting that NO₃⁻ uptake rates by smaller microorganisms in the Combined treatment were not significantly higher than in the Control, since rates significantly increased when CO₂ alone was manipulated. The smaller microorganisms in the Combined treatment may have preferred other nitrogen substrates (i.e. urea) when both temperature and CO₂ were elevated.

The results for nitrogen and carbon uptake in the Combined treatment indicate that temperature, more so than CO₂, has the greater potential to change the cycling of urea and DIC in California coastal waters in the future. Uptake of NO₃⁻ may increase as a function of elevated CO₂; however, this relationship may only be important during the winter, when temperatures are cooler. Our results for dissolved and particulate carbon as well as $\rho_{\text{DIC}}:\rho_{\text{NO}_3}$ ratios in the Combined treatment further suggest that C:N ratios are likely to be different under future climate scenarios.

Nitrate versus urea

Within the same experimental conditions, uptake rates for urea were significantly greater than uptake rates for NO₃⁻ (Figs. 2 & 3). This result was somewhat counterintuitive, given that dissolved concentrations of NO₃⁻ were consistently lower than urea (Tables 1 & 2). One possible explanation is active urea regeneration, which would result in higher urea concentrations than if just uptake was taking place. Regenera-

tion of urea can occur on the order of minutes to days (Bronk et al. 2007), and rates of regeneration can be high in natural waters (Bronk et al. 1998). Sources of water column urea regeneration include zooplankton, bacteria, and phytoplankton (reviewed by Solomon et al. 2010). With respect to phytoplankton, some diatoms possess the genes that are necessary to produce urea within their cells (Armbrust et al. 2004, Bowler et al. 2008). It is definitely possible that urea was being rapidly used and regenerated within the ecostats.

Kinetics and different half saturating constants (K_s) for NO₃⁻ and urea uptake might also explain why urea uptake was higher than NO₃⁻ uptake even though dissolved concentrations of NO₃⁻ were lower than urea. For genera that were present in this study, including *Leptocylindrus* spp. and *Pseudo-nitzschia* spp., K_s values for NO₃⁻ uptake typically fall between 1 and 3 $\mu\text{mol N l}^{-1}$ (Eppley et al. 1969, Cochlan et al. 2008, Seeyave et al. 2009). Less is known about urea uptake kinetics, but K_s values for diatom urea uptake can range considerably (Bronk & Flynn 2006, Solomon et al. 2010 and references therein), falling lower and higher than K_s values for NO₃⁻ uptake. Cochlan et al. (2008) found that K_s values for NO₃⁻ uptake by *P. australis* were lower than K_s values for uptake of other nitrogen substrates (NH₄⁺ and glutamine), and that urea uptake did not follow typical Michaelis-Menten kinetics because uptake was non-saturating. They suggested that some species of diatoms have a relatively higher affinity for NO₃⁻, and are therefore able to more effectively take it up at low concentrations. In our study, it is possible that the microorganisms were taking up urea more quickly, and yet were not as efficient at drawing urea down to the very low concentrations observed for NO₃⁻.

We chose to assess NO₃⁻ and urea utilization in this study; however, another substrate of interest is NH₄⁺. Although the microorganisms in the ecostat bottles received considerably less NH₄⁺ than either NO₃⁻ or urea, which were added to the diluent, concentrations of NH₄⁺ were higher than concentrations of both NO₃⁻ and urea at the SCMI site (Table 1). Given the reduced nature of NH₄⁺, this substrate is often thought of as the preferred substrate for phytoplankton nitrogen uptake (McCarthy 1981, Glibert et al. 2016 and references therein), and it can inhibit and/or reduce NO₃⁻ uptake by phytoplankton depending on a variety of factors including ambient concentrations, environmental conditions, and species composition (Dortch 1990). In our study, it is possible that NO₃⁻ uptake rates were reduced due to the presence of NH₄⁺; however, any inhibitory effects were likely

minimal. First, concentrations of NH₄⁺ would be low in the ecostats themselves as the diluent was added slowly. Lack of NH₄⁺ inhibition can also be inferred by the continually low NO₃⁻ concentrations observed throughout the experiment (Table 2), indicating that NO₃⁻ was readily being used. It would be useful to include NH₄⁺ as a substrate in future ecostats, because of the dynamic interactions that occur between NO₃⁻, NH₄⁺, and urea with respect to microbial utilization and availability. For example, concentrations of NH₄⁺ can also increase as urea undergoes hydrolysis (Solomon et al. 2010). It would also be worthwhile to repeat the study and measure urea uptake in the treatments that received NO₃⁻ diluent and vice versa. We were unable to do this due to water limitations; however, background concentrations of both NO₃⁻ and urea were present in the diluent, and the low concentrations of both NO₃⁻ and urea measured at the end of the experiment in all treatments, regardless of the diluent type, indicate that the microorganisms in our study were using all of these nutrients.

In the Southern California Bight, it is particularly important to consider possible future changes in the nature and magnitude of nitrogen utilization by diatoms. The production of toxins by HAB species, such as the *Pseudo-nitzschia* diatoms that frequently bloom off the coast of California, is influenced by nutrient availability. *Pseudo-nitzschia* are capable of utilizing both inorganic and organic forms of nitrogen, including urea, as a source for growth (Cochlan et al. 2008, Loureiro et al. 2009, Auro & Cochlan 2013), and several factors including nutrient limitation, nitrogen source, and CO₂ can influence domoic acid production (Bates et al. 1998, Pan et al. 1998, Bates & Trainer 2006 and references therein, Sun et al. 2011, Sahraoui et al. 2012, Tatters et al. 2012). In this study, *Pseudo-nitzschia* spp. were present throughout the experiment, and domoic acid was detected in all treatments, with varying cellular toxin levels (A. O. Tatters et al. unpubl.).

CONCLUSIONS

Here we provide evidence that uptake of NO₃⁻, urea, and DIC is enhanced by changes in temperature and/or CO₂. Our results suggest that uptake of urea and DIC by phytoplankton communities dominated by diatoms will increase as sea surface temperatures rise in temperate nearshore environments. Although smaller microorganisms were a minor component of the community that we studied, we found that rates of NO₃⁻ uptake by smaller microorganisms

increased as a function of elevated CO₂ concentration. This relationship may be even more important during the winter, or if elevations in temperature and CO₂ are decoupled in the future, or when smaller microorganisms are more dominant. The relationship that we found between elevated temperature and urea uptake rates indicate that urea will be more rapidly utilized if warming continues. Additionally, the increased DIC to NO₃⁻ uptake ratios that occurred at elevated temperature indicate that C:N stoichiometry may also be altered in the future. Finally, this study demonstrates that uptake trends of nitrogen substrates under climate change conditions cannot be generalized, and it is important to study multiple substrates to gain a more comprehensive view of potential change.

Acknowledgements. We acknowledge Q. N. Roberts and M. P. Sanderson for their assistance with nutrient analysis, and E. A. Canuel, M. W. Lomas, D. K. Steinberg, and 3 anonymous reviewers for constructive feedback on the manuscript. This research was supported by grant numbers 1043635, 1043671, and 1043748, awarded to D.A.B., A.A.A., and D.A.H., respectively, from the National Science Foundation; D.A.H. received additional support from USC Urban Ocean Sea Grant. J.L.S. also received support from NSF GK-12 (DGE-0840804). This paper is Contribution No. 3650 of the Virginia Institute of Marine Science, College of William & Mary.

LITERATURE CITED

- ✦ Armbrust EV, Berges JA, Bowler C, Green BR and others (2004) The genome of the diatom *Thalassiosira pseudonana*: ecology, evolution, and metabolism. *Science* 306: 79–86
- ✦ Auro ME, Cochlan WP (2013) Nitrogen utilization and toxin production by two diatoms of the *Pseudo-nitzschia pseudodelicatissima* complex: *P. cuspidata* and *P. fryxelliana*. *J Phycol* 49:156–169
- Bates SS, Trainer VL (2006) The ecology of harmful diatoms. In: Granéli E, Turner JT (eds) *Ecology of harmful algae. Ecological studies (analysis and synthesis)*, Vol 189. Springer, Berlin, p 81–93
- ✦ Bates SS, Bird CJ, de Freitas ASW, Foxall R and others (1989) Pennate diatom *Nitzschia pungens* as the primary source of domoic acid, a toxin in shellfish from eastern Prince Edward Island, Canada. *Can J Fish Aquat Sci* 46: 1203–1215
- Bates SS, Garrison DL, Horner RA (1998) Bloom dynamics and physiology of domoic acid-producing *Pseudo-nitzschia* species. In: Anderson DM, Cembella AD, Hallegraeff GM, (eds) *Physiological ecology of harmful algal blooms*. Springer-Verlag, Heidelberg, p 267–292
- ✦ Beardall J, Stojkovic S, Larsen S (2009) Living in a high CO₂ world: impacts of global climate change on marine phytoplankton. *Plant Ecol Divers* 2:191–205
- ✦ Berges JA, Varela DE, Harrison PJ (2002) Effects of temperature on growth rate, cell composition and nitrogen metabolism in the marine diatom *Thalassiosira pseudonana* (Bacillariophyceae). *Mar Ecol Prog Ser* 225:139–146

- ✦ Bond NA, Cronin MF, Freeland H, Mantua N (2015) Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophys Res Lett* 42:3414–3420
- ✦ Bowler C, Allen AE, Badger JH, Grimwood J and others (2008) The *Phaeodactylum* genome reveals the evolutionary history of diatom genes. *Nature* 456:239–244
- Boyd PW, Doney SC (2002) Modelling regional responses by marine pelagic ecosystems to global climate change. *Geophys Res Lett* 29:53-1–53-4
- ✦ Boyd PW, Hutchins DA (2012) Understanding the responses of ocean biota to a complex matrix of cumulative anthropogenic change. *Mar Ecol Prog Ser* 470:125–135
- ✦ Boyd PW, Rynearson TA, Armstrong EA, Fu FX and others (2013) Marine phytoplankton temperature versus growth responses from polar to tropical waters – outcome of a scientific community-wide study. *PLOS ONE* 8:e63091
- ✦ Bradley PB, Sanderson MP, Frischer ME, Brofft J, Booth MG, Kerkhof LJ, Bronk DA (2010a) Inorganic and organic nitrogen uptake by phytoplankton and heterotrophic bacteria in the stratified Mid-Atlantic Bight. *Estuar Coast Shelf Sci* 88:429–441
- ✦ Bradley PB, Lomas MW, Bronk DA (2010b) Inorganic and organic nitrogen use by phytoplankton along Chesapeake Bay, measured using a flow cytometric sorting approach. *Estuaries Coasts* 33:971–984
- Bronk DA, Flynn KJ (2006) Algal cultures as a tool to study the cycling of dissolved organic nitrogen. In: Subba Rao DV (ed) *Algal cultures, analogues of blooms and applications*, Vol I. Science Publishers, Enfield, NH, p 301–341
- ✦ Bronk DA, Glibert PM, Malone TC, Banahan S, Sahlsten E (1998) Inorganic and organic nitrogen cycling in Chesapeake Bay: autotrophic versus heterotrophic processes and relationships to carbon flux. *Aquat Microb Ecol* 15: 177–189
- ✦ Bronk DA, See JH, Bradley PB, Killberg L (2007) DON as a source of bioavailable nitrogen for phytoplankton. *Biogeosciences* 4:283–296
- ✦ Bronk DA, Roberts QN, Sanderson MP, Canuel EA and others (2010) Effluent organic nitrogen (EON) bioavailability and photochemical and salinity-mediated release. *Environ Sci Technol* 44:5830–5835
- ✦ Bronk DA, Killberg-Thoreson L, Sipler RE, Mulholland MR and others (2014) Nitrogen uptake and regeneration (ammonium regeneration, nitrification and photoproduction) in waters of the West Florida Shelf prone to blooms of *Karenia brevis*. *Harmful Algae* 38:50–62
- ✦ Burkhardt S, Amoroso G, Riebesell U, Sultemeyer D (2001) CO₂ and HCO₃⁻ uptake in marine diatoms acclimated to different CO₂ concentrations. *Limnol Oceanogr* 46: 1378–1391
- California Department of Food and Agriculture (2015) California agricultural production statistics. www.cdffa.ca.gov/statistics/ (accessed on 31 May 2017)
- ✦ Capone DG, Hutchins DA (2013) The microbial biogeochemistry of coastal upwelling regimes in a changing ocean. *Nat Geosci* 6:711–715
- ✦ Cochlan WP, Herndon J, Kudela RM (2008) Inorganic and organic nitrogen uptake by the toxigenic diatom *Pseudo-nitzschia australis* (Bacillariophyceae). *Harmful Algae* 8: 111–118
- Cubasch U, Wuebbles D, Chen D, Facchini MC and others (2014) Working Group I contribution to the IPCC Fifth Assessment Report, Climate Change 2013: the physical science basis. IPCC, Geneva
- ✦ Cullen JJ, Reid FMH, Stewart E (1982) Phytoplankton in the surface and chlorophyll maximum off southern California in August, 1978. *J Plankton Res* 4:665–694
- Cupp EE (1943) *Marine plankton diatoms of the West Coast of North America*. University of California Press, Berkeley, CA
- ✦ Doney SC, Fabry VJ, Feely RA, Kleypas JA (2009) Ocean acidification: the other CO₂ problem. *Annu Rev Mar Sci* 1:169–192
- ✦ Dortch Q (1990) The interaction between ammonium and nitrate uptake in phytoplankton. *Mar Ecol Prog Ser* 61: 183–201
- ✦ Du X, Peterson W, Fisher J, Hunter M, Peterson J (2016) Initiation and development of a toxic and persistent *Pseudo-nitzschia* bloom off the Oregon coast in spring/summer 2015. *PLOS ONE* 11:e0163977
- ✦ Dugdale RC, Wilkerson FP (1986) The use of ¹⁵N to measure nitrogen uptake in eutrophic oceans; experimental considerations. *Limnol Oceanogr* 31:673–689
- ✦ Engel A, Zondervan I, Aerts K, Beaufort L and others (2005) Testing the direct effect of CO₂ concentration on a bloom of the coccolithophorid *Emiliania huxleyi* in mesocosm experiments. *Limnol Oceanogr* 50:493–507
- Eppley R (1972) Temperature and phytoplankton growth in the sea. *Fish Bull* 70:1063–1085
- ✦ Eppley RW, Rogers JN, McCarthy JJ (1969) Half-saturation constants for uptake of nitrate and ammonium by marine phytoplankton. *Limnol Oceanogr* 14:912–920
- ✦ Fabry VJ, Seibel BA, Feely RA, Orr JC (2008) Impacts of ocean acidification on marine fauna and ecosystem processes. *ICES J Mar Sci* 65:414–432
- ✦ Fan C, Glibert PM, Alexander J, Lomas MW (2003) Characterization of urease activity in three marine phytoplankton species, *Aureococcus anophagefferens*, *Prorocentrum minimum*, and *Thalassiosira weissflogii*. *Mar Biol* 142: 949–958
- ✦ Feng Y, Hare CE, Leblanc K, Rose J and others (2009) Effects of increased pCO₂ and temperature on the North Atlantic spring bloom. I. The phytoplankton community and biogeochemical response. *Mar Ecol Prog Ser* 388: 13–25
- ✦ Feng Y, Hare CE, Rose JM, Handy SM and others (2010) Interactive effects of iron, irradiance and CO₂ on Ross Sea phytoplankton. *Deep-Sea Res I* 57:368–383
- ✦ Gao Y, Smith G, Alberte R (2000) Temperature dependence of nitrate reductase activity in marine phytoplankton: biochemical analysis and ecological implications. *J Phycol* 36:304–313
- ✦ Giordano M, Beardall J, Raven JA (2005) CO₂ concentrating mechanisms in algae: mechanisms, environmental modulation, and evolution. *Annu Rev Plant Biol* 56:99–131
- ✦ Glibert PM, Conley DJ, Fisher TR, Harding LW Jr, Malone TC (1995) Dynamics of the 1990 winter/spring bloom in Chesapeake Bay. *Mar Ecol Prog Ser* 122:27–43
- ✦ Glibert P, Harrison J, Heil C, Seitzinger S (2006) Escalating worldwide use of urea—a global change contributing to coastal eutrophication. *Biogeochemistry* 77:441–463
- ✦ Glibert PM, Wilkerson FP, Dugdale RC, Raven JA and others (2016) Pluses and minuses of ammonium and nitrate uptake and assimilation by phytoplankton and implications for productivity and community composition, with emphasis on nitrogen-enriched conditions. *Limnol Oceanogr* 61:165–197
- ✦ Goldman JC (1977) Temperature effects on phytoplankton growth in continuous culture. *Limnol Oceanogr* 22: 932–936

- Hama T, Miyazaki T, Ogawa Y, Iwakuma T, Takahashi M, Otsuki A, Ichimura S (1983) Measurement of photosynthetic production of a marine phytoplankton population using a stable ¹³C isotope. *Mar Biol* 73:31–36
- Hansell DA (2005) Dissolved organic carbon reference material program. *EOS Trans Am Geophys Union* 86:318
- Hare CE, Leblanc K, DiTullio GR, Kudela RM and others (2007) Consequences of increased temperature and CO₂ for phytoplankton community structure in the Bering Sea. *Mar Ecol Prog Ser* 352:9–16
- Hauri C, Gruber N, Plattner GK, Alin S, Feely RA, Hales B, Wheeler PA (2009) Ocean acidification in the California Current System. *Oceanography* 22:60–71
- Hofmann LC, Straub S, Bischof K (2013) Elevated CO₂ levels affect the activity of nitrate reductase and carbonic anhydrase in the calcifying rhodophyte *Corallina officinalis*. *J Exp Bot* 64:899–908
- Howard MDA, Cochlan WP, Ladizinsky N, Kudela RM (2007) Nitrogenous preference of toxigenic *Pseudo-nitzschia australis* (Bacillariophyceae) from field and laboratory experiments. *Harmful Algae* 6:206–217
- Howard MDA, Sutula M, Caron DA, Chao Y and others (2014) Anthropogenic nutrient sources rival natural sources on small scales in the coastal waters of the Southern California Bight. *Limnol Oceanogr* 59:285–297
- Hutchins DA, Pustizzi F, Hare CE, DiTullio GR (2003) A shipboard natural community continuous culture system for ecologically relevant low-level nutrient enrichment experiments. *Limnol Oceanogr Methods* 1:82–91
- Hutchins DA, Fu FX, Zhang Y, Warner ME, Portune K, Bernhardt PW, Mulholland MR (2007) CO₂ control of *Trichodesmium* N₂ fixation, photosynthesis, growth rates, and elemental ratios: implications for past, present, and future ocean biogeochemistry. *Limnol Oceanogr* 52:1293–1304
- IPCC (2014) Climate Change 2014: synthesis report. Contribution of Working Groups I, II and III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change. IPCC, Geneva
- Jacox MG, Bograd SJ, Hazen EL, Feichter J (2015) Sensitivity of the California Current nutrient supply to wind, heat, and remote ocean forcing. *Geophys Res Lett* 42:5950–5957
- Kaplan A, Reinhold L (1999) CO₂ concentrating mechanisms in photosynthetic microorganisms. *Annu Rev Plant Physiol Plant Mol Biol* 50:539–570
- Kim JM, Lee K, Shin K, Yang EJ, Engel A, Karl DM, Kim HC (2011) Shifts in biogenic carbon flow from particulate to dissolved forms under high carbon dioxide and warm ocean conditions. *Geophys Res Lett* 38:L08612
- King AL, Sanudo-Wilhelmy SA, Leblanc K, Hutchins DA, Fu FX (2011) CO₂ and vitamin B₁₂ interactions determine bioactive trace metal requirements of a subarctic Pacific diatom. *ISME J* 5:1388–1396
- Koroleff F (1983) Determination of nutrients. In: Grasshoff K, Ehrhardt M, Kremling K (eds) *Methods of seawater analysis*. Verlag Chemie, New York, NY, p 125–187
- Kristiansen S (1983) The temperature optimum of the nitrate reductase assay for marine phytoplankton. *Limnol Oceanogr* 28:776–780
- Kroeker KJ, Kordas RL, Crim R, Hendriks IE and others (2013) Impacts of ocean acidification on marine organisms: quantifying sensitivities and interaction with warming. *Glob Change Biol* 19:1884–1896
- Landry MR, Ohman MD, Goericke R, Stukel MR, Tsyrvlevich K (2009) Lagrangian studies of phytoplankton growth and grazing relationships in a coastal upwelling ecosystem off Southern California. *Prog Oceanogr* 83:208–216
- Lefebvre S, Claquin P, Orvain F, Veron B, Charpy L (2012) Spatial and temporal dynamics of size structured photosynthetic parameters (PAM) and primary production (¹³C) of pico- and nano-phytoplankton in an atoll lagoon. *Mar Pollut Bull* 65:478–489
- Leinweber A, Gruber N (2013) Variability and trends of ocean acidification in the Southern California Current System: a time series from Santa Monica Bay. *J Geophys Res* 118:3622–3633
- Lewis E, Wallace DWR (1998) Program developed for CO₂ system calculations (Internet), ORNL/CDIAC-105. <http://cdiac.ornl.gov/ftp/co2sys>
- Lewitus AJ, Horner RA, Caron DA, Garcia-Mendoza E and others (2012) Harmful algal blooms along the North American west coast: history, trends, causes, and impacts. *Harmful Algae* 19:133–159
- Li QP, Franks PJS, Landry MR, Goericke R, Taylor AG (2010) Modeling phytoplankton growth rates and chlorophyll to carbon ratios in California coastal and pelagic ecosystems. *J Geophys Res* 115:G04003
- Lomas MW, Glibert PM (1999) Temperature regulation of nitrate uptake: a novel hypothesis about nitrate uptake and reduction in cool-water diatoms. *Limnol Oceanogr* 44:556–572
- Loureiro S, Jauzein C, Garces E, Collos Y, Camp J, Vague D (2009) The significance of organic nutrients to *Pseudo-nitzschia delicatissima* (Bacillariophyceae) nutrition. *J Plankton Res* 31:399–410
- MacIntyre HL, Cullen JJ (2005) Using cultures to investigate the physiological ecology of microalgae. In: Anderson R (ed) *Algal culturing techniques*. Academic Press, Burlington, MA, p 287–326
- Magnusson G, Larsson C, Lennart A (1996) Effects of high CO₂ treatment on nitrate and ammonium uptake by *Ulva lactuca* grown in different nutrient regimes. *Sci Mar* 60:179–189
- Martin-Jézéquel V, Calu G, Candela L, Amzil Z, Jauffrais T, Séchet V, Weigel P (2015) Effects of organic and inorganic nitrogen on the growth and production of domoic acid by *Pseudo-nitzschia multiseriata* and *P. australis* (Bacillariophyceae) in culture. *Mar Drugs* 13:7067–7086
- McCabe RM, Hickey BM, Kudela RM, Lefebvre KA and others (2016) An unprecedented coastwide toxic algal bloom linked to anomalous ocean conditions. *Geophys Res Lett* 43:10366–10376
- McCarthy JJ (1981) The kinetics of nutrient utilization. *Can J Fish Aquat Sci* 210:211–213
- Meehl G, Stocker T, Collins W, Friedlingstein P and others (2007) Global climate projections. In: Solomon S, Qin D, Manning M, Chen Z and others (eds) *Climate change 2007: the physical science basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge
- Mercado JM, Javier F, Gordillo L, Niell FX, Figueroa FL (1999) Effects of different levels of CO₂ on photosynthesis and cell components of the red alga *Porphyra leucosticta*. *J Appl Phycol* 11:455–461
- Mérigout P, Lelandais M, Bitton F, Renou JP, Briand X, Meyer C, Daniel-Vedele F (2008) Physiological and transcriptomic aspects of urea uptake and assimilation in *Arabidopsis* plants. *Plant Physiol* 147:1225–1238

- Nezlin NP, Oram JJ, DiGiacomo PM, Gruber N (2004) Sub-seasonal to interannual variations of sea surface temperature, salinity, oxygen anomaly, and transmissivity in Santa Monica Bay, California from 1987 to 1997. *Cont Shelf Res* 24:1053–1082
- Pan Y, Bates SS, Cembella AD (1998) Environmental stress and domoic acid production by *Pseudo-nitzschia*: a physiological perspective. *Nat Toxins* 6:127–135
- Parsons T, Maita Y, Lalli C (1984) A manual of chemical and biological methods for seawater analysis. Pergamon Press, New York, NY
- Pickell LD, Wells ML, Trick CG, Cochlan WP (2009) A sea-going continuous culture system for investigating phytoplankton community response to macro- and micro-nutrient manipulations. *Limnol Oceanogr Methods* 7:21–32
- Price N, Harrison P (1987) A comparison of methods for the measurement of dissolved urea concentrations in seawater. *Mar Biol* 94:307–317
- R Development Core Team (2010) R: a language and environment for statistical computing. www.r-project.org
- Raven JA (1991) Physiology of inorganic carbon acquisition and implications for resource use efficiency by marine phytoplankton: relation to increased CO₂ and temperature. *Plant Cell Environ* 14:774–794
- Raven JA, Geider RJ (1988) Temperature and algal growth. *New Phytol* 110:441–461
- Raven JA, Ball LA, Beardall J, Giordano M, Maberly SC (2005) Algae lacking carbon-concentrating mechanisms. *Can J Bot* 83:879–890
- Reid FMH, Fuglister E, Jordan JB (1970) Part V: phytoplankton taxonomy and standing crop. In: Strickland JDH (ed) *The ecology of the plankton off La Jolla, California, in the period April through September, 1967*. Scripps Institution of Oceanography, UC San Diego, San Diego, CA, p 51–66
- Reid FMH, Stewart E, Eppley RW, Goodman D (1978) Spatial distribution of phytoplankton species in chlorophyll maximum layers off southern California. *Limnol Oceanogr* 23:219–226
- Riebesell U (2004) Effects of CO₂ enrichment on marine phytoplankton. *J Oceanogr* 60:719–729
- Riebesell U, Zondervan I, Rost B, Tortell PD, Zeebe RE, Morel FMM (2000) Reduced calcification of marine plankton in response to increased atmospheric CO₂. *Nature* 407:364–367
- Riebesell U, Schulz KG, Bellerby RGJ, Botros M and others (2007) Enhanced biological carbon consumption in a high CO₂ ocean. *Nature* 450:545–548
- Rost B, Riebesell U, Burkhardt S, Sultemeyer D (2003) Carbon acquisition of bloom-forming marine phytoplankton. *Limnol Oceanogr* 48:55–67
- Rost B, Zondervan I, Wolf-Gladrow D (2008) Sensitivity of phytoplankton to future changes in ocean carbonate chemistry: current knowledge, contradictions and research directions. *Mar Ecol Prog Ser* 373:227–237
- Rykaczewski RR, Dunne JP (2010) Enhanced nutrient supply to the California Current Ecosystem with global warming and increased stratification in an earth system model. *Geophys Res Lett* 37:L21606
- Sahraoui I, Grami B, Bates SS, Bouchouicha D, Chikhaoui MA, Mabrouk HH, Hlaili AS (2012) Response of potentially toxic *Pseudo-nitzschia* (Bacillariophyceae) populations and domoic acid to environmental conditions in a eutrophied, SW Mediterranean coastal lagoon (Tunisia). *Estuar Coast Shelf Sci* 102-103:95–104
- Sarmiento H, Montoya JM, Vázquez-Domínguez E, Vaqué D, Gasol JM (2010) Warming effects on marine microbial food web processes: How far can we go when it comes to predictions? *Philos Trans R Soc Lond B Biol Sci* 365: 2137–2149
- Schnetzer A, Miller PE, Schaffner RA, Stauffer BA and others (2007) Blooms of *Pseudo-nitzschia* and domoic acid in the San Pedro Channel and Los Angeles harbor areas of the Southern California Bight, 2003–2004. *Harmful Algae* 6: 372–387
- Seeyave S, Probyn TA, Pitcher GC, Lucas MI, Purdie DA (2009) Nitrogen nutrition in assemblages dominated by *Pseudo-nitzschia* spp., *Alexandrium catenella* and *Dinophysis acuminata* off the west coast of South Africa. *Mar Ecol Prog Ser* 379:91–107
- Sharp JH, Bearegard AY, Burdige D, Cauwet G and others (2004) A direct instrument comparison for measurement of total dissolved nitrogen in seawater. *Mar Chem* 84: 181–193
- Slawyk G, Collos Y, Auclair JC (1977) The use of the ¹³C and ¹⁵N isotopes for the simultaneous measurement of carbon and nitrogen turnover rates in marine phytoplankton. *Limnol Oceanogr* 22:925–932
- Solomon CM, Collier JL, Berg GM, Glibert PM (2010) Role of urea in microbial metabolism in aquatic systems: a biochemical and molecular review. *Aquat Microb Ecol* 59:67–88
- Sun J, Hutchins DA, Feng Y, Seubert EL, Caron DA, Fu FX (2011) Effects of changing pCO₂ and phosphate availability on domoic acid production and physiology of the marine harmful bloom diatom *Pseudo-nitzschia multiseries*. *Limnol Oceanogr* 56:829–840
- Tatters AO, Fu FX, Hutchins DA (2012) High CO₂ and silicate limitation synergistically increase the toxicity of *Pseudo-nitzschia fraudulenta*. *PLOS ONE* 7:e32116
- Taucher J, Schulz KG, Dittmar T, Sommer U, Oschlies A, Riebesell U (2012) Enhanced carbon overconsumption in response to increasing temperatures during a mesocosm experiment. *Biogeosciences* 9:3531–3545
- Taucher J, Jones J, James A, Brzezinski MA, Carlson CA, Riebesell U, Passow U (2015) Combined effects of CO₂ and temperature on carbon uptake and partitioning by the marine diatoms *Thalassiosira weissflogii* and *Dactyliosolen fragilissimus*. *Limnol Oceanogr* 60:901–919
- Thomas MK, Kremer CT, Klausmeier CA, Litchman E (2012) A global pattern of thermal adaptation in marine phytoplankton. *Science* 338:1085–1088
- Tomas C (ed) (1997) *Identifying marine phytoplankton*. Academic Press, San Diego, CA
- Tortell PD, DiTullio GR, Sigman DM, Morel FMM (2002) CO₂ effects on taxonomic composition and nutrient utilization in an Equatorial Pacific phytoplankton assemblage. *Mar Ecol Prog Ser* 236:37–43
- Tortell PD, Payne C, Gueguen C, Strzepek RF, Boyd PW, Rost B (2008) Inorganic carbon uptake by Southern Ocean phytoplankton. *Limnol Oceanogr* 53:1266–1278
- Trimborn S, Wolf-Gladrow D, Richter KU, Rost B (2009) The effect of pCO₂ on carbon acquisition and intracellular assimilation in four marine diatoms. *J Exp Mar Biol Ecol* 376:26–36
- Utermöhl H (1931) Neue Wege in der quantitativen Erfassung des Planktons. (Mit besonderer Berücksichtigung des Ultraplanktons). *Verh Int Verein Limnol* 5:567–596
- Venrick EL (2015) Phytoplankton species in the California Current System off Southern California: the spatial

- dimensions. Calif Coop Ocean Fish Invest Rep 56: 168–184
- ✦ Verity PG (1982) Effects of temperature, irradiance, and daylength on the marine diatom *Leptocylindrus danicus* Cleve. III. Dark respiration. J Exp Mar Biol Ecol 60: 197–207
- ✦ Wohlers J, Engel A, Zöllner E, Breithaupt P and others (2009) Changes in biogenic carbon flow in response to sea surface warming. Proc Natl Acad Sci USA 106:7067–7072
- ✦ Wu Y, Gao K, Riebesell U (2010) CO₂-induced seawater acidification affects physiological performance of the marine diatom *Phaeodactylum tricornutum*. Biogeosciences 7:2915–2923
- ✦ Zou D (2005) Effects of elevated atmospheric CO₂ on growth, photosynthesis and nitrogen metabolism in the economic brown seaweed, *Hizikia fusiforme* (Sargassaceae, Phaeophyta). Aquaculture 250:726–735
- ✦ Zou D, Gao K, Luo H (2011) Short- and long-term effects of elevated CO₂ on photosynthesis and respiration in the marine macroalga *Hizikia fusiformis* (Sargassaceae, Phaeophyta) grown at low and high N supplies. J Phycol 47:87–97

Editorial responsibility: Antonio Bode,
A Coruña, Spain

Submitted: June 20, 2016; Accepted: June 28, 2017
Proofs received from author(s): July 28, 2017