

2011

Precise Extraction of the Induced Polarization in the He-4(e,e(l)(p)over-right-arrow)H-3 Reaction

S. P. Malace

M. Paolone

S. Strauch

L. Pentchev

William & Mary

C. F. Perdrisat

William & Mary, perdrisa@jlab.org

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

Recommended Citation

Malace, S. P., Paolone, M., Strauch, S., Albayrak, I., Arrington, J., Berman, B. L., ... & Christy, M. E. (2011). Precise Extraction of the Induced Polarization in the He 4 (e, e' p-) H 3 Reaction. *Physical review letters*, 106(5), 052501.

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Precise Extraction of the Induced Polarization in the ${}^4\text{He}(e, e'\vec{p}){}^3\text{H}$ Reaction

S. P. Malace,¹ M. Paolone,¹ S. Strauch,¹ I. Albayrak,² J. Arrington,³ B. L. Berman,⁴ E. J. Brash,⁵ B. Briscoe,⁴ A. Camsonne,⁶ J.-P. Chen,⁶ M. E. Christy,² E. Chudakov,⁶ E. Cisbani,⁷ B. Craver,⁸ F. Cusanno,⁷ R. Ent,⁶ F. Garibaldi,⁷ R. Gilman,^{9,6} O. Glamazdin,¹⁰ J. Glister,^{11,12} D. W. Higinbotham,⁶ C. E. Hyde-Wright,¹³ Y. Ilieva,¹ C. W. de Jager,⁶ X. Jiang,⁹ M. K. Jones,⁶ C. E. Keppel,² E. Khrosinkova,¹⁴ E. Kuchina,⁹ G. Kumbartzki,⁹ B. Lee,¹⁵ R. Lindgren,⁸ D. J. Margaziotis,¹⁶ D. Meekins,⁶ R. Michaels,⁶ K. Park,⁶ L. Pentchev,¹⁷ C. F. Perdrisat,¹⁷ E. Piassetzky,¹⁸ V. A. Punjabi,¹⁹ A. J. R. Puckett,²⁰ X. Qian,²¹ Y. Qiang,²⁰ R. D. Ransome,⁹ A. Saha,⁶ A. J. Sarty,¹¹ E. Schulte,⁹ P. Solvignon,³ R. R. Subedi,¹⁴ L. Tang,² D. Tedeschi,¹ V. Tvaskis,² J. M. Udias,²² P. E. Ulmer,¹³ J. R. Vignote,²³ F. R. Wesselmann,¹⁹ B. Wojtsekhowski,⁶ and X. Zhan²⁰

(E03-104 Collaboration)

¹University of South Carolina, Columbia, South Carolina 29208, USA

²Hampton University, Hampton, Virginia 23668, USA

³Argonne National Laboratory, Argonne, Illinois 60439, USA

⁴The George Washington University, Washington, D.C. 20052, USA

⁵Christopher Newport University, Newport News, Virginia 23606, USA

⁶Thomas Jefferson National Accelerator Facility, Newport News, Virginia 23606, USA

⁷INFN Rome, Sanitá Group and Istituto Superiore di Sanitá, I-00161 Rome, Italy

⁸University of Virginia, Charlottesville, Virginia 22904, USA

⁹Rutgers, The State University of New Jersey, Piscataway, New Jersey 08854, USA

¹⁰Kharkov Institute of Physics and Technology, Kharkov 310108, Ukraine

¹¹Saint Mary's University, Halifax, Nova Scotia, Canada

¹²Dalhousie University, Halifax, Nova Scotia, Canada

¹³Old Dominion University, Norfolk, Virginia 23529, USA

¹⁴Kent State University, Kent, Ohio 44242, USA

¹⁵Seoul National University, Seoul, Korea

¹⁶California State University, Los Angeles, Los Angeles, California 90032, USA

¹⁷College of William and Mary, Williamsburg, Virginia 23187, USA

¹⁸Tel Aviv University, Tel Aviv 69978, Israel

¹⁹Norfolk State University, Norfolk, Virginia 23504, USA

²⁰Massachusetts Institute of Technology, Cambridge, Massachusetts 02139, USA

²¹Duke University, Durham, North Carolina 27708, USA

²²Universidad Complutense de Madrid, E-28040 Madrid, Spain

²³Instituto de Estructura de la Materia, CSIC, E-28006 Madrid, Spain

(Received 22 November 2010; published 31 January 2011)

We measured with unprecedented precision the induced polarization P_y in ${}^4\text{He}(e, e'\vec{p}){}^3\text{H}$ at $Q^2 = 0.8$ and 1.3 (GeV/c)². The induced polarization is indicative of reaction-mechanism effects beyond the impulse approximation. Our results are in agreement with a relativistic distorted-wave impulse approximation calculation but are overestimated by a calculation with strong charge-exchange effects. Our data are used to constrain the strength of the spin-independent charge-exchange term in the latter calculation.

DOI: 10.1103/PhysRevLett.106.052501

PACS numbers: 25.30.Fj, 21.65.-f, 25.10.+s, 27.10.+h

Why and to what extent the nucleon changes its structure while embedded in nuclear medium has been a long-standing question in nuclear physics, attracting experimental and theoretical attention. In this context, one of the hotly debated topics has been the interpretation of the quenching in the polarization-transfer double ratio, $(P'_x/P'_z)_{\text{He}}/(P'_x/P'_z)_{\text{H}}$ extracted from measurements of the polarization-transfer coefficients, P'_x and P'_z , in elastic $\vec{e}p$ scattering and quasielastic scattering on ${}^4\text{He}$ [1–4]. In elastic $\vec{e}p$ scattering P'_x/P'_z is directly proportional to

the ratio of the electric and magnetic form factors of the proton, G_E/G_M [5]. In $A(\vec{e}, e'\vec{p})B$ quasielastic scattering, the polarization-transfer ratio is expected to be sensitive to the form-factor ratio of the proton embedded in the nuclear medium. The polarization double ratio is then taken to emphasize differences between the in-medium and free values. For a ${}^4\text{He}$ nucleus this double ratio was found to be quenched by 10% [1,4]. This quenching could be due to conventional nuclear medium effects like nucleon off shellness, meson-exchange currents, final-state

interactions (FSI), but also to unconventional effects like modifications of the electric and magnetic form factors of the proton in the nuclear medium [6]. However, an interpretation of a small quenching in the polarization-transfer double ratio as evidence of unconventional nuclear effects requires excellent control of conventional reaction mechanisms and hence remains, to some degree, model dependent. The induced polarization P_y , experimentally accessible along with P'_x and P'_z , is a measure of conventional nuclear effects and offers vital constraints for the interpretation of the polarization-transfer double ratio.

The polarization-transfer double ratio from Jefferson Lab experiments E93-049 [1] and E03-104 [4] has been successfully modeled by two competing theoretical predictions: the relativistic distorted-wave impulse approximation (RDWIA) calculation by the Madrid group [2] with medium-modified form factors from the quark-meson coupling (QMC) model [6], and the calculation of Schiavilla *et al.* [3] which assumes free nucleon form factors but has different modeling of nuclear conventional effects, in particular, the FSIs. Whether the two models give an accurate description of P_y has become the key in the interpretation of the polarization-transfer double ratio. The large uncertainties of E93-049 P_y measurements precluded any definite conclusion. Experiment E03-104 provides the most precise measurements to date of the induced polarization in ${}^4\text{He}(e, e'\vec{p}){}^3\text{H}$, and this Letter presents the results.

We report measurements of the induced polarization P_y in the quasielastic reaction ${}^4\text{He}(e, e'\vec{p}){}^3\text{H}$, at four-momentum transfer Q^2 of 0.8 and 1.3 $(\text{GeV}/c)^2$ and missing momentum, p_m ranging from 0 to 160 MeV/c . A longitudinally polarized electron beam with flipping polarization direction and a current of 80 μA was incident on ${}^4\text{He}$ and ${}^1\text{H}$ targets, and the scattered electron and recoil proton were detected in coincidence in two high-resolution spectrometer arms. The ${}^4\text{He}$ target was chosen because its relative simplicity allows for realistic microscopic theoretical calculations while its high nuclear density increases the sensitivity to nuclear medium effects. The proton arm central momenta for the ${}^1\text{H}(\vec{e}, e'\vec{p})$ reaction were adjusted in 2% increments from -8% to $+8\%$ so that protons in elastic $\vec{e}p$ scattering had a similar coverage of the focal plane as in the ${}^4\text{He}(\vec{e}, e'\vec{p}){}^3\text{H}$ reaction [4]. These $\vec{e}p$ measurements provided a baseline for the comparison of in-medium to free proton polarizations and were also used to check for possible instrumental asymmetries.

The polarized recoil protons traveled through the magnetic field of the spectrometer to the detector package used to measure the polarizations, the focal plane polarimeter (FPP) [7]. The spin precession of the protons was calculated using a well-established model of the spectrometer's magnetic field [4]. In the FPP, the polarized protons scattered in a carbon block leading to azimuthal asymmetries. These asymmetries in combination with information on the proton spin precession and the carbon analyzing

power were analyzed by means of a maximum likelihood method to obtain the induced polarization [8].

The extraction of the induced polarization P_y is complicated by the presence of instrumental asymmetries. For the particular reaction that we studied, P_y was expected to be small, $<6\%$ [1]. Thus even small instrumental asymmetries could constitute a significant background. The ${}^1\text{H}$ data have been used to check for the presence of instrumental asymmetries. In the one-photon-exchange approximation P_y in ${}^1\text{H}(e, e'\vec{p})$ is expected to be zero. The two-photon-exchange processes could yield a nonzero but rather small induced polarization, theoretical calculations predicting a value below 1% [9,10] at our kinematics. When taking into account the analyzing power and the recoil proton spin transport, this will translate into an expectation for the physics azimuthal asymmetries of $<0.4\%$, making any significant instrumental asymmetries easy to detect.

We performed an extensive study to identify and correct for these asymmetries. The azimuthal distributions of the polarized protons are reconstructed from the track information provided by the FPP straw chambers located before (front) and after (rear) the carbon analyzer [7]. We engaged in a thorough check of the performance of the chambers, and we found that inefficient regions and misalignments of the front and rear chambers lead to a contamination of the physics asymmetries. We devised a new tracking algorithm to allow track reconstruction even in inefficient regions, and we developed a more precise alignment procedure to correct for misalignments. As a result, we see only small variations, at the subpercent level, in the experimental azimuthal asymmetries for ${}^1\text{H}(e, e'\vec{p})$, with few outliers up to 1% in the few inefficient regions and at the edges of the FPP acceptance. On average, the experimental azimuthal asymmetries for ${}^1\text{H}(e, e'\vec{p})$ are at the subpercent level.

To cancel out these residual instrumental asymmetries, we obtain P_y in ${}^4\text{He}(e, e'\vec{p}){}^3\text{H}$, P_y in Table I, as the difference of P_y extracted from ${}^4\text{He}$, $P_y(\text{raw})$ in Table I, and ${}^1\text{H}$ data. Our systematic studies found the induced polarization thus extracted to be very robust on average and within the acceptance of the detector when binned in various kinematic variables. We reduced the systematic uncertainty on the P_y extraction by a factor of 4 when compared to previous, similar measurements from

TABLE I. The induced polarization P_y from E03-104. $P_y(\text{raw})$ is the experimental value of the induced polarization in ${}^4\text{He}(e, e'\vec{p}){}^3\text{H}$. P_y is the difference between the experimental values of P_y in ${}^4\text{He}(e, e'\vec{p}){}^3\text{H}$ and ${}^1\text{H}(e, e'\vec{p})$, which gives, in the absence of the induced polarization in ${}^1\text{H}$, the induced polarization in ${}^4\text{He}$. Stat and Syst represent the statistical and systematic uncertainties, respectively.

Q^2 ($\text{GeV}/c)^2$	$P_y(\text{raw})$	Stat	P_y	Stat	Syst
0.8	-0.0366	0.0042	-0.0415	0.0050	0.0050
1.3	-0.0394	0.0039	-0.0373	0.0043	0.0058

E93-049 [1]. Our data are used to put to stringent test state-of-the-art theoretical calculations, and such comparisons are presented in what follows.

In Fig. 1 we show the induced polarization P_y in ${}^4\text{He}(e, e'\bar{p}){}^3\text{H}$ extracted from E03-104 (upper panel) together with earlier results from E93-049 [1] (lower panel) and theoretical calculations from the Madrid group [2] (curves and band) which were averaged over the spectrometer acceptance. In the RDWIA, the nuclear current is calculated with relativistic wave functions for the initial bound and outgoing proton. The nuclear current operator can be of $cc1$ or $cc2$ forms [11] depending on the prescription used to enforce current conservation. The final outgoing proton wave function is a solution of a Dirac equation with global optical potentials to account for FSIs. The optical potential models used are McNeil-Ray-Wallace (MRW) [12] and Love-Franey (RLF) [13]. In




FIG. 1 (color online). The acceptance-averaged induced polarization P_y in ${}^4\text{He}(e, e'\bar{p}){}^3\text{H}$ as a function of Q^2 from E03-104 (upper panel) and E93-049 (lower panel) [1]. The inner and outer error bars represent the statistical and total uncertainties, respectively. The green band displays the Madrid calculation [2] when using MRW as optical potential and $cc1$ or $cc2$ for the form of the nuclear current operator. For the solid and dashed curves RLF was used with $cc1$ (solid) and $cc2$ (dashed). The dash-dotted and dotted curves were obtained from the dashed and solid ones, respectively, by including medium-modified form factors via the QMC model [6]. Lines connect the acceptance-averaged theory calculations.

Fig. 1 the green band represents the Madrid calculation when MRW is used, and the width of the band depends on the form of the nuclear current operator, $cc1$ or $cc2$, with $cc1$ giving a larger P_y in absolute value. The blue solid and dashed curves represent the Madrid calculation when using the RLF optical potential and $cc1$ (solid line) or $cc2$ (dashed line). The older E93-049 experiment [1] averaged over a larger range in missing momentum than E03-104, see Fig. 1, and P_y is predicted to increase, in absolute value, with increasing missing momentum. This causes the apparent drop in the theory results at the kinematics of E93-049 when compared to those of E03-104. The considerably reduced systematic uncertainties of the new results make possible a clear distinction between various theoretical prescriptions: the best description of the data is given by RDWIA (RLF, $cc1$). The inclusion of medium-modified form factors via the QMC model [6] slightly increases P_y , in absolute value, more so at large Q^2 , as shown by the dash-dotted and dotted curves.

In Fig. 2 we present the distribution of the induced polarization P_y as a function of the missing momentum p_m . Our results are compared to the Madrid RDWIA calculation [2]. Overall, there is good agreement between data and the theoretical prediction. Both data and calculation show an increase in P_y (in absolute value) with




FIG. 2 (color online). P_y in ${}^4\text{He}(e, e'\bar{p}){}^3\text{H}$ as a function of missing momentum p_m from E03-104 (solid circles). Also shown are calculations from the Madrid group [2] (notations as in Fig. 1).

increasing p_m . The calculation predicts a stronger variation of P_y in the range of p_m from -0.15 to 0 GeV/ c compared to 0 to 0.15 GeV/ c . Although there is some hint in the data that supports this behavior, especially at $Q^2 = 0.8$ (GeV/ c)², the size of the statistical uncertainties precludes any definite conclusion.

Another state-of-the-art theoretical calculation is the computation of Schiavilla *et al.* which uses variational wave functions for the bound three- and four-nucleon systems, nonrelativistic meson-exchange currents (2-body currents) and free nucleon form factors. The FSIs are treated within the optical potential framework and include both spin-independent and spin-dependent charge-exchange terms which play a crucial role in the prediction of P_y [3] for this calculation. The spin-independent charge-exchange term is constrained by $p + {}^3\text{H} \rightarrow n + {}^3\text{He}$ charge-exchange cross section data while the spin-dependent one is largely unconstrained [3].

In Fig. 3 our P_y results are compared to the calculation of Schiavilla *et al.* [3]. To facilitate this comparison our data have been corrected for the spectrometer acceptance as this theoretical calculation is only available at $p_m \approx 0$. This correction ($< 20\%$ for this experiment) was determined using the Madrid RDWIA (RLF, *cc1*) model because it offers a very good qualitative description of the P_y dependence on p_m . The stability of the acceptance correction was studied by using other prescriptions within the Madrid RDWIA calculation, and the variations were negligible when compared to the experimental uncertainties. The calculation from the Madrid group is also shown at $p_m \approx 0$. The remaining small but visible variation in the induced polarization between E93-049 and E03-104 kinematics is due to the higher beam energies used in E93-049. Although the prediction of Schiavilla *et al.*, Schiavilla (2005) in Fig. 3, offers a good description of the




FIG. 3 (color online). The induced polarization P_y in ${}^4\text{He}(e, e'\bar{p}){}^3\text{H}$ as a function of Q^2 extrapolated at missing momentum $p_m \approx 0$ from E03-104 (solid circles) and E93-049 (empty circles) [1]. The inner and outer error bars represent the statistical and total uncertainties, respectively. Calculations from the Madrid group [2] (continuous and dashed curves) and from Schiavilla *et al.* [3,14] (bands) are also shown.

polarization-transfer double ratio [4], it overpredicts, in absolute value, our measurements of P_y , especially at larger Q^2 . This evident discrepancy prompted a revision of this calculation. The new calculation [14], Schiavilla (2010) in Fig. 3, uses our P_y results as additional constraints for the modeling of the charge-exchange terms. The calculation proved insensitive to variations of the spin-dependent charge-exchange term (especially at larger Q^2), and this remains largely unconstrained [14]. However, the spin-independent charge-exchange contribution has been modified to provide a good fit to our data. It remains to be verified whether the agreement with the charge-exchange cross section data from $p + {}^3\text{H} \rightarrow n + {}^3\text{He}$ is still maintained. This 2010 version of the calculation is also in good agreement with the polarization-transfer double ratio.

The role of the charge exchange in ${}^4\text{He}(e, e'\bar{p}){}^3\text{H}$ still needs to be clarified. P_y in Schiavilla's 2010 calculation [14] proves to be mostly sensitive to the charge-exchange spin-independent term, leaving the spin-dependent one still unconstrained; the Madrid group deems P_y largely insensitive to both terms within the RDWIA framework. On the other hand, the charge-exchange cross sections as predicted by the Madrid RDWIA calculation need to be gauged against data. Possibly, a comparison of the induced polarization in ${}^4\text{He}(e, e'\bar{p}){}^3\text{H}$ and ${}^4\text{He}(e, e'\bar{n}){}^3\text{He}$ could cast some light on the role of charge exchange in this reaction.

To summarize, we measured with unprecedented precision the induced polarization P_y in ${}^4\text{He}(e, e'\bar{p}){}^3\text{H}$ at $Q^2 = 0.8$ and 1.3 (GeV/ c)². For the first time the systematic uncertainties on this observable were reduced to a size comparable to the statistical uncertainties. We compared our results with theoretical calculations from the Madrid group [2] and Schiavilla *et al.* [3,14]. The Madrid RDWIA prediction describes well our data when RLF is used as optical potential and *cc1* as form for the nuclear current operator. The 2005 prediction from Schiavilla *et al.* overestimates our P_y results (in absolute value) but gives a good description of the polarization-transfer double ratio [4]. Our induced polarization data have then been used to constrain the charge-exchange spin-independent term in the calculation, and this 2010 version describes well both the induced polarization and the polarization-transfer double ratio. Our high-precision data point to the need to carefully consider both the charge-exchange spin-independent and spin-dependent terms in realistic calculations, to settle the extent of medium modifications in nucleon structure.

The collaboration wishes to acknowledge the Hall A technical staff and the Jefferson Lab Accelerator Division for their support. This work was supported by the U.S. Department of Energy and the U.S. National Science Foundation. Jefferson Science Associates operates the Thomas Jefferson National Accelerator Facility under DOE Contract No. DE-AC05-06OR23177.

- [1] S. Strauch *et al.*, *Phys. Rev. Lett.* **91**, 052301 (2003).
- [2] J.M. Udias *et al.*, *Phys. Rev. C* **48**, 2731 (1993); J.M. Udias *et al.*, *Phys. Rev. C* **51**, 3246 (1995); J.M. Udias *et al.*, *Phys. Rev. C* **53**, R1488 (1996); J.M. Udias *et al.*, *Phys. Rev. Lett.* **83**, 5451 (1999); J.M. Udias and J.R. Vignote, *Phys. Rev. C* **62**, 034302 (2000).
- [3] R. Schiavilla *et al.*, *Phys. Rev. Lett.* **94**, 072303 (2005).
- [4] M. Paolone *et al.*, *Phys. Rev. Lett.* **105**, 072001 (2010).
- [5] A.I. Akhiezer and M.P. Rekalov, *Sov. J. Part. Nucl.* **4**, 277 (1974).
- [6] D.H. Lu *et al.*, *Phys. Lett. B* **417**, 217 (1998); , *Phys. Rev. C* **60**, 068201 (1999); M.R. Frank, B.K. Jennings, and G.A. Miller, *Phys. Rev. C* **54**, 920 (1996); U.T. Yakhshiev, U-G. Meissner, and A. Wirzba, *Eur. Phys. J. A* **16**, 569 (2003); J.R. Smith and G.A. Miller, *Phys. Rev. C* **70**, 065205 (2004); T. Horikawa and W. Bentz, *Nucl. Phys. A* **762**, 102 (2005); S. Liuti, [arXiv:hep-ph/0601125v2](https://arxiv.org/abs/hep-ph/0601125v2); V. Guzey, A.W. Thomas, and K. Tsushima, *Phys. Rev. C* **79**, 055205 (2009).
- [7] V. Punjabi *et al.*, *Phys. Rev. C* **71**, 055202 (2005).
- [8] D. Besset *et al.*, *Nucl. Instrum. Methods* **166**, 515 (1979).
- [9] P.G. Blunden, W. Melnitchouk, and J.A. Tjon, *Phys. Rev. C* **72**, 034612 (2005).
- [10] Y.C. Chen *et al.*, *Phys. Rev. Lett.* **93**, 122301 (2004); A.V. Afanasev *et al.*, *Phys. Rev. D* **72**, 013008 (2005).
- [11] T. DeForest, *Nucl. Phys.* **A392**, 232 (1983).
- [12] J.A. McNeil, L. Ray, and S.J. Wallace, *Phys. Rev. C* **27**, 2123 (1983).
- [13] C.J. Horowitz, *Phys. Rev. C* **31**, 1340 (1985); D.P. Murdock and C.J. Horowitz, *Phys. Rev. C* **35**, 1442 (1987).
- [14] R. Schiavilla (private communication).