

W&M ScholarWorks

Arts & Sciences Articles

Arts and Sciences

2011

Dendritic Calcium Activity Precedes Inspiratory Bursts in preBotzinger Complex Neurons

Christopher A. Del Negro William & Mary, cadeln@wm.edu

John A. Hayes *William & Mary*

Jens C. Rekling

Follow this and additional works at: https://scholarworks.wm.edu/aspubs

Recommended Citation

Del Negro, Christopher A.; Hayes, John A.; and Rekling, Jens C., Dendritic Calcium Activity Precedes Inspiratory Bursts in preBotzinger Complex Neurons (2011). *Journal of Neuroscience*, 31(3), 1017-1022. 10.1523/JNEUROSCI.4731-10.2011

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Dendritic Calcium Activity Precedes Inspiratory Bursts in preBötzinger Complex Neurons

Christopher A. Del Negro,¹ John A. Hayes,¹ and Jens C. Rekling²

¹Department of Applied Science, McGlothlin-Street Hall, The College of William and Mary, Williamsburg, Virginia 23187, and ²Department of Neuroscience and Pharmacology, Panum Institute, Copenhagen DK-2200, Denmark

Medullary interneurons of the preBötzinger complex assemble excitatory networks that produce inspiratory-related neural rhythms, but the importance of somatodendritic conductances in rhythm generation is still incompletely understood. Synaptic input may cause Ca²⁺ accumulation postsynaptically to evoke a Ca²⁺-activated inward current that contributes to inspiratory burst generation. We measured Ca²⁺ transients by two-photon imaging dendrites while recording neuronal somata electrophysiologically. Dendritic Ca²⁺ accumulation frequently precedes inspiratory bursts, particularly at recording sites 50–300 μ m distal from the soma. Preinspiratory Ca²⁺ transients occur in hotspots, not ubiquitously, in dendrites. Ca²⁺ activity propagates orthodromically toward the soma (and antidromically to more distal regions of the dendrite) at rapid rates (300–700 μ m/s). These high propagation rates suggest that dendritic Ca²⁺ activates an inward current to electrotonically depolarize the soma, rather than propagate as a regenerative Ca²⁺ wave. These data provide new evidence that respiratory rhythmogenesis may depend on dendritic burst-generating conductances activated in the context of network activity.

Introduction

Breathing behavior in mammals depends on inspiratory neural rhythms that originate in the preBötzinger complex (preBötC) of the ventral medulla. Glutamatergic interneurons comprise its rhythmogenic core and a key unsolved question pertains to the spatiotemporal distribution of active conductances in the somatodendritic membrane and their role in rhythmogenesis (Ramirez et al., 2004; Feldman and Del Negro, 2006). One possible mechanism posits that recurrent synaptic interactions evoke postsynaptic conductances to generate inspiratory bursts (Rekling et al., 1996; Rekling and Feldman, 1998). Specifically, recurrent excitation may elevate postsynaptic Ca²⁺ and thus activate nonspecific cationic current (I_{CAN}) (Mironov, 2008; Rubin et al., 2009). Studies in a slice model of respiratory rhythm show that $I_{\rm CAN}$ is one important charge carrier underlying inspiratory bursts (Thoby-Brisson and Ramirez, 2001; Peña et al., 2004) and that the synaptic recruitment of I_{CAN} takes place in the dendrites (Pace et al., 2007b; Mironov, 2008).

Glutamatergic preBötC neurons have the potential to excite their postsynaptic targets and thus cause Ca²⁺ to accumulate in the dendrites before inspiratory bursts. This has been

Received Sept. 9, 2010; revised Oct. 18, 2010; accepted Nov. 7, 2010.

J. A. Hayes' present address: Institut de Neurobiologie Alfred Fessard, Neurobiologie et Développement Centre National de la Recherche Scientifique, 91198 Gif-sur-Yvette, France. E-mail: john.hayes@inaf.cnrs-gif.fr. DOI: 10.1523/JNEUROSCI.4731-10.2011

Copyright © 2011 the authors 0270-6474/11/311017-06\$15.00/0

examined in slices and organotypic cultures but recording sites were limited to proximal dendrites (~50 μ m from the soma) and many data were acquired from nonrhythmic cultures (Mironov, 2008, 2009). Therefore, the extent to which these data apply to preBötC neuron dendrites in the context of rhythmogenesis remains uncertain. We measured postsynaptic Ca²⁺ dynamics in soma-close and soma-distal dendritic locations, using acute slices that retain respiratory function *in vitro*. We show that endogenous synaptic drive evokes dendritic Ca²⁺ transients before somatic inspiratory bursts, which suggests that synaptically activated dendritic conductances may influence respiratory rhythm generation.

Materials and Methods

The Institutional Animal Care and Use Committee at the College of William and Mary and the Danish National Committee for Ethics in Animal Research approved all procedures. Neonatal mice [postnatal days (P) 0-6] were dissected in artificial CSF (ACSF) containing the following (in mM): 124 NaCl, 3 KCl, 1.5 CaCl₂, 1 MgSO₄, 25 NaHCO₃, 0.5 NaH₂PO₄, and 30 dextrose, equilibrated with 95% O₂/5% CO₂, pH 7.4. CaCl₂ was adjusted to 1.1 mM in some experiments (see below). We cut 550- μ m-thick transverse slices that exposed the preBötC at the rostral face (Ruangkittisakul et al., 2010) Slices were perfused with 27°C oxygenated ACSF at 4 ml/min. The K^+ concentration ($[K^+]_0$) was raised to 9 mM in ACSF containing 1.5 mM CaCl₂ and to 5 mM in ACSF containing 1.1 mM CaCl₂. In both cases, the tradeoff of raising $[K^+]_0$ toward a maximum of 9 mM and lowering [Ca²⁺]_o toward a minimum of 1 mM maintained excitability and sustained rhythmic function (Ruangkittisakul et al., 2007). Motor output was recorded from hypoglossal (XII) nerve roots.

Infrared-enhanced differential interference contrast videomicroscopy was performed using a Zeiss fixed-stage microscope with Köhler illumination and a 20×/1.0 numerical aperture (NA) objective (Carl Zeiss).

This work was supported by Grants R01-HL104127-01 (to C.D.N.), The Danish Medical Council, the NovoNordisk Foundation, Den Owensenske Fond, Lægeforeningens forskningsfond, and Agnes og Pouls Friis Fond (to J.C.R.).

Correspondence should be addressed to Dr. Christopher A. Del Negro. Present address (until May 2011): Institut de Neurobiologie Alfred Fessard, Neurobiologie et Développement, Centre National de la Recherche Scientifique, Avenue de la terrasse, 91198 Gif-sur-Yvette, France. Permanent address: McGlothlin-Street Hall, Room 318, The College of William and Mary, Williamsburg, Virginia 23187-8795. E-mail: cadeln@wm.edu.

We performed whole-cell recordings using an EPC-10 amplifier (HEKA). Data were filtered at 1 kHz and acquired at 10 kHz. Standard patch solution contained the following (in mM): 140 K-Gluconate, 10 NaCl, 0.5 CaCl₂, 10 HEPES, 1 EGTA, 2 Mg-ATP, and 0.3 Na3-GTP, pH 7.3. We added 200 μ M Oregon Green 488 BAPTA-2 (OGB-2; K_d = 580 nM) and 50 μ M Alexa 568 hydrazide (Invitrogen) before use.

We performed two-photon imaging using a Ti:Sapphire laser (Spectra Physics) in conjunction with a laser-scanning microscope (Zeiss). Images were acquired by non-descanned detectors at 20 Hz.

We gated image-acquisition protocols via logic pulses from the EPC-10. Gating pulses were recorded at 10 kHz in a separate channel of the chart recorder along with logic pulses returned by the LSM 510 upon the initiation of the first (and last) scans of the time series. Return pulses were used to sync the electrophysiology data with the image stack.

We report 54 neurons recorded with 200 μ M OGB-2. We varied the concentration and the affinity of the Ca²⁺ dye to discern optimal recording conditions, using 400 µM Oregon Green 488 BAPTA-1 (OGB-1; $K_d = 190 \text{ nM}$) and 100–500 μM Oregon Green 488 BAPTA-6F (OGB-6F; $K_d = 3 \mu M$) in preliminary experiments. OGB-6F yielded fast-decaying somatic signals, but the low affinity of the dye diminished dendritic fluorescence. OGB-1 was readily detectable in dendrites, but its long-lasting decay caused spurious temporal summation of fluorescence changes (ΔF). We report OGB-1 experiments wherein ΔF decayed to baseline during each respiratory cycle. The rise time was equally rapid for OGB-1, OGB-2, and OGB-6F.

Alexa 568 hydrazide dye in the patch solution enabled us to visualize the dendritic arbor within 5 min of establishing the whole-cell configuration. Three to six principle dendrites were always visible in preBötC neurons. OGB

diffuses more slowly than Alexa 568 hydrazide, so we waited at least 20 min before measuring dendritic Ca^{2+} activity. After 20 min, we recorded one principle dendrite per 10 min until whole-cell recording conditions deteriorated.

In eight experiments, we imaged the soma and proximal dendrites of preBötC neurons loaded with membrane-permeable Ca²⁺ dye *in vitro*. Slices were loaded with Fluo-8 AM (K_d = 389 nm; AAT Bioquest) via submersion for 60–90 min while bubbling with 95% O₂/5% CO₂. We dissolved 50 μ g of Fluo-8 AM in 50 μ l of DMSO and then combined 20 μ l of the Fluo-8 AM solution with 5 μ l of cremophore EL (Fluka) and 5 μ l of 20% Pluronic F-127 in DMSO. We added the 30 μ l of Fluo-8 AM solution to 1 ml of ACSF with 100 mM mannitol (Sigma-Aldrich) and 100 μ M MK-571 (Enzo Life Sciences). The final Fluo-8 AM concentration was 20 μ M.

We imaged Fluo-8 AM-labeled neurons using an Olympus BX51 microscope with a $63 \times /0.95$ NA objective. Image stacks were acquired at 70 Hz using a CCD camera (Andor Technology).

Images were analyzed with ImageJ (NIH). For each bout, we created a composite image showing the SD of fluorescence through the image stack. This composite shows dendritic regions of interest (ROIs) with large ΔF , which often indicates periodicity. For each ROI, we plotted a histogram of fluorescence measurements and fitted a Gaussian function to determine the mean and SD. The mean reflects baseline fluorescence (F_0) and was used to normalize the ΔF signal, which was subsequently displayed as $\Delta F/F_0$. The SD was used to determine the onset of fluores-



Figure 1. Ca²⁺ measurements in preBötC neurons recorded with QX-314. *A*, Infrared-enhanced differential interference contrast videomicroscopy (*a*), Alexa 568 (*b*), and OGB-2 (*c*) images with on-cell recording (top) and XII output (bottom). *B*, Imaging and electrophysiology from cell in *A*. *C*, Cycle-triggered average of Ca²⁺ and membrane potential from cell in *A*. Vertical line indicates inspiratory onset. Numbered traces correspond to ROIs in the image. *D*, In a different cell, dendritic imaging 91 μ m from the soma using OGB-1. Cycles marked with † show positive dendritic Ca²⁺ latency, cycle marked with * shows negative drive latency. *E*, Cycle marked * displayed at higher sweep speed. *F*, Dendritic and somatic responses to current injection. Broken lines indicate the *T*_{DEN} and *T*_{SOMA}.

cence bursts. The first point in the $\Delta F/F_0$ signal to exceed baseline by ≥ 2 SD was considered statistically significant at p < 0.05 and was defined as the dendritic fluorescence onset time (T_{DEN}). The onset time of the somatic burst (T_{SOMA}) was defined as the absolute time of maximum rate of rise (dV_M/dt) of the voltage, or the maximum rate of rise of somatic Ca²⁺ fluorescence. The inherent disparity in sampling rate and signal-to-noise ratio was the basis for defining T_{DEN} and T_{SOMA} . Cycle-triggered averages of inspiratory bursts were computed from ≥ 7 consecutive cycles based on a threshold-crossing algorithm applied to the XII output. Dendritic Ca²⁺ latency (ΔT) was defined as the difference between T_{DEN} and T_{SOMA} , ($\Delta T = T_{\text{DEN}} - T_{\text{SOMA}}$).

Results

We recorded preBötC interneurons showing evidence of a build-up of presynaptic excitatory input and strong inspiratory bursts, including summating EPSPs, ramp-like depolarization, and spike discharge at a low rate (2-10 Hz) during the expiratory phase, particularly 300-900 ms before XII output. Robust inspiratory bursts experience depolarization block (i.e., voltage-dependent spike inactivation), which attenuates intraburst spiking. We probed for these characteristics, which may indicate rhythmogenic function, during on-cell recording (Figs. 1A, 2A) as selection criteria for subsequent whole-cell recording.



Figure 2. Dendritic Ca²⁺ measurements in preBötC neurons recorded with standard K-gluconate patch solution. *A*, On-cell recording and XII output. *B*, Image of the cell in *A* (Alexa 568). *C*, *D*, Dendritic Ca²⁺ dynamics in ROI₁ (*C*) and ROI₂ (*D*) with whole-cell and XII recordings. Step commands were applied while imaging ROI₁ in *C*. Vertical calibration bars apply to *C*–*F*. *E*, The first cycles from *C* and *D* replotted and superimposed at higher sweep speed. *F*, Second evoked response from *C* replotted at higher sweep speed. *G*, Image of a different preBötC neuron (Alexa 568). *H*, Cycle-triggered average of somatodendritic activity. Imaging sweeps for ROIs 1– 4 with membrane potential and XII output. Asterisks indicate somatic action potentials that evoked Ca²⁺ transients. Arrowheads (top traces) indicate negative dendritic Ca²⁺ latency. *I*–*K*, Ca²⁺ imaging from soma and dendrites. *I*, Epifluorescence image. *J*, Imaging sweeps showing raw and differentiated traces. *K*, Dendritic Ca²⁺ latency versus recording distance for 20 dendritic recording sites. Horizontal error bars indicate the length of the dendritic ROI.

The patch solution contained 2 mM QX-314 to prevent voltage-gated Ca²⁺ channels evoked during action potentials from obscuring synaptic–dendritic Ca²⁺ activity. QX-314 blocks spiking intracellularly, but has no effect on network properties such as recurrent excitation, which can be detected as preinspiratory ramp-like voltage trajectory and temporal summation of EPSPs (Fig. 1*B*). Nevertheless, Ca²⁺ transients in the soma and proximal dendrites (20–50 μ m) coincided with the somatic inspiratory drive potential (Fig. 1*B*, *C*).

We hypothesized that postsynaptic Ca²⁺ accumulation before somatic burst generation might be detectable in more distal dendrites. To quantify the temporal relationship of dendritic Ca²⁺ and somatic inspiratory bursts, we subtracted the onset time of the inspiratory drive potential (T_{SOMA}) from the fluorescence onset time in the dendrite (T_{DEN}) to obtain the dendritic Ca²⁺ latency (ΔT), which we dub drive latency. Figure 1, *D* and *E*, shows dendritic $\Delta F/F_0$ measured 90 µm from the soma. ΔT was <0 in 75% of cycles measured; thus, dendritic Ca²⁺ generally increased before the inspiratory drive potential. The average dendritic Ca²⁺ latency measured -12 ± 8 ms, with a range of -6 to -34 ms (n = 7 cycles at 90 µm), and -28 ± 15 ms at an average distance of 76 ± 15 µm in five dendritic branches (n = 24 cycles in total). These latency times are reliable, even though they are smaller than the sampling interval (49 ms/frame), because T_{DEN} was determined based on the first dendritic measurement to exceed baseline fluorescence by two or more standard deviations; we did not interpolate between points to estimate T_{DEN} . This method of computing ΔT may underestimate the dendritic Ca²⁺ latency because if the sampling rate was higher, a significant increase in dendritic Ca²⁺ fluorescence would be detectable earlier. The dendritic sampling rate has no effect on T_{SOMA} , which is determined by electrophysiology (10 kHz).

 ΔT exceeded zero in two cycles (5 and 28 ms, respectively) (Fig. 1*D*), which may reflect dendritic Ca²⁺ channel activity evoked electrotonically by somatic depolarization. To evaluate this possibility, we applied step-current commands 500 ms after the inspiratory burst, when network synaptic activity is minimal and few spontaneous synaptic potentials can be detected. The step command caused a significant dendritic fluorescence response within 38 ms (Fig. 1*F*), which was commensurate with inspiratory bursts in which ΔT was also positive and the dendritic $\Delta F/F_0$ occurred 5–28 ms after the inspiratory drive potential.

The peak dendritic fluorescence always occurred \sim 200 ms (or more) after the onset of the drive potential, even during respiratory cycles in which initial dendritic Ca²⁺ preceded the somatic

burst ($\Delta T < 0$) (Fig. 1*E*). The dendritic fluorescence similarly peaked ~240 ms after the step command (Fig. 1*F*). These data suggest that the maximum dendritic Ca²⁺ response is attributable to voltage-gated Ca²⁺ channels evoked electrotonically by somatic depolarization.

A caveat is that QX-314 also enhances the amplitude of the inspiratory drive potential (Pace et al., 2007a). AMPA receptors ordinarily evoke Ca^{2+} channels (Pace and Del Negro, 2008), but the QX-314-amplified drive potential may artificially increase Ca^{2+} channel recruitment and cause one to overestimate the synaptically triggered Ca^{2+} activity. Therefore, we subsequently omitted QX-314 from the patch solution and concentrated on measuring dendritic Ca^{2+} latency.

Figure 2, A-F, shows a preBötC neuron recorded with K-gluconate solution. Preinspiratory spiking was robust on-cell (Fig. 2A), so in whole-cell we applied negative bias to prevent intraburst spiking that would otherwise obscure our ability to measure synaptic Ca²⁺ activity. During inspiratory cycles, ΔT measured 87 \pm 13 ms at a site 220 μ m from the cell body (n = 22cycles; ROI_1) (Fig. 2B). The latency at ROI_1 was significantly longer in response to step-current injection through the somatic patch electrode (182 \pm 35 ms, n = 5 pulse injections, p < 0.05) (Fig. $2C_{r}F$). This disparity suggests that a proximal dendritic site triggered Ca²⁺ accumulation at ROI₁ during inspiratory cycles. To test this, we recorded contiguous regions of the dendrite between ROI₁ and the soma. At a site 81 μ m from the cell body, ΔT measured $-90 \pm 14 \text{ ms}$ ($n = 23 \text{ cycles from ROI}_2$) (Fig. 2D, E). Because ROI₁ and ROI₂ were separated by 139 μ m, and T_{DEN} occurred 177 ms later at ROI₁, the data suggest that Ca²⁺ activity originating at or near ROI2 influenced adjacent regions of the dendrite at a rate of \sim 785 μ m/s. We detected Ca²⁺ transients in the dendrite occurring both

before and after the inspiratory burst in four experiments. Figure 2G shows a cycle-triggered average in a representative preBötC neuron. Distal sites such as ROI_1 and ROI_2 (154 and 94 μ m from the soma, respectively) showed Ca²⁺ transients before the inspiratory phase, which propagated orthodromically at a rate in the range of 660–1000 μ m/s. ROI₃ (57 μ m from the soma) showed no evidence of Ca²⁺ activity before the inspiratory burst. Preinspiratory Ca^{2+} transients detected in the soma (ROI₄) were most likely attributable to preinspiratory spiking activity that we could not fully suppress via direct current bias (Fig. 2H, asterisks). These data suggest that not all sites in the dendrite are excitable and that preinspiratory Ca²⁺ activity occurs in specialized hotspots. Peak fluorescence always occurred after the onset of the inspiratory phase, which is equally clear in Figures 1 and 2. Ca²⁺ transients that accompany the inspiratory burst occur synchronously throughout the dendrite, not in hotspots, and always

peaked within ~200 ms of T_{SOMA} (Fig. 2*H*). We measured dendritic Ca²⁺ latency at 20 dendritic ROIs in eight preBötC neurons loaded by simultaneously imaging the dendrites and soma, which circumvents whole-cell dialysis that can affect the Ca²⁺ dynamics in the soma and dendrites. A representative experiment shows ΔT of -34 ms for a dendritic site located 71 μ m from the soma (Fig. 2*I*,*J*). ΔT measured <0 in 17 dendritic sites 24–74 μ m away from the soma and was ~0 in three cases (Fig. 2*K*).

We plotted the entire set of dendritic Ca²⁺ latency measurements for endogenous inspiratory bursts and responses to stepcurrent injection as a function of dendritic recording distance. These data include 36 recordings with K-gluconate patch solution (Fig. 3A, red), 17 recordings with QX-314 included (Fig. 3A, green), 20 different dendritic sites in eight neurons imaged at the



Figure 3. Plots of dendritic Ca²⁺ latency versus recording distance for the entire dataset. *A*, Drive latency for endogenous cycles of respiratory activity (red, K-gluconate recordings; green, QX-314 experiments). Inset reproduces Figure 2 *K*. *B*, Dendritic Ca²⁺ latency in response to step commands (blue). Vertical error bars correspond to SE of ΔT . Horizontal error bars indicate the length of the dendritic ROI.

soma and dendrite (Figs. 2*K*, 3*A*, inset), and Ca²⁺ latency in response to step-current injection from 20 neurons (Fig. 3*B*). Points below the origin reflect dendritic Ca²⁺ activity that precedes inspiratory burst generation. Points above the origin reflect Ca²⁺ activity evoked electrotonically as a result of somatic or local dendritic depolarization. Ca²⁺ activity was detected before somatic burst generation in most experiments. However, at recording sites <60 μ m from the soma, ΔT often exceeded zero, suggesting dendritic Ca²⁺ attributable to the electrotonic effects of somatic depolarization. The step-current injections universally evoked positive dendritic Ca²⁺ latency, as expected.

Discussion

The physiological properties of putatively rhythmogenic preBötC neurons—observed in vivo and in vitro over ~30 years—include a preinspiratory firing pattern and robust inspiratory bursts that experience intraburst depolarization block, which suggests the massive recruitment of inward current during inspiration (Richter, 1982; Rekling et al., 1996; Rekling and Feldman, 1998; Guyenet and Wang, 2001; Stornetta et al., 2003; Hayes and Del Negro, 2007; Rubin et al., 2009; Gray et al., 2010). We examined the Ca²⁺ signaling in dendrites to elucidate how the build-up of excitatory synaptic input evokes burst-generating conductances. In this study, the membrane properties of the presynaptic neurons remain unknown and could include a wide variety of discharge phenotypes, including neurons with bursting-pacemaker properties and intrinsic spiking activity, as well as neurons that participate in recurrent excitatory clusters. However, all the presynaptic neurons share a propensity to increase their activity during the early inspiratory phase.

We recorded preBötC neurons based on preinspiratory firing pattern and depolarization block (Figs. 1*A*, 2*A*), which strongly suggests glutamatergic phenotype (Gray et al., 2010). The other major subpopulations in the preBötC are inhibitory interneurons that display low-amplitude inspiratory bursts (<5 mV) without preinspiratory firing (Kuwana et al., 2006; Winter et al., 2009).

Mironov (2008, 2009) documented dendritic waves of Ca²⁺induced Ca²⁺-release (CICR) propagating at 72 μ m/s, triggered by metabotropic glutamate receptors in preBötC neurons. The ability to generate CICR waves may develop in culture or it may be limited to the proximal dendrite; we never detected similar Ca²⁺ waves in acute rhythmically active slices. We also recorded dendritic Ca²⁺ transients before somatic bursts, but not in every case. Ca²⁺ transients preceded somatic inspiratory bursts in 55 of 74 experiments (74%). In general, the more distal the recording site, the earlier we detected Ca²⁺ signals. What propagates from dendrite to soma: Ca²⁺ or voltage? CICR generally propagates at ~70 μ m/s in dendrites (Augustine et al., 2003). Alternatively, Ca²⁺ may evoke I_{CAN} at or near the synapse to depolarize the soma (and neighboring regions of dendrite) electrotonically. Cable theory shows that voltage changes propagate faster than CICR by orders of magnitude in neuronal dendrites. Since we did not detect Ca²⁺ waves per se, propagation reflects a quotient wherein the numerator is the distance between two recording sites and the denominator is the temporal disparity of burst onset at each site. According to this definition, we recorded propagation rates of 300–700 μ m/s, which is consistent with electrotonic propagation rather than CICR.

Dendritic Ca²⁺ transients originated after or in sync with somatic bursts or drive potentials in 19 of 74 experiments (26%). Here too, the antidromic propagation rates were 300–700 μ m/s. Similar antidromic propagation rates were obtained in response to somatic step-current injections (Fig. 3*B*). The fact that their propagation rate is of similar magnitude (but opposite sign) to cycles with negative dendritic Ca²⁺ latency suggests that electrotonic factors govern the dendritic–somatic interactions in the preBötC in both orthodromic and antidromic responses. The data also suggest that some dendrites may receive very little or no drive at all, but still show Ca²⁺ activity by back propagating electrotonic effects.

We also detected dendritic sites with negative drive latency, whereas neighboring regions of the dendrite, both proximal and distal, showed either no Ca^{2+} activity (ROI₃) (Fig. 2*G*,*H*) or positive drive latency (ROI₁) (Fig. 2*B*,*C*,*E*). These data indicate that synaptic drive leading to postsynaptic Ca^{2+} elevation occurs in hotspots, which presumably indicates points at or near excitatory synaptic contacts. Sites with positive drive latency reflect Ca^{2+} channels activated electrotonically from the soma or neighboring regions of the dendrite, which again suggests electrotonic interactions among the dendrites and soma rather than CICR.

According to the group-pacemaker model, preBötC interneurons mutually interact to elevate postsynaptic Ca²⁺, activate the inward current I_{CAN}, and periodically generate inspiratory bursts (Rekling et al., 1996; Rekling and Feldman, 1998; Mironov, 2008; Rubin et al., 2009). The relative timing of dendritic Ca²⁺ transients and somatic voltage changes is one measurable link between network-mediated synaptic drive and postsynaptic mechanisms that activate I_{CAN} (Morgado-Valle et al., 2008). Although these data affirm the key prediction that postsynaptic Ca²⁺ transients be detectable before somatic bursts, they do not prove that respiratory rhythm is group-pacemaker driven. Nor can these data rule out alternative models that emphasize voltagedependent conductances such as persistent Na⁺ current in rhythm generation (Koizumi and Smith, 2008). Regardless of whether emergent network properties constitute the core mechanism of respiratory rhythmogenesis, as the group-pacemaker model postulates, the synaptic activation of I_{CAN} contributes substantially to inspiratory burst generation (Johnson et al., 2007; Pace et al., 2007b) and is relevant to understanding the neural origins of respiratory rhythm.

We conclude that synaptic input typically triggers Ca²⁺ accumulation in dendrites before burst generation. Our measurements are inconsistent with regenerative Ca²⁺ propagation from dendrite to soma, which is a relatively slow process (\sim 70 µm/s) and unlikely to promote synaptic integration in the context of network rhythms. For example, consider an excitatory input 250 µm from the cell body. A CICR-dependent Ca²⁺ wave would require 3.5 s to influence the soma. In slices or en bloc preparations *in vitro*, the typical respiratory period is \sim 5 s. Synaptic inputs located 500 μ m (or farther) from the soma would be impractically sited to influence somatic integration during a typical respiratory cycle. Moreover, the slow propagation rate of CICR would prevent spatial and temporal summation of excitatory synaptic inputs that occur simultaneously at multiple positions in the dendritic arbor. Instead, synaptic inputs may elevate Ca²⁺, evoke I_{CAN} , and depolarize the dendrite locally. These hotspots could thus act electrotonically to influence other regions of the dendrite and the soma in support of inspiratory burst generation. With propagation rates of $300-700 \,\mu$ m/s, it would be reasonable for multiple dendritic sites to undergo temporal and spatial summation of inputs during a typical respiratory cycle in vitro, and even during breathing cycles in vivo in which cycle period is typically <1 s in rodents.

References

- Augustine GJ, Santamaria F, Tanaka K (2003) Local calcium signaling in neurons. Neuron 40:331–346.
- Feldman JL, Del Negro CA (2006) Looking for inspiration: new perspectives on respiratory rhythm. Nat Rev Neurosci 7:232–242.
- Gray PA, Hayes JA, Ling GY, Llona I, Tupal S, Picardo MC, Ross SE, Hirata T, Corbin JG, Eugenín J, Del Negro CA (2010) Developmental origin of preBötzinger complex respiratory neurons. J Neurosci 30:14883–14895.
- Guyenet PG, Wang H (2001) Pre-Bötzinger neurons with preinspiratory discharges "in vivo" express NK1 receptors in the rat. J Neurophysiol 86:438-446.
- Hayes JA, Del Negro CA (2007) Neurokinin receptor-expressing preBötzinger complex neurons in neonatal mice studied in vitro. J Neurophysiol 97:4215–4224.
- Johnson SM, Wiegel LM, Majewski DJ (2007) Are pacemaker properties required for respiratory rhythm generation in adult turtle brain stems in vitro? Am J Physiol Regul Integr Comp Physiol 293:R901–R910.
- Koizumi H, Smith JC (2008) Persistent Na+ and K+-dominated leak currents contribute to respiratory rhythm generation in the pre-Bötzinger complex in vitro. J Neurosci 28:1773–1785.
- Kuwana S, Tsunekawa N, Yanagawa Y, Okada Y, Kuribayashi J, Obata K (2006) Electrophysiological and morphological characteristics of GABAergic respiratory neurons in the mouse pre-Botzinger complex. Eur J Neurosci 23:667–674.
- Mironov S (2009) Respiratory circuits: function, mechanisms, topology, and pathology. Neuroscientist 15:194–208.
- Mironov SL (2008) Metabotropic glutamate receptors activate dendritic calcium waves and TRPM channels which drive rhythmic respiratory patterns in mice. J Physiol 586:2277–2291.
- Morgado-Valle C, Beltran-Parrazal L, DiFranco M, Vergara JL, Feldman JL (2008) Somatic Ca2+ transients do not contribute to inspiratory drive in preBötzinger complex neurons. J Physiol 586:4531–4540.
- Pace RW, Del Negro CA (2008) AMPA and metabotropic glutamate receptors cooperatively generate inspiratory-like depolarization in mouse respiratory neurons in vitro. Eur J Neurosci 28:2434–2442.
- Pace RW, Mackay DD, Feldman JL, Del Negro CA (2007a) Role of persistent sodium current in mouse preBotzinger complex neurons and respiratory rhythm generation. J Physiol 580:485–496.
- Pace RW, Mackay DD, Feldman JL, Del Negro CA (2007b) Inspiratory bursts in the preBötzinger complex depend on a calcium-activated nonspecific cationic current linked to glutamate receptors. J Physiol 582:113–125.
- Peña F, Parkis MA, Tryba AK, Ramirez JM (2004) Differential contribution of pacemaker properties to the generation of respiratory rhythms during normoxia and hypoxia. Neuron 43:105–117.
- Ramirez JM, Tryba AK, Peña F (2004) Pacemaker neurons and neuronal networks: an integrative view. Curr Opin Neurobiol 14:665–674.
- Rekling JC, Feldman JL (1998) PreBötzinger complex and pacemaker neurons: hypothesized site and kernel for respiratory rhythm generation. Annu Rev Physiol 60:385–405.
- Rekling JC, Champagnat J, Denavit-Saubi é M (1996) Electroresponsive

properties and membrane potential trajectories of three types of inspiratory neurons in the newborn mouse brain stem in vitro. J Neurophysiol 75:795–810.

- Richter DW (1982) Generation and maintenance of the respiratory rhythm. J Exp Biol 100:93–107.
- Ruangkittisakul A, Secchia L, Bornes TD, Palathinkal DM, Ballanyi K (2007) Dependence on extracellular Ca2+/K+ antagonism of inspiratory centre rhythms in slices and en bloc preparations of newborn rat brainstem. J Physiol 584:489–508.
- Ruangkittisakul A, Panaitescu B, Ballanyi K (2010) K+ and Ca2+ dependence of inspiratory-related rhythm in novel "calibrated" mouse brainstem slices. Respir Physiol Neurobiol. Advance online publication. Retrieved September 15, 2010. doi:10.1016/j.resp.2010.09.004.

- Rubin JE, Hayes JA, Mendenhall JL, Del Negro CA (2009) Calciumactivated nonspecific cation current and synaptic depression promote network-dependent burst oscillations. Proc Natl Acad Sci USA 106:2939–2944.
- Stornetta RL, Sevigny CP, Guyenet PG (2003) Inspiratory augmenting bulbospinal neurons express both glutamatergic and enkephalinergic phenotypes. J Comp Neurol 455:113–124.
- Thoby-Brisson M, Ramirez JM (2001) Identification of two types of inspiratory pacemaker neurons in the isolated respiratory neural network of mice. J Neurophysiol 86:104–112.
- Winter SM, Fresemann J, Schnell C, Oku Y, Hirrlinger J, Hülsmann S (2009) Glycinergic interneurons are functionally integrated into the inspiratory network of mouse medullary slices. Pflugers Arch 458:459–469.