

2012

## Polymorphisms of the acid sensing histidine kinase gene *arsS* in *Helicobacter pylori* populations from anatomically distinct gastric sites

Daniel R. Hallinger  
*William & Mary*

Mark H. Forsyth  
*William & Mary*, mhfors@wm.edu

Judith Romero-Gallo

Richard M. Peek

Richard M. Peek

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

---

### Recommended Citation

Hallinger, Daniel R.; Forsyth, Mark H.; Romero-Gallo, Judith; Peek, Richard M.; and Peek, Richard M., Polymorphisms of the acid sensing histidine kinase gene *arsS* in *Helicobacter pylori* populations from anatomically distinct gastric sites (2012). *Microbial Pathogenesis*, 53(6-May), 227-233.  
10.1016/j.micpath.2012.08.002

This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).



## Polymorphisms of the acid sensing histidine kinase gene *arsS* in *Helicobacter pylori* populations from anatomically distinct gastric sites

Daniel R. Hallinger<sup>a</sup>, Judith Romero-Gallo<sup>b</sup>, Richard M. Peek Jr.<sup>b,c,d</sup>, Mark H. Forsyth<sup>a,\*</sup>

<sup>a</sup> Department of Biology, The College of William and Mary, Integrated Science Center 3051, Williamsburg, VA 23185, USA

<sup>b</sup> Department of Gastroenterology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

<sup>c</sup> Departments of Medicine and Cancer Biology, Vanderbilt University Medical Center, Nashville, TN 37232, USA

<sup>d</sup> Veterans Affairs Tennessee Valley Healthcare System, Nashville, TN 37212, USA

### ARTICLE INFO

#### Article history:

Received 20 July 2012

Accepted 14 August 2012

Available online 30 August 2012

#### Keywords:

*Helicobacter pylori*

*arsS*

Polymorphism

Homopolymeric tract

### ABSTRACT

Phase variation is frequently utilized by bacterial species to affect gene expression such that phenotypic variants are maintained within populations, ensuring survival as environmental or host conditions change. Unusual among *Helicobacter pylori* phase variable or contingency genes is *arsS*, encoding a sensory histidine kinase involved in the acid acclimation of the organism. The presence of a 3' homopolymeric cytosine tract of variable length in *arsS* among *Helicobacter pylori* strains allows for the expression of various functional ArsS isoforms, differing in carboxy-terminal protein domains. In this study, we analyzed this 3' *arsS* region via amplified fragment length polymorphism (AFLP) and sequencing analyses for *H. pylori* populations from 3 different gastric sites of 12 patients. Our data indicate the presence of multiple *arsS* alleles within each population of *H. pylori* derived from the gastric antrum, cardia, or corpus of these patients. We also show that *H. pylori*, derived from the same anatomical site and patient, are predicted to express multiple ArsS isoforms in each population investigated. Furthermore, we identify a polymorphic deletion within *arsS* that generates another alternate ArsS C-terminal end. These findings suggest that four C-terminal variations of ArsS adds to the complexity of the ArsRS acid adaptation mechanism as a whole and may influence the ability of *H. pylori* to persist in the gastric niche for decades.

© 2012 Elsevier Ltd. All rights reserved.

### 1. Introduction

*Helicobacter pylori* is a spiral shaped, Gram-negative, micro-aerophilic, highly motile bacterium [1–4]. It is a neutrophile that is ecologically restricted to the mucus layer overlaying the human gastric epithelium and an etiologic agent of peptic ulcer disease, chronic active gastritis, and non-ulcer dyspepsia [1–5]. When untreated, infection may persist for decades, increasing the risk for the development of gastric malignancies such as mucosal-associated lymphoid tissue (MALT) lymphoma and gastric adenocarcinoma [5–7]. *H. pylori* encounters marked pH fluctuations during its decades long persistence within the gastric environment as pH levels in the stomach lumen can vary from 5 to 1 depending if the host is in a fed or fasting state [7,8].

Many studies show that *H. pylori* possesses unique acid acclimation mechanisms to maintain its periplasmic and cytoplasmic pH levels near neutrality [4,7,9–11]. These buffering mechanisms

allow *H. pylori* to withstand severe acid shock and to grow at moderately low pH levels [2,7]. One important acid acclimation mechanism is the two-component signal transduction system ArsRS, which is composed of a sensory histidine kinase, ArsS, and its cognate response regulator, ArsR [2,7,12]. In response to acidic conditions, ArsS dimerizes to promote autophosphorylation at a conserved histidine residue within its C-terminal transmitter domain [2,13]. Phosphorylated ArsS can serve as a phosphoryl donor for ArsR, which subsequently regulates gene expression [2,12]. Interestingly, *arsS* mutants are viable while *arsR* mutants are not, suggesting an essential function for non-phosphorylated ArsR [2,12,14].

*H. pylori* strains exhibit remarkable genetic variation and one mechanism promoting this genetic heterogeneity is slipped-strand mispairing during DNA replication [15,16]. Slipped-strand mispairing occurs more frequently at repetitive sequences, resulting in misalignment of template and nascent DNA strands [15,16]. Thus, insertion and deletion mutations (indels) are more frequent in sequence repeats of single nucleotides (homopolymeric tracts) or multiple nucleotides (heteropolymeric tracts). Short sequence repeats and the associated indels result in phase variation,

\* Corresponding author. Tel.: +1 757 221 2489.

E-mail address: [mhfors@wm.edu](mailto:mhfors@wm.edu) (M.H. Forsyth).

a mechanism utilized by bacteria for transcriptional or translational regulation in response to alterations in environmental or host conditions [17].

*arsS* exhibits unusual sequence variation due to a 3' homopolymeric tract that differs in the number of repeated cytosine nucleotides among *H. pylori* strains [12,18,19]. Different *arsS* alleles possessing cytosine repeat length polymorphisms are predicted to be expressed as alternate, yet functional, ArsS isoforms *in vitro* and, thus far, these ArsS variants have been considered to be strain specific [12]. In this study, we hypothesized that slipped-strand mispairing at the *arsS* cytosine tract would lead to the generation of alternate alleles within populations of *H. pylori*. Here, we demonstrate substantial polymorphic variation in the 3' *arsS* homopolymeric cytosine tract of all *H. pylori* populations examined. We also determined that a predominant ArsS isoform is predicted in *H. pylori* populations from different patients and, in some cases, different populations of the same patient. We additionally document a previously unidentified deletion polymorphism in the 3' *arsS* region, leading to a novel alternate ArsS C-terminal domain.

## 2. Materials and methods

### 2.1. Patients, bacterial populations, and growth conditions

Populations of clinical *H. pylori* strains used in this study were cultured from gastric biopsies harvested from the antrum, cardia, and corpus (A, Ca, and Cs) of 12 patients (patients 1–12) who were scheduled for endoscopy at the Veterans Administration Hospital in Nashville, Tennessee. All but one patient was male, and patient age ranged from 50 to 74 years with a median age of 62.5 years. *H. pylori* populations were collected by sweep culture of all resulting *H. pylori* colonies. To minimize selection of mutants more fit to *in vitro* conditions, *H. pylori* populations were generated from cryopreserved, low passage stocks for each DNA extraction. All populations were cultured on Trypticase Soy Agar II plates with 5% sheep blood (BD) and incubated in a humidified environment at 37 °C and 5% CO<sub>2</sub>. Specific information on patients' age, gender, and clinical findings as well as documented virulence factor genes *cagA*, *vacA*, and *iceA* status of the *H. pylori* isolates is located in Supplemental Table 1.

### 2.2. *arsS* region AFLP

For each gastric population, 3 separate genomic DNA isolations were performed via CTAB extraction as previously described [20,21]. Each DNA extraction was used in triplicate to amplify a ~300 base pair region encoding the 3' end of *arsS*, including the homopolymeric cytosine tract, and part of the 5' end of downstream gene *hemB*. Amplicons were generated according to manufacturer's protocol with Expand High Fidelity PCR kit (Roche) with 6-carboxymethyl fluorescein (FAM)-labeled primer *arsS* F-1 FAM (CTTCTAACCCAGCCAAGCCCATGG) and unlabeled *hemB* R-1 (CGTGCTTCGTAATCTTCTCAATCG). Amplification conditions consisted of a hot start at 94°C/2 min, 30 cycles of 94°C/30 s, 60°C/30 s, and 72°C/30 s, and a final extension at 72°C/7 min. PCR was performed 3 times for each DNA extraction for a total of 9 PCR reactions for each population.

One µL of 1:100 diluted PCR samples was added to 12 µL Hi-Di Formamide (Applied Biosystems, ABI) plus 0.25 µL GeneScan ROX500 Size Standard (ABI) in 96-well plates. Samples were denatured at 95 °C for 3 min and analyzed with an ABI3130 Genetic Analyzer. Amplicon frequency was quantified in Microsoft Excel with data generated from GeneMapper version 4.0 (ABI). For each PCR run, individual area under the curve for each amplicon was summed to determine total area under the curve for all amplicons.

Individual amplicon frequency was calculated by dividing individual area by total area. To integrate AFLP data from the 9 PCR runs for each gastric population, frequencies of corresponding amplicons were used to calculate average and standard error of the mean (SEM).

Each amplicon length was considered to be representative of an individual *arsS* allele. Amplicons that differed in length by multiples of 3bp were considered to be alleles encoding in the same *arsS* open reading frame (ORF). Thus, allele (amplicon) frequency data was used to calculate ORF frequency by taking the sum of the frequencies of amplicons that differed in length by multiples of 3bp. ORF SEM was also calculated with corresponding allele (amplicon) SEM values.

### 2.3. Statistics

To determine the significance of different amplicon lengths within individual populations (i.e. antral, cardia, or corpus populations), one-way ANOVAs with Dunnett post-tests were performed (GraphPad Prism 5.0). The significance of amplicon length between these individual gastric regional populations of the same patient was determined with one-way ANOVAs with Bonferroni post-tests (GraphPad Prism 5.0). One-way ANOVAs with Bonferroni post-tests were also used to calculate significance for the predicted ORFs within and among populations of an individual patient. Thus, comparisons were made with *H. pylori* antrum, cardia, and corpus populations of individual patients. Comparisons among *H. pylori* populations of different patients were not considered.

### 2.4. *arsS* region sequencing

Unlabeled *arsS* region amplicons were generated with primers *arsS* F-1 and *hemB* R-1 as described above. Amplicons were cloned using pCR-Blunt II-TOPO (Invitrogen) according to manufacturer's protocols. Plasmid DNA was purified with QIAprep Spin Miniprep Kit or IBI High-Speed Plasmid Mini Kit and 10 cloned 3' *arsS* amplicons from each *H. pylori* gastric population were sequenced.

Both strands of each cloned amplicon were sequenced using Big Dye Terminator version 3.1 (ABI). In general, 20 µL sequencing reactions were prepared with 1X Big Dye Terminator v1.1/v3.1 Sequencing Buffer, 0.25X Big Dye Terminator v3.1 Sequencing Reaction Mix, 10 µM of T7 (20mer; Promega) or Sp6 (19mer; Promega) promoter primer, 10 µL of purified plasmid DNA, and sdH<sub>2</sub>O. Reactions were achieved with thermal cycling conditions of 94°C/5 min and 26 cycles of 94°C/45 s, 50°C/30 s, and 60°C/4 min. Reactions were purified with Performa Dye Terminator Removal (DTR) Gel Filtration Cartridges (Edge Biosystems) according to manufacturer's protocol. Purified sequencing reactions were vacuum-dried and resuspended in 12 µL Hi-Di Formamide, denatured for 3 min at 95 °C, and analyzed with an ABI3130 Genetic Analyzer. SequencingAnalysis version 5.2 (ABI) software was used for base calling and sequencing analyses were performed with MacVector version 9.0 (Accelrys, Inc.), 4Peaks version 1.7.2 (Mek&Tosj.com), and WebLogo version 3.0 (Threplusone.com).

Using nine complete *H. pylori* genomes accessible from National Center for Biotechnology Information (NCBI) databases as of July 2010, we determined that no two completely sequenced *H. pylori* strains were identical in nucleotide sequence within the 3' *arsS* region amplified in our analyses [18,19,22–28]. Based on sequence alignments for the amplified *arsS* region, the two most highly related *arsS* sequences are *H. pylori* strains 26695 and HPAG1. These strains differed by only two nucleotide substitutions when disregarding length differences in their *arsS* cytosine tracts [19,27]. To make our analyses more stringent, we considered one base pair differences, disregarding polycytosine tract length polymorphisms,

as a metric to denote individual *H. pylori* sequence types within each gastric population.

### 3. Results

#### 3.1. *arsS* polymorphisms

Amplified fragment length polymorphism (AFLP) PCR was utilized to determine whether amplicons of variable base pair (bp) length could be generated from the 3' *arsS* region of *H. pylori* populations isolated from various sites within the stomachs of 12 patients. Resulting AFLP data show six to nine amplicons of variable length from each of the populations investigated in this study (Table 1). The 3' *arsS* region of interest was predicted to be approximately 300 bp in length and AFLP data showed that amplicons of 296 bp through 307 bp were generated from the various populations. We quantified each amplicon as a proportion of the total amplicon yield. This indicated that 33 of 36 *H. pylori* populations produced an amplicon whose quantity was significantly greater than other amplicons from the same gastric

population from the same patient ( $p < 0.01$ ) (Table 1). Amplicons of 302 bp, 301 bp, and 300 bp predominated in 33%, 25%, and 19.4% of the populations, respectively.

*H. pylori* populations from different gastric sites of patients 1, 2, 3, 8, 11, and 12 produced a common quantitatively predominant amplicon from all gastric regions that was significantly different than any other amplicon length ( $p < 0.0001$ ) for five of these six patients (Table 1). However, one of the three gastric *H. pylori* populations of patients 4, 5, 6, 7, 9, and 10 produced a quantitatively predominant amplicon of different length when compared to the predominant amplicon length for the other two *H. pylori* gastric populations of the individual patients. Difference in predominant amplicon length among populations of the same patient was significant ( $p < 0.01$ ) in 5 of these 6 patients (Table 1).

For each of the 36 *H. pylori* gastric populations examined in this study, ten cloned amplicons containing 3' *arsS* region amplicons were generated to confirm that AFLP results were due to differences in homopolymeric cytosine tract length. Our data show that at least two to as many as six different cytosine tract lengths could be documented from the amplicons generated from each gastric *H. pylori*

**Table 1**

Summary of AFLP and sequencing data for 3' *arsS* region amplicons generated from each *H. pylori* population of this study.

Patient designation	Strain designation	Gastric site	<i>arsS</i> amplicon AFLP data					<i>arsS</i> amplicon sequence data		
			Total # of amplicon lengths detected	Amplicon length range (bp)	Predominant amplicon length (bp)	Significance of predominant amplicon <sup>a</sup>	Common predominant amplicon <sup>b</sup>	Total # of cytosine tract lengths detected	Lengths of cytosine tracts (bp) detected	Presence of thymine deletion <sup>c</sup>
Patient 1	B215	Antrum	7	298–304	301	***	Yes	2	12–13	Yes
		Cardia	7	297–303	301	***		2	12–13	
		Corpus	6	298–303	301	***		3	12–14	
Patient 2	B221	Antrum	6	299–304	302	***	Yes	3	11–13	No <sup>d</sup>
		Cardia	6	299–304	302	***		5	10–14	
		Corpus	6	299–304	302	***		4	11–14	
Patient 3	B253	Antrum	7	297–303	301	***	Yes	4	11–14	Yes
		Cardia	7	297–303	301	***		3	11–13	
		Corpus	6	298–303	301	***		4	11–14	
Patient 4	B256	Antrum	7	299–305	303	***	No	4	12–15	No
		Cardia	7	299–305	303	***		3	12–14	
		Corpus	7	299–305	302	***		4	11–14	
Patient 5	B266	Antrum	6	297–302	300	***	No	4	10–13	Yes
		Cardia	6	297–302	300	***		4	11–13, 15	
		Corpus	8	296–303	301	ns		4	11–14	
Patient 6	B268	Antrum	6	299–304	302	***	No	2	12–13	No
		Cardia	7	299–305	303	***		11, 13–14		
		Corpus	6	299–304	302	***		3	11–13	
Patient 7	B284	Antrum	8	299–306	304	**	No	5	13–17	Yes
		Cardia	9	298–306	303	***		6	12–17	
		Corpus	9	299–307	304	***		5	12–16	
Patient 8	B292	Antrum	7	298–304	302	***	Yes	5	10–14	Yes
		Cardia	8	298–305	302	***		4	12–15	
		Corpus	7	298–304	302	***		4	12–15	
Patient 9	B294	Antrum	7	296–302	299	***	No	5	9, 11–14	Yes
		Cardia	6	297–302	300	***		3	11–13	
		Corpus	6	297–302	300	***		3	11–13	
Patient 10	B295	Antrum	9	298–306	301	ns	No	6	8, 11–15	Yes
		Cardia	7	300–306	304	***		5	10, 11, 14–16	
		Corpus	6	298–303	301	***		2	12–13	
Patient 11	B300	Antrum	6	297–302	300	***	Yes	3	11–13	Yes
		Cardia	7	297–303	300	ns		3	12–14	
		Corpus	6	297–302	300	***		3	11–13	
Patient 12	B301	Antrum	6	299–304	302	***	Yes	5	9–13	No
		Cardia	6	299–304	302	***		3	9, 12–13	
		Corpus	7	298–304	302	***		4	10–13	

<sup>a</sup> The statistical significance of the predominant amplicon versus other amplicon lengths was tested for individual *H. pylori* populations via one-way ANOVAs with Dunnett post-tests; \*,  $p < 0.05$ , \*\*,  $p < 0.01$ , \*\*\*,  $p < 0.001$ , \*\*\*\*,  $p < 0.0001$ .

<sup>b</sup> *H. pylori* populations that shared the same predominant amplicon within a single patient were designated as having a common predominant amplicon. Commonality is only valid for populations within patients.

<sup>c</sup> Sequenced amplicons from each *H. pylori* population of individual patients either encoded for a functional *arsS* stop codon (no deletion) or had a deletion of a thymine nucleotide within the *arsS* stop codon (deletion).

<sup>d</sup> The thymine deletion was detected in a single sequence derived from the cardia *H. pylori* population of patient 2. The other 29 sequences from the populations of this patient were negative for the deletion.

population (Table 1). Amplicons from 88.9% of the populations possessed three or more different cytosine tract lengths. Only two different cytosine tract lengths were detected in four of the 36 gastric *H. pylori* populations. In total, sequence data indicated that homopolymeric tract lengths ranged from 8 to 17 tandem cytosine nucleotides, depending on the source population (Table 1). Cytosine tract lengths of 13, 12, and 14 represented 40%, 24.7%, and 15.8% of all sequenced amplicons, respectively. Due to the relatively small number of sequences analyzed, relationships between AFLP amplicon length and sequenced cytosine tract length were not considered. In sum, AFLP and sequencing analyses suggest that variant 3' *arsS* homopolymeric tract lengths may be generated *in vivo* and suggested that predominant amplicon lengths could differ among *H. pylori* populations colonizing different regions of the stomach (Table 1).

Our sequence analysis of 360 cloned amplicons generated from the 36 *H. pylori* populations revealed a thymine deletion in 241 of the cloned sequences. This thymine deletion is not associated with a repeated sequence (Fig. 1). This deletion eliminates the stop codon of one predicted *arsS* reading frame. Interestingly, 240 of these 241 sequences were found in the *H. pylori* populations of 8 patients (Table 1). Only one of the remaining 120 sequenced amplicons from populations of the other 4 patients possessed this stop codon thymine deletion. This sequence was generated from the cardia population of patient 2 and the deletion was not observed in sequences from the other 29 amplicons generated from populations from that patient. Outside this study, the NCBI sequence database indicates that *H. pylori* strains B38 and J99 are the only sequenced strains to possess this thymine deletion. Thus, differences in homopolymeric cytosine tract length as well as the presence or absence of this deletion appear to be common polymorphisms for this region of *arsS*.


### 3.2. Variant *ArsS* C-terminal regions

AFLP and sequence data indicated that the *arsS* amplicons could be considered as different alleles distinct in their 3' regions due to differences in homopolymeric cytosine tract length and the presence/absence of a thymine deletion. Cytosine tract length differences and presence/absence of the thymine deletion can shift the *arsS* open reading frame (ORF) to alter the *ArsS* C-terminal amino

acid sequence. Thus, AFLP amplicon lengths that varied by multiples of 3bp were presumed to be of the same *arsS* ORF and used to calculate total ORF frequencies. For each population, *arsS* amplicons of 301 bp, 302 bp, and 303 bp were designated to be of ORFs 1, 2, and 3, respectively. These variant alleles are predicted to express *ArsS* isoforms differing in their carboxy-terminal domains. The ORFs of all other amplicons were determined with the multiple of three bp parameter. A predominant *ArsS* ORF was apparent for each *H. pylori* population and ORFs 1, 2, and 3 were predominant in 33.3%, 41.7%, and 25% of all populations.

A predominant *ArsS* isoform was significant ( $p < 0.05$ ) for 29 of 36 populations, which may suggest a potential selective pressure on *ArsS* isoform expression dependent on the gastric environment. Overall, five of the 12 patients had *H. pylori* populations predicted to express a uniform predominant ORF regardless of the gastric site of origin. However, seven patients had *H. pylori* populations encoding for different predominant ORFs among their gastric regions, and the predominance difference amongst regions was significant ( $p < 0.05$ ) for the populations of five of these seven patients (Table 2). We speculate that local conditions in the cardia, corpus, or antrum of these five patients may differ sufficiently such that selection has allowed populations with differing *ArsS* isoforms to attain numeric predominance within the stomach of individual patients (Table 2).

As the ORF of *arsS* is shifted due to differences in homopolymeric cytosine tract length and the presence/absence of the thymine deletion, the translated *ArsS* amino acid sequence will be affected. *arsS* ORFs that differ by three bp multiples of cytosines in the homopolymeric tract should only differ by the number of proline amino acids translated in *ArsS*. However, differences in cytosine tract length paired with the presence/absence of the thymine deletion can considerably alter the translated *ArsS* C-terminal region. With the 3' *arsS* polymorphisms now documented for each *H. pylori* population of this study, we performed *in silico* translation of the 360 cloned amplicon sequences to demonstrate that the polymorphic differences lead to the translation of variant *ArsS* C-terminal ends. Again, because of the relatively small number of sequenced amplicons, comparisons between AFLP-predicted *arsS* ORFs and sequence-predicted *ArsS* C-terminal ends were not considered. Although *ArsS* C-terminal domains vary from two to 31 amino acids past the poly proline regions, for simplicity, alternate



**Fig. 1.** *arsS* polymorphisms and the *arsRS* operon. The *arsRS* operon consists of *arsR*, *arsS*, *hemB*, and *hp0162*, a gene encoding a hypothetical protein. DNA sequence of the 3' terminus of *arsS* and the 5' region of *hemB* is shown. The *arsS* homopolymeric cytosine tract and associated stop codons of the different *arsS* ORFs are indicated. As the homopolymeric cytosine tract varies in length ( $C_n$ ), different C-terminal regions of *ArsS* (PKI\*, LWG\*, EKQ\*, and SND\*) are encoded due to different ORFs. Many of the *H. pylori* populations in this study possessed a thymine deletion in the third stop codon (in italics), allowing for the alternate SND\* C-terminal region to be predicted. The predicted stop codon of this C-terminal end is located within *hemB* gene.

**Table 2**Summary of AFLP and sequencing data for predicted *arsS* ORFs from each *H. pylori* population of this study.

Patient designation	Strain designation	Gastric site	AFLP <i>arsS</i> amplicon data				Sequenced <i>arsS</i> amplicon data	
			ORF range	Predominant ORF	Significance of predominant ORF <sup>a</sup>	Common predominant ORF <sup>b</sup>	C-terminal regions detected	Thymine deletion <sup>c</sup>
Patient 1	B215	Antrum	1–3	1	****	Yes	LWG* SND*	Yes
		Cardia	1–3	1	****		LWG* SND*	
		Corpus	1–3	1	****		PKI* LWG* SND*	
Patient 2	B221	Antrum	1–3	2	****	Yes	PKI* LWG* EKQ*	No <sup>d</sup>
		Cardia	1–3	2	****		PKI* LWG* EKQ*	
		Corpus	1–3	2	****		PKI* LWG* EKQ*	
Patient 3	B253	Antrum	1–3	1	****	Yes	PKI* LWG* SND*	Yes
		Cardia	1–3	1	****		PKI* LWG* SND*	
		Corpus	1–3	1	****		PKI* LWG* SND*	
Patient 4	B256	Antrum	1–3	3	****	No	PKI* LWG* EKQ*	No
		Cardia	1–3	3	****		PKI* LWG* EKQ*	
		Corpus	1–3	2	ns		PKI* LWG* EKQ*	
Patient 5	B266	Antrum	1–3	3	*	No	PKI* LWG* SND*	Yes
		Cardia	1–3	3	**		PKI* LWG* SND*	
		Corpus	1–3	1	ns		PKI* LWG* SND*	
Patient 6	B268	Antrum	1–3	2	****	No	PKI* LWG* EKQ*	No
		Cardia	1–3	3	****		PKI* LWG*	
		Corpus	1–3	2	****		PKI* LWG* EKQ*	
Patient 7	B284	Antrum	1–3	1	***	No	PKI* LWG* SND*	Yes
		Cardia	1–3	2	ns		PKI* LWG* SND*	
		Corpus	1–3	1	****		PKI* LWG* SND*	
Patient 8	B292	Antrum	1–3	2	****	Yes	PKI* LWG* SND*	Yes
		Cardia	1–3	2	****		PKI* LWG* SND*	
		Corpus	1–3	2	****		PKI* LWG* SND*	
Patient 9	B294	Antrum	1–3	2	ns	No	PKI* LWG* SND*	Yes
		Cardia	1–3	3	****		PKI* LWG* SND*	
		Corpus	1–3	3	****		PKI* LWG* SND*	
Patient 10	B295	Antrum	1–3	2	ns	No	PKI* LWG* SND*	Yes
		Cardia	1–3	1	ns		PKI* LWG* SND*	
		Corpus	1–3	1	***		LWG* SND*	
Patient 11	B300	Antrum	1–3	3	****	Yes	PKI* LWG* SND*	Yes
		Cardia	1–3	1	ns		PKI* LWG* SND*	
		Corpus	1–3	3	****		PKI* LWG* SND*	
Patient 12	B301	Antrum	1–3	2	****	Yes	PKI* LWG* EKQ*	No
		Cardia	1–3	2	****		LWG* EKQ*	
		Corpus	1–3	2	****		PKI* LWG* EKQ*	

<sup>a</sup> The statistical significance of the predominant amplicon versus other amplicon lengths was tested for individual *H. pylori* populations via one-way ANOVAs with Dunnett post-tests; \*,  $p < 0.05$ , \*\*,  $p < 0.01$ , \*\*\*,  $p < 0.001$ , \*\*\*\*,  $p < 0.0001$ .

<sup>b</sup> *H. pylori* populations that shared the same predominant ORF within a single patient were designated as having a common predominant ORF. Commonality is only valid for populations within patients.

<sup>c</sup> Sequenced amplicons from each *H. pylori* population of individual patients either encoded for a functional *arsS* stop codon (no deletion) or had a deletion of a thymine nucleotide within the *arsS* stop codon (deletion).

<sup>d</sup> The thymine deletion was detected in a single sequence derived from the cardia *H. pylori* population of patient 2. The other 29 sequences from the populations of this patient were negative for the deletion.

C-terminal domains are abbreviated as the last 3 predicted amino acids prior to their respective stop codons, symbolized by \*.


In agreement with NCBI *arsS* ORF designations, we designated the 0 and -1 *arsS* ORFs as ArsS C-terminal ends PKI\* and LWG\* (Fig. 1). Since the polymorphic thymine deletion was located within the stop codon of the +1 *arsS* ORF, two alternative C-terminal ends are predicted (Fig. 1). These ends are designated as SND\* or EKQ\* depending on the thymine deletion presence or absence, respectively. The stop codon for the SND\* C-terminal end overlaps the sequence encoding the downstream gene *hemB* (Fig. 1). The peptide sequences of the alternative C-terminal ends vary in length by two to as many as 31 amino acids beyond the final proline encoded by the homopolymeric cytosine tract. These four alternate ArsS C-terminal ends also differ in the composition of their amino acid sequences (Fig. 2). However, the predicted EKQ\* and SND\* ends appear to be homologous until the SND\* sequence is extended due to the thymine deletion in the EKQ\* stop codon (Fig. 2).

From amplicon sequence data, the PKI\* ArsS C-terminal end was predicted to be expressed by 31 of the 36 gastric *H. pylori* populations, but was not detected in 5 individual gastric regional populations from four patients (Table 2). The LWG\* C-terminal end

was predicted in all *H. pylori* populations investigated. The EKQ\* C-terminal end was predicted in 11 populations of four patients and the SND\* alternative +1 end was detected in 24 populations from the remaining eight patients (Table 2). PKI\* and LWG\* were predicted to be expressed by 93 (25.8%) and 158 (43.9%) of the 360 cloned 3' *arsS* amplicons, respectively. Collectively, the alternative EKQ\* and SND\* ends, represented 107 (29.7%) of these sequences. Individually, the EKQ\* and SND\* C-terminal ends represented 30 (8.3%) and 77 (21.4%) of all sequenced amplicons. The remaining two (0.6%) cloned sequences possessed deletions within the *arsS* coding sequence, which may represent naturally occurring *arsS* null mutants or deletion mutations.

### 3.3. *arsS* point mutations

Our analyses of cloned *arsS* amplicon sequences generated from individual *H. pylori* populations from various gastric anatomical regions indicated that sequence variation due to point mutations outside the homopolymeric cytosine was present. Disregarding differences in hypermutable cytosine tract length, we considered single nucleotide mutations as a determinant of variant 3' *arsS*



**Fig. 2.** Variable *ArsS* C-terminal regions. Sequence logos were generated based on the 360 cloned 3' *arsS* region sequences and depict the 4 variable *ArsS* C-terminal regions past the homopolymeric cytosine tract. Amino acid sequence differences and similarities are apparent. The x-axis represents the amino acid sequence distal to the final proline (P) Encoded by the homopolymeric cytosine tract. The y-axis represents the probability of the amino acid present. Amino acids are also colored due to their hydropathy index values, where hydrophobic amino acids are black, hydrophilic are blue, and neutral are green. (A) Sequence logo for PKI\* was generated with 93 cloned 3' *arsS* region sequences from 31 *H. pylori* populations. (B) Sequence logo for LWC\* was derived from 158 sequences from 36 *H. pylori* populations. (C) Sequence logo EKQ\* was generated with 30 sequences from 11 populations. (D) Sequence logo for SND\* was derived from 76 3' *arsS* region sequences from 24 populations.

sequence types. Thus, nucleotide sequences that were identical within a gastric population were considered to be of the same sequence type and sequences that possessed at least one nucleotide mutation were considered to be of an alternate sequence type. We frequently observed sequenced amplicons that possessed no nucleotide substitutions. At least 5 to as many as 10 sequenced amplicons had identical sequence types within each population. We also observed amplicons that differed in nucleotide sequence from 1 to as many as 11 nucleotides within some *H. pylori* gastric populations. In all, 29 of the 36 *H. pylori* gastric regional populations examined in the current study contained distinct 3' *arsS* region sequence types, while only seven of 36 *H. pylori* populations showed identical *arsS* sequence types. In total, eight of twelve patients in the study showed evidence of polyclonal *H. pylori* infection.

#### 4. Discussion

*Helicobacter pylori* genome sequences deposited in NCBI databases suggest that individual strains possess a homopolymeric cytosine tract of specific length near the 3' terminus of *arsS*

[18,19,22–28]. In the present study, AFLP and sequence data generated with 3' *arsS* amplicons indicate that individual *H. pylori* populations encode multiple 3' *arsS* regions that can differ greatly in their cytosine tract lengths. AFLP data suggested that 6 to as many as 9 different 3' *arsS* amplicons of variable length were present in each population investigated. Furthermore, sequencing data indicated that 8 to 17 repeated cytosine nucleotides could be detected in this 3' *arsS* tract, depending on the population. The large number of cytosine tract lengths may be due in part to the fact that we queried entire *H. pylori* populations and not isolated, clonal strains as in previous studies [18,19,22–28]. However, Alm *et al.* observed 3 different 3' *arsS* homopolymeric cytosine tract lengths among J99 sequencing reads, and suggested that slipped-strand mispairing or *in vitro* conditions could influence the generation of multiple tract lengths. Hence in the current study, *H. pylori* populations were harvested from gastric biopsy specimens and passed minimally *in vitro* to limit nonselective mutations and maintain original *arsS* homopolymeric cytosine tract length frequencies.

Using AFLP and sequence analyses to probe cytosine tract length differences and the presence/absence of the thymine deletion, we

found that each *H. pylori* population of this study encoded various ArsS C-terminal regions. Beier *et al.* showed previously that different *H. pylori* strains with presumably strain-dependent ArsS C-terminal ends are each capable of ArsS autophosphorylation and phosphotransfer to the cognate response regulator, ArsR [12]. However, their study assumed that ArsS with a single C-terminal end was expressed for each of the *H. pylori* strains investigated. Our data indicate that multiple ArsS isoforms, variant at their C-terminal end, are expressed within each population investigated. Our study makes clear that *H. pylori* populations possess multiple *arsS* alleles leading to the expression of multiple ArsS isoforms with alternate C-termini. Thus, our data indicate that *arsS* is a contingency gene. It is interesting to speculate that ArsS isoforms, variant in their C-terminal end and still functional, may differ in regards to phosphotransfer capacity or efficiency; or perhaps each isoform or its encoding mRNA may have variable longevity in *H. pylori*. To test such hypotheses, we attempted to produce *H. pylori* strains capable of expressing ArsS with a single C-terminal end. However, these studies have been unsuccessful perhaps due to the importance of the *hemB* gene present in the *arsRS* operon.

The complexity of ArsS isoform functionality is increased with the discovery of a fourth (the SND\*) C-terminal end. Interestingly, the thymine deletion that facilitates SND\* translation is present in the sequenced *H. pylori* strains J99 and B38 [18,19,22–28]. In this study, we found that 8 of the 12 patients were infected with *H. pylori* that possessed this thymine deletion. We entertained the possibility that this was due to a bias in geographic locale as all patients in this study were evaluated in Nashville, Tennessee and strain J99 was also isolated from a patient in this geographic region. However, strain B38 was isolated from a patient in France [28]. Yet, the published sequences of *H. pylori* J99 and B38 do not suggest that the SND\* C-terminal end of ArsS would be translated due to the apparent length of the homopolymeric cytosine tract [18,28]. Thus, the acknowledgment of the SND\* C-terminal end or its frequency has not been considered in prior studies.

We also detected nucleotide substitutions within regions of *arsS* outside the homopolymeric cytosine tract and the thymine deletion and considered each variant to be evidence of additional sequence types. Interestingly, Alm *et al.* also noted nucleotide substitutions in sequence reads of J99, but at low frequency [18]. Comparison of the 9 *H. pylori* genome sequences within the NCBI database at the time of this study demonstrate that at least two, and frequently more than two, nucleotide substitutions exist within the 3' *arsS* region examined in our study [18,19,22–28]. At least 5 to as many as 10 *arsS* amplicons generated from each population had sequence types that possessed no nucleotide substitutions. However, there were also 3' *arsS* amplicons that had between 1 and 11 nucleotide substitutions within certain populations. While polyclonal *H. pylori* infections are frequent in developing nations, our data showed that identical sequence types were generated from only 7 gastric *H. pylori* populations in 4 of the 12 patients examined [29]. Due to the somewhat limited scope of our study, we cannot contend that the remaining 29 populations are truly polyclonal. However, the notion that 8 of the 12 patients in our study may have multiple sequence types of *H. pylori* concurrently is interesting to consider, especially considering the well documented proclivity of *H. pylori* for horizontal gene transfer.

## Acknowledgments

This work was supported by NIH grant R15 AI053062 to MHF and P01 CA116087, R01 DK58587, R01 CA77955 to RMP Jr. This research was also supported in part by a Howard Hughes Medical Institute grant through the Undergraduate Biological Sciences Education Program to the College of William and Mary.

## Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.micpath.2012.08.002>.

## References

- [1] Pattison CP, Combs MJ, Marshall BJ. *Helicobacter pylori* and peptic ulcer disease: evolution to revolution to resolution. *Am J Roentgenol* 1996;168:1415–20.
- [2] Pflock M, Finsterer N, Joseph B, Mollenkopf H, Meyer TF, Beier D. Characterization of the ArsRS regulon of *Helicobacter pylori*, involved in acid adaptation. *J Bacteriol* 2006;188:3449–62.
- [3] Schär J, Sickmann A, Beier D. Phosphorylation-independent activity of atypical response regulators of *Helicobacter pylori*. *J Bacteriol* 2005;187:3100–9.
- [4] Wen Y, Feng J, Scott DR, Marcus EA, Sachs G. Involvement of the HP0165-HP0166 two-component system in expression of some acidic-pH-upregulated genes of *Helicobacter pylori*. *J Bacteriol* 2006;188:1750–61.
- [5] Pflock M, Kennard S, Delany I, Scarlato V, Beier D. Acid-induced activation of the urease promoters is mediated directly by the ArsRS two-component system of *Helicobacter pylori*. *Infect Immun* 2005;73:6437–45.
- [6] Parsonnet J, Vandersteen D, Goates J, Sibley RK, Pritikin J, Chang Y. *Helicobacter pylori* infection in intestinal- and diffuse-type gastric adenocarcinomas. *J Natl Cancer* 1991;83:640–3.
- [7] Pflock M, Kennard S, Finsterer N, Beier D. Acid-responsive gene regulation in the human pathogen. *Helicobacter Pylori J Biotechnol* 2006;126:52–60.
- [8] Pflock M, Müller S, Beier D. The CrdRS (HP1365-HP1364) two-component system is not involved in pH-responsive gene regulation in the *Helicobacter pylori* strains 26695 and G27. *Curr Microbiol* 2006;54:320–4.
- [9] Sachs G, Weeks DL, Wen Y, Marcus EA, Scott DR. Acid acclimation by *Helicobacter pylori*. *Physiology* 2005;20:429–38.
- [10] Scott DR, Marcus EA, Wen Y, Oh J, Sachs G. Gene expression *in vivo* shows that *Helicobacter pylori* colonizes an acidic niche on the gastric surface. *P Natl Acad Sci USA* 2007;104:7235–40.
- [11] Scott DR, Marcus EA, Wen Y, Singh S, Feng J, Sachs G. Cytoplasmic histidine kinase (HP0244)-regulated assembly of urease with UreI, a channel for urea and its metabolites, CO<sub>2</sub>, NH<sub>3</sub>, and NH<sub>4</sub><sup>+</sup> is necessary for acid survival of *Helicobacter pylori*. *J Bacteriol* 2010;192:94–103.
- [12] Beier D, Frank R. Molecular characterization of two-component systems of *Helicobacter pylori*. *J Bacteriol* 2000;182:2068–76.
- [13] Stock AM, Robinson VL, Goudreau PN. Two-component signal transduction. *Annu Rev Biochem* 2000;69:183–215.
- [14] Dietz P, Gerlach G, Beier D. Identification of target genes regulated by the two-component system HP166-HP165 of *Helicobacter pylori*. *J Bacteriol* 2002;184:350–62.
- [15] Achtman M, Azuma T, Berg DE, Ito Y, Morelli G, Pan Z, et al. Recombination and clonal groupings within *Helicobacter pylori* from different geographical regions. *Mol Microbiol* 1999;32:459–70.
- [16] Han S, Zschausch HE, Meyer HW, Schneider T, Loos M, Bhakdi S, et al. *Helicobacter pylori*: clonal population structure and restricted transmission within families revealed by molecular typing. *J Clin Microbiol* 2000;38:3646–51.
- [17] Henderson IR, Owen P, Nataro JP. Molecular switches - the ON and OFF of bacterial phase variation. *Mol Microbiol* 1999;33:919–32.
- [18] Alm RA, Ling LS, Moir DT, King BL, Brown ED, Doig PC, et al. Genomic-sequence comparison of two unrelated isolates of the human gastric pathogen *Helicobacter pylori*. *Nature* 1999;397:176–80.
- [19] Tomb JF, White O, Kerlavage AR, Clayton RA, Sutton GG, Fleischmann RD, et al. The complete genome sequence of the gastric pathogen *Helicobacter pylori*. *Nature* 1997;388:539–47.
- [20] McNulty SL, Mole BM, Dailidienė D, Segal I, Ally R, Mistry R, et al. Novel 180- and 480-base-pair insertions in African and African-American strains of *Helicobacter pylori*. *J Clin Microbiol* 2004;42:5658–63.
- [21] Orsi R, Bowen BM, Wiedmann M. Homopolymeric tracts represent a general regulatory mechanism in prokaryotes. *BMC Genomics* 2010;11:1471–2164.
- [22] Baltus DA, Amieva MR, Covacci A, Lowe TM, Merrell DS, Ottemann KM, et al. The complete genome sequence of *Helicobacter pylori* strain G27. *J Bacteriol* 2009;191:447–8.
- [23] Fischer, W., L. Windhager, A. Karnholz, M. Zeiller, R. Zimmer, and R. Haas. 2008. The complete genome sequence of *Helicobacter Pylori* Strain P12. Unpublished.
- [24] Kersulyte, D., A. Kalia, R.H. Gilman, and D.E. Berg. 2010. Genome sequence of *Helicobacter pylori* from the remote Amazon: traces of Asian ancestry of the first Americans. Unpublished.
- [25] Kim S, Lee WK, Choi SH, Kang S, Park HS, Kim YS, et al. *Helicobacter Pylori* 2004;51, <http://www.ncbi.nlm.nih.gov/nucore/CP000012.1>.
- [26] Kim S, Lee WK, Choi SH, Kang S, Park HS, Kim YS, et al. *Helicobacter Pylori* 2009;52, <http://www.ncbi.nlm.nih.gov/nucore/261838873>.
- [27] Oh JD, Kling-Bäckhed H, Giannakis M, Xu J, Fulton RS, Fulton LA, et al. The complete genome sequence of a chronic atrophic gastritis *Helicobacter pylori* strain: evolution during disease progression. *P Natl Acad Sci USA* 2006;103:9999–10004.
- [28] Thiberge J, Boursaux-Eude C, Lehours P, Dillies M, Creno S, Coppee J, et al. Array-based hybridization of *Helicobacter pylori* isolates to the complete genome sequence of an isolate associated with MALT lymphoma. *BMC Genomics* 2010;11:368.
- [29] Logan Robert PH, Walker MM. Epidemiology and diagnosis of *Helicobacter pylori* infection. *Br Med J* 2001;323:920–2.