

2012

## Every invertible matrix is diagonally equivalent to a matrix with distinct eigenvalues

Zejun Huang

Nung-Sing Sze

Chi-Kwong Li

*William & Mary*, [ckli@wm.edu](mailto:ckli@wm.edu)

Man-Duen Choi

Follow this and additional works at: <https://scholarworks.wm.edu/aspubs>

---

### Recommended Citation

Choi, M. D., Huang, Z., Li, C. K., & Sze, N. S. (2012). Every invertible matrix is diagonally equivalent to a matrix with distinct eigenvalues. *Linear algebra and its applications*, 436(9), 3773-3776.


This Article is brought to you for free and open access by the Arts and Sciences at W&M ScholarWorks. It has been accepted for inclusion in Arts & Sciences Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).



Contents lists available at SciVerse ScienceDirect

# Linear Algebra and its Applications

journal homepage: [www.elsevier.com/locate/laa](http://www.elsevier.com/locate/laa)



## Every invertible matrix is diagonally equivalent to a matrix with distinct eigenvalues

Man-Duen Choi<sup>a</sup>, Zejun Huang<sup>b</sup>, Chi-Kwong Li<sup>c,1</sup>, Nung-Sing Sze<sup>b,\*</sup>

<sup>a</sup> Department of Mathematics, University of Toronto, Toronto, Ontario, Canada M5S 2E4

<sup>b</sup> Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong

<sup>c</sup> Department of Mathematics, College of William & Mary, Williamsburg, VA 23187-8795, USA

### ARTICLE INFO

#### Article history:

Received 6 November 2011

Accepted 7 December 2011

Available online 20 January 2012

Submitted by R.A. Brualdi

#### AMS classification:

15A18

#### Keywords:

Invertible matrices

Diagonal matrices

Distinct eigenvalues

### ABSTRACT

We show that for every invertible  $n \times n$  complex matrix  $A$  there is an  $n \times n$  diagonal invertible  $D$  such that  $AD$  has distinct eigenvalues. Using this result, we affirm a conjecture of Feng, Li, and Huang that an  $n \times n$  matrix is not diagonally equivalent to a matrix with distinct eigenvalues if and only if it is singular and all its principal minors of size  $n - 1$  are zero.

© 2011 Elsevier Inc. All rights reserved.

## 1. Introduction

Denote by  $M_n$  the set of  $n \times n$  complex matrices. In [1], the authors pointed out that matrices with distinct eigenvalues have many nice properties. They then raised the question whether every invertible matrix in  $M_n$  is diagonally equivalent to a matrix with distinct eigenvalues, and conjectured that a matrix in  $M_n$  is not diagonally equivalent to a matrix with distinct eigenvalues if and only if it is singular and every principal minor of size  $n - 1$  is zero. They provided a proof for matrices in  $M_n$  with

\* Corresponding author.

E-mail addresses: [choi@math.toronto.edu](mailto:choi@math.toronto.edu) (M.-D. Choi), [huangzejun@yahoo.cn](mailto:huangzejun@yahoo.cn) (Z. Huang), [ckli@math.wm.edu](mailto:ckli@math.wm.edu) (C.-K. Li), [raymond.sze@inet.polyu.edu.hk](mailto:raymond.sze@inet.polyu.edu.hk) (N.-S. Sze).

<sup>1</sup> This research was done while he was visiting The Hong Kong University of Science & Technology in 2011 under the support of a Fulbright Fellowship. Li is an honorary professor of The University of Hong Kong, Taiyuan University of Technology, and Shanghai University.

$n \leq 3$ , and demonstrated the complexity of the problem for matrices in  $M_4$  using their approach. In this note, we affirm their conjecture by proving the following theorem.

**Theorem 1.1.** *Suppose  $A \in M_n$  is invertible. There is an invertible diagonal  $D \in M_n$  such that  $AD$  has distinct eigenvalues.*

Once this result is proved, we have the following corollary.

**Corollary 1.2.** *Let  $A \in M_n$ . The following are equivalent.*

- (a)  *$A$  is not diagonally equivalent to a matrix with distinct eigenvalues.*
- (b) *There is no diagonal matrix  $D$  such that  $AD$  has distinct eigenvalues.*
- (c) *The matrix  $A$  is singular and all principal minors of size  $n - 1$  are zero.*

**Proof.** The implication (a)  $\Rightarrow$  (b) is clear. Suppose condition (c) does not hold. Then either  $A$  is invertible or  $A$  has an invertible principal submatrix of size  $n - 1$ . Assume the former case holds. There is an invertible diagonal matrix  $D$  such that  $AD$  has distinct eigenvalues by Theorem 1.1. If the latter case holds, we may assume without loss of generality that the leading principal submatrix  $A_1 \in M_{n-1}$  is invertible. By Theorem 1.1, there is an invertible diagonal matrix  $D_1 \in M_{n-1}$  such that  $A_1D_1$  has distinct (nonzero) eigenvalues. Let  $D = D_1 \oplus [0]$ . Then  $AD$  has distinct eigenvalues including zero as an eigenvalue. Thus, (b) cannot hold. So, we have proved (a)  $\Rightarrow$  (b)  $\Rightarrow$  (c).

Recall that the characteristic polynomial of a matrix  $B \in M_n$  has the form  $\det(xI_n - B) = x^n + b_{n-1}x^{n-1} + b_{n-2}x^{n-2} + \dots + b_1x + b_0$ , where  $(-1)^j b_{n-j}$  is the sum of  $j \times j$  principal minors of  $B$ . Suppose condition (c) holds. Since the principal minors of  $D_1AD_2$  are scalar multiples of the corresponding principal minors of  $A$ , then  $D_1AD_2$  has characteristic polynomial of the form  $\det(xI_n - D_1AD_2) = x^n + a_{n-1}x^{n-1} + \dots + a_2x^2$  so that 0 is a root with multiplicity at least two. Thus,  $D_1AD_2$  cannot have  $n$  distinct eigenvalues. So, the implication (c)  $\Rightarrow$  (a) is proved.  $\square$

Note that the set of diagonal matrices is an  $n$ -dimensional subspace in  $M_n$ . We can extend Theorem 1.1 to the following.

**Corollary 1.3.** *Suppose  $\mathcal{V}$  is a subspace of matrices in  $M_n$ .*

- (a) *If there are invertible matrices  $R$  and  $S$  such that  $R\mathcal{V}S = \{RXS : X \in \mathcal{V}\}$  contains the subspace of diagonal matrices, then for any invertible  $A \in M_n$  there is  $X \in \mathcal{V}$  such that  $AX$  has distinct eigenvalues.*
- (b) *If there are invertible matrices  $R$  and  $S$  such that  $RXS$  has zero first row and zero last column for every  $X \in \mathcal{V}$ , then  $A = SR$  is invertible and  $AX$  is similar to  $RXS$  which cannot have distinct eigenvalues for any  $X \in \mathcal{V}$ .*

**Proof.** (a) Suppose  $A$  is invertible. Then there is a diagonal matrix  $D$  such that  $S^{-1}AR^{-1}D$  has distinct eigenvalues by Theorem 1.1. Set  $X = R^{-1}DS^{-1} \in \mathcal{V}$ . Notice that  $AX$  has distinct eigenvalues as  $S^{-1}(AX)S = S^{-1}(AR^{-1}DS^{-1})S = (S^{-1}AR^{-1})D$ .

Assertion (b) can be verified readily.  $\square$

## 2. Proof of Theorem 1.1

We will prove Theorem 1.1 by induction on  $n$ . The result is clear if  $A \in M_1$ . Assume that the result holds for all  $k \times k$  invertible matrices with  $1 \leq k < n$ . Suppose  $A \in M_n$  is invertible. We consider two cases.

**Case 1.** If all  $k \times k$  principal minors of  $A$  are singular for  $k = 1, \dots, n - 1$ , then the characteristic polynomial of  $A$  has the form  $x^n - a_0$  and has  $n$  distinct roots. So, the result holds with  $D = I_n$ .

**Case 2.** Suppose  $A$  has an invertible  $k \times k$  principal minor. Without loss of generality, we may assume that  $A = \begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}$  such that  $A_{11} \in M_k$  is invertible for some  $1 \leq k < n$ . Then the

Schur complement of  $A_{22}$  equals  $B = A_{22} - A_{21}A_{11}^{-1}A_{12}$  which is invertible; see [2, pp. 21–22]. By induction assumption, there are diagonal invertible  $D_1 \in M_k$  and  $D_2 \in M_{n-k}$  such that each of  $A_{11}D_1$  and  $BD_2$  has distinct nonzero eigenvalues, say,  $\lambda_1, \dots, \lambda_k$  and  $\lambda_{k+1}, \dots, \lambda_n$ , respectively. Thus,  $A_{11}D_1$  and  $BD_2$  are diagonalizable and there are invertible  $S_1 \in M_k$  and  $S_2 \in M_{n-k}$  such that  $S_1A_{11}D_1S_1^{-1} = \Lambda_1 = \text{diag}(\lambda_1, \dots, \lambda_k)$  and  $S_2BD_2S_2^{-1} = \Lambda_2 = \text{diag}(\lambda_{k+1}, \dots, \lambda_n)$ . Let  $D_{r,s} = rD_1 \oplus sD_2$ . The proof is complete if one can find some suitable  $r$  and  $s$  so that  $AD_{r,s}$  has distinct eigenvalues. Notice that  $AD_{r,s}$  has the same eigenvalues as

$$\begin{aligned} \tilde{A} &= \begin{bmatrix} S_1 & 0 \\ 0 & sS_2 \end{bmatrix} \begin{bmatrix} I_k & 0 \\ -A_{21}A_{11}^{-1} & I_{n-k} \end{bmatrix} AD_{r,s} \begin{bmatrix} I_k & 0 \\ A_{21}A_{11}^{-1} & I_{n-k} \end{bmatrix} \begin{bmatrix} S_1^{-1} & 0 \\ 0 & s^{-1}S_2^{-1} \end{bmatrix} \\ &= \begin{bmatrix} r\Lambda_1 + sS_1A_{12}D_2A_{21}A_{11}^{-1}S_1^{-1} & S_1A_{12}D_2S_2^{-1} \\ s^2S_2BD_2A_{21}A_{11}^{-1}S_1^{-1} & s\Lambda_2 \end{bmatrix}. \end{aligned}$$

Denote by  $D(a, d)$  the closed disk in  $\mathbb{C}$  centered at  $a$  with radius  $d \geq 0$ . Suppose the  $k \times k$  matrix  $S_1A_{12}D_2A_{21}A_{11}^{-1}S_1^{-1}$  has diagonal entries  $\mu_1, \dots, \mu_k$  and let

$$d_1 = k\|S_1A_{12}D_2A_{21}A_{11}^{-1}S_1^{-1}\|, \quad d_2 = (n-k)\|S_1A_{12}D_2S_2^{-1}\|, \quad \text{and} \quad d_3 = k\|S_2BD_2A_{21}A_{11}^{-1}S_1^{-1}\|,$$

where  $\|\cdot\|$  is the operator norm. By Geršgorin disk result (see [2, pp. 344–347]), the eigenvalues of  $\tilde{A}$  must lie in the union of the  $n$  Geršgorin disks, which is a subset of the union of  $n$  disks

$$D(r\lambda_1 + s\mu_1, sd_1 + d_2), \dots, D(r\lambda_k + s\mu_k, sd_1 + d_2), D(s\lambda_{k+1}, s^2d_3), \dots, D(s\lambda_n, s^2d_3).$$

We can choose sufficiently large  $r > 0$  and sufficiently small  $s > 0$  so that these disks are disjoint, and hence  $\tilde{A}$  has  $n$  disjoint Geršgorin disks. Then  $\tilde{A}$  has distinct eigenvalues.  $\square$

We thank Editor Zhan for sending us the two related Refs. [3,4]. In these papers, the author proved following. Suppose  $A$  is an  $n \times n$  matrix and  $a_1, \dots, a_n$  are complex numbers. Then there is a diagonal matrix  $E$  such that  $A + E$  has eigenvalues  $a_1, \dots, a_n$ . Moreover, if all principal minors of  $A$  are nonzero, then there is a diagonal matrix  $D$  such that  $AD$  has eigenvalues  $a_1, \dots, a_n$ .

Note that the assumption on the principal minors of  $A$  is important in the second assertion. Obviously, if  $\det(A) = 0$ , then one cannot find diagonal  $D$  such that  $AD$  has  $n$  nonzero eigenvalues. Even if we remove this obvious obstacle and assume that  $A$  is invertible, one may not be able to find diagonal  $D$  so that  $AD$  has prescribed eigenvalues. For example, if  $\{E_{1,1}, E_{1,2}, \dots, E_{n,n}\}$  is the standard basis for  $M_n$  and  $A = E_{1,2} + \dots + E_{n-1,n}$ , then the eigenvalues of  $AD$  always have the form  $z, zw, \dots, zw^{n-1}$  for some  $z \in \mathbb{C}$ , where  $w$  is the primitive  $n$ th root of unity.

It is interesting to determine the condition on  $A$  so that for any complex numbers  $a_1, \dots, a_n$ , one can find a diagonal  $D$  such that  $AD$  has  $a_1, \dots, a_n$  as eigenvalues.

**Acknowledgments**

Research of Choi was supported by a NSERC grant. Research of Huang was supported by a HK RGC grant. Research of Li was supported by a USA NSF grant, and a HK RGC grant. Research of Sze was supported by HK RGC grants.

## References

- [1] X.-L. Feng, Z. Li, T.-Z. Huang, Is every nonsingular matrix diagonally equivalent to a matrix with all distinct eigenvalues? *Linear Algebra Appl.*, 436 (2012) 120–125.
- [2] R.A. Horn, C.R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 1985.
- [3] S. Friedland, On inverse multiplicative eigenvalue problems for matrices, *Linear Algebra Appl.* 12 (1975) 127–137.
- [4] S. Friedland, Inverse eigenvalue problems, *Linear Algebra Appl.* 17 (1977) 15–51.