

Reports

11-4-1985

Fall 1985 Potomac River Oyster Bar Survey

Elgin A. Dunnington
Virginia Institute of Marine Science

James P. Whitcomb
Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Dunnington, E. A., & Whitcomb, J. P. (1985) Fall 1985 Potomac River Oyster Bar Survey. Marine Resource Report No. 85-6. Virginia Institute of Marine Science, College of William and Mary. <http://dx.doi.org/doi:10.21220/m2-nddy-2n39>

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

CENTER FOR ENVIRONMENTAL AND
ESTUARINE STUDIES
UNIVERSITY OF MARYLAND

Ref. No. [UMCEES]CBL 85-126
VMRR 85-6

November 4, 1985

Chesapeake Biological Laboratory
Solomons, MD 20688

FALL 1985 POTOMAC RIVER OYSTER BAR SURVEY

Elgin A. Dunnington
James P. Whitcomb

Selected oyster bars in the Potomac River were sampled on 21, 22, and 23 October 1985. Bottom material was dredged aboard the MISS KAY, and counts and observations were made by personnel from MD/DNR, UMCEES/CBL, VIMS, PRFC, and VMRC. Standard bar composition data were recorded. A summary of the collected information is appended.

There has been a good general set of oyster spat this year. Reproduction has not been confined, as it often is, to the area near the mouth of the river; it has extended up through the middle bars to some upper bars, and into the tributaries. This is the most widely distributed recruitment since 1963. Only the most upriver bottoms, which are nearly devoid of broodstock, failed to share in this year's set. Planted shells which were disappointingly clean of spat (and fouling) last year received a good spatfall this year. Most important however, are the spat attached to "natural" cultch such as oysters, oystershell fragments, clamshells, mussel shells, and stones, because most of them can grow to maturity where they are without being transplanted. Maturation of the 1985 set will yield more widely dispersed oyster populations, resulting in less concentrated harvest efforts. Although this year's spat counts may seem low when compared to what are usually regarded as "seed quality" numbers in other areas, they are nevertheless significant quantities in the Potomac, where survival is usually good.

The maximum yield from this fortuitous reproduction will only be realized if these spat are allowed to grow to market size. Because of the general scarcity of cultch, a substantial portion of the 1985 spat are attached to market size oysters. Spat which are difficult to cull off market oysters now without killing them will be more easily separated without injury next year.

Mortalities were generally low except at Cornfield and in the 1985 Cole's Point seed.

The condition of oysters in the middle and upper river was good, and in the lower river it was fair. These levels of "fatness" are normal for this time of year.

df
attachment
11/4/85

SUMMARY, FALL 1985 POTOMAC RIVER OYSTER BAR SURVEY

Box	Market Oysters	Volume % Markets	Small Oysters	Volume % Small	Spat	Old Boxes	Recent Boxes	Gapers	Volume % Shells	% Recent Boxes and Gapers	Remarks
Cedar Point ('81 seed)	56	20	20	5	0	8	0	0	74	0	
Cedar Point (natural)	40	25	2	<1	0	0	0	0	75	0	One spat in dredge lick
Watsons ('81 seed)	20	15	4	<1	0	14	0	0	84	0	All boxes were smalls
Gum Bar (natural)	68	30	38	8	0	24	0	0	42	0	Old brood stock planting
Swan Point (outside)	76	60	4	1	8	10	0	0	30	0	
Swan Point (inside)	12	8	0	0	0	16	0	0	82	0	One spat in dredge lick
Cobb Bar (lower)	102	50	8	1	18	32	0	0	30	0	
Cobb Bar ('81 seed)	130	64	4	<1	16	26	0	0	22	0	
Dukelhart Channel ('82 seed)	108	55	2	<1	20	14	0	0	46	0	
Sheepshead (lower inside)	64	36	12	2	16	8	6	0	49	6	
Huggins (tug damage)	60	30	8	3	72	16	2	0	49	1	
Knott's Hollow (natural)	30	22	2	1	85	12	6	0	63	5	
Posey's Bluff ('81 & '83 seed)	106	60	30	7	24	12	0	0	25	0	
Kingcopsico ('81 seed)	86	40	16	3	4	20	0	0	47	0	
Coles Point ('85 seed)	104	32	118	18	4	44	22	0	35	9	Most recent boxes are transplanted seed
Ragged Point (upper)	82	60	4	<1	30	2	4	0	35	3	

Box	Market Oysters	Volume % Markets	Small Oysters	Volume % Small	Spat	Old Boxes	Recent Boxes	Gapers	Volume % Shells	% Recent Boxes and Gapers	Remarks
Ragged Point (outside deep)	38	30	10	3	18	36	8	0	46	11	
Ragged Point (lower inside)	80	35	10	1	108	12	8	0	47	4	
St. Georges Is. ('85 gr. shell)	20	10	4	1	312	4	2	0	48	<1	
St. Georges Is. ('84 gr. shell)	6	1	10	2	176	0	6	0	66	3	
St. Georges Is. (natural)	92	35	42	10	112	14	2	0	42	<1	
Bonum (nat. & gr. shell)	32	30	6	1	122	8	2	0	50	1	
Bonum ('84 shells, dr. & gr.)	0	0	0	0	46	0	2	0	92	4	
Bonum (natural)	28	20	0	0	112	2	6	0	67	4	
Lynch Point ('79 shells)	46	20	10	1	204	4	8	0	56	1	
Hog Island ('85 gr. shells)	20	12	12	4	40	4	8	0	61	1	
Hog Island (natural)	146	44	40	9	60	4	4	0	41	2	Probable work site. About 12% of markets have spat.
Great Neck ('85 gr. shells)	4	1	6	1	132	0	4	0	70	3	
Great Neck (natural)	62	20	14	2	132	4	4	0	60	2	Probable work site. About 20% of markets have spat.
Kitts Point ('85 dr. shell)	0	0	4	2	180	12	4	0	63	2	
Kitts Point ('85 dr. shell)	0	0	0	0	296	2	12	0	75	4	
Kitts Point ('79 dr. shell)	72	35	52	11	266	36	12	0	21	3	
PRFC/DNR add.#2 ('84 dr. shell)	0	0	28	10	786	0	10	0	30	1	

Box	Market Oysters	Volume % Markets	Small Oysters	Volume % Small	Spat	Old Boxes	Recent Boxes	Gapers	Volume % Shells	% Recent Boxes and Gapers	Remarks
Jones Shore Lot #1, outside ('83 shell)	66	22	234	40	166	12	14	0	12	3	
Jones Shore Lot #1, inside ('83 shell)	104	40	126	22	58	6	12	0	31	4	
Jones Shore Lot #2, outside ('84 shell)	16	7	28	6	442	2	2	0	57	<1	
Jones Shore Lot #2, inside ('84 shell)	32	18	8	1	210	22	12	0	31	5	
Jones Shore Lot #12 ('82 shells)	48	21	48	11	528	14	8	0	35	1	
Cornfield Harbor (natural)	80	30	60	13	470	18	38	0	16	6	
Cornfield Harbor ('85 dr. shell)	8	2	6	1	222	0	0	0	17	0	
Cornfield Harbor (natural)	88	30	252	30	296	32	28	0	17	4	