

# Identification and partial characterisation of metalloproteases secreted by a *Mesanophrys*-like ciliate parasite of the Norway lobster *Nephrops norvegicus*

H. J. Small<sup>1,3,\*</sup>, D. M. Neil<sup>1</sup>, A. C. Taylor<sup>1</sup>, G. H. Coombs<sup>2</sup>

<sup>1</sup>Division of Environmental and Evolutionary Biology, and <sup>2</sup>Division of Infection and Immunity, Institute of Biomedical & Life Sciences, University of Glasgow, Glasgow G12 8QQ, UK

<sup>3</sup>Present address: Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, Virginia 23062, USA

**ABSTRACT:** A ciliate parasite, tentatively identified as *Mesanophrys* sp. of Norway lobsters *Nephrops norvegicus*, is demonstrated to secrete several proteases into the culture medium (modified *Nephrops* saline). Analyses using substrate-impregnated sodium dodecylsulfate-polyacrylamide gel electrophoresis (SDS-PAGE) revealed 12 activity bands differing greatly in mobility in the gels. The complete inhibition of proteolytic activity by 1,10-phenanthroline indicated that the proteases are of the metallo class. The proteases were active at the physiological temperature (8°C) and haemolymph pH (7.8) of the host. The proteases were selective in the degradation of several host proteins, including the myosin heavy chain, which is a major structural component of lobster muscle. Consequently, these proteases may have important roles in several aspects of the host–parasite interaction including invasion, nutrient uptake by the ciliate, and pathogenesis.

**KEY WORDS:** Ciliate · Parasite · Proteases · Secreted · Pathology

Resale or republication not permitted without written consent of the publisher

## INTRODUCTION

Histophagous ciliates of the genus *Mesanophrys* have previously been described parasitising several crustaceans, including the shore crab *Carcinus maenas* (Cattaneo 1888), the edible crab *Cancer pagurus* (Grolhier & Leglise 1977), the Dungeness crab *C. magister* (Morado & Small 1994) and the blue crab *Callinectes sapidus* (Messick & Small 1996). We recently discovered a histophagous ciliate infecting Norway lobsters *Nephrops norvegicus* from the Clyde Sea area in Scotland that appears to belong to the genus *Mesanophrys* by morphological feature comparison (Small et al. 2005).

The invasion of these histophagous ciliates into their crustacean host is poorly understood. Morado et al. (1999) suggested that *Mesanophrys pugettensis* enters the Dungeness crab via lesions in the epidermal surface associated with the moult cycle; inside they spread through the body and multiply to high densities in tis-

sues and haemolymph. Hibbits & Sparks (1983) also suggested that the cuticle of the isopod *Gnorimosphaeroma oregonensis* is the location of initial ciliate invasion. In several other parasitic infections, proteases released by the pathogen have been shown to play various roles in the establishment and progression of disease. For example, some parasite proteases degrade host extracellular matrix proteins, thus facilitating spread of the parasite within host tissues (Schulte & Scholze 1989). Many marine bacterial, fungal and protozoan pathogens secrete proteases during *in vitro* culture, suggesting that protease release could be important in pathogenesis. Examples include the salmon pathogen *Aeromonas salmonicida* (Rockey et al. 1988), the crayfish pathogen *Aphanomyces astaci* (Söderhäll & Unestam 1975), and the oyster parasite *Perkinsus marinus* (La Peyre & Faisal 1995, La Peyre et al. 1995). *Uronema marinum*, a histophagous parasitic ciliate of the olive flounder *Paralichthys olivaceus*, has also been reported to secrete proteases during *in vitro* culture

\*Email: hamish@vims.edu

(Lee et al. 2003); in this case, secreted proteases may be directly responsible for many of the clinical symptoms associated with the disease, such as tissue necrosis.

Several reports of *Mesanothryx* spp. infections in crustaceans have indicated that these ciliates infiltrate and consume host tissues and haemocytes (Poisson 1930, Bang et al. 1972, Messick & Small 1996). The discovery of a *Mesanothryx*-like parasitic ciliate infecting Norway lobsters *Nephrops norvegicus*, and isolation of the ciliate via *in vitro* cultivation, has allowed proteolytic factors that may play a role in the parasite's establishment and proliferation to be investigated. This paper reports on the characterisation of these proteases, and suggests ways in which they may be involved in the pathogenesis of ciliate disease in lobsters.

## MATERIALS AND METHODS

**Isolation and cultivation of the ciliate parasite.** Routine screening of *Nephrops norvegicus* from the Clyde Sea area for the parasitic dinoflagellate *Hematodinium* sp. led to the discovery and identification of a ciliate infection in several Norway lobsters (Small et al. 2005). Haemolymph samples were withdrawn aseptically from the base of the fifth pereopod using a 1 ml disposable syringe and 25-gauge needle following sterilisation of the cuticle with 70% (v/v) ethanol. Ciliate infections were diagnosed by viewing haemolymph smears using light microscopy. Parasitic ciliates were isolated in 3.5 cm well plates with 0.2 ml infected haemolymph added to 5 ml culture medium in each well. The culture medium comprised 10% (v/v) heat-inactivated foetal calf serum (FCS) in autoclaved balanced *Nephrops* saline (Appleton & Vickerman 1998), containing NaCl, 27.99 g l<sup>-1</sup>; KCl 0.95 g l<sup>-1</sup>; CaCl<sub>2</sub> 2.014 g l<sup>-1</sup>; MgSO<sub>4</sub> 2.465 g l<sup>-1</sup>; Na<sub>2</sub>SO<sub>4</sub> 0.554 g l<sup>-1</sup>; HEPES 1.92 g l<sup>-1</sup>, adjusted to pH 7.8. Penicillin G (10 U ml<sup>-1</sup>) and streptomycin (10 µg ml<sup>-1</sup>) were added to inhibit bacterial contamination. The medium was filter-sterilised (0.2 µm) after the addition of all constituents. Cultures were incubated at 8°C. Serial subculturing gave rise to axenic cultures.

Experimental subcultures for protease analyses were initiated with the addition of 1 × 10<sup>5</sup> ciliates to 5 ml culture medium in 3.5 cm well plates, in quadruplicate, and were maintained at 8°C. Cultures were checked by light microscopy daily for bacterial contamination. On Days 3 and 7, a 50 µl sample of the ciliate suspension was removed and added to 10 µl of phosphate-buffered saline (PBS) containing 20 µg ml<sup>-1</sup> 4',6'-diamidino-2-phenylindole (DAPI, Sigma). We placed 20 µl of this suspension on a glass slide and covered it with a coverslip. The ciliate nuclei were observed under oil immersion using fluorescence microscopy

(Zeiss Axioplan 2 at an excitation wavelength of 350 nm). No fluorescent signal from contaminating protists or bacteria was detectable. Cell viability was assessed by viewing cultures under phase contrast microscopy. The total volume of medium in each well was collected on Days 1, 3, 5 and 7. The samples were centrifuged at 400 × *g* for 10 min at 4°C, and 4.9 ml culture medium was collected, passed through a 0.2 µm filter and frozen at -80°C until analysed. The cell pellet was resuspended in 100 µl culture medium and 10 µl of this was mixed with 10 µl 1% (v/v) formaldehyde (in PBS) to immobilise the ciliates. Ciliate densities were estimated using an Improved Neubauer counting chamber following standard procedures.

**Preparation of ciliate lysates.** Ciliates from an *in vitro* culture were collected, and cell density was estimated as described above. The resulting ciliate suspension (5 × 10<sup>4</sup> cells 100 µl<sup>-1</sup>) was washed and centrifuged (400 × *g* for 10 min at 4°C) 3 times with *Nephrops* saline before resuspension in 100 µl *Nephrops* saline. Cell lysates were prepared by freezing and thawing the sample 3 times, followed by centrifugation (10 000 × *g* for 10 min) at 4°C. The supernatant fraction of the cell lysate was carefully removed and stored at -80°C until analysed.

**Spectrophotometric assay for protease activity.** Protease activities in ciliate culture medium samples were assayed following the method of Sarath et al. (1989). Azocasein substrate (3% w/v) was prepared by dissolving azocasein (Sigma) in 0.1 M Tris-HCl, pH 8.0, followed by centrifugation at 10 000 × *g* for 10 min at 4°C. The supernatant (substrate) was removed and stored on ice prior to use. The assay was initiated by the addition of 75 µl of ciliate culture medium sample to 125 µl substrate solution. Samples were incubated for 18 h at 37°C. The reaction was terminated by the addition of 600 µl ice-cold 10% (w/v) trichloroacetic acid (TCA), after which the mixtures were kept at 4°C for 30 min. The samples were centrifuged at 9000 × *g* for 5 min to pellet the precipitated proteins (including undigested azocasein). Then 0.6 ml of the supernatant was removed and mixed with 0.7 ml 1.0 M NaOH, and the absorbance of the mixture read at 440 nm on a spectrophotometer. We defined 1 U of protease activity as the enzyme activity resulting in an absorbance of 1.0 in a 1 cm cuvette, under the conditions of the assay. Controls comprised cell culture media without the addition of ciliates. Each sample was assayed in triplicate.

**Substrate-impregnated SDS-PAGE.** The proteases in the ciliate culture medium and in ciliate lysates were separated using gelatin-impregnated polyacrylamide gel electrophoresis according to the method of La Peyre et al. (1995), with minor modifications. Gelatin was added to the 8% (w/v) acrylamide resolving gel at a final concentration of 0.2% protein (w/v). Ciliate cul-

ture medium/lysate samples (5  $\mu$ l) were mixed with 15  $\mu$ l of electrophoresis sample buffer (62.5 mM Tris-HCl, pH 6.8, containing SDS (2% w/v),  $\beta$ -mercaptoethanol (5% w/v), glycerol (10% w/v) and 0.004% pyronin Y) before loading onto gels containing copolymerised gelatin. Electrophoresis was performed using a BioRad mini-Protean system at 30 mA constant current, containing the Tris-glycine buffer for 90 min at 4°C. After electrophoresis, the resolved gels were incubated with agitation in Triton X-100 (2.5% v/v in distilled H<sub>2</sub>O) for 30 min at 4°C to remove SDS and reactivate the resolved proteases. To detect proteolytic activity, the gels were incubated in 0.1 M Tris-HCl, pH 8.0, for 4 to 12 h at 37°C. The gels were fixed and stained with Coomassie blue, then destained until clear bands were apparent where hydrolysis of embedded substrate had occurred. Apparent molecular masses of the proteases were determined from their mobility relative to known protein standards, although these were used only as means of identifying individual bands. Proteins did not migrate according to their molecular masses under the conditions used.

**Effect of inhibitors on protease activity.** Stock solutions of phenylmethylsulphonyl fluoride (PMSF, 200 mM), Pepstatin A (1 mM), and 1,10-phenanthroline (200 mM) were prepared in 100% methanol. A stock solution of N-(N-(L-3-trans-carboxirane-2-carbonyl)-L-leucyl)-agmatine (E-64, 10 mM) was prepared in 100% ethanol. Stock solutions of leupeptin (10 mM) and ethylenediaminetetraacetic acid (EDTA, 500 mM) were prepared in 0.1 M Tris-HCl, pH 8.0. Stock solutions of all the inhibitors were diluted in 0.1 M Tris-HCl, pH 8.0, to the appropriate working concentration (see list in Table 1). Ciliate culture medium samples with the maximum protease activity (after 7 d culture) were used in the inhibitor studies. To quantify the effects of the inhibitors, each of the appropriately diluted inhibitors was incubated for 1 h at 37°C with 100  $\mu$ l of ciliate culture medium. To these, 125  $\mu$ l of azocasein substrate solution was then added, and the samples were incubated for 18 h at 37°C. The proteolytic activity was measured spectrophotometrically as described in the foregoing subsection. The samples were compared with assays run in the absence of inhibitors. The effect of each inhibitor was assayed in triplicate.

To investigate the possible differences in inhibitor susceptibility between proteolytic bands, ciliate culture medium samples were resolved by gelatin-SDS-PAGE as in the foregoing subsection with minor alterations. The supernatant-sample buffer mixture was loaded across all wells of the gelatin-SDS-PAGE minigel. After electrophoresis and Triton X-100 treatment, the gel was sectioned into six 1.5 cm vertical strips, and the strips were incubated individually in 5 ml of 0.1 M Tris-HCl, pH 8.0, containing the appropriate inhibitor

for 6 h at 37°C. The gels were fixed and stained with Coomassie blue, then destained (inhibitor concentrations are detailed in Table 1).

**Effect of temperature and pH on protease activity.** The effect of temperature on protease activity in ciliate cell culture medium samples was examined spectrophotometrically as described above. The substrate-sample mixtures were incubated at 8, 25 and 37°C for 18 h. At each temperature, samples were assayed in triplicate.

The effect of pH on protease activity in ciliate culture medium samples was examined spectrophotometrically and by gelatin-SDS-PAGE. For the spectrophotometric assay, azocasein substrate (3% w/v) was dissolved in 0.1 M Tris-HCl pH 7.0, followed by centrifugation at 10 000  $\times g$  for 10 min at 4°C. The supernatant was removed and the pH increased by the addition of 10 M NaOH. Samples of substrate at pH 7.0 and 8.0 were removed and stored on ice prior to use. Ciliate culture medium samples (75  $\mu$ l) were mixed with 125  $\mu$ l substrate solution at the different pH values, and incubated for 18 h at 37°C. Protease activity was measured as described earlier. Controls comprised culture media without the addition of ciliates. Each substrate sample was assayed in triplicate. Ciliate culture medium samples were subjected to gelatin-SDS-PAGE, and washed in Triton X-100 as described earlier. Each gel was sectioned into respective sample lanes and incubated individually in 5 ml of 0.1 M Tris-HCl at pH 7.0 and 8.0 for 6 h at 37°C. The gels were then stained and destained.

**Effect of proteases on host muscle proteins.** The effect of proteases present in ciliate culture medium on host tissue proteins was investigated using a sample of abdominal flexor muscle from the Norway lobster. Briefly, 100 mg of abdominal flexor muscle was homogenised in 500  $\mu$ l 0.1 M Tris-HCl, pH 8.0, followed by centrifugation at 3000  $\times g$  for 10 min at 4°C. The muscle protein supernatant was removed and placed on ice prior to use. Samples of protein supernatant (20  $\mu$ l) were mixed with 20  $\mu$ l ciliate culture medium (Day 7) and 20  $\mu$ l 0.1 M Tris-HCl pH 8.0. Reaction mixtures were incubated at 8°C for 15 h. Controls included the incubation of 20  $\mu$ l of 0.1 M Tris-HCl pH 8.0 with equal volumes of protein supernatant, ciliate culture medium, and uninoculated culture medium under the same conditions. Protein degradation was evaluated by standard SDS-PAGE (Sambrook et al. 1989).

## RESULTS

### Protease activity and ciliate growth *in vitro*

Ciliate density increased approximately 10-fold over 7 d in culture. Ciliates were always observed to be

motile when cultures were examined daily using phase contrast microscopy. Protease activity in samples of culture medium was detected as early as 1 d post-inoculation, and increased over Days 3, 5 and 7. This increase in activity correlated closely with the increase in ciliate density (Fig. 1). No proteolytic activity could be detected in uninoculated culture medium.

#### Detection of proteases in ciliate culture medium and lysates by gelatin-SDS-PAGE

The presence of multiple protease activities in ciliate culture medium and lysate samples was confirmed using gelatin-SDS-PAGE. We detected 12 zones of hydrolysis with apparent molecular masses ranging from 20 to 220 kDa in the samples of culture medium, while only 8 of the zones of hydrolysis were clearly evident in lysate samples (Fig. 2). The profile of protease activities in the ciliate lysate differed from that in the medium not only in a reduction in the number of bands of gelatin hydrolysis (notably the absence of the 50 kDa band, Band 7, observed in the samples of ciliate culture medium), but also in the relative intensity of gelatin digestion of the various bands. The low molecular mass bands (20 to 50 kDa) appeared faint, while the high molecular mass bands (70 to 220 kDa) were very intense.

#### Effect of inhibitors on proteases present in ciliate culture medium

Pre-incubation of samples of the ciliate culture medium with protease inhibitors before determining the proteolytic activity by the spectrophotometric assay revealed marked inhibition of proteolytic activity when samples were incubated in the presence of 10 mM 1,10-phenanthroline (only  $2.5 \pm 8.8\%$  activity remained) (Table 1). Partial inhibition was observed with the addi-




Fig. 1. Changes in ciliate density and protease activity in cultures over 7 d. Mean  $\pm$  SD,  $n = 3$




Fig. 2. Detection of proteases in (A) ciliate culture medium, and (B) ciliate crude lysates by gelatin-SDS-PAGE. Positions of molecular mass standards are indicated on the left; arrows indicate zones of hydrolysis

tion of EDTA ( $42.9 \pm 3.3\%$  activity remained with 10 mM EDTA). However, addition of the inhibitors PMSF, Pepstatin A, E-64 and leupeptin had no effect on the proteolytic activity of samples of ciliate culture medium. The activities of all proteases separated by gelatin-SDS-PAGE were completely inhibited when the post-electrophoresis activation buffer contained 10 mM 1,10-phenanthroline (Table 1). The inclusion of 10 mM EDTA in the incubation buffer inhibited the majority of proteases apart from the 3 largest proteases, which were partially inhibited in apparent activity. Addition of the inhibitors PMSF, Pepstatin A, E-64 and leupeptin did not detectably inhibit hydrolysis of gelatin.

#### Effect of temperature and pH on protease activity in ciliate culture medium

To investigate the effect of temperature on enzyme activity, protease activity in samples of ciliate culture medium was assayed by the spectrophotometric assay at different temperatures. Protease activity could be detected at  $8^\circ\text{C}$  ( $2.2 \pm 0.5 \text{ U ml}^{-1}$ ), and increased 2.9-fold to  $6.2 \pm 0.3$  at  $25^\circ\text{C}$ , and 4.5-fold to  $9.8 \pm 0.2 \text{ U ml}^{-1}$  when the assay was performed at  $37^\circ\text{C}$ . Spectrophotometric measurements of protease activity indicated that the hydrolysis of azocasein was highest at pH 8.0 ( $7.4 \pm 0.2 \text{ U ml}^{-1}$ ) with 0.1 M Tris-HCl as the buffer. Decreasing the pH to 7.0 resulted in a reduction of protease activity ( $6.5 \pm 0.2 \text{ U ml}^{-1}$ ). Gelatin-SDS-PAGE analysis of the ciliate culture medium samples revealed 12 distinct zones of hydrolysis both at pH 7.0 and 8.0 (data not shown). However, the hydrolysis was more intense at pH 8.0.

Table 1. Effect of inhibitors on activity of proteases present in ciliate culture medium. 100% activity was  $7.08 \pm 0.33 \text{ U ml}^{-1}$  (mean  $\pm$  SD,  $n = 3$ )

Inhibitor	Conc.	Protease activity (% control)	Inhibition of gelatin-SDS-PAGE digestion
Control (0.1 M Tris-HCl) pH 8.0		100 $\pm$ 4.6	None
Control (ethanol)		94.5 $\pm$ 6.1	None
Control (methanol)		95.9 $\pm$ 2.9	None
<b>Serine protease/cysteine protease inhibitors:</b>			
PMSF	1 mM	94.6 $\pm$ 2.4	None
	10 mM	92.0 $\pm$ 2.8	
Leupeptin	1 $\mu$ M	94.4 $\pm$ 2.9	None
	10 $\mu$ M	91.1 $\pm$ 3.7	
<b>Cysteine protease inhibitor:</b>			
E-64	1 $\mu$ M	97.5 $\pm$ 5.2	None
	10 $\mu$ M	94.2 $\pm$ 2.5	
<b>Aspartic protease inhibitor:</b>			
Pepstatin A	1 $\mu$ M	92.5 $\pm$ 4.4	None
<b>Metallo-protease inhibitors:</b>			
1,10 phenanthroline	1 mM	41.4 $\pm$ 3.8	Complete
	10 mM	2.5 $\pm$ 8.8	
EDTA	1 mM	87.0 $\pm$ 3.6	Partial
	10 mM	42.9 $\pm$ 3.3	

### Host protein degradation by proteases present in ciliate culture medium

The multiple proteases present in samples of the ciliate culture medium had a selective effect on several abdominal flexor muscle proteins (Lane C, Fig. 3). In particular, the myosin heavy chain and one of the Troponin I family of proteins were completely degraded. There was also degradation of several unidentified muscle proteins with masses in the range of 100 to 120 kDa. However, paramyosin, actin and other muscle proteins remained unaffected. The FCS protein component of the culture medium occurred in smaller amounts in samples of ciliate culture medium (Lane D, Fig. 3) compared with the uninoculated control culture medium (Lane E, Fig. 3).

### DISCUSSION

Proteolytic enzymes from parasites are thought to be involved in parasite nutrition, anticoagulation and evasion of the host immune system (McKerrow 1989). They are also believed to play key roles in the processes of penetration and migration of parasites through host tissues (Klemba & Goldberg 2002). The extensive tissue infiltration and damage observed during *Mesanoophrys* spp. infections in crustaceans (Armstrong et al. 1981, Sparks et al. 1982, Hibbits & Sparks 1983, Messick & Small 1996) indicates a likely role for proteases in such processes.

The results of this study show that the *Mesanoophrys*-like ciliate multiplied during *in vitro* culture, with density increasing 10-fold over the 7 d period of the incubations. Comparable increases in ciliate numbers occurred in experimental infections of *Cancer pagurus* and *C. magister* with *Mesanoophrys* sp. (Bang et al. 1972, Cain & Morado 2001). Densities of the scuticociliate fish pathogen *Philasterides dicentrarchi* have also been observed to multiply 5-fold over 7 d (reaching a plateau phase after 3 d) during *in vitro* culture using a similar seeding density to that used in the present study (Inglesias et al. 2003).

Associated with the increase in growth of the ciliate *in vitro*, was the release of multiple extracellular proteases. As the ciliates were apparently always motile, and were still increasing in number when the experiment was terminated (Day 7), we concluded that the extracellular proteases detected in

the culture medium had been actively secreted. The relatively close correlation between the amount of secreted proteases and number of ciliates (Fig. 1) suggests that the secreted proteases did not accumulate in the medium, or at least not in active forms.




Fig. 3. Digestion of deep abdominal flexor (DAF) muscle proteins by proteases present in ciliate culture medium; sizes are indicated on the left. Lane A: molecular mass standards; Lane B: DAF protein sample incubated in absence of ciliate culture medium; Lane C: DAF protein sample incubated in presence of ciliate culture medium; Lane D: ciliate culture medium sample; Lane E: uninoculated (control) culture medium sample. Note degradation of DAF muscle proteins in Lane C (arrows), and also reduced intensity of FCS proteins in ciliate culture medium sample (Lane D) compared to control medium sample (Lane E). MHC: myosin heavy chain; P: paramyosin; Ac: actin; TnI: Troponin I family



Inhibitory studies using both spectrophotometry and substrate-impregnated SDS-PAGE methods (Table 1) indicated that the proteases secreted by the ciliate are of the metallo class, with complete and partial inhibition of activity being observed using the inhibitors 1,10-phenanthroline and EDTA, respectively. Metalloproteases produced by several protozoan parasites are suggested to contribute to the invasion and degradation of host tissues (McKerrow et al. 1993), and are important in the pathogenesis of parasitic infections. Metalloproteases have been documented from several marine pathogens during *in vitro* culture, including the piscine haemoflagellate *Cryptobia salmositica* (Zuo & Woo 1998), the bivalve pathogen *Vibrio alginolyticus* (Nottage & Birkbeck 1987), and the fish pathogen *Aeromonas salmonicida* (Arnesen et al. 1995). In the scuticociliate parasite of fish *Uronema marinum*, Lee et al. (2003) detected metalloprotease secretion as early as 10 min in ciliate cell suspensions using fluorescence polarisation. They suggested that the metalloproteases are involved in the pathogenesis of this ciliate infection in fishes. Other ciliate species such as *Tetrahymena pyriformis* and *T. thermophila* have previously been shown to contain and secrete proteases (Banno et al. 1983, Straus et al. 1992). In these studies, proteases were suggested to be involved in both intracellular and extracellular digestion. In addition, lysosomal-released enzymes from *T. thermophila* have been implicated in the process of extracellular digestion and parasite nutrition, with proteolytic enzymes facilitating the release of host particulate and soluble nutrients (Florin-Christensen et al. 1985, Tiedtke & Rasmussen 1988). Thus, the proteases released by the present ciliate under investigation may also play a similar role in nutrient uptake, both *in vitro* and *in vivo*.

During ciliate infections of other crustaceans, extensive host tissue penetration and destruction are common observations (Armstrong et al. 1981, Sparks et al. 1982, Hibbits & Sparks 1983, Messick & Small 1996, Athanassopoulou et al. 2004). The present study has shown that proteases present in the ciliate culture medium selectively degrade the myosin heavy chain and one of the Troponin I family of proteins from *Nephrops norvegicus* abdominal flexor muscle *in vitro* (Fig. 3). Several unidentified muscle proteins with masses in the range of 100 to 120 kDa were also degraded, but the paramyosin, actin and other muscle proteins remained unaffected by the proteases. The myosin heavy chain proteins are present in all muscle groups within the Norway lobster, and it seems likely that the same proteases secreted by the ciliate *in vivo* will degrade tissues containing these muscle groups and so be mediators of extensive tissue penetra-

tion and digestion. Similarly, the multiple proteases secreted by the apicomplexan parasite *Perkinsus marinus* appear to degrade the gut epithelium and basement membranes of its host, the eastern oyster *Crassostrea virginica*, assisting parasitic invasion and spread through the oyster tissues (La Peyre et al. 1996).

In addition to tissue penetration/digestion, haemocytopenia is also a classic sign of advanced ciliate infections in crustaceans (Cattaneo 1888, Armstrong et al. 1981, Sparks et al. 1982, Cain & Morado 2001). The Norway lobsters that were infected with the ciliate under study also had very few haemocytes present in the haemolymph samples taken to initiate *in vitro* cultures (H. J. Small pers. obs.). Released proteins, including proteases, can modulate the immune system of a pathogen's host. For example, the 64 kDa protease and extracellular proteins secreted by *Aeromonas salmonicida*, which suppress the humoral immune response of Atlantic salmon *Salmo salar*, result in leucopenia and allow the bacteria to spread via the circulatory system and invade other tissues (Ellis et al. 1981, Hussain et al. 2000). Released proteins from *A. salmonicida* are also reported to induce degranulation of eosinophilic granular cells, releasing vasoactive compounds leading to acute shock in Atlantic salmon (Ellis et al. 1981). Similarly, the pathogenic haemoflagellate *Cryptobia salmositica* secretes a metalloprotease which can directly lyse fish erythrocytes causing anaemia (Zuo & Woo 2000), while the proteases secreted by the apicomplexan oyster parasite *Perkinsus marinus* compromise the oyster's host defence mechanisms by inhibiting haemocyte motility and degrading plasma lysozyme and haemagglutinins (Garreis et al. 1996). Therefore, the reduction in circulating haemocytes during ciliate infections of crustaceans, previously attributed to consumption (Bang et al. 1972, Armstrong et al. 1981, Messick & Small 1996), may also be due in part to the effect of secreted proteases.

In conclusion, the histophagous ciliate under study, thought to be a species of the genus *Mesanoophrys*, secreted several metalloproteases during *in vitro* culture. These may well be important in the invasion of host tissues by the ciliate and also in the parasite's nutrition *in vivo*. Thus these metalloproteases may be key virulence factors and responsible for the tissue pathologies observed in infected crustaceans. Further studies are required to ascertain the exact contribution of these enzymes to the host-parasite relationship.

*Acknowledgements.* H.J.S. received a Natural Environment Research Council Studentship (NER/S/A/2000/03368). Thanks to Dr. Jeffrey Shields, who critically read the manuscript.

## LITERATURE CITED

- Appleton PL, Vickerman KV (1998) *In vitro* cultivation and development cycle in culture of a parasitic dinoflagellate (*Hematodinium* sp.) associated with mortality of the Norway lobster (*Nephrops norvegicus*) in British waters. *Parasitology* 116:115–130
- Armstrong DA, Burren EM, Sparks AK (1981) A ciliate infection (*Paranophrys* sp.) in laboratory-held Dungeness crabs, *Cancer magister*. *J Invertebr Pathol* 37:201–209
- Arnesen JA, Eggset G, Jørgensen TO (1995) Partial purification and characterization of extracellular metalloproteases from *Aeromonas salmonicida* ssp. *J Fish Dis* 18:283–295
- Athanassopoulou F, Speare D, Cawthorn RJ, MacMillan R, Despres B (2004) Pathology of *Anophryoides haemophila* (Scuticociliatida: Orchitophryidae), parasite of American lobster *Homarus americanus* kept under experimental conditions. *Aquaculture* 236:103–117
- Bang FB, Audouin J, Leglise M (1972) Ciliate infection of the blood of the edible crab, *Cancer pagurus*, in holding tanks in Brittany, France. *J Invertebr Pathol* 20:226–227
- Banno Y, Yano K, Nozawa Y (1983) Purification and characterization of a secreted proteinase from *Tetrahymena pyriformis*. *Eur J Biochem* 132:563–568
- Cain TA, Morado JF (2001) Changes in total hemocyte and differential counts in Dungeness crabs infected with *Mesanoophrys pugettensis*, a marine facultative parasitic ciliate. *J Aquat Anim Health* 13:310–319
- Cattaneo G (1888) Su di un infusorio ciliato, parassito del sangue del *Carcinus maenas*. *Zool Anz* 11:456–459
- Ellis AE, Hastings TS, Munro ALS (1981) The role of *Aeromonas salmonicida* extracellular products in the pathology of furunculosis. *J Fish Dis* 4:41–51
- Florin-Christensen J, Florin-Christensen M, Knudsen J, Rasmussen L (1985) Cytolytic activity released from *Tetrahymena*. *J Protozool* 32:657–660
- Garreis KA, La Peyre JF, Faisal M (1996) The effects of *Perkinsus marinus* extracellular products and purified proteases on oyster defence parameters *in vitro*. *Fish Shellfish Immunol* 6:581–597
- Grolier CA, Leglise M (1977) *Paranophrys carcini* n. sp., cilié *Philasterina* récolté dans l'hémolymphe du crabe *Cancer pagurus* Linné. *Protistologica* 13:503–507
- Hibbits J, Sparks AK (1983) Observations on the histopathology caused by a parasitic ciliate (*Paranophrys* sp.?) in the isopod *Gnorimosphaeroma oregonensis*. *J Invertebr Pathol* 41:51–56
- Hussain I, Mackie C, Cox D, Alderson R, Birkbeck TH (2000) Suppression of the humoral immune response of Atlantic salmon, *Salmo salar* L. by the 64 kDa serine protease of *Aeromonas salmonicida*. *Fish Shellfish Immunol* 10:359–373
- Inglesias R, Paramá A, Alvarez MF, Leiro J, Aja C, Sanmartín ML (2003) *In vitro* growth requirements for the fish pathogen *Philasterides dicentrarchi* (Ciliophora, Scuticociliatida). *Vet Parasitol* 111:19–30
- Klemba M, Goldberg DE (2002) Biological roles of proteases in parasitic protozoa. *Annu Rev Biochem* 71:275–305
- La Peyre JF, Faisal M (1995) *Perkinsus marinus* produces extracellular proteolytic factors *in vitro*. *Bull Assoc Fish Pathol* 15:28–31
- La Peyre JF, Schafhauser DY, Rizkalla EH, Faisal M (1995) Production of serine proteases by the oyster pathogen *Perkinsus marinus* (Apicomplexa) *in vitro*. *J Eukaryot Microbiol* 42:544–551
- La Peyre JF, Yarnall HA, Faisal M (1996) Contribution of *Perkinsus marinus* extracellular products in the infection of Eastern oysters (*Crassostrea virginica*). *J Invertebr Pathol* 68:312–313
- Lee EH, Kim CS, Cho JB, Ahn KJ, Kim KH (2003) Measurement of protease activity of live *Uronema marinum* (Ciliata: Scuticociliatida) by fluorescence polarisation. *Dis Aquat Org* 54:85–88
- McKerrow JH (1989) Parasite proteinases. *Exp Parasitol* 68:111–115
- McKerrow JH, Sun E, Rosenthal PJ, Bouvier J (1993) The proteases and pathogenicity of parasitic protozoa. *Annu Rev Microbiol* 47:821–853
- Messick GA, Small EB (1996) *Mesanoophrys chesapeakeensis* n.sp., a histophagous ciliate in the blue crab, *Callinectes sapidus*, and associated histopathology. *Invertebr Biol* 115:1–12
- Morado JF, Small EB (1994) Morphology and stomatogenesis of *Mesanoophrys pugettensis* n. sp. (Scuticociliatida: Orchitophryidae), a facultative parasitic ciliate of the Dungeness crab, *Cancer magister* (Crustacea: Decapoda). *Trans Am Microsc Soc* 113:343–364
- Morado JF, Giesecke RH, Syrjala SE (1999) Molt related mortalities of the Dungeness crab *Cancer magister* caused by a marine facultative ciliate *Mesanoophrys pugettensis*. *Dis Aquat Org* 38:143–150
- Nottage AS, Birkbeck TH (1987) Purification of a proteinase produced by the bivalve pathogen *Vibrio alginolyticus* NCMB 1339. *J Fish Dis* 10:211–220
- Poisson R (1930) Observations sur *Anophrys sarcophaga* Cohn (= *A. maggii* Cattaneo) infusoire holotriche marin et sur son parasitisme possible chez certains crustacés. *Bull Biol Fr Belg* 64:288–331
- Rockey DD, Fryer JL, Rohovec JS (1988) Separation and *in vivo* analysis of two extracellular proteases and the T-hemolysin from *Aeromonas salmonicida*. *Dis Aquat Org* 5:197–204
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular cloning: a laboratory manual*, 2nd edn. Cold Spring Harbor Laboratory (CSH), Cold Spring Harbor, NY
- Sarath G, De la Motte RS, Wagner FW (1989) Protease assay methods. In: Benyon RJ, Bond JS (eds) *Proteolytic enzymes: a practical approach*, Vol 3. IRL Press, Oxford
- Schulte W, Scholze H (1989) Action of major protease from *Entamoeba histolytica* on proteins of the extracellular matrix. *J Protozool* 36:538–543
- Small HJ, Neil DM, Taylor AC, Batman K, Coombs GH (2005) A parasitic scuticociliate infection in the Norway lobster (*Nephrops norvegicus*). *J Invertebr Pathol* 90:108–117
- Söderhäll K, Unestam T (1975) Properties of extracellular enzymes from *Aphanomyces astaci* and their relevance in the penetration process of crayfish cuticle. *Physiol Plant* 35:140–146
- Sparks AK, Hibbits J, Fegley J (1982) Observations on the histopathology of a systemic ciliate (*Paranophrys* sp.?) disease in the Dungeness crab, *Cancer magister*. *J Invertebr Pathol* 39:219–228
- Straus JW, Migaki G, Finch MT (1992) Assessment of proteolytic enzymes in *Tetrahymena thermophila*. *J Protozool* 39:655–662
- Tiedtke A, Rasmussen L (1988) Lysosomal enzymes in extracellular digestion in the unicellular eukaryote *Tetrahymena*. *J Cell Physiol* 136:554–556
- Zuo X, Woo PTK (1998) *In vitro* secretion of the metalloprotease by the pathogenic haemoflagellate *Cryptobia salmositica* Katz, 1951 and stimulation of the protease production by collagen. *J Fish Dis* 21:249–255
- Zuo X, Woo PTK (2000) *In vitro* haemolysis of piscine erythrocytes by purified metalloprotease from the pathogenic haemoflagellate *Cryptobia salmositica* Katz. *J Fish Dis* 23:227–230