

Photochemical production of dissolved inorganic nitrogen and primary amines from dissolved organic nitrogen in waters of two estuaries and adjacent surficial groundwaters

Dirk J. Koopmans¹, Deborah A. Bronk^{2,*}

¹Department of Marine Sciences, University of Georgia, Athens, Georgia 30602-3636, USA

²Virginia Institute of Marine Science, College of William and Mary, Gloucester Point, Virginia 23062-1346, USA

ABSTRACT: Recent studies have shown that in a number of humic-rich surface waters in North America, NH_4^+ is released when dissolved organic matter (DOM) is exposed to sunlight. However, photochemical NH_4^+ production has not been observed in all surface waters, and factors that contribute to it are not well understood. We hypothesized that the presence or absence of NH_4^+ photo-production may be affected by the light exposure history of DOM. The present study was undertaken to determine whether DOM from surficial groundwaters, with minimal light exposure history, would produce labile nitrogen (N) photoproducts more consistently. In this study, estuarine surface waters and surficial groundwaters, collected just adjacent to estuaries in Georgia and South Carolina, USA, were exposed to sunlight to quantify the photochemical production of NH_4^+ , dissolved primary amines (DPA), and NO_2^- . The photoproduction of NH_4^+ was observed in 4 of 5 irradiated estuarine surface water samples but in only 2 of 13 groundwater samples. In contrast, NH_4^+ concentrations decreased in 5 of 13 groundwater samples when exposed to sunlight. The results indicate that a small amount of NH_4^+ may be lost from waters in which groundwater-derived DOM is first exposed to sunlight. No consistent trends were observed in the photoproduction or loss of DPA and NO_2^- .

KEY WORDS: UV radiation · Dissolved organic nitrogen · Groundwater · Ammonium · Dissolved primary amines · Nitrite

Resale or republication not permitted without written consent of the publisher

INTRODUCTION

Dissolved organic nitrogen (DON) accounts for 40 to 90% of the nitrogen (N) in rivers of the southeastern United States (Alberts & Takács 1999). Recent research has shown that microbes can utilize estuarine DON at rates that suggest it is a significant source of bioavailable N to estuaries and the coastal ocean (Seitzinger & Sanders 1997; reviewed in Bronk 2002). One process that may contribute to the bioavailability of riverine DON is photochemical decomposition. Photochemical processes have been shown to result in the release of ammonium (NH_4^+), dissolved primary amines (DPA), nitrite (NO_2^-), and unidentified labile N compounds

when more refractory DON compounds are exposed to sunlight (Bushaw et al. 1996, Bushaw-Newton & Moran 1999, Kieber et al. 1999, reviewed in Moran & Zepp 1997). Of these, the observed rates of photoproduction of NH_4^+ are generally the greatest, between 0.04 and 0.4 $\mu\text{M h}^{-1}$ (Bushaw et al. 1996, Gao & Zepp 1998, Gardner et al. 1998). This production occurs at a wide range of sites rich in humic substances, including boreal ponds in Manitoba, a swamp and estuary in Georgia (Bushaw et al. 1996, Gao & Zepp 1998), a river and bayou in Louisiana (Wang et al. 2000), and a humic-rich lake in Venezuela (Gardner et al. 1998). However, photochemical release of labile N is not always observed. For example, no NH_4^+ release was observed during the irradiation of river, lake, and groundwater samples from boreal catchments in Sweden (Jørgensen et al. 1998, Bertilsson et al. 1999).

*Corresponding author. E-mail: bronk@vims.edu

The propensity for dissolved organic matter (DOM) to release N photoproducts is likely dependent on the source of the DOM. An additional factor that may contribute to the variability observed in photoproduction is light exposure history. For example, surface water DOM collected from stratified lakes and slow moving estuaries during the dry season would presumably have had a long exposure to light. In contrast, DOM collected from low order streams and ponds during a wet season would likely have experienced a shorter exposure time to light. These differences in light exposure, as well as variability in DOM source, likely contribute to the variability observed in the photoproduction of NH_4^+ and DPA in different studies (Bushaw et al. 1996, Bertilsson et al. 1999, Bushaw-Newton & Moran 1999).

A source of DON that has received little attention in photoproduction studies, with the exception of Bertilsson et al. (1999), is groundwater. In surficial groundwaters, DOM is leached from overlying soils, vegetation, and anthropogenic sources. In surficial groundwaters in the southeastern United States, DON concentrations have been studied only rarely, but dissolved organic carbon (DOC) concentrations are generally 400 to 800 $\mu\text{M C}$ (Thurman 1985). As a rough extrapolation, even if the average atomic C:N ratio of this DOM were high at 40:1 (i.e., Satilla River humic substances; Alberts & Filip 1994), there would be between 10 and 20 $\mu\text{M DON}$ present in this groundwater. We hypothesized that groundwater-derived DON may play a proportionally greater role in the photoproduction of labile N because of its limited exposure to sunlight.

In this study, the photoproduction of labile N (NH_4^+ , DPA, and NO_2^-) from DON collected from surficial groundwaters and from the epilimnion of estuaries was quantified. Groundwater samples were collected from residential and pristine sites in coastal Georgia and South Carolina and exposed to sunlight. To improve detection of photochemically produced labile N, the




Fig. 1. Map of the South Atlantic Bight with 3 sampling sites: Site I, North Inlet, SC; Site II, Brunswick and St. Simon's Island, GA; Site III, Satilla River Estuary, GA

ratio of DON to dissolved inorganic nitrogen (DIN) was increased in a subset of surface water and groundwater samples collected from the Satilla River Estuary, which has been the focus of previous photochemical studies (Bushaw et al. 1996, Gao & Zepp 1998).

MATERIALS AND METHODS

Surficial groundwater samples (<10 m depth) were collected from 3 sites: North Inlet, South Carolina; Brunswick/St. Simon's Island, Georgia; and the Satilla River, Georgia. Water column samples were collected at North Inlet and the Satilla River sites (Fig. 1, Table 1).

Site I: North Inlet, South Carolina. North Inlet is a pristine estuary 90 km northeast of Charleston, South

Table 1. Summary of sites where groundwater and surface water samples were collected including the treatments used to increase the dissolved organic nitrogen (DON) to NH_4^+ ratio of samples from Site III, and the light source used for irradiating samples

Site no. and location	Sample date	Site description	Organic matter treatments	Light source
I. North Inlet, SC	Feb 1999	<i>Spartina alterniflora</i> salt marsh	None	Sunlight
II. Brunswick and St. Simon's, GA	Sep 1999	Suburban communities	None	Sunlight
III. Satilla River Estuary, GA	May 2000, Jul 2000	Low population density, pine forests	Rotary evaporation, ultrafiltration	Xenon arc lamp

Carolina, surrounded by *Spartina alterniflora* salt marsh. Groundwater samples were collected in February 1999 from 5 polyvinyl chloride (PVC) groundwater monitoring wells (1.2 to 2.4 m deep): 2 in an upland pine forest, 1 mid-marsh, and 2 in the marsh edge on the bank of a tidal creek. The marsh wells drew water from a layer of sand beneath the marsh mud. Approximately 10 l of water was removed and discarded from each well before 1.5 l of sample was collected. One water column sample was also collected from just below the surface (<0.5 m depth) of Town Creek, a creek that drains the study area. Within 7 h after collection, the samples were filtered through a pre-combusted (500°C for 2 h) GF/F filter (nominal cutoff of 0.7 µm) and a Supor filter (0.2 µm cutoff). All groundwater samples collected in this study were initially sulfidic. Oxygen was reintroduced to groundwaters prior to photooxidation by gently shaking samples in bottles with headspace. The samples were stored at 4°C until photooxidized.

Site II: Brunswick and St. Simon's Island, Georgia.

Brunswick is a town of approximately 17 000 inhabitants, 100 km south of Savannah, Georgia. St. Simon's Island is adjacent to Brunswick and has been developed primarily for vacation homes. Groundwater was collected from 1 PVC-lined residential well (the well screen depth was approximately 3 to 5 m) in a suburban community in Brunswick and 3 similar wells on St. Simon's Island in September 1999. The surficial aquifer in this area extends to >40 m depth (Clarke et al. 1990), so all the groundwater collected was from this surficial aquifer. Water was pumped from the wells until temperature and dissolved oxygen readings stabilized (over 20 l of water was discarded) before collecting samples. Each sample was stored on ice for up to 2 wk until they were filtered through GF/F and Supor filters and photooxidized.

Site III: Satilla River Estuary, Georgia. The Satilla River is a 'black water' river that receives DOM from extensive swamps in the drainage area of Georgia's coastal plain. Two groundwater and 2 water column samples were collected from the Satilla River Estuary during the summer of 2000. Groundwater samples were drawn from surficial aquifers (1.5 m depth) through a stainless steel well point sampler (Valiela & Costa 1988), 20 m or less from Umbrella Creek, a tidal creek that drains into the Satilla River Estuary. Water column samples were collected just below the water surface (<0.5 m) in May 2000 from the head of the estuary (non-saline) and from Umbrella Creek (salinity of 25 parts per thousand). The surface and groundwater samples were filtered within 14 h of collection through GF/F and Supor filters and stored at 4°C until subdivision into fractions for photooxidation pre-treatments 1 to 2 wk later.

Treatments of Site III samples prior to photo-oxidation. To improve the detection of photochemically produced NH_4^+ , rotary evaporation and tangential flow ultrafiltration were used to increase the concentration of DON relative to the background concentration of NH_4^+ , which was greater than 20 µM in some groundwater samples. In the rotary evaporation treatment, 1 l of sample was brought to pH 9.5 and rotary evaporated (Büchi Rotavapor R110, Büchi Analytical Ltd, Manchester, UK) at a vacuum of 71 cm Hg at room temperature for 45 min. The change in pH resulted in the conversion of ambient NH_4^+ to volatile ammonia (NH_3) gas and the loss of NH_3 during evaporation. As a result, the background NH_4^+ concentration, which might obscure a small amount of photo-production, was reduced. Immediately following rotary evaporation, the sample was diluted to its original volume with deionized water and acidified to the original pH. The volumes of the rotary evaporated Satilla River surface water, Umbrella Creek surface water, and Umbrella groundwater (GW-A; replicate samples are designated A, B or C) samples were diluted an additional 10 to 14% to provide sufficient sample volume for the analyses. All rotary evaporated samples were gently shaken and then stored with headspace for at least 24 h to allow oxygen to dissolve into solution.

In the tangential flow ultrafiltration treatment, 10 l of the unmanipulated sample from each site was filtered through a series of two 1 kDa molecular weight cut off cellulose acetate ultrafiltration membranes (Millipore) using a DC10L tangential flow ultrafiltration unit (Amicon Inc, Beverly, MA, USA). Before the sample was ultrafiltered, the membranes were cleaned with 20 l of 0.2 N NaOH and rinsed with 60 to 80 l of deionized water. The sample was brought to pH 7 and forced across the 2 membranes, at inlet and outlet pressures of 3.5 kg cm². The volume of the retentate, containing the fraction of the DON that did not pass through the membranes, was brought to one-third the initial sample volume. Then the outlet pressure was briefly reduced to 0.4 kg cm² to improve recovery of DOM adsorbed to the membranes, as recommended by Benner et al. (1997). The retentate was brought back to the original pH and stored at 4°C for later photooxidation experiments. The permeate, the fraction of the DON that passed through the membrane, was discarded because the concentration of DON was low (approximately 5 µM) such that detecting photoproduction of labile N from it would have been unlikely.

Photooxidation of samples. To reduce the number of bacteria in irradiated and control samples and to minimize microbial uptake of photoproducts, all samples from Sites I, II, and III were filtered through GF/F and 0.2 µm Supor filters and stored at 4°C until they were

exposed to artificial sunlight. Samples from Sites I and II were GF/F and 0.2 μm Supor filtered only once, 24 to 80 h before photooxidation, to minimize the possibility of introducing DON artifacts of filtration. Samples from Site III were GF/F and 0.2 μm Supor filtered twice, immediately after collection and again 20 to 40 h before photooxidation, to reduce bacterial regrowth that may have occurred during rotary evaporation and ultrafiltration. Each sample was poured into 6 quartz tubes, 3 of which were wrapped in aluminum foil as experimental controls. The experimental and control tubes of samples from Sites I and II were placed in a cool water bath (10 to 15°C) and exposed to sunlight in Athens, Georgia (latitude 34° N) for 5 to 10 h. The experimental and control tubes from Site III were placed in adjacent cold water baths and exposed to 8 h of artificial sunlight at an intensity of 1000 $\mu\text{E m}^{-2} \text{s}^{-1}$ generated by a xenon arc lamp (Suntest CPS solar simulator, Atlas Electric Devices, Chicago, IL, USA). The spectrum of UV irradiance had been adjusted to closely match midday sunlight in June in Athens, Georgia. The heat of the lamp brought the temperatures of both water baths from 5 to 10°C. Following irradiation the samples were subdivided and frozen for NH_4^+ , NO_2^- , and DPA analyses. The absorbance of samples from Site III, before and after irradiation, was determined at a wavelength of 350 nm by a Beckman DU 640 spectrophotometer.

Chemical analyses. For irradiated and control treatments, the reported nutrient concentrations are the means of triplicate analyses performed on each of the 3 control and irradiated experimental replicates. Ammonium was analyzed spectrophotometrically by the manual phenol-hypochlorite method, which responds negligibly to amino acids (Koroleff 1983, detection limit 0.05 μM). Nitrite concentrations in waters from Sites I and III were measured by the chemiluminescent method (Garside 1982; detection limit of 0.1 μM). Nitrite concentrations in waters from Site II were determined by the colorimetric method (Grasshoff 1983; detection limit of 0.25 μM in highly colored water). Concentrations of DPA were measured spectrofluorometrically (Parsons et al. 1984); DPA concentrations were reported as glycine equivalents after correction for the fluorescence of ambient NH_4^+ (Liebezeit & Behrends 1999). Samples for DOC analyses were acidified to pH 3, sparged for 15 min with carbon-free air, and analyzed using a Shimadzu TOC-5000 (Moran et al. 1999).

The concentration of DON was determined by subtracting the concentration of DIN (NO_3^- , NO_2^- , and NH_4^+) from the concentration of total dissolved nitrogen (TDN) determined by persulfate oxidation (Bronk et al. 2000). The humic-N and C concentrations were determined by humic adsorption to Supelite DAX-8 macroporous resin (the current replacement for

Amberlite XAD-8, Supelco, Bellefonte, Pennsylvania; Aiken 1985).

Data analyses. The concentrations of a given compound determined in the 3 irradiated replicates were compared to the concentrations in the corresponding controls with a 2-tailed Student's *t*-test ($\alpha = 0.05$). Equal variance was assumed for irradiated and control concentrations. When the mean concentration of a constituent in the irradiated water was significantly greater than the mean concentration in the controls ($p \leq 0.05$), the difference was interpreted as photo-production. When the reverse was true, the difference was interpreted as a photochemically mediated loss.

Rate normalization. To compare the net rates of NH_4^+ photoproduction between samples and treatments, rates were normalized to DON concentrations by dividing the NH_4^+ production rate by the initial DON concentration in the sample. For Site III, NH_4^+ photoproduction was normalized to absorbance by dividing net production rates by the sample absorption coefficient at 350 nm (a_{350}), defined as $2.303 \times A_{350}/b$, where A_{350} is the measured absorbance at 350 nm and b is the path length of light through the sample in meters (Miller & Zepp 1995). Photoproduction rates were corrected for self-shading at 350 nm by a light screening factor calculated using the derivation of a formula presented by Zepp (1982) given in Bushaw-Newton & Moran (1999): $(1 - e^{-a_{350} \times b})/(a_{350} \times b)$, where b is the average path-length of light through the irradiation vessel in meters (0.009 m in this study). This is a correction factor for the amount of light attenuated within a sample by DOM.

RESULTS

Water samples collected from 2 tidal creeks, the Satilla River, and shallow groundwaters were exposed to natural or artificial sunlight to examine photo-production of labile N. The initial DIN and DON concentrations and the effect of irradiation on NH_4^+ , NO_2^- , and DPA concentrations are presented below.

Inorganic and organic N concentrations

In surface water samples, most of the dissolved N (77 to 97%) was organic in nature. In the Satilla River and Umbrella Creek, most of that organic N (83 and 63%, respectively) was associated with humic substances (Table 2). The predominant forms of DIN were NO_3^- in the Satilla River and NH_4^+ in Umbrella and Town Creeks. In contrast, most of the dissolved N in groundwater samples was inorganic, specifically NH_4^+ . About 80% of the dissolved N in Marsh Creek GW-A and B samples and the Brunswick GW sample was NH_4^+ . In

Table 2. Concentration of dissolved organic nitrogen (DON; mean \pm SD), percentage of DON eluted with humic substances, the dissolved organic carbon to DON (C:N) ratio, and the absorptivity of surface and groundwater samples. Concentrations were determined immediately before the samples were photooxidized, except where noted. See Table 1 for site definitions. Two treatments were used at Site III: NH_4^+ was removed by rotary evaporation (RE) or DON was concentrated with ultrafiltration (UF). GW: groundwater; a_{350} : absorption coefficient at 350 nm; na: not available. Replicate samples taken from the same location are designated by A, B or C

Site no. and sample	DON ($\mu\text{M N}$)	% DON in humic-N	C:N	a_{350} (m^{-1})
Surface water				
I. Town Creek	12 \pm 0.9	65	na	na
III. Satilla River (RE)	37 \pm 0.8	83 ^a	39	43.8
III. Satilla River (UF)	107 \pm 1.1	na	38	128
III. Umbrella Creek (RE)	22 \pm 0.5	63 ^a	23	9.1
III. Umbrella Creek (UF)	50 \pm 1.3	na	25	21.2
Shallow groundwater				
I. Upland GW-A	27 \pm 1.0	57	na	na
I. Upland GW-B	64 \pm 1.6	66	na	na
I. Marsh GW	16 \pm 1.7	>90	na	na
I. Creek GW-A	1.9 \pm 2.6	>90	na	na
I. Creek GW-B	12 \pm 0.9	54	na	na
II. Brunswick GW	1 \pm 0.1	54	56	na
II. St. Simon's GW-A	35 \pm 0.9	57	43	na
II. St. Simon's GW-B	14 \pm 0.8	62	47	na
II. St. Simon's GW-C	21 \pm 0.5	51	43	na
III. Umbrella GW-A (RE)	26 \pm 0.4	71 ^a	46	21.8
III. Umbrella GW-A (UF)	67 \pm 1.8	na	43	56.8
III. Umbrella GW-B (RE)	9 \pm 0.3	63 ^a	74	3.7
III. Umbrella GW-B (UF)	25 \pm 3.3	na	48	10.4

^aAnalysis performed before RE treatment

Table 3. Effect of rotary evaporation (RE) at pH 9.5 for 45 min on the concentration of NH_4^+ and DON in surface and groundwater samples collected at Site III, the Satilla River Estuary. Following rotary evaporation the Satilla River, Umbrella Creek and Umbrella GW-A samples were diluted with deionized water (DI) to provide sufficient volume for photooxidation (Dilution with DI)

	Initial NH_4^+ (μM)	Final NH_4^+ (μM)	Initial DON (μM)	DON loss (%)	Dilution with DI (%)
Satilla River	0.33	0.26	46.4	6	14
Umbrella Creek	6.29	0.30	22.4	0	13
Umbrella GW-A	8.62	0.14	32.9	13	10
Umbrella GW-B	21.9	1.14	11.8	19	0

all groundwater samples except Upland GW-B, St. Simon's GW-A and B, and Umbrella GW-A, over 50% of dissolved N was NH_4^+ . The concentration of NO_3^- was highest in the Brunswick GW sample, where it accounted for 18% of the dissolved N. In all other groundwater samples NO_3^- accounted for 3% or less of the N. Most of the DON in groundwater samples, similar to DON in surface waters, was associated with humic substances (Table 2).

Effect of rotary evaporation and tangential flow ultrafiltration

As a result of the rotary evaporation treatment, NH_4^+ concentrations were reduced to low levels (1.2 to 0.1 μM), a reduction of up to 98% from ambient concentrations (Table 3). There was a concurrent loss of up to 19% of the DON during rotary evaporation, perhaps due to the volatilization of amine groups. The ultrafiltration treatment only slightly reduced the average NH_4^+ concentrations, while increasing the DON concentration approximately 2-fold (Table 4). There was some loss of DON through the 1 kDa membranes, but the overall retention was about 70%. At less than a 10-fold concentration factor, tangential flow ultrafiltration may not completely separate high and low molecular weight (LMW) compounds (Benner et al. 1997), so a significant fraction of the retentate DON in these samples may have been LMW compounds. Concentrations of DOC, consistent with DON, increased 1.7- to 2.3-fold during ultrafiltration, with 60 to 75% of the DOC retained by the membranes (data not shown).

Rates of photoproduction

The photoproduction of NH_4^+ was observed in river and creek samples. However, there was more photochemically mediated loss of NH_4^+ than production in the groundwater samples. Statistically significant (*t*-test, $p < 0.05$) photoproduction of NH_4^+ was observed in 4 of 5 surface water experiments (Table 5). In groundwaters from Sites I and II, the photochemically mediated loss of NH_4^+ was observed in 3 of 9 irradiated samples ($p < 0.01$), but photoproduction was observed in only 2 of 9 irradiated samples ($p < 0.05$; Table 5). No net photochemical production or loss of NH_4^+ was observed in the rotary evaporation treatment of irradiated groundwater samples from Site III, but NH_4^+ was lost from both ultrafiltration treatments of the same samples ($p < 0.01$; Table 5).

Table 4. Effect of tangential flow UF (1 kDa molecular weight cut-off membrane) on the retentate concentrations of NH_4^+ and DON in samples from Site III, the Satilla River Estuary. Percentages of DON that passed through the UF membranes (permeate) and that were unaccounted for are also given

Sample	Initial NH_4^+ (μM)	Final NH_4^+ (μM)	Initial DON (μM)	DON conc. factor	DON in retentate (%)	DON in permeate (%)	Missing DON (%)
Satilla River	0.3	0.4	47.7	2.2	73	13	14
Umbrella Creek	6.3	5.5	24.3	2.0	70	24	6
Umbrella GW-A	8.6	7.7	31.2	2.2	76	16	8
Umbrella GW-B	21.9	20.5	13.0	2.0	69	23	8

Table 5. Mean NH_4^+ concentrations ($\pm\text{SD}$) in surface and groundwaters, unexposed to light (controls) and irradiated. Samples from Site III were RE at pH 9.5 to reduce NH_4^+ concentrations or concentrated by tangential flow UF before irradiation. P and L mark statistically significant production and loss of NH_4^+ , respectively, in irradiated samples ($\alpha = 0.05$, Student's *t*-test). The DON normalized (norm.) rate is the rate of photochemical NH_4^+ production divided by the DON concentration. –: no significant change

Site no. and sample	Control NH_4^+ (μM)	Irradiated NH_4^+ (μM)	Production or loss	DON norm. $\times 10^{-3} \text{ h}^{-1}$
Surface water				
I. Town Creek	0.21 \pm 0.02	0.45 \pm 0.02	P	2.9 \pm 0.4
III. Satilla River (RE)	0.29 \pm 0.01	0.44 \pm 0.02	P	0.50 \pm 0.1
III. Satilla River (UF)	0.33 \pm 0.04	0.44 \pm 0.03	P	0.1 \pm 0.1
III. Umbrella Creek (RE)	0.41 \pm 0.11	0.54 \pm 0.09	–	
III. Umbrella Creek (UF)	5.65 \pm 0.06	5.88 \pm 0.08	P	0.6 \pm 0.2
Shallow groundwater				
I. Upland GW-A	26.08 \pm 0.12	25.05 \pm 0.10	–	
I. Upland GW-B	27.89 \pm 0.05	25.69 \pm 0.15	L	–4.9 \pm 0.4
I. Marsh GW	19.48 \pm 0.05	19.50 \pm 0.09	–	
I. Creek GW-A	21.18 \pm 0.31	20.53 \pm 0.07	–	
I. Creek GW-B	14.52 \pm 0.15	14.96 \pm 0.15	P	7.3 \pm 3.4
II. Brunswick GW	4.53 \pm 0.12	4.68 \pm 0.11	–	
II. St. Simon's GW-A	18.93 \pm 0.14	16.84 \pm 0.11	L	–6.4 \pm 0.5
II. St. Simon's GW-B	6.74 \pm 0.09	7.91 \pm 0.05	P	9.1 \pm 0.8
II. St. Simon's GW-C	30.11 \pm 0.11	28.45 \pm 0.05	L	–8.1 \pm 0.6
III. Umbrella GW-A (RE)	0.42 ^a	0.54 \pm 0.08	–	
III. Umbrella GW-A (UF)	7.64 \pm 0.03	7.29 \pm 0.06	L	–0.6 \pm 0.1
III. Umbrella GW-B (RE)	1.21 \pm 0.06	1.30 \pm 0.07	–	
III. Umbrella GW-B (UF)	20.65 \pm 0.04	20.49 \pm 0.04	L	–0.8 \pm 0.3

^an = 1

The DON-normalized rates of NH_4^+ photoproduction and loss were greater among samples from Sites I and II than the manipulated samples from Site III (Table 5). The greatest DON-normalized rate of NH_4^+ photoproduction in surface waters was observed in Town Creek, from Site I ($2.9 \times 10^{-3} \text{ h}^{-1}$). The DON-normalized rates of NH_4^+ photochemical production observed in the 2 groundwater samples were 7.3×10^{-3} and $9.1 \times 10^{-3} \text{ h}^{-1}$. As in surface waters, the magnitude of DON-normalized rates of NH_4^+ loss among groundwaters from Sites I and II (-4.9×10^{-3} to $-8.1 \times 10^{-3} \text{ h}^{-1}$) were greater than rates observed in manipulated samples (Table 5).

Few significant differences in DPA and NO_2^- concentrations between irradiated and control treat-

ments were observed (Table 6). No statistically significant changes ($p < 0.05$) in DPA concentrations were observed in irradiated water column samples. Among groundwaters, the photochemical production of DPA ($p < 0.05$) was observed in only 1 of 9 irradiated samples from Sites I and II. The photochemically mediated loss of DPA ($p < 0.05$) was observed in both rotary evaporation and ultrafiltration treatments of Umbrella Creek GW-A but not in GW-B. Normalized to the concentration of DON in the samples, DPA production and loss rates were low (-1.6×10^{-3} , -0.6×10^{-3} and $1.3 \times 10^{-3} \text{ h}^{-1}$) but comparable to rates of NH_4^+ production and loss.

The photochemically mediated loss of NO_2^- ($p < 0.05$) was observed in 1 of 4 treatments of surface

Table 6. Mean dissolved primary amines (DPA) and NO_2^- concentrations (\pm SD) in surface and groundwaters, unexposed to light (controls) and irradiated. Samples from Site III were RE at pH 9.5 to reduce NH_4^+ concentrations or concentrated by tangential flow UF before irradiation. The DON normalized rates of photochemical DPA and NO_2^- production are presented for samples in which statistically significant ($p < 0.05$, Student's t -test) production or loss was observed. bd: below detection; -:no significant change

Site sample	Control DPA (μM)	Irradiated DPA (μM)	DON norm. DPA prod. $\times 10^{-3} \text{ h}^{-1}$	Control NO_2^- (μM)	Irradiated NO_2^- (μM)	DON norm. NO_2^- prod. $\times 10^{-3} \text{ h}^{-1}$
Surface water						
I. Town Creek	0.14 \pm 0.07	0.14 \pm 0.08	-	bd	bd	-
III. Satilla River (RE)	0.82 \pm 0.04	0.81 \pm 0.10	-	0.24 \pm 0.01	0.24 \pm 0.00	-
III. Satilla River (UF)	2.68 \pm 0.07	2.54 \pm 0.08	-	0.63 \pm 0.01	0.56 \pm 0.01	-0.08 \pm 0.02
III. Umbrella Creek (RE)	0.30 \pm 0.03	0.34 \pm 0.02	-	0.36 \pm 0.02	0.37 \pm 0.05	-
III. Umbrella Creek (UF)	0.66 \pm 0.05	0.66 \pm 0.05	-	0.49 \pm 0.02	0.48 \pm 0.01	-
Shallow groundwater						
I. Upland GW-A	0.67 \pm 0.07	0.53 \pm 0.07	-	bd	bd	-
I. Upland GW-B	1.29 \pm 0.02	1.88 \pm 0.25	1.31 \pm 0.77	bd	bd	-
I. Marsh GW	1.82 \pm 0.11	1.80 \pm 0.10	-	bd	bd	-
I. Creek GW-A	1.32 \pm 0.06	1.24 \pm 0.05	-	bd	bd	-
I. Creek GW-B	1.28 \pm 0.35	1.28 \pm 0.25	-	bd	bd	-
II. Brunswick GW	0.07 \pm 0.02	0.12 \pm 0.01	-	0.05 \pm 0.01	0.03 \pm 0.01	-1.05 \pm 0.01
II. St. Simon's GW-A	bd	0.16 \pm 0.08	-	0.08 \pm 0.00	0.08 \pm 0.01	-
II. St. Simon's GW-B	0.14 \pm 0.03	0.22 \pm 0.09	-	0.06 \pm 0.01	0.04 \pm 0.00	-
II. St. Simon's GW-C	bd	bd	-	0.04 \pm 0.00	0.07 \pm 0.01	0.16 \pm 0.01
III. Umbrella GW-A (RE)	1.19 \pm 0.08	0.86 \pm 0.10	-1.61 \pm 0.63	0.09 \pm 0.02	0.09 \pm 0.00	-
III. Umbrella GW-A (UF)	2.11 \pm 0.04	1.78 \pm 0.07	-0.6 \pm 0.15	0.23 \pm 0.02	0.24 \pm 0.01	-
III. Umbrella GW-B (RE)	0.25 \pm 0.02	0.25 \pm 0.04	-	0.09 \pm 0.02	0.07 \pm 0.00	-
III. Umbrella GW-B (UF)	0.48 \pm 0.09	0.37 \pm 0.02	-	0.09 \pm 0.00	0.08 \pm 0.01	-

water samples and in 1 of the 8 groundwater samples. Photochemical NO_2^- production ($p < 0.01$) was also observed in 1 of 8 groundwater samples. Normalized to the concentration of DON in the samples, NO_2^- production and loss rates were low relative to those observed for NH_4^+ and DPA (-1.1×10^{-3} , -0.1×10^{-3} and $0.2 \times 10^{-3} \text{ h}^{-1}$).

The absorbances of samples collected at Site III were compared before and after irradiation to measure the fading of chromophoric DOM caused by exposure to sunlight. In both the rotary evaporation and ultrafiltration treatments, the irradiated groundwater samples were slightly more faded (11 to 12%) relative to the fading of irradiated surface water samples (6 to 9%; Table 7). Normalized to either absorbance or DON, the NH_4^+ photoproduction rates of manipulated (Site III) samples indicated the same relative DOM photoreactivity. The greatest rate of NH_4^+ photoproduction was observed in concentrated ultrafiltered tidal creek water, and the loss observed in groundwaters was comparable to, or greater than, production observed in surface waters (Tables 5 & 7). DOC concentrations were also determined in samples from Site III, but there were no detectable differences in DOC between irradiated and control replicates (not shown).

DISCUSSION

In this study, 3 primary results were observed. First, there was photochemical NH_4^+ production in surface samples. Second, there was more photochemical loss than photoproduction of NH_4^+ in groundwater

Table 7. Effect of irradiation on the absorbance of samples collected at Site III, the Satilla River Estuary. The light screening factor is a measure of the self-shading of light within a sample by colored dissolved organic matter. Absorbance-normalized NH_4^+ production, corrected for light screening, is also presented for samples in which statistically significant ($p < 0.05$, Student's t -test) production or loss was observed. -: no significant change

Sample	Fading 350 nm (%)	Light screening factor	Absorbance-normalized production (nM m h^{-1})
Surface water			
Satilla River (RE)	8.3	0.83	0.52 \pm 0.07
Satilla River (UF)	7.0	0.59	0.17 \pm 0.09
Umbrella Creek (RE)	6.4	0.96	-
Umbrella Creek (UF)	9.3	0.91	1.45 \pm 0.63
Shallow groundwater			
Umbrella GW-A (RE)	11.5	0.91	-
Umbrella GW-A (UF)	12.4	0.78	-0.98 \pm 0.20
Umbrella GW-B (RE)	11.6	0.98	-
Umbrella GW-B (UF)	11.8	0.95	-2.00 \pm 0.67

samples. Third, consistent photoproduction of DPA and NO_2^- was not observed. Each of these results is discussed below.

Photochemical production of NH_4^+ in surface waters

DON in the surface waters of many systems, including the Satilla River, is known to release NH_4^+ when exposed to sunlight (Bushaw et al. 1996, Gao & Zepp 1998). The results of the present study support that conclusion. Ammonium was photochemically produced in most (4 of 5) river and creek samples and treatments, although photoproduction rates were low in manipulated samples, where NH_4^+ was removed or DON was concentrated. Normalized to DON concentrations, the rate of whole water photochemical NH_4^+ production observed in the unmanipulated surface water sample ($2.9 \times 10^{-3} \text{ h}^{-1}$) was similar to DON-normalized rates of NH_4^+ production observed from Satilla River Estuary fulvic acids and whole water from the Suwanee River (2.5×10^{-3} and 2.9 to $4.9 \times 10^{-3} \text{ h}^{-1}$; Bushaw et al. 1996), but less than rates of photochemical production from DOM in a Louisiana Bayou (12×10^{-3} to $26 \times 10^{-3} \text{ h}^{-1}$; Wang et al. 2000). Normalized to light absorbance at 350 nm, NH_4^+ photochemical production in manipulated samples at Site III was also less than absorbance-normalized production rates observed from whole DOM in the Satilla River and from the Louisiana Bayou (Bushaw et al. 1996, Wang et al. 2000).

The low rates of photochemical NH_4^+ production in samples from Site III may have been the result of the experimental manipulations. In a study of NH_4^+ photochemical production from DOM fractions in Bayou Trepagnier, Louisiana, Wang et al. (2000) observed the greatest DON-normalized rate of photochemical NH_4^+ production from LMW (<1000 Da) DOM. The ultrafiltration treatment used in the present study preferentially removed that LMW DOM. The NH_4^+ removal treatment may have also reduced photochemical NH_4^+ production in manipulated groundwater samples, in which 13 to 19% of the DON was lost, possibly due to volatilization of amine N.

Photochemically mediated loss of NH_4^+ in groundwaters

Statistically significant (*t*-test, $p < 0.05$) photochemical production of NH_4^+ from groundwater-derived DON was observed in only 2 of the 13 groundwater samples in this study. This suggests that shallow groundwater DON in these systems, despite limited previous exposure to sunlight, is not more likely to

photochemically produce NH_4^+ than surface water DON, as initially hypothesized.

In contrast, the photochemically mediated loss of NH_4^+ was observed in 5 of 13 groundwater samples (Table 5). There are a few potential sinks of NH_4^+ or NH_3 in these samples that could explain the observed loss of NH_4^+ (as NH_4^+ and NH_3 are in equilibrium at a given pH, a loss of NH_3 would lead to the loss of NH_4^+). First, the loss of NH_4^+ may have been the result of NH_4^+ oxidation; however, no parallel increase in the concentration of oxidized forms of inorganic N, NO_3^- or NO_2^- was observed during the experiment. Second, NH_4^+ may have been volatilized (as NH_3) due to microscale increases in the concentration of hydroxide produced during the photooxidation of DOM (Stumm & Morgan 1996). Third, NH_4^+ loss may have been the result of photochemical incorporation of NH_3 into DOM. Abiotic, oxidative incorporation of NH_3 into DOM has been observed in other systems, including the incorporation of NH_3 into triglycerides via a photochemically mediated process (Kieber et al. 1997). Aldehydes and ketoacids, products of the photooxidation of humic substances, also readily form imines from NH_3 and DPA, which can hydrolyze abiotically, releasing NH_4^+ gradually (Cohen & Ojanpera 1975). Ammonia also appears to react with keto and quinone groups in humic substances, in a process associated with oxidation, and is incorporated into biologically unavailable forms, such as pyrrole and indole N (Thorn & Mikita 1992). These mechanisms suggest that both biologically available (imines) and biologically unavailable (pyrrole and indole N) organic N forms are sinks for NH_3 in the process of photooxidation.

The tendency for NH_4^+ to be lost from photooxidized groundwater samples appeared to be influenced by NH_4^+ concentrations. The concentration of NH_4^+ in many of these samples was high, but this is common in surficial groundwaters in coastal Georgia and South Carolina (Joye et al. unpubl.). A linear regression of the change in NH_4^+ concentration after exposure to light against the initial NH_4^+ concentration indicated that groundwater samples with higher ambient NH_4^+ concentrations lost more NH_4^+ during irradiation (Fig. 2; $r^2 = 0.50$, $n = 12$). This suggests that in surface waters with low ambient NH_4^+ concentrations, losses of NH_4^+ due to the photooxidation of this DOM would be slight. In support of this possibility, no photochemical loss of NH_4^+ was observed in the groundwater samples in which NH_4^+ concentrations were experimentally reduced through rotary evaporation (Table 5).

While the photochemical production of NH_4^+ from aquatic DOM has been observed in many studies, the photochemically mediated loss of NH_4^+ has also been observed, generally in non-estuarine waters with

higher NH_4^+ concentrations ($>1 \mu\text{M}$). In humic-rich river, lake and groundwater samples from the River Öre and Svartberget catchments in northern Sweden, exposure to sunlight slightly reduced NH_4^+ concentrations (Bertilsson et al. 1999). In pasture and forest runoff waters from New Brunswick and Stanton, New Jersey, exposure to sunlight also slightly reduced NH_4^+ concentrations (Wiegner & Seitzinger 2001). The abiotic losses of NH_4^+ in groundwater samples exposed to sunlight in the present study were consistent with the abiotic losses of NH_4^+ above and suggest that the source of DON and the ambient NH_4^+ concentration affect the net direction of photochemical NH_4^+ production or loss.

Photochemical losses of NH_4^+ may have important implications for microbial decomposition of DOM at these sites. The irradiation of DOM produces an array of compounds that enhance bacterial production (Moran & Zepp 1997), of which NH_4^+ is only one. However, N often limits bacterial productivity in aquatic systems, so the photochemical production or loss of NH_4^+ and other forms of bioavailable N may have a significant effect on the microbial decomposition of DOM. Ammonium was photochemically produced by the irradiation of fulvic and humic acids from a boreal pond in northern Canada and from rivers and an estuary in the southeastern USA. In samples where the photochemical production of NH_4^+ was observed, a parallel release of bacteria from N limitation was also observed (Bushaw et al. 1996, Bushaw-Newton & Moran 1999). In studies where no photochemical NH_4^+ production or a net loss of NH_4^+ was observed, bacterial productivity was not enhanced (Bertilsson et al. 1999, Wiegner & Seitzinger 2001).

Photoproduction of DPA and NO_2^-

In the present study, the photochemical production of DPA and NO_2^- were not consistently observed. However, organic matter from the Skidaway River, Satilla River Estuary, and Suwanee Rivers in the southeastern USA and Lake Skärshult in southern Sweden has been shown to release DPA when exposed to sunlight (Jørgensen et al. 1998, Bushaw-Newton & Moran 1999, Tarr et al. 2001). This release, in addition to the release of NH_4^+ , may increase the productivity of N-limited bacteria as they degrade humic substances. Nitrite is also produced by the photolysis of humic substances (Kieber et al. 1999) and from NO_3^- by a number of reactions energized by UVB light (Zafiriou & True 1979). However, NO_2^- production was observed in only 1 of the 8 irradiated samples in the present study in which NO_2^- concentrations were above the limit of detection.




Fig. 2. Change in NH_4^+ concentration after an exposure of unmanipulated (◆), NH_4^+ removed (□), and DON concentrated (△) groundwater samples to natural or artificial sunlight (xenon arc lamp, $1000 \mu\text{E m}^{-2} \text{s}^{-1}$), plotted against the initial NH_4^+ concentration. Means \pm SD shown, $n = 3$

The lack of consistency in the photoproduction of DPA and NO_2^- may be due to the relatively low concentrations of DOM used in this study. Both of the photochemical studies mentioned above (Bushaw-Newton & Moran 1999, Kieber et al. 1999) were conducted on isolated and concentrated humic substances. The higher concentrations of humic substances in those studies may have allowed more consistent detection of the low rates of photochemical production that have been observed.

CONCLUSION

While the photooxidation of surface water DOM appears to be a source of NH_4^+ to many estuaries, the photooxidation of groundwater DOM at these sites does not appear to be. At high concentrations of NH_4^+ , as in the groundwater samples collected for this study, the photooxidation of groundwater DOM may slightly reduce NH_4^+ concentrations. This finding suggests an additional role for the photooxidation of DOM in the regulation of NH_4^+ concentrations in surface waters. The change in NH_4^+ concentrations in irradiated, unmanipulated groundwater samples ranged from a production of $0.1 \mu\text{M h}^{-1}$ to a loss of $0.3 \mu\text{M h}^{-1}$. These rates are low compared to summer rates of biotic NH_4^+ uptake (1.2 to $1.6 \mu\text{M h}^{-1}$; Bronk unpubl.) in the Altamaha Estuary, a major estuary between the study sites. During the winter and spring, however, biotic NH_4^+ uptake rates are significantly lower (0.11 to $0.12 \mu\text{M h}^{-1}$), only slightly greater than the median rate of loss observed in this study. This suggests that following events that introduce a large volume of groundwater to surface waters, fresh DOM may be a significant sink for NH_4^+ in the near surface, but this effect may be reduced by low NH_4^+ concentrations.

Acknowledgements. This research was supported by NOAA grant NAO6 RG0029 from Georgia Sea Grant to D.A.B. and NOAA grants NA87OZ0231 and NA97OZ0155 through a Georgia Sea Grant Coastal Management Fellowship to D.J.K. We thank Dr Moran of the University of Georgia and Dr Zepp of the US Environmental Protection Agency for helpful discussions and the use of laboratory equipment. We also thank M. Sanderson of VIMS for assistance with laboratory analyses, Dr Gardner at the University of South Carolina for use of groundwater monitoring wells at North Inlet (NSF EAR9218972), and the University of Georgia Marine Extension in Brunswick, Georgia, for assistance with field work.

LITERATURE CITED

- Aiken GR (1985) Isolation and concentration techniques for aquatic humic substances. In: Aiken GR, McKnight DM, Wershaw RL, MacCarthy P (eds) Humic substances in soil, sediment and water. Wiley, Chichester, p 363–385
- Alberts JJ, Filip Z (1994) Humic substances in rivers and estuaries of Georgia, USA. *Trends Chem Geol* 1:143–162
- Alberts JJ, Takács M (1999) Importance of humic substances for carbon and nitrogen transport into southeastern United States estuaries. *Org Geochem* 30:385–395
- Benner R, Biddanda B, Black B, McCarthy M (1997) Abundance, size distribution, and stable carbon and nitrogen isotopic compositions of marine organic matter isolated by tangential-flow ultrafiltration. *Mar Chem* 57:243–263
- Bertilsson S, Stepanauskas R, Cuadros-Hansson R, Granéli W, Wikner J, Tranvik L (1999) Photochemically induced changes in bioavailable carbon and nitrogen pools in a boreal watershed. *Aquat Microb Ecol* 19:47–56
- Bronk DA (2002) Dynamics of organic nitrogen. In: Hansell DA, Carlson CA (eds) Biogeochemistry of marine dissolved organic matter. Academic Press, San Diego, p 153–247
- Bronk DA, Lomas MW, Glibert PM, Schukert KJ, Sanderson MP (2000) Total dissolved nitrogen analysis: comparisons between the persulfate, UV and high temperature oxidation methods. *Mar Chem* 69:163–178
- Bushaw KL, Zepp RG, Tarr MA, Schulz-Jander D, Bourbonniere RA, Hodson RE, Miller WL, Bronk DA, Moran MA (1996) Photochemical release of biologically available nitrogen from aquatic dissolved organic matter. *Nature* 381:404–407
- Bushaw-Newton KL, Moran MA (1999) Photochemical formation of biologically available nitrogen from dissolved humic substances in coastal marine systems. *Aquat Microb Ecol* 18:285–292
- Clarke JS, Hacke CM, Peck MF (1990) Geology and groundwater resources of the coastal area of Georgia. *Georgia Geol Surv Bull* 113
- Cohen SG, Ojanpera SJ (1975) Photooxidation of methionine and related compounds. *J Am Chem Soc* 97:5633–5634
- Gao HZ, Zepp RG (1998) Factors influencing photoreactions of dissolved organic matter in a coastal river of the southeastern United States. *Environ Sci Technol* 32:2940–2946
- Gardner WS, Cavaletto JF, Bootsma HA, Lavrentyev PJ, Troncone F (1998) Nitrogen cycling rates and light effects in tropical Lake Maracaibo, Venezuela. *Limnol Oceanogr* 43:1814–1825
- Garside C (1982) A chemi-luminescent technique for the determination of nanomolar concentrations of nitrate and nitrite in sea-water. *Mar Chem* 11:159–167
- Grasshoff K (1983) Determination of nitrate. In: Grasshoff K, Ehrhardt M, Kremling K (eds) *Methods of seawater analysis*. Verlag Chemie, Weinheim, p 143–150
- Jørgensen NOG, Tranvik L, Edling H, Granéli W, Lindell M (1998) Effects of sunlight on occurrence and bacterial turnover of specific carbon and nitrogen compounds in lake water. *FEMS Microbiol Ecol* 25:217–227
- Kieber RJ, Hydro LH, Seaton PJ (1997) Photooxidation of triglycerides and fatty acids in seawater: implication toward the formation of marine humic substances. *Limnol Oceanogr* 42:1454–1462
- Kieber RJ, Li A, Seaton PJ (1999) Production of nitrite from the photodegradation of dissolved organic matter in natural waters. *Environ Sci Technol* 33:993–998
- Koroleff F (1983) Determination of ammonia. In: Grasshoff K, Ehrhardt M, Kremling K (eds) *Methods of seawater analysis*. Verlag Chemie, Weinheim, p 150–157
- Liebezeit G, Behrends B (1999) Determination of amino acids and carbohydrates. In: Grasshoff K, Kremling K, Ehrhardt M (eds) *Methods of seawater analysis*. Wiley-VCH, Weinheim, p 541–544
- Miller WL, Zepp RG (1995) Photochemical production of dissolved inorganic carbon from terrestrial organic-matter: significance to the oceanic organic-carbon cycle. *Geophys Res Lett* 22:417–420
- Moran MA, Zepp RG (1997) Role of photoreactions in the formation of biologically labile compounds from dissolved organic matter. *Limnol Oceanogr* 42:1307–1316
- Moran MA, Sheldon WM, Sheldon JE (1999) Biodegradation of riverine dissolved organic carbon in five estuaries of the southeastern United States. *Estuaries* 22:55–64
- Parsons TR, Maita Y, Lalli CM (1984) *A manual of chemical and biological methods for seawater analysis*. Pergamon Press, New York
- Seitzinger SP, Sanders RW (1997) Contribution of dissolved organic nitrogen from rivers to estuarine eutrophication. *Mar Ecol Prog Ser* 159:1–12
- Stumm W, Morgan JJ (1996) *Aquatic chemistry: chemical equilibria and rates in natural waters*. John Wiley & Sons Inc, New York
- Tarr MA, Wang W, Bianchi TS, Engelhaupt E (2001) Mechanisms of ammonia and amino acid photoproduction from aquatic humic and colloidal matter. *Water Res* 15:3688–3696
- Thorn KA, Mikita MA (1992) Ammonia fixation by humic substances: a N-15 and C-13 NMR-study. *Sci Total Environ* 113:67–87
- Thurman EM (1985) *Organic geochemistry of natural waters*. John Wiley and Sons, New York
- Valiela I, Costa JE (1988) Eutrophication of Buttermilk Bay, a Cape-Cod coastal embayment: concentrations of nutrients and watershed nutrient budgets. *Environ Manage* 12:539–553
- Wang WW, Tarr MA, Bianchi TS, Engelhaupt E (2000) Ammonium photoproduction from aquatic humic and colloidal matter. *Aquat Geochem* 6:275–292
- Wiegner TN, Seitzinger SP (2001) Photochemical and microbial degradation of external dissolved organic matter inputs to rivers. *Aquat Microb Ecol* 24:27–40
- Zafriou OC, True MB (1979) Nitrate photolysis in seawater by sunlight. *Mar Chem* 8:33–42
- Zepp RG (1982) Experimental approaches to environmental chemistry. In: Hutzinger O (ed) *The handbook of environmental chemistry*, Vol 2B. Springer-Verlag, New York, p 19–41