

9-11-2019

Bacterial and Archaeal Specific-Predation in the North Atlantic Basin

LM Seyler

S Tuorto

LR McGuinness

DL Gong

Virginia Institute of Marine Science

LJ Kerkof

Follow this and additional works at: <https://scholarworks.wm.edu/vimsarticles>



Part of the [Marine Biology Commons](#)

Recommended Citation

Seyler, LM; Tuorto, S; McGuinness, LR; Gong, DL; and Kerkof, LJ, "Bacterial and Archaeal Specific-Predation in the North Atlantic Basin" (2019). *VIMS Articles*. 1735.

<https://scholarworks.wm.edu/vimsarticles/1735>

This Article is brought to you for free and open access by the Virginia Institute of Marine Science at W&M ScholarWorks. It has been accepted for inclusion in VIMS Articles by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



Bacterial and Archaeal Specific-Predation in the North Atlantic Basin

Lauren M. Seyler^{1,2*}, Steve Tuorto³, Lora R. McGuinness⁴, Donglai Gong⁵ and Lee J. Kerkhof⁴

¹ Department of Marine Chemistry & Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA, United States,

² Blue Marble Space Institute of Science, Seattle, WA, United States, ³ The Watershed Institute, Pennington, NJ,

United States, ⁴ Department of Marine and Coastal Sciences, Rutgers University, New Brunswick, NJ, United States,

⁵ Virginia Institute of Marine Science, College of William & Mary, Gloucester Point, VA, United States

OPEN ACCESS

Edited by:

Fengping Wang,
Shanghai Jiao Tong University, China

Reviewed by:

Yu Zhang,
Shanghai Jiao Tong University, China
Qiang Zheng,
Xiamen University, China

*Correspondence:

Lauren M. Seyler
lmseyler@gmail.com

Specialty section:

This article was submitted to
Marine Molecular Biology
and Ecology,
a section of the journal
Frontiers in Marine Science

Received: 02 May 2019

Accepted: 23 August 2019

Published: 11 September 2019

Citation:

Seyler LM, Tuorto S,
McGuinness LR, Gong D and
Kerkhof LJ (2019) Bacterial
and Archaeal Specific-Predation
in the North Atlantic Basin.
Front. Mar. Sci. 6:555.
doi: 10.3389/fmars.2019.00555

Stable isotope probing (SIP) was used to track prokaryotic and eukaryotic carbon uptake along a meridional transect (Long. 52°W) in the North Atlantic to assess if ¹³C-resource partitioning between bacteria and archaea and ¹³C-labeled eukaryotic predators could be detected. One-liter SIP microcosms were amended with ¹³C-acetate or ¹³C-urea and incubated for 48 h. Our data indicated archaea often outcompeted bacteria for ¹³C-urea while both archaea and bacteria could incorporate ¹³C-acetate. This ¹³C label could also be tracked into eukaryotic microbes. The largest number of ¹³C-labeled eukaryotic OTUs, and the greatest percentage of eukaryotic ¹³C signal, were observed in conjunction with both archaeal and bacterial ¹³C incorporation, suggesting that most eukaryotic predators do not distinguish between archaeal and bacterial prey. However, other ¹³C-eukaryotic OTUs were exclusively associated with either ¹³C-archaeal or ¹³C-bacterial OTUs. These archaeal-specific and bacterial-specific ¹³C-eukaryotic OTUs were related to known bacterivorous predators including *Ancyromonas*, *Amastigomonas*, *Cafeteria*, and *Caecitellus*. Our SIP findings suggest both resource partitioning between bacteria and TACK (Thaumarchaeota, Aigarchaeota, Crenarchaeota, and Korarchaeota) archaea and selective predation by eukaryotic predators. Determining the equalizing mechanisms for co-existence in the marine environment can help map predator/prey interactions to better estimate carbon flow in the deep ocean.

Keywords: archaea, bacteria, predation, competition, stable isotope probing, deep ocean

INTRODUCTION

G. E. Hutchinson confronted a major contradiction between ecological theory and the natural world in “The Paradox of the Plankton” (1961). According to the competitive exclusion principle, two species competing for the same resources cannot stably coexist. However, hundreds/thousands of phytoplankton species can be found within small volumes of water in lakes, despite limited room for niche specialization and intense competition for the same resources (Scheffer and Carpenter, 2003). Hutchinson surmised that because environmental conditions can change very rapidly in aquatic systems, no one species has the advantage long enough to exclude others (Hutchinson, 1961;

Richardson et al., 1970). Additionally, stabilizing forces may also exist that reduce intra- and inter-specific competition (Chesson and Huntley, 1997; Chesson, 2000). Multiple mechanisms have been proposed to account for this co-occurrence of highly similar species, including temporal separation of activity, resource partitioning, immigration/emigration, selective predation, and dormancy.

These ecological theories of coexistence can also be tested in deep ocean environments containing both archaea and bacteria. For many years archaea were believed to be solely extremophiles. Specifically, these microorganisms were thought to grow under conditions of very high temperature, high salinity, or extreme anaerobiosis where CO₂ is used as a terminal electron acceptor, yielding methane. Because archaea occurred in habitats that would not support the growth of eukaryotes or prokaryotes, no apparent conflict was perceived between these various life forms. However, the concept of archaea being obligate extremophiles changed when these microbes were found to inhabit coastal oceans (DeLong, 1992; Fuhrman et al., 1992). Since then, mesothermal archaeal diversity has been revealed in open ocean areas around the globe (Massana et al., 2000) and in deep-sea sediments (Vetriani et al., 1999; Teske et al., 2002; Sørensen and Teske, 2006; Biddle et al., 2008). Archaea have also been found in waters between 4 and 8°C (Massana et al., 1997), polar seas (Murray et al., 1998), soils (Birtrim et al., 1997), caves (Gonzalez et al., 2006), salt marshes (Nelson et al., 2009), and in estuarine settings (Abreu et al., 2001). The discovery of marine mesophilic archaea represents an entire clade of organisms that, up until 30 years ago, had gone completely unnoticed because of their resistance to laboratory culturing, yet are apparently ubiquitous and frequently abundant in the ocean (DeLong et al., 1994; Stein and Simon, 1996; Karner et al., 2001).


Exactly what strategies marine archaea are using to remain relatively abundant and ubiquitous remains largely unknown. Surface ocean prokaryotic communities tend to be dominated by bacteria and a small percentage of euryarchaea, but below the euphotic zone and in sediments, thaumarchaea and bathyarchaea of the TACK superphylum (Guy and Ettema, 2011) equal or outnumber bacteria in abundance (Karner et al., 2001; Herndl et al., 2005; Amano-Santo et al., 2013). Both thaumarchaea (Ouverney and Fuhrman, 2000; Seyler et al., 2014, 2018) and bathyarchaea (Seyler et al., 2014; Lazar et al., 2016) appear to be capable of organic carbon uptake, competing with bacteria for the same organic substrates. Metagenomic evidence suggests that thaumarchaea may also be capable of both autotrophy and mixotrophy (Hallam et al., 2006), and radiocarbon data has suggested that autotrophy is the dominant lifestyle of thaumarchaea below the photic zone (Ingalls et al., 2006). Thaumarchaea, however, are most frequently studied with respect to their role in ammonia oxidation in the ocean. In 2004, a unique ammonia monooxygenase (*amoA*) gene was discovered on an archaeal-associated genomic scaffold (Venter et al., 2004); a year later *amoA* genes were found in thaumarchaea (Treusch et al., 2005) and examples of archaeal *amoA* genes were observed in a variety of other marine environments (Francis et al., 2005). Currently, few thaumarchaea have been cultured, and all are ammonia oxidizers (Könneke et al., 2005; Park et al., 2012;

Qin et al., 2014). However, a nitrifying lifestyle for archaea would also directly compete with nitrifying bacteria in the marine environment. It has been suggested that ammonia-oxidizing archaea are of particular importance in low-oxygen environments such as sediments and oxygen minimum zones (Francis et al., 2005; Mosier and Francis, 2008; Moin et al., 2009; Park et al., 2010). These nitrifying archaea may also be well-adapted to oligotrophic environments where the amount of available nitrogen is comparatively low (Martens-Habbena et al., 2009) which could be considered as a stabilizing force regarding preference of low rather than high ammonium concentrations.

All this conflicting potential metabolic data indicates that if the TACK-archaeota and bacteria are living in the same environment, using similar carbon and energy sources, there should be competitive exclusion and one group would eventually out-compete the other. Yet, we find both living together in the same oceanic samples. In this report, resource partitioning and selective predation were investigated as two possible mechanisms that may allow archaea and bacteria to stably coexist in the deep ocean. We chose to utilize DNA stable-isotope probing techniques (SIP; Radajewski et al., 1999) as a means to create different pools of ¹³C-microbial prey items in marine samples (by labeling either bacteria or archaea with different ¹³C-carbon sources). Prior research had demonstrated that archaea could successfully outcompete bacteria for ¹³C-urea in a coastal salt marsh at low concentrations (<30 μM) during short term incubations (5 days) (Seyler et al., 2014). However, it is unclear if deep ocean archaea would also outcompete bacteria under comparable conditions to our salt marsh study. We initially set out to determine if similar resource partitioning results could also be observed in oceanic water column samples. If resource partitioning could be demonstrated, we could then test a second mechanism of co-existence (selective predation) by determining if eukaryotic predators consumed ¹³C-archaeal or ¹³C-bacterial prey in a specific fashion. Our findings suggested there was specific uptake of ¹³C-urea by TACK archaea in many of our SIP experiments. Furthermore, particular ¹³C-eukaryotes were associated exclusively with ¹³C-archaea (archaeovores), with ¹³C-bacteria (bactivores) or detected in samples without any archaea or bacterial ¹³C uptake (osmotrophs). This information on ¹³C-resource partitioning and specific predation by eukaryotes provides insight into the competitive relationships between archaea and bacteria. Finally, this SIP approach also demonstrates how the deep sea can be a model system for studying competition between other microbial groups in natural settings to further our understanding of microbial ecology.

MATERIALS AND METHODS

In April 2012, water samples were collected onboard the R/V Atlantis as part of the US CLIVAR/CO₂ Repeat Hydrography program, along World Ocean Circulation Experiment (WOCE) line A20. This was a cruise of opportunity, done in conjunction with a larger effort to characterize water chemistry and marine microbial productivity. Therefore, the volumes of water available for sampling were often limited (generally <5 L at each



station/depth per cast). As such, a total of 36 SIP microcosms were established at three stations (**Figure 1A**) from three water column zones: the mixed layer (euphotic zone, 65–115 m), the local oxygen minimum (mesopelagic zone, 365–835 m), and the bathypelagic zone (2335–2585 m; **Figure 1B**), as part of a larger effort to determine autotrophy, heterotrophy and mixotrophy in N. Atlantic waters (Seyler et al., 2018). Duplicate 1-L samples of water from each depth were amended with ^{13}C -acetate/ ^{12}C -urea or ^{13}C -urea/ ^{12}C -acetate at equimolar concentrations (20 μM), to stimulate the same populations of microbes in all microcosms from the same station/depth and maximize the number of ^{13}C -carbon sources that could be tested. Microcosms were incubated in covered containers on deck for 48 h, with a constant inflow of surface seawater to maintain stable temperature ($\sim 22^\circ\text{C}$). Early in the cruise, duplicate ^{12}C -acetate and ^{13}C -acetate microcosms were established and incubated for 12–72 h to verify the minimum time necessary to obtain a positive ^{13}C -signal (Seyler et al., 2018). Negative SIP controls using ^{12}C sodium acetate were incubated for 72 h. Biomass was collected on a 0.2- μm filter using vacuum filtration.

DNA was purified using phenol-chloroform methods and $^{12}\text{C}/^{13}\text{C}$ -DNA was separated by isopycnic cesium chloride gradient ultracentrifugation at $200,000 \times g$ for 36 h, using ^{13}C -labeled *H. salinarum* or *E. coli* DNA as a carrier (Gallagher et al., 2005). Carrier DNA produces a visible ^{13}C -lower band in the gradient, removing the need to fractionate the gradient and increasing sensitivity (Gallagher et al., 2010; Kerkhof et al., 2011; Seyler et al., 2014, 2018; Tuorto et al., 2014). Fluorescently-labeled (using ethidium bromide) ^{12}C -(upper) and ^{13}C -(lower) bands were collected by pipette. Upper and lower bands (1 μl each) were amplified

using nested PCR with 5'-fluorescently-labeled, 16S rRNA archaea-specific 21F (5'-TCCGGTTGATCCYGCCGG)/958R (5'-YCCGCGTTGAMTCCAATT) primers using the following protocol: 94°C for 5 min, 23 cycles of (94°C for 1 min, 55°C for 2 min, 72°C for 3 min). A 1 μl aliquot was then taken from this reaction and amplified using thaum/cren/bathyarchaeota-specific 7F (5'-TTCCGGTTGATCCYGCCGGACC)/518R (5'-GCTGGTWTACC GCGCGGCTGA) primers using the following protocol: 94°C for 5 min, 30 cycles of (94°C for 1 min, 73°C for 2 min, 72°C for 3 min), 72°C for 10 min (Perevalova et al., 2003; Ionescu et al., 2008; Seyler et al., 2018). Upper and lower bands were also amplified using bacteria-specific 27F (5'-AGAGTTTGATCMTGGCTCAG)/1100R (5'-GGGTTGCGCTCGTTG) primers (Ludwig, 2007) in a 2-step process: step 1 at 94°C for 1 min 30 s, 20 cycles of (94°C for 30 s, 57°C for 30 s, 72°C for 1 min 10 s); 1 μl aliquot amplified in step 2 at 94°C for 1 min 30 s, 25 cycles of (94°C for 30 s, 57°C for 30 s, 72°C for 1 min 10 s), 72°C for 10 min. Finally, upper and lower bands were amplified for eukaryotes using 18S rRNA universal eukaryote 328F (5'-ACCTGGTTGATCCTGCCAG) (Medlin et al., 1988)/516R (5'-ACCAGACTTGCCCTCC) (Amann et al., 1990) primers in a 2-step PCR: 94°C for 1 min 30 s, 15 cycles of [94°C for 45 s, 56°C for 45 s, 72°C for 2 min], followed by 94°C for 1 min 30 s, 30 cycles of [94°C for 45 s, 56°C for 45 s, 72°C for 2 min], 72°C for 10 min. Amplicons were digested with *MnII* (prokaryotic amplicons) or *HaeIII* (eukaryotic amplicons) in 20 μl volumes for 6 h at 37°C , then precipitated using sodium acetate and 95% ethanol (McGuinness et al., 2006). Precipitated DNA was dried and re-suspended in 19.7 μl de-ionized formamide with 0.3 μl ROX 500 size standard (Applied Biosystems). TRFLP fingerprinting

(Avaniss-Aghajani et al., 1994) was carried out on an ABI 310 genetic analyzer (Applied Biosystems, Foster City, CA, United States) using GENESCAN software. Peak detection was set at 25 arbitrary fluorescent units. Each resulting peak was considered an operational taxonomic unit (OTU).

An 18S amplicon clonal library was constructed using the Topo TA cloning kit according to the manufacturer's instructions (Invitrogen, Carlsbad, CA, United States). 192 recombinant clones from the ^{13}C -labeled bands were screened in a multiplex format (Babcock et al., 2007), to determine portions of the 18S rRNA genes which corresponded to the TRF of the ^{13}C -eukaryotic profile. Thirteen 18S rRNA genes that matched TRFs of interest were sequenced via Sanger methods using M13 primers (GENEWIZ, Inc., NJ, United States), producing 9 unique sequences (<99% similarity to each other) between 520 and 603 bp. These sequences were compared to the Silva-ARB Eukaryotic database by discontinuous MEGABLAST. Best hits were used to construct a phylogenetic tree using 460 unambiguously aligned base pairs and maximum likelihood methods (Guindon et al., 2010) in Geneious 11.1.4. Clone sequences were submitted to GenBank (accession numbers KT749976–KT749984).

RESULTS

A minimum incubation time of 48 h was required to unambiguously detect ^{13}C -archaeal DNA synthesis in the ^{13}C -carrier band (Seyler et al., 2018). In 7 microcosms (6 epipelagic, 1 bathypelagic), no signal could be detected in the ^{13}C -band (no PCR amplicon could be detected by electrophoresis gel, and the resulting digests produced no TRFLP peaks). In 7 microcosms from mesopelagic or bathypelagic samples, signal from ^{13}C -urea was only detected in archaea. In one bathypelagic microcosm (Station 63), signal from ^{13}C -urea was only detected in bacteria. Three mesopelagic microcosms had both archaea and bacteria ^{13}C signals resulting from ^{13}C -urea amendments. In all twelve ^{13}C -acetate mesopelagic and bathypelagic microcosms, both archaea and bacteria incorporated ^{13}C , except bathypelagic microcosms at Station 63, where only bacterial uptake of ^{13}C -acetate was detected (Figure 2). No ^{13}C uptake was detected in ^{12}C controls.

TRFLP analysis detected 37 eukaryotic OTUs, grouped based on whether the ^{13}C signal was also detected in the ^{13}C -archaea and ^{13}C -bacteria, in the ^{13}C -archaea alone, in the ^{13}C -bacteria alone, or without a ^{13}C signal from either prokaryotic group (Figure 3). Seventeen eukaryotic OTUs were detected in SIP microcosms with both bacterial and archaeal ^{13}C uptake. Ten eukaryotic OTUs were associated exclusively with ^{13}C -archaea (archaevores), and 7 eukaryotic OTUs were only observed with ^{13}C -bacteria (bacterivores). Three eukaryotic OTUs were detected in samples without any archaea or bacterial ^{13}C uptake (osmotrophs). The total number of observed archaeal-specific predator OTUs was higher than that of bacterial-specific predators, however, if the total adjusted TRFLP peak area was considered a proxy for relative abundance, the non-specific predators tended to constitute a

	Station 33		Station 48		Station 63	
	1	2	1	2	1	2
^{13}C -Urea						
Mesopelagic	A B	A —	A B	A B	A —	A —
Bathypelagic	A —	—	A —	A —	— B	A —

	Station 33		Station 48		Station 63	
	1	2	1	2	1	2
^{13}C -Acetate						
Mesopelagic	A B	A B	A B	A B	A B	A B
Bathypelagic	A B	A B	A B	A B	— B	— B


FIGURE 2 | Patterns of ^{13}C -incorporation by archaea (A) and bacteria (B) based on ^{13}C -urea or ^{13}C -acetate additions in meso-bathypelagic samples. No incorporation of ^{13}C was detected in any microcosms established with water from the epipelagic zone.

greater percentage of the overall ^{13}C -eukaryotic community profile (Figure 3).

Nine rRNA genes with ^{13}C -eukaryotic TRF peaks were identified in the clonal library. Of these, three were found exclusively in microcosms where bacteria incorporated ^{13}C signal (TRFs 182, 201, 269), three were found only in microcosms where archaea incorporated ^{13}C signal (TRFs 329, 333, 339), two were found in both (TRFs 273, 305), and one was found in the microcosm where neither bacteria nor archaea incorporated ^{13}C signal (TRF 293). An 18S phylogenetic tree (Figure 4) indicated bacterial uptake-associated eukaryotes were associated with relatives of the parasitic dinoflagellate *Syndiniales* group, a flagellate (*Ancyromonas* sp.), and an apuzoan (*Amastigomonas* sp.). The archaea uptake-associated eukaryotes were associated with radiolarians or a dinoflagellate group (*Heterocapsa* sp.). Non-specific prokaryotic predators were related to a bicosoecida group (*Cafeteria* sp.) and a marine fungus (*Paraphysoderma sedebokerense*). The 18S rRNA gene from the eukaryote believed to be osmotrophic aligned to another flagellate group (*Caecitellus paraparvulus*).

DISCUSSION


Interactions between prokaryotes, protist predators, and dissolved organic carbon pools in the world's oceans may have far-reaching effects on global biogeochemical cycles (Kujawinski, 2011). Deep ocean waters represent the largest pool of dissolved organic carbon (Hansel and Carlson, 1998), the respiration of which accounts for up to one third of oceanic biological CO_2 production (Del Giorgio and Duarte, 2002; Aristegui et al., 2005). The factors which control microbial community dynamics, carbon respiration, and mortality in the deep ocean are therefore of great importance to the global carbon cycle. Archaea, particularly thaumarchaea belonging to the TACK superphylum, have been found to dominate prokaryotic communities in the deep ocean (Herndl et al., 2005; Varela et al., 2008). Yet, archaeal



utilization of dissolved and particulate organic matter in the deep sea remains poorly constrained, and even less is known about the relationship of predator grazing to archaeal function in the deep sea.

“Bottom-up” controls such as the availability of nutrients (Wilkins et al., 2012) and the character of dissolved organic

matter (Smriga et al., 2016) may have significant effects on the community structure of prokaryotes. Past studies demonstrated that archaea take up certain carbon sources at an equal or higher rate than bacteria in deep ocean waters (e.g., Ouverney and Fuhrman, 2000; Teira et al., 2006; Kirchman et al., 2007). Thaumarchaea are known to incorporate urea as a carbon source



(Seyler et al., 2014, 2018) and they may use it as an alternative nitrogen source to fuel nitrification (Alonso-Sáez et al., 2012). In many of our mesopelagic and bathypelagic microcosms, ^{13}C -urea uptake was detected in archaea but not bacteria, suggesting that archaea can outcompete bacteria for urea at those particular samples, at the concentration tested here. Competition for substrates in oligotrophic environments where nutrients and energy are limiting seems to favor archaea over bacteria (Jardillier et al., 2005; Vuillemin et al., 2019). Additionally, PCR amplification of key genes in the Thaumarchaeota suggest that ecological functioning of this group may shift with depth (Hu et al., 2011). One possibility of our observed resource partitioning for ^{13}C -urea by archaea may be our SIP incubation conditions. That is, by conducting the SIP incubations at sea surface temperature, rather than at *in situ* mesopelagic and bathypelagic temperatures, we may have stimulated or accelerated activity in some clades. However, an increase in temperature cannot bestow metabolic capability where none had previously existed; we therefore feel confident in stating that archaea are capable of outcompeting bacteria for urea under the conditions provided in our microcosms [For further discussion, see Seyler et al. (2018)].

In addition to resource partitioning, our SIP experiment indicated that selective predation may partly account for archaeal/bacterial co-existence. The main causes of mortality in marine microorganisms are viruses and grazing by phagotrophic protists (Jardillier et al., 2005; Arístegui et al., 2009). Phagotrophic

grazing- “top-down” population control- can structure the microbial community, activity, and physiology of prokaryotic prey (Šimek et al., 1997; Lin et al., 2007; Saleem et al., 2012), including shifts in community assemblages as a result of the prey preferences of protists (Jardillier et al., 2005; Jezbera et al., 2006). Protistan species can act as bacterivore, herbivore, carnivore of other protozoa and/or metazoans (e.g., nauplii, copepodite, and metazoan larvae), and can even consume dissolved organic matter through saprotrophy or osmotrophy (Swanberg and Caron, 1991; Caron et al., 2009; Worden et al., 2015). Specific predation of archaea has been addressed in very few studies, e.g., at marine oxic-anoxic interfaces (Anderson et al., 2012) and in freshwater (Ballen-Segura et al., 2017). Here, we demonstrate both archaeal and bacterial-specific predation by a subset of eukaryotic predators in the mesopelagic and bathypelagic water column. However, our data suggest the archaea-specific predators represent a small percentage of the overall the eukaryotic predatory community compared to non-specific microbial predators.

This study demonstrates that certain ^{13}C -eukaryotic OTUs are associated with either ^{13}C -archaeal or ^{13}C -bacterial signals. This transfer of ^{13}C -carbon from prokaryote to eukaryote could result from predation or by exudation/episymbiosis. Often ciliates or flagellates have surface associations with bacteria for detoxification or defense (Petroni et al., 2000; Edgcomb et al., 2011). Although we did not test for the presence of prokaryotes in food vacuoles or on the external surfaces of deep-sea eukaryotes

by microscopy, an assessment of whether predation or symbiosis is the predominant mechanism for carbon transfer can be made using the SIP results directly. To a first order, if predation is the dominant mechanism, we can presume that in many microcosms, the ^{13}C -carbon will pass directly from substrate to prokaryote to eukaryote predator. Therefore, a much higher ^{13}C -eukaryotic signal compared to a ^{13}C -prokaryotic signal will be observed in the PCR product from the ^{13}C -labeled DNA. On the other hand, if growth of the eukaryote is predicated on growth of an epibiont, there should always be a comparable PCR signal from both prokaryote and eukaryote within the ^{13}C -band. All of our SIP microcosms had significantly higher eukaryotic signal than prokaryotic signal in the ^{13}C -band (**Supplementary Figure S1**), suggesting a tight coupling between these groups that is likely a predation signal.

If differential predation influences the coexistence of archaea and bacteria, the mechanisms used by predators to discern between the archaea and bacteria are largely unexplored. It is known that prey item size, surface properties, concentration, and particle attachment can be major controlling elements selecting for prey items by eukaryotic grazers (Lampert, 1987; Gonzalez et al., 1990, 1993; Jürgens and Matz, 2002; Thurman et al., 2010; Tuorto and Taghon, 2014). Recent studies suggest that marine archaea are significantly smaller than bacteria on average (Stepanauskas et al., 2017; Santoro et al., 2018). Aggregation on detrital particles, as observed in euryarchaea (Orsi et al., 2015), may provide refuge from predation (Langenheder and Jürgens, 2001; Šimek et al., 2001), as well as a host of biochemical interactions, exopolymers, and toxicity (Jürgens and Matz, 2002). Further studies will be necessary to explore the mechanisms of selective predation in deep ocean settings.

Predation by protists is also a key mechanism for the transfer of dissolved organic carbon to higher trophic levels in the microbial loop (Azam et al., 1983; Jürgens and Massana, 2008; Worden et al., 2015). The structure of the marine microbial food web and efficiency of carbon transfer are major determinants in the rate of oceanic carbon sequestration and the type/amount of higher consumers in the ecosystem (Limardo and Worden, 2015). Although prokaryotes mobilize dissolved carbon into biomass, it is the heterotrophic strategies of small eukaryotic consumers that determine whether the net flow of organic matter will ultimately end up as a trophic “link” (i.e., trophic transfer to top predators and export) versus a trophic “sink” (i.e., be remineralized as dissolved organic matter or CO_2). The archaeal contribution to the microbial loop is still poorly understood and, consequently, often easy to ignore. SIP studies such as described here can map trophic interactions between microbial predators and prey to predict the fate of organic carbon in marine ecosystems.

REFERENCES

Abreu, C., Jurgens, G., De Marco, P., Saana, A., and Bordalo, A. A. (2001). Crenarchaeota and euryarchaeota in temperate estuarine sediments. *J. Appl. Microbio.* 90, 713–718. doi: 10.1046/j.1365-2672.2001.01297.x

DATA AVAILABILITY

The datasets generated for this study can be found in GenBank, KT749976–KT749984.

AUTHOR CONTRIBUTIONS

LS and LK wrote the original manuscript. LS, LM, and LK designed the SIP experiments. LS set up microcosms and performed all the laboratory work. ST and LM performed the data analysis and interpretation of data. DG was co-chief scientist on R/V Atlantis during CLIVAR leg A20 and supervised all onboard research. LK conceived the original idea, oversaw all laboratory work, and provided data analysis.

FUNDING

This research was made possible through the support of the U.S Global Ocean Carbon and Repeat Hydrography Program and NSF Ocean Technology and Interdisciplinary Program grant (#1131022) to LK.

ACKNOWLEDGMENTS

The authors wish to thank Dr. Michael McCartney for allowing LS to participate as a graduate student researcher onboard the R/V Atlantis during CLIVAR leg A20.

SUPPLEMENTARY MATERIAL

The Supplementary Material for this article can be found online at: <https://www.frontiersin.org/articles/10.3389/fmars.2019.00555/full#supplementary-material>

FIGURE S1 | Top half of gel showing amplification of 18S rRNA genes from either the ^{12}C -band or the ^{13}C -band within the SIP microcosms indicated. Lanes are (A) empty; (B) lambda HinDIII ladder; (C) negative PCR control; (D) ^{13}C -*Halobacterium* carrier; (E) station 63–2585 m, ^{13}C -acetate amended, ^{12}C -band; (F) station 63–2585 m, ^{13}C -acetate amended, ^{13}C -band; (G) station 48–2335 m, ^{13}C -acetate amended, ^{12}C -band; (H) station 48–2335 m, ^{13}C -acetate amended, ^{13}C -band; (I) station 48–2335 m, ^{13}C acetate-amended, ^{13}C -band (duplicate Niskin cast); (J) empty; (K) positive control; (L) empty. Bottom half gel of indicates amplification of 16S rRNA genes from the identical samples as the 18S rRNA amplification above with the exception that the ^{12}C -bands (Lanes E, G) were diluted 1:10 prior to amplification of the 16S rRNA genes. All ^{13}C -bands were all amplified for 16S rRNA genes without dilution as above. The concentration of PCR product from each amplification is indicated above the band.

Alonso-Sáez, L., Waller, A. S., Mende, D. R., Bakker, K., Farnelid, H., Yager, P. L., et al. (2012). Role for urea in nitrification by polar marine Archaea. *PNAS* 109, 17989–17994. doi: 10.1073/pnas.1201914109

Amann, R. I., Binder, B. J., Olson, R. J., Chisholm, S. W., Devereux, R., and Stahl, D. A. (1990). Combination of 16S rRNA-targeted oligonucleotide probes

- with flow cytometry for analyzing mixed microbial populations. *Appl. Environ. Microbiol.* 56, 1919–1925.
- Amano-Santo, C., Akiyama, S., Uchida, M., Shimada, K., and Utsumi, M. (2013). Archaeal distribution and abundance in water masses of the Arctic Ocean, Pacific sector. *Aquat. Microb. Ecol.* 69, 101–112. doi: 10.3354/ame01624
- Anderson, R., Winter, C., and Jürgens, K. (2012). Protist grazing and viral lysis as prokaryotic mortality factors at Baltic Sea oxic-anoxic interfaces. *MEPS* 467, 1–14. doi: 10.3354/meps10001
- Aristegui, J., Duarte, C. M., Gasol, J. M., and Alonso-Sáez, L. (2005). Active mesopelagic prokaryotes support high respiration in the subtropical northeast Atlantic Ocean. *Geophys. Res. Lett.* 32, 1–4.
- Aristegui, J., Gasol, J. M., Duarte, C. M., and Herndl, G. J. (2009). Microbial oceanography of the dark ocean's pelagic realm. *Limnol. Oceanogr.* 54, 1501–1529. doi: 10.4319/lo.2009.54.5.1501
- Avaniss-Aghajani, E., Jones, K., Chapman, D., and Brunk, C. (1994). A molecular technique for identification of bacteria using small subunit ribosomal RNA sequences. *Biotechniques* 17, 144–146.
- Azam, F., Fenichel, T., Field, J., Gray, J., Meyer-Reil, L. A., and Thingstad, F. (1983). The ecological role of water-column microbes in the sea. *Mar. Ecol. Prog. Ser.* 10, 257–263. doi: 10.3354/meps010257
- Babcock, D. A., Wawrik, B., Paul, J. H., McGuinness, L., and Kerkhof, L. J. (2007). Rapid screening of a large insert BAC library for specific 16S rRNA genes using TRFLP. *J. Microbiol. Methods* 71, 156–161. doi: 10.1016/j.mimet.2007.07.015
- Ballen-Segura, M., Felip, M., and Catalan, J. (2017). Some mixotrophic flagellate species selectively graze on archaea. *Appl. Environ. Microbiol.* 83:e2317-16. doi: 10.1128/AEM.02317-16
- Biddle, J. F., Fitz-Gibbon, S., Schuster, S. C., Brenchley, J. E., and House, C. H. (2008). Metagenomic signatures of the Peru margin seafloor biosphere show a genetically distinct environment. *PNAS* 105, 10583–10588. doi: 10.1073/pnas.0709942105
- Birtrim, S. B., Donohue, T. J., Handelsman, J., Roberts, G. P., and Goodman, R. M. (1997). Molecular phylogeny of Archaea from soil. *PNAS* 94, 277–282. doi: 10.1073/pnas.94.1.277
- Caron, D. A., Worden, A. Z., Countway, P. D., Demir, E., and Heidelberg, K. B. (2009). Protists are microbes too: a perspective. *ISME J.* 3, 4–12. doi: 10.1038/ismej.2008.101
- Chesson, P. (2000). Mechanisms of maintenance of species diversity. *Annu. Rev. Ecol. Syst.* 31, 343–366. doi: 10.1146/annurev.ecolsys.31.1.343
- Chesson, P., and Huntley, N. (1997). The roles of harsh conditions in the dynamics of ecological communities. *Am. Nat.* 150, 519–553. doi: 10.1086/286080
- Del Giorgio, P. A., and Duarte, C. M. (2002). Respiration in the open ocean. *Nature* 420, 379–384. doi: 10.1038/nature01165
- DeLong, E. F. (1992). Archaea in coastal marine environments. *PNAS* 89, 5685–5689. doi: 10.1073/pnas.89.12.5685
- DeLong, E. F., Wu, K. Y., Prézélin, B. B., and Jovine, R. V. M. (1994). High abundance of Archaea in Antarctic marine picoplankton. *Nature* 37, 695–697. doi: 10.1038/371695a0
- Edgcomb, V. P., Breglia, S. A., Yubuki, N., Beaudoin, D., Patterson, D. J., Leander, B. S., et al. (2011). Identity of epibiotic bacteria on symbiont euglenozoans in O₂-depleted marine sediments: evidence for symbiont and host co-evolution. *ISME J.* 5, 231–243. doi: 10.1038/ismej.2010.121
- Francis, C. A., Roberts, K. J., Beman, J. M., Santoro, A. E., and Oakley, B. B. (2005). Ubiquity and diversity of ammonia-oxidizing archaea in water columns and sediments of the ocean. *PNAS* 102, 14683–14688. doi: 10.1073/pnas.0506625102
- Fuhrman, J. A., McCallum, K., and Davis, A. A. (1992). Novel major archaeobacterial group from marine plankton. *Nature* 356, 148–149. doi: 10.1038/356148a0
- Gallagher, E., McGuinness, L., Phelps, C., Young, L. Y., and Kerkhof, L. J. (2005). ¹³C-carrier DNA shortens the incubation time needed to detect benzoate-utilizing denitrifying bacteria by stable-isotope probing. *Appl. Environ. Microbiol.* 71, 5192–5196. doi: 10.1128/aem.71.9.5192-5196.2005
- Gallagher, E. M., Young, L. Y., McGuinness, L. M., and Kerkhof, L. J. (2010). Detection of 2,4,6-trinitrotoluene anaerobic bacteria by ¹⁵N and ¹³C incorporation. *Appl. Environ. Microbiol.* 76, 1695–1698. doi: 10.1128/AEM.02274-09
- Gonzalez, J. M., Portilo, M. C., and Saiz-Jimenez, C. (2006). Metabolically active Crenarchaeota in Altamira cave. *Naturwissenschaften* 93, 42–45. doi: 10.1007/s00114-005-0060-3
- Gonzalez, J. M., Sherr, E. B., and Sherr, B. F. (1990). Size-selective grazing on bacteria by natural assemblages of estuarine flagellates and ciliates. *Appl. Environ. Microbiol.* 56, 583–589.
- Gonzalez, J. M., Sherr, E. B., and Sherr, B. F. (1993). Differential feeding by marine flagellates on growing versus starving, and on motile versus nonmotile, bacterial prey. *Mar. Ecol. Prog. Ser.* 102, 257–267. doi: 10.3354/meps095257
- Guindon, S., Dufayard, J. F., Lefort, V., Anisimova, M., Hordijk, W., and Gascuel, O. (2010). New algorithms, and methods to estimate maximum-likelihood phylogenies: assessing the performance of PhyML 3.0. *Syst. Biol.* 59, 307–321. doi: 10.1093/sysbio/syq010
- Guy, L., and Ettema, T. J. G. (2011). The archaeal ‘TACK’ superphylum and the origin of eukaryotes. *Trends Microbiol.* 19, 580–587. doi: 10.1016/j.tim.2011.09.002
- Hallam, S. J., Mincer, T. J., Schleper, C., Preston, C. M., Roberts, K., Richardson, P. M., et al. (2006). Pathways of carbon assimilation and ammonia oxidation suggested by environmental genomic analyses of marine Crenarchaeota. *PLoS Biol.* 4:e95. doi: 10.1371/journal.pbio.0040095
- Hansel, D. A., and Carlson, C. A. (1998). Deep-ocean gradients in the concentration of dissolved organic carbon. *Nature* 395, 263–266. doi: 10.1038/26200
- Herndl, G. J., Reinthaler, T., Teira, E., van Aken, H., Veth, C., Pernthaler, A., et al. (2005). Contribution of Archaea to total prokaryotic production in the deep Atlantic Ocean. *Appl. Environ. Microbiol.* 71, 2303–2309. doi: 10.1128/aem.71.5.2303-2309.2005
- Hu, A., Jiao, N., Zhang, R., and Yang, Z. (2011). Niche partitioning of marine group I Crenarchaeota in the euphotic and upper mesopelagic zones of the East China Sea. *Appl. Environ. Microbiol.* 77, 7469–7478. doi: 10.1128/AEM.00294-11
- Hutchinson, G. E. (1961). The paradox of the plankton. *Am. Nat.* 45, 137–145.
- Ingalls, A. E., Shah, S. R., Hansman, R. L., Aluwihare, L. L., Santos, G. M., Druffel, E. R. M., et al. (2006). Quantifying archaeal community autotrophy in the mesopelagic ocean using natural radiocarbon. *PNAS* 103, 6442–6447. doi: 10.1073/pnas.0510157103
- Ionescu, D., Penno, S., Haimovich, M., Rihtman, B., Goodwin, A., Schwartz, D., et al. (2008). Archaea in the Gulf of Aqaba. *FEMS Microbiol.* 69, 425–438. doi: 10.1111/j.1574-6941.2009.00721.x
- Jardillier, L., Bettarel, Y., Richardot, M., Bardot, C., Amblard, C., Sime-Ngando, T., et al. (2005). Effects of viruses and predators on prokaryotic community composition. *Microbiol. Ecol.* 50, 557–569. doi: 10.1007/s00248-005-5030-y
- Ježbera, J., Hornák, K., and Šimek, K. (2006). Prey selectivity of bacterivorous protists in different size fractions of reservoir water amended with nutrients. *Environ. Microbiol.* 8, 1330–1339. doi: 10.1111/j.1462-2920.2006.01026.x
- Jürgens, K., and Massana, R. (2008). “Protist grazing on marine bacterioplankton,” in *Microbial Ecology of the Oceans*, 2 Edn, ed. D. L. Kirchman, (Hoboken, NJ: Wiley), 383–441. doi: 10.1002/9780470281840.ch11
- Jürgens, K., and Matz, C. (2002). Predation as a shaping force for the phenotypic and genotypic composition of planktonic bacteria. *Antonie Leeuwenhoek* 81, 413–434.
- Karner, M. B., DeLong, E. F., and Karl, D. M. (2001). Archaeal dominance in the mesopelagic zone of the Pacific Ocean. *Nature* 409, 507–510. doi: 10.1038/35054051
- Kerkhof, L. J., Williams, K. H., Long, P. E., and McGuinness, L. R. (2011). Phase preference by active, acetate-utilizing bacteria at the rifle, CO integrated field research challenge site. *Environ. Sci. Technol.* 45, 1250–1256. doi: 10.1021/es102893r
- Kirchman, D. L., Elifantz, H., Dittel, A. I., Malmstrom, R. R., and Cottrell, M. T. (2007). Standing stocks and activity of Archaea and Bacteria in the western Arctic Ocean. *Limnol. Oceanogr.* 52, 495–507. doi: 10.4319/lo.2007.52.2.0495
- Könneke, M., Bernhard, A. E., de la Torre, J. R., Walker, C. B., Waterbury, J. B., and Stahl, D. A. (2005). Isolation of an autotrophic ammonia-oxidizing marine archaeon. *Nature* 437, 543–546. doi: 10.1038/nature03911
- Kujawinski, E. B. (2011). The impact of microbial metabolism on marine dissolved organic matter. *Annu. Rev. Mar. Sci.* 3, 567–599. doi: 10.1146/annurev-marine-120308-081003
- Lampert, W. (1987). “Predictability in lake ecosystems: the role of biotic interactions,” in *Potential and Limitations of Ecosystem Analysis (Ecological Studies 61)*, eds E. D. Schultz, and H. Zwölfer, (Berlin: Springer-Verlag), 333–346. doi: 10.1007/978-3-642-71630-0_16

- Langenheder, S., and Jürgens, K. (2001). Regulation of bacterial biomass and community structure by metazoan and protozoan predation. *Limnol. Oceanogr.* 46, 121–134. doi: 10.4319/lo.2001.46.1.0121
- Lazar, C. S., Baker, B. J., Seitz, K., Hyde, A. S., Dick, G. J., Hinrichs, K. U., et al. (2016). Genomic evidence for distinct carbon substrate preferences and ecological niches of Bathyarchaeota in estuarine sediments. *Environ. Microbiol.* 18, 1200–1211. doi: 10.1111/1462-2920.13142
- Limardo, A. J., and Worden, A. Z. (2015). Microbiology: exclusive networks in the sea. *Nature* 522, 36–37. doi: 10.1038/nature14530
- Lin, X., Scranton, M. I., Varela, R., Chistoserdov, A., and Taylor, G. T. (2007). Compositional responses of bacterial communities to redox gradients and grazing in the anoxic Cariaco Basin. *Aquat. Microb. Ecol.* 47, 57–72. doi: 10.3354/ame047057
- Ludwig, W. (2007). Nucleic acid techniques in bacterial systematics and identification. *Int. J. Food Microbiol.* 120, 225–236. doi: 10.1016/j.ijfoodmicro.2007.06.023
- Martens-Habbena, W., Berube, P. M., Urakawa, H., de la Torre, J. R., and Stahl, D. A. (2009). Ammonia oxidation kinetics determine niche separation of nitrifying Archaea and bacteria. *Nature* 461, 976–979. doi: 10.1038/nature08465
- Massana, R., DeLong, E. F., and Pedrós-Alió, C. (2000). A few cosmopolitan phylotypes dominate planktonic archaeal assemblages in widely different oceanic provinces. *Environ. Microbiol.* 66, 1777–1787. doi: 10.1128/aem.66.5.1777-1787.2000
- Massana, R., Murray, A. E., Preston, C. M., and DeLong, E. F. (1997). Vertical distribution and phylogenetic characterization of marine planktonic Archaea in the Santa Barbara channel. *Appl. Environ. Microbiol.* 63, 50–56.
- McGuinness, L. M., Salganik, M., Vega, L., Pickering, K. D., and Kerkhof, L. J. (2006). Replicability of bacterial communities in denitrifying bioreactors as measured by PCR/T-RFLP analysis. *Environ. Sci. Technol.* 40, 509–515. doi: 10.1021/es050900l
- Medlin, L., Elwood, H. J., Stickel, S., and Sogin, M. L. (1988). The characterization of enzymatically amplified eukaryotic 16S-like rRNA-coding regions. *Gene* 71, 491–499. doi: 10.1016/0378-1119(88)90066-2
- Moin, N. S., Nelson, K. A., Bush, A., and Bernhard, A. E. (2009). Distribution and diversity of archaeal and bacterial ammonia oxidizers in salt marsh sediments. *Appl. Environ. Microbiol.* 75, 7461–7468. doi: 10.1128/AEM.01001-09
- Mosier, A. C., and Francis, C. A. (2008). Relative abundance and diversity of ammonia-oxidizing archaea and bacteria in the San Francisco Bay estuary. *Environ. Microbiol.* 10, 3002–3016. doi: 10.1111/j.1462-2920.2008.01764.x
- Murray, A. E., Preston, C. M., Massana, R., Taylor, L. T., Blakis, A., Wu, K., et al. (1998). Seasonal and spatial variability of bacterial and archaeal assemblages in the coastal waters near Anvers Island, Antarctica. *Appl. Environ. Microbiol.* 64, 2585–2595.
- Nelson, K. A., Moin, N. S., and Bernhard, A. E. (2009). Archaeal diversity and the prevalence of Crenarchaeota in salt marsh sediments. *Appl. Environ. Microbiol.* 75, 4211–4215. doi: 10.1128/AEM.00201-09
- Orsi, W. D., Smith, J. M., Wilcox, H. M., Swallow, J. E., Carini, P., Worden, A. Z., et al. (2015). Ecophysiology of uncultivated marine euryarchaea is linked to particulate organic matter. *ISME J.* 9, 1747–1763. doi: 10.1038/ismej.2014.260
- Ouverney, C. C., and Fuhrman, J. A. (2000). Marine planktonic archaea take up amino acids. *Appl. Environ. Microbiol.* 66, 4829–4833. doi: 10.1128/aem.66.11.4829-4833.2000
- Park, B. J., Park, S. J., Yoon, D. N., Schouten, S., Sinninghe Damsté, J. S., and Rhee, S. K. (2010). Cultivation of autotrophic ammonia-oxidizing archaea from marine sediments in coculture with sulfur-oxidizing bacteria. *Appl. Environ. Microbiol.* 76, 7575–7587. doi: 10.1128/AEM.01478-10
- Park, S. J., Kim, J. G., Jung, M. Y., Kim, S. J., Cha, I. T., Kwon, K. K., et al. (2012). Draft genome sequence of an ammonia-oxidizing archaeon, “*Candidatus Nitrosopumilus koreensis*” AR1, from marine sediment. *J. Bacteriol.* 194, 6940–6941. doi: 10.1128/jb.01857-12
- Perevalova, A. A., Lebedinsky, A. V., Bonch-Osmolovskaya, E. A., and Chernykh, N. A. (2003). Detection of hyperthermophilic Archaea of the genus *Desulfurococcus* by hybridization with oligonucleotide probes. *Microbiology* 72, 340–346.
- Petroni, G., Spring, S., Schleifer, K. H., Verni, F., and Roasati, G. (2000). Defensive extrusive ectosymbionts of *Euplotidium* (Ciliophora) that contain microtubule-like structures are bacteria related to Verrucomicrobia. *PNAS* 97, 1813–1817. doi: 10.1073/pnas.030438197
- Qin, W., Amin, S. A., Martins-Habbena, W., Walker, C. B., Urakawa, H., Devol, A. H., et al. (2014). Marine ammonia-oxidizing archaeal isolates display obligate mixotrophy and wide ecotypic variation. *PNAS* 111, 12504–12509. doi: 10.1073/pnas.1324115111
- Radajewski, S., Ineson, P., Parekh, N. R., and Murrell, J. C. (1999). Stable-isotope probing as a tool in microbial ecology. *Nature* 403, 646–649. doi: 10.1038/35001054
- Richardson, P., Armstrong, R., and Goldman, C. R. (1970). Contemporaneous disequilibrium, a new hypothesis to explain the “Paradox of the Plankton.” *PNAS* 67, 1710–1714. doi: 10.1073/pnas.67.4.1710
- Saleem, M., Fetzer, I., Dormann, C. F., Harms, H., and Chatzinotas, A. (2012). Predator richness increases the effect of prey diversity on prey yield. *Nat. Commun.* 3:1305. doi: 10.1038/ncomms2287
- Santoro, A. E., Richter, R. A., and Dupont, C. L. (2018). Planktonic marine archaea. *Ann. Rev. Mar. Sci.* 11, 131–158. doi: 10.1146/annurev-marine-121916-063141
- Scheffer, M., and Carpenter, S. R. (2003). Catastrophic regime shifts in ecosystems: linking theory to observation. *Trends Ecol. Evol.* 18, 648–656. doi: 10.1016/j.tree.2003.09.002
- Seyler, L. M., McGuinness, L. M., and Kerkhof, L. J. (2014). Crenarchaeal heterotrophy in salt marsh sediments. *ISME J.* 8, 1534–1543. doi: 10.1038/ismej.2014.15
- Seyler, L. M., McGuinness, L. R., Gilbert, J. A., Biddle, J. F., Gong, D., and Kerkhof, L. J. (2018). Discerning autotrophy, mixotrophy, and heterotrophy in marine TACK archaea from the North Atlantic. *FEMS Microbiol. Ecol.* 94:fy014. doi: 10.1093/femsec/fy014
- Šimek, K., Pernthaler, J., Weinbauer, M. G., Hornák, K., Dolan, J. R., Nedoma, J., et al. (2001). Changes in bacterial community composition and dynamics and viral mortality rates associated with enhanced flagellate grazing in a mesoeutrophic reservoir. *Appl. Environ. Microbiol.* 67, 2723–2733. doi: 10.1128/aem.67.6.2723-2733.2001
- Šimek, K., Vrba, J., Pernthaler, J., Posch, T., Hartman, P., Nedoma, J., et al. (1997). Morphological and compositional shifts in an experimental bacterial community influenced by protists with contrasting feeding modes. *Appl. Environ. Microbiol.* 63, 587–595.
- Smruga, S., Fernandez, V. I., Mitchell, J. G., and Stocker, R. (2016). Chemotaxis toward phytoplankton drives organic matter partitioning among marine bacteria. *PNAS* 113, 1576–1581. doi: 10.1073/pnas.1512307113
- Soresonson, K. B., and Teske, A. (2006). Stratified communities of active archaea in deep marine subsurface sediments. *Appl. Environ. Microbiol.* 72, 4596–4603. doi: 10.1128/aem.00562-06
- Stein, J. L., and Simon, M. I. (1996). Archaeal ubiquity. *PNAS* 93, 6228–6230. doi: 10.1073/pnas.93.13.6228
- Stepanouskas, R., Fergusson, E. A., Brown, J., Poulton, N., Tupper, B., Labonté, J. M., et al. (2017). Improved genome recovery and integrated cell-size analyses of individual uncultured microbial cells and vial particles. *Nat. Commun.* 8:84. doi: 10.1038/ncomms13287
- Swanberg, N. R., and Caron, D. A. (1991). Patterns of sardine feeding in epipelagic oceanic plankton. *J. Plankton Res.* 13, 287–312. doi: 10.1093/plankt/13.2.287
- Teira, E., van Aken, H., Veth, C., and Herndl, G. J. (2006). Archaeal uptake of enantiomeric amino acids in the meso- and bathypelagic waters of the North Atlantic. *Limnol. Oceanogr.* 51, 60–69. doi: 10.4319/lo.2006.51.1.0060
- Teske, A., Hinrichs, K.-U., Edgcomb, V., Gomez, A. D. V., Kysela, D., Sylva, S. P., et al. (2002). Microbial diversity of hydrothermal sediments in the guaymas basin: evidence for anaerobic methanotrophic communities. *Appl. Environ. Microbiol.* 68, 1994–2007. doi: 10.1128/aem.68.4.1994-2007.2002
- Thurman, J., Parry, J. D., Hill, P. J., and Laybourn-Parry, J. (2010). The filter-feeding ciliates *Colpidium striatum* and *Tetrahymena pyriformis* display selective feeding behaviors in the presence of mixed, equally-sized, bacterial prey. *Protist* 161, 577–588. doi: 10.1016/j.protis.2010.04.001
- Treusch, A. H., Leininger, S., Kletzin, A., Schuster, S. C., Klenk, H. P., and Schleper, C. (2005). Novel genes for nitrite reductase and Amo-related proteins indicate a role of uncultivated mesophilic crenarchaeota in nitrogen cycling. *Environ. Microbiol.* 7, 1985–1995. doi: 10.1111/j.1462-2920.2005.00906.x

- Tuorto, S. J., Darias, P., McGuinness, L. R., Panikov, N., Zhang, T., Häggblom, M. M., et al. (2014). Bacterial genome replication at subzero temperatures in permafrost. *ISME J.* 8, 139–149. doi: 10.1038/ismej.2013.140
- Tuorto, S. J., and Taghon, G. L. (2014). Rates of benthic bacterivory of marine ciliates as a function of prey concentration. *J. Exp. Mar. Biol. Ecol.* 460, 129–134. doi: 10.1016/j.jembe.2014.06.014
- Varela, M. M., van Aken, H. M., Sintès, E., and Herndl, G. J. (2008). Latitudinal trends of Crenarchaeota and bacteria in the meso- and bathypelagic water masses of the Eastern North Atlantic. *Environ. Microbiol.* 10, 110–124. doi: 10.1111/j.1462-2920.2007.01437.x
- Venter, J. C., Remington, K., Heidelberg, J. F., Halpern, A. L., Rusch, D., Eisen, J. A., et al. (2004). Environmental genome shotgun sequencing of the Sargasso Sea. *Science* 304, 66–74. doi: 10.1126/science.1093857
- Vetriani, C., Jannasch, H. W., MacGregor, B. J., Stahl, D. A., and Reysenbach, A. L. (1999). Population structure and phylogenetic characterization of marine benthic archaea in deep-sea sediments. *Appl. Environ. Microbiol.* 65, 4375–4384.
- Vuillemin, A., Wankel, S. D., Coskun, O. K., Magriths, T., Vargas, S., Estes, E. R., et al. (2019). Archaea dominate oxic subseafloor communities over multi-million-year time scales. *Sci. Adv.* 5:eaa4108. doi: 10.1126/sciadv.aaw4108
- Wilkins, D., Lauro, F. M., Williams, T. J., Demaree, M. Z., Brown, M. V., Hoffman, J. M., et al. (2012). Biogeographic partitioning of southern ocean microorganisms revealed by metagenomics. *Environ. Microbiol.* 15, 1318–1333. doi: 10.1111/1462-2920.12035
- Worden, A. Z., Follows, M. J., Giovannoni, S. J., Wilken, S., Zimmerman, A. E., and Keeling, P. J. (2015). Rethinking the marine carbon cycle: factoring in the multifarious lifestyles of microbes. *Science* 347:1257594. doi: 10.1126/science.1257594

Conflict of Interest Statement: The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Copyright © 2019 Seyler, Tuorto, McGuinness, Gong and Kerkhof. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.