

Reports

---

1979

## Estuarine fluid mud : its behavior and accumulation

Maynard M. Nichols  
*Virginia Institute of Marine Science*

Richard Faas  
*Virginia Institute of Marine Science*

Galen Thompson  
*Virginia Institute of Marine Science*

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Geology Commons](#)

---

### Recommended Citation

Nichols, M. M., Faas, R., & Thompson, G. (1979) Estuarine fluid mud : its behavior and accumulation. Special scientific report; no. 98.. Virginia Institute of Marine Science, College of William and Mary. <http://dx.doi.org/doi:10.21220/m2-bcyc-3d73>

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact [scholarworks@wm.edu](mailto:scholarworks@wm.edu).

ESTUARINE FLUID MUD: ITS BEHAVIOR AND ACCUMULATION

Final Report

by Maynard Nichols, Richard Faas and Galen Thompson

April 1979

For: U.S. Army Research Office  
P.O. Box 12221  
Research Triangle Park  
North Carolina 27709

Under Grant: DAAG29-76-G-0086

To: Virginia Institute of Marine Science  
Gloucester Point, Virginia 23062

APPROVED FOR PUBLIC RELEASE;  
DISTRIBUTION UNLIMITED.

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER	2. GOVT ACCESSION NO.	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) Estuarine Fluid Mud: Its Behavior and Accumulation.		5. TYPE OF REPORT & PERIOD COVERED Final 1 Nov. 1975 - 28 Feb. 1979
7. AUTHOR(s) Maynard Nichols, Richard Faas and Galen Thompson		6. PERFORMING ORG. REPORT NUMBER
9. PERFORMING ORGANIZATION NAME AND ADDRESS Virginia Institute of Marine Science Gloucester Point, Virginia 23062		8. CONTRACT OR GRANT NUMBER(s) DAAG29-76-G-0086
11. CONTROLLING OFFICE NAME AND ADDRESS U.S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS R & D Project No. 1T161102BH57-01 Atmospheric & Terrestrial
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		12. REPORT DATE April 1979
		13. NUMBER OF PAGES 6 pages
		15. SECURITY CLASS. (of this report) Unclassified
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE NA
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)  NA		
18. SUPPLEMENTARY NOTES  The findings in this report are not to be construed as an official Department of the Army position, unless so designated by other authorized documents.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Estuaries Sediment transport Turbidity Suspended sediment		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  A study of fluid mud in Virginia estuaries was conducted to determine how the mud accumulates in a dynamic tidal flow regime. The mud occurs as lenses and blanket deposits in zones of fast sedimentation, i.e. on channel floors and in the turbidity maximum zone. Viscosity measurements indicate resuspension potential of the mud is greater in the turbidity maximum than elsewhere.		

ESTUARINE FLUID MUD: ITS BEHAVIOR AND ACCUMULATION

Final Report

by Maynard Nichols, Richard Faas and Galen Thompson

April 1979

For: U.S. Army Research Office  
P.O. Box 12221  
Research Triangle Park  
North Carolina 27709

Under Grant: DAAG29-76-G-0086

To: Virginia Institute of Marine Science  
Gloucester Point, Virginia 23062

APPROVED FOR PUBLIC RELEASE;  
DISTRIBUTION UNLIMITED.

## FORWARD

This report summarizes results of a study concerning the behavior and accumulation of fluid mud in estuaries. It is prepared to conform with ARO 70-31 and instruction 18 of 31 July 1975 and contains the requested information: (1) a statement of the problem, (2) a summary of important results, (3) a list of publications, (4) a list of participating scientific personnel.

## 1. Introduction

Masses of soft mud are observed on the floor of many estuaries. These dense suspensions of sediment, variously called "fluff", sludge, "slingmud" or "creme de vase" occur as transient layers, ephemeral pools and lenses 0.01 to 10.0 meters thick. The term "fluid mud" is a descriptor to describe mud of high water content with densities in the range 1.005 to 1.30 g/cc corresponding to concentrations of 10 to 480 g/l.

## 2. Statement of the Problem

Whereas much sediment in estuaries is deposited directly by settling out from suspension in the water column, a substantial amount of sediment may undergo repeated resuspension and settling. This fraction then accumulates as dense suspensions of fluid mud despite fast currents that exceed speeds normally required to erode the mud. Because fluid mud forms in dynamic flow regimes, it is logical to ask: What processes are responsible for its accumulation? In turn, how does the mud maintain its integrity and resist shear under stress of tidal currents and intense turbulence? These questions were approached by examining (1) the fluid stress on the mud surface, (2) the cohesive properties of the mud, and (3) the dynamic interaction between the mud and water.

According to a model developed by McCave (1970), deposition of suspended sediment is accomplished by trapping in a viscous sublayer of the boundary layer. Trapping occurs below a limiting shear stress for deposition. Consequently, deposition does not depend on the critical flow velocity but is controlled by the balance between input to, and ejection from, the sublayer. Deposition is essentially a function of the settling velocity. The model is valid at low shear values ( $U_* < 1.2$  cm/sec) and neglects the influence of organisms and bed roughness. The rate of deposition is given by:

$$dm/dt = C_s W$$

where  $C_s$  is the concentration of suspended sediment of settling velocity  $W$ , just<sup>s</sup> above an assumed plane which is taken as the edge of the viscous sublayer.

Another model postulates periodic settling at each slack tide. Krone (1962) deduced a linear relation between rate of deposition and shear stress given by:

$$R = C_s W (1 - \tau_0/\tau_1)$$

where  $\tau_1$  is the limiting shear stress for deposition and  $\tau_0$  the bottom shear stress. The occurrence of fluid mud is an indication of the long-term balance between the factors  $C_s$ ,  $W$  and  $p$ , i.e. the probability of deposition which depends on the time for<sup>s</sup> which  $\tau_0 < \tau_1$ .




Figure 1. Schematic diagram of the sediment-water interface probe and sensors.

#### 4. Site Description

The James and Rappahannock estuaries and the Upper Chesapeake Bay are transitional zones between fresh-water flows from rivers and the marine environment. The tide, which ranges 35 to 51 cm, produces unsteady quasi-periodic flows with speeds varying from nearly zero at slack water to 65 cm per sec at maximum current. In the transition from river to tide-induced flow, salinity ranges from nearly zero to about 16 ppt and produces a neutrally stratified bottom boundary layer. Wave action was not important at the time of observations.

Sediments transported into landward parts of these estuaries are mainly derived from the river. They are fine-grained, 2 to 16 $\mu$  particle size,

Field Observations. Fluid mud was recorded in vertical density profiles throughout channels of the James and Rappahannock estuaries from both freshwater and saline zones. It attains greatest thickness, up to 62 cm, in shipping channels where sedimentation is fast. With a rate of 10 cm per year, it takes 3 to 4 years for fluid mud to dewater and consolidate to a density greater than 1.30 g per cc. Relative large "blanket" deposits 10 to 20 cm thick persist in the turbidity maximum zone near the inner limit of salty water. Most of the deposits examined are static suspensions or settled mud. However, the top millimeter or less may move in response to storm waves or strong tidal currents. The resuspension potential partly depends on mud viscosity.

Viscosity measurements and resulting rheograms reveal that the mud exhibits both a pseudoplastic and a dilatant behavior. During accelerating rates of shear, the apparent viscosity profile decreases to its lowest value and becomes highly variable. By contrast, during decelerating shear, the profile displayed a hysteresis effect. This indicates thixotropic behavior whereby the mud changes its properties, i.e. yield stress, and viscosity. Viscosity decreased more rapidly in mud from low salinity zones (2 to 8 ppt), i.e. the zone of the turbidity maximum, than in mud from high salinity zones or from fresh water. Yield values increased with settling time of the mud being greatest in mud from low salinity zones. The observations show that the resuspension potential of mud in the turbidity maximum zone is greater than in either more saline or in freshwater zones. These data suggest the bed is responsive to shear stresses, either ejecting sediment or resisting shear with time.

Reynolds stress and bed shear stress were derived from times series tidal current measurements of a two component electromagnetic meter. Measurements were made at 6, 15 and 100 cm above the mud-water interface at four stations in the Rappahannock over one tidal cycle. The data were analyzed to obtain mean longitudinal ( $\mu$ ) and vertical ( $\omega$ ) currents and corresponding turbulent fluctuations ( $\mu$  and  $\omega$ ). The Reynolds stress ranged from nearly zero to about 2.0 dynes per sq cm generated by currents reaching 30 cm per sec. Stress increased linearly with acceleration of mean current. Like bed shear stress, it lagged the maximum current velocity by one to two hours, reflecting the increased turbulence intensity during deceleration of the tidal current. However, there is a high sampling variability. Intermittency of the fluctuations extends upward one meter above the bed where most turbulence is produced.

It is concluded that deposition of fluid mud is controlled by turbulent processes active near the bed. Fresh mud of low viscosity and relatively low shear strength will fail when stress exceeds about 0.8 dynes per sq cm. It is broken into small units which are resuspended by turbulent lift forces. When tidal currents are weak during 3 to 5 days of neap tide range, the mud develops sufficient strength, by attachment of cohesive bonds, to withstand current stress up to 0.8 dynes per sq cm. The mud may be eroded quickly, but deposition of mud masses is a slow process.

## 6. Publications

"Sticky Muds": Viscosity of Estuarine Sediments, by R. Faas, Abstr. Annual Meet AAPG-SEPM 1979, p. 84.