

W&M ScholarWorks

Reports

7-31-2010

# 2010 Annual Report Estimating Relative Juvenile Abundance of Ecologically Important Finfish in the Virginia Portion of Chesapeake Bay Project Number: F-104-R-14

Troy D. Tuckey Virginia Institute of Marine Science

Mary C. Fabrizio Virginia Institute of Marine Science

Follow this and additional works at: https://scholarworks.wm.edu/reports

Part of the Aquaculture and Fisheries Commons

#### **Recommended Citation**

Tuckey, T. D., & Fabrizio, M. C. (2010) 2010 Annual Report Estimating Relative Juvenile Abundance of Ecologically Important Finfish in the Virginia Portion of Chesapeake Bay Project Number: F-104-R-14. Virginia Institute of Marine Science, William & Mary. https://doi.org/10.25773/P0JD-VH42

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

## **2010 ANNUAL REPORT**

## Estimating Relative Juvenile Abundance of Ecologically Important Finfish in the Virginia Portion of Chesapeake Bay

Project Number: F-104-R-14

Submitted to:

Virginia Marine Resources Commission P.O. Box 756 Newport News, VA 23607-0756

Prepared by:

Troy D. Tuckey and Mary C. Fabrizio

School of Marine Science, Virginia Institute of Marine Science The College of William and Mary, Gloucester Point, Virginia 23072



31 July 2010

ACKNOWLEDGMENTS	3
LIST OF TABLES	4
LIST OF FIGURES	6
EXECUTIVE SUMMARY	8
INTRODUCTION	9
METHODS	11
RESULTS	16
DISCUSSION	24
LITERATURE CITED	27
TABLES	31
FIGURES	56
APPENDICES	76

#### ACKNOWLEDGMENTS

Thanks to the many individuals who participated in the field collections, often under difficult and arduous circumstances, especially Captains Wendy Lowery and Hank Brooks, and the scientific and technical staff including Aimee Halvorson, Jennifer Greaney, Ashleigh Rhea, Karen Capossela, Ryan Schloesser, and Leonard Machut. Appreciation is expressed to Chris Bonzek for data management assistance.

We thank the marinas that provided monthly mooring facilities for the *R/V Fish Hawk*: Sunset Marina and Kingsmill Marina on the James River, and Norview Marina on the Rappahannock River.

This project was supported by the Virginia Marine Resources Commission, Project No. F-104-R-14.

Cover: Juvenile spot captured by trawl.

#### DISCLAIMER

Some of the results contained in this report have recently been completed and may contain errors and/or need further refinement.

## LIST OF TABLES

Table 1. Yearly comparison of substrate (habitat type) 1998 to 2010	
Table 2. Spatial, temporal, and length criteria used to calculate indices	33
Table 3. Summary of samples collected, 1955 - May 2010	
Table 4. VIMS Trawl Survey pooled catch from June 2009 to May 2010	
Table 5. American eel indices (1988–2009).	37
Table 6. Spring Atlantic croaker indices (1988–2009).	
Table 7. Bay anchovy indices (1988–2009).	
Table 8. Black sea bass indices (1988–2008)	40
Table 9. Juvenile blue catfish indices (1988–2009)	41
Table 10. Age 1+ blue catfish indices (1988–2009)	42
Table 11. Juvenile channel catfish indices (1988–2009)	43
Table 12. Age 1+ channel catfish indices (1988–2009).	44
Table 13. Scup indices (1988–2008)	45
Table 14. Silver perch indices (1988–2009)	46
Table 15. Spot indices (1988–2009)	47
Table 16. Striped bass indices (1988–2009).	48
Table 17. Summer flounder indices (1988–2009)	49
Table 18. Weakfish indices (1988–2009)	50
Table 19. Juvenile white catfish indices (1988–2009)	51
Table 20. Age 1+ white catfish indices (1988–2009).	
Table 21. Juvenile white perch indices (1988–2009)	53
Table 22. Age 1+ white perch indices (1988–2009)	54

Appendix Table 1.	VIMS Trawl Survey Advisory Service Requests	76
Appendix Table 2.	Mobjack Bay pooled catch for March to May 2010	77

## LIST OF FIGURES

Figure 1.	The VIMS Trawl Survey random stratified design
Figure 2.	American eel random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of American eel from index strata and months
Figure 3.	Spring juvenile Atlantic croaker random stratified (RSI, 95% C.I.), fixed transect (Rivers only - RO), and Bay and fixed river station (BRI) indices, and distribution of juvenile Atlantic croaker from index strata and months
Figure 4.	Bay anchovy random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of index-sized bay anchovy from index strata and months
Figure 5.	Juvenile black sea bass random stratified (RSI, 95% C.I.), fixed transect (Rivers only - RO), and Bay and fixed river station (BRI) indices, and distribution of juvenile black sea bass from index strata and months
Figure 6.	Juvenile blue catfish random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of juvenile blue catfish from index strata and months62
Figure 7.	Age 1+ blue catfish random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of age 1+ blue catfish from index strata and months
Figure 8.	Juvenile channel catfish random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of juvenile channel catfish from index strata and months64
Figure 9.	Age 1+ channel catfish random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of age 1+ channel catfish from index strata and months65
Figure 10	. Scup random stratified index (RSI, 95% C.I.), and distribution of index-sized scup from index strata and months
Figure 11	. Juvenile silver perch random stratified (RSI, 95% C.I.), fixed transect (Rivers only - RO), and Bay and fixed river station (BRI) indices, and distribution of index-sized juvenile silver perch from index strata and months
Figure 12.	Juvenile spot random stratified (RSI, 95% C.I.), fixed transect (Rivers only - RO), and Bay and fixed river station (BRI) indices, and distribution of index-sized juvenile spot from index strata and months
Figure 13	. Juvenile striped bass random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of index-sized juvenile striped bass from index strata and months

Figure 14.	Juvenile summer flounder random stratified (RSI, 95% C.I.), fixed transect (Rivers only - RO), and Bay and fixed river station (BRI) indices, and distribution of index-sized juvenile summer flounder from index strata and months
Figure 15.	Juvenile weakfish random stratified (RSI, 95% C.I.), fixed transect (Rivers only - RO), and Bay and fixed river station (BRI) indices, and distribution of index-sized juvenile weakfish from index strata and months
Figure 16.	Juvenile white catfish random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of index-sized juvenile white catfish from index strata and months
Figure 17.	Age 1+ white catfish random stratified (RSI) and fixed transect (Rivers only - RO) indices, and distribution of age 1+ white catfish from index strata and months73
Figure 18.	Juvenile white perch random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of index-sized juvenile white perch from index strata and months
Figure 19.	Age 1+ white perch random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices, and distribution of age 1+ white perch from index strata and months
Appendix	Figure 1. Length frequency distributions by species78

#### **EXECUTIVE SUMMARY**

#### June 2009 – May 2010

The juvenile fisheries trawl survey conducted by the Virginia Institute of Marine Science (VIMS) is the oldest continuing monitoring program (55 years) for marine and estuarine fishes in the United States. This survey provides a monthly assessment of abundance of juvenile marine and estuarine fishes and crustaceans in the tidal rivers and main stem of Chesapeake Bay. The survey provides crucial data to state, regional, and national fisheries management agencies, including the Virginia Marine Resources Commission (VMRC), the Atlantic States Marine Fisheries Commission (ASMFC), the Mid-Atlantic Fisheries Management Council (MAFMC), and the National Marine Fisheries Service (NMFS). The MAFMC recognizes the VIMS Juvenile Trawl Survey as one of the key predictors of summer flounder recruitment, and the American eel index was a vital component of the assessment that led to the 2006 ASMFC American Eel Management Plan.

Several annual indices of juvenile abundance have been generated from Trawl Survey data for species of key recreational, ecological, and commercial importance in the Virginia portion of Chesapeake Bay. These include spot, Atlantic croaker, weakfish, summer flounder, black sea bass, scup, striped bass, white perch, white catfish, channel catfish, blue catfish, silver perch, American eel, and bay anchovy. Historically, four different estimates of relative abundance were developed and reported for juvenile finfish in the survey. However, only the unconverted indices (Random Stratified Index – RSI, 1988 to present) for the target species are the focus of this report. We chose to use this index because it is based on data collected from a random stratified survey design with consistent spatial and temporal domains. Furthermore, gear changes since 1988 were minor and inconsequential (replacement of standard trawl doors with China-V doors in 1991), thus, the index can be calculated without the use of gear conversion factors.

In 2009, abundance indices for American eel, black sea bass, channel and white catfish (YOY and age 1+), spot, striped bass, summer flounder, and white perch age 1+ were below the time series average (1988 – 2009, based on the RSI). Abundance indices for YOY Atlantic croaker, bay anchovy, blue catfish age 1+, scup, and silver perch indicated above average recruitment, while YOY blue catfish, weakfish, and white perch experienced average recruitment.

In May 2010 a scale model of the trawl survey gear was tested in a flume tank at Memorial University, Centre for Sustained Aquatic Resources, Newfoundland, Canada. Results from the gear testing project will allow us to optimize the attachment of gear mensuration sensors to the trawl net to measure the wing spread without affecting how the trawl net currently fishes. With the measured wing spread and GPS starting and ending positions, relative abundance data can be calculated on an area swept basis instead of assuming a constant area is sampled by each tow as is the current practice.

Routine sampling in Mobjack Bay, located on the western side of Chesapeake Bay and north of the York River entrance, was initiated in March 2010. Mobjack Bay sampling stations consist of a mix of seven fixed and ten random stratified stations each month that encompass the main portion of the bay and the four river systems that empty into it. Extensive seagrass beds occur in Mobjack bay and likely serve as important nursery habitat for juvenile fishes. Results from Mobjack Bay sampling were analyzed separately to maintain consistency with spatial coverage discussed in previous reports. In three months of sampling (51 stations), 26 species were captured totaling 34,593 individuals. Eleven species were captured in greater relative abundance in Mobjack Bay compared with a subset of data for the same time period (March – May) from the full trawl survey.

#### **INTRODUCTION**

Relative abundance estimates of early juvenile (age 0) fishes and invertebrates generated from fishery-independent survey programs provide a reliable and early indicator of year-class strength (Goodyear, 1985), and may be used to evaluate the efficacy of management actions. The Chesapeake Bay Stock Assessment Committee (CBSAC) reviewed available indices of juvenile abundance for important fishery resources in Chesapeake Bay (hereafter referred to as "Bay") and recommended that "a unified, consistent trawl program should be one of the primary monitoring tools for finfish and crab stock assessment" (Chesapeake Bay Program Stock Assessment Plan, Chesapeake Executive Council, 1988). Subsequently, pilot studies directed at developing a comprehensive trawl survey for Chesapeake Bay began at VIMS with monthly trawl sampling in the main stem of the lower Bay. This effort complemented and expanded the monthly trawl sampling conducted in major Virginia tributaries (James, York, and Rappahannock rivers) by the Virginia Institute of Marine Science (VIMS).

The present sampling program, which includes the Bay and its tributaries, ensures that data are of sufficient geographic coverage to generate relative abundance indices for recreationally, commercially, and ecologically important finfishes and invertebrates. The National Marine Fisheries Service Marine Recreational Fisheries Statistics Survey shows that recreational catches in Virginia are dominated by Atlantic croaker (*Micropogonias undulatus*), summer flounder (*Paralichthys dentatus*), spot (*Leiostomus xanthurus*), striped bass (*Morone saxatilis*), black sea bass (*Centropristis striata*), bluefish (*Pomatomus saltatrix*), pigfish (*Orthopristis chrysoptera*), weakfish (*Cynoscion regalis*), and kingfishes (*Menticirrhus* spp.). These species depend on the lower Chesapeake Bay and its tributaries as nursery areas and, with the exception of bluefish, are highly vulnerable to bottom trawls. Additional species of recreational interest, such as scup (*Stenotomus chrysops*), white perch

(*Morone americana*), silver perch (*Bairdiella chrysoura*), white catfish (*Ameiurus catus*), channel catfish (*Ictalurus punctatus*) and blue catfish (*I. furcatus*), are also taken with sufficient regularity during trawling operations to provide information suitable for the generation of juvenile abundance indices. Although annual juvenile indices are the primary focus of this project, survey results can be used to address other aspects of finfish population biology, such as habitat utilization, early growth and survival, environmental effects on recruitment, or disease prevalence. For example, episodic climatic events, such as hurricanes, affect recruitment of shelf spawning species such as Atlantic croaker (Montane and Austin, 2005).

The development of juvenile indices requires a continuous time series of data to determine the most appropriate area-time sequences for index calculations. Provisional annual juvenile abundance indices were developed for spot, weakfish, Atlantic croaker, summer flounder, and black sea bass (Colvocoresses and Geer, 1991), followed by scup (Colvocoresses et al., 1992), white perch and striped bass (Geer et al., 1994), and white catfish, channel catfish, and silver perch (Geer and Austin, 1994). More recently, blue catfish, American eel (*Anguilla rostrata*), and bay anchovy (*Anchoa mitchilli*) indices were developed. Through the use of gear conversion factors and post stratification, a time series of index values can be produced back to 1955 for most species (Geer and Austin, 1997).

Many species of interest are captured in significant numbers across several year classes. As a result, both juvenile and age 1+ (i.e., all fish older than age 0) indices are reported for white perch, white catfish, channel catfish, and blue catfish.

This report summarizes the activity of the VIMS Juvenile Finfish Trawl Survey from June 2009 through May 2010. Abundance indices are provided from 1988 to the present, along with the mean value estimated across the time series; indices for years prior to 1988 are available in previous reports.

#### **METHODS**

#### Field Sampling

The field sampling protocol is described in detail in Lowery and Geer (2000). In brief, a 30' (9.14m) semi-balloon otter trawl, with 1.5" (38.1mm) stretched mesh and 0.25" (6.35mm) cod-end liner, is towed along the bottom for 5 minutes during daylight hours. Sampling in the Bay occurs monthly except during January and March, when few target species are available. Sampling in the tributaries also occurs monthly, at both the random stratified and historical fixed (mid-channel) stations. The stratification system is based on depth and latitudinal regions in the Bay, or depth and longitudinal regions in the rivers. Each Bay region spans 15 latitudinal minutes and consists of six strata: western and eastern shore shallow (4-12 ft), western and eastern shoal (12-30 ft), central plain (30- 42 ft), and deep channel ( $\geq$  42 ft). Each tributary is partitioned into four regions of approximately ten longitudinal minutes, with four depth strata in each (4-12 ft, 12-30 ft, 30-42 ft, and  $\geq$  42 ft; Figure 1). Strata are collapsed in areas where certain depths are limited. Fixed stations were assigned to a stratum according to their location and depth.

With the exception of the fixed river stations, trawling sites within strata are selected randomly from the National Ocean Service's Chesapeake Bay bathymetric grid, a database of depth records measured or calculated at 15-cartographic-second intervals. Between two and four trawling sites are randomly selected for each Bay stratum each month, and the number varies seasonally. Exceptions include the shallow water strata where only a single station is sampled each month. For most river strata, one to two random stations are selected per month. Sampling in the York River has been altered slightly as of 1991 to make depth strata similar to those in the James and Rappahannock rivers and main stem Bay. The stratification scheme for the tributaries was modified in January 1996 to create separate depth strata of 30-42 ft and  $\geq$  42 ft (Geer and Austin, 1996). Because tributary

sampling had occurred at these depths prior to 1996, samples collected previously were reassigned to the new strata established in 1996.

Fixed stations were sampled monthly (nearly continuously) since 1980 with sites in each tributary spaced at approximately 5-mile intervals from the river mouth up to the freshwater interface. From the mid-1950's (York River) and early-1960's (James and Rappahannock rivers) to 1972, fixed stations were sampled monthly using an unlined 30' trawl (gear code 010). During 1973-79, semi-annual random stratified sampling was performed by the VIMS Ichthyology Department, while the VIMS Crustaceology Department continued monitoring the fixed tributary stations on a limited monthly basis (May - November). Area-based weightings for the tributaries were previously assigned by dividing each river into two approximately equal length 'strata' by assuming that the stations in each stratum were representative of the channel areas in those reaches (see Lowery and Geer, 2000). As of 1996, all three tributaries were sampled with a random stratified design; the fixed stations were assigned to a stratum based on location and depth. The current design (combined fixed and random stations) provides greater spatial coverage and a long-term historical reference.

At the completion of each tow, all fishes are identified to species, counted, and measured to the nearest millimeter fork length (FL), total length (TL), or total length centerline (TLC, black sea bass only). Species that have varying size ranges are measured and counted by size class and large catches of a particular species are randomly subsampled, measured, and the remaining unmeasured catch is counted. In instances of extremely large catches (e.g., bay anchovy), subsampling is performed volumetrically.

Since May 1998, habitat or substrate type sampled by the trawl has been recorded (Table 1). Fish distribution and abundance may be influenced by various substrates such as shell, sponge, hydroids, and sea squirts that may be used as shelter, spawning habitat or for feeding. Substrates are

measured at each trawling site based on the quantity (volume in a standard container) observed in the net. Volumetric measurements of gelatinous zooplankton are also recorded for each trawl site because large catches of jellyfish and ctenophores may affect the catch (e.g., changes in gear saturation or efficiency).

#### Juvenile Index Computations

Many of the target species of this study are migratory and abundance estimation presents special difficulties, particularly if the timing and duration of migration varies annually. Juvenile fishes that use estuarine nursery areas are especially vulnerable to the vagaries of the environment, as many rely on wind-driven and tidal circulation patterns for transport into the estuaries as larvae and early juveniles (Norcross, 1983; Bodolus, 1994; Wood, 2000). The outward migration of some species from the nursery area may follow annually variable environmental cues (e.g., temperature changes). Ideally, juvenile abundance should be measured when young fish are fully recruited to the nursery area under study. In practice, however, this can only be accomplished if the time of maximal abundance and size at recruitment to the gear can be predicted (and surveys timed accordingly), or if surveys can be conducted with high frequency over the season of potential maximal abundance. Neither of these two approaches is practical. The period of maximal abundance and the scope of the area occupied by juvenile fish have proven to be variable among years and among species. This observation, coupled with multi-species monitoring objectives, precludes temporally intense surveys. Consequently, the survey is operated with a regular periodicity (monthly) and sample-site selection is performed using a standard sampling design for multispecies resource surveys.

A monthly size threshold value is applied to the length frequency information collected for each target species to partition the catch data into young-of-year and older components for juvenile index calculation (Table 2). Threshold values vary among months for each species and are based on

modal analyses of historical, composite length-frequency data and on reviews of ageing studies (Colvocoresses and Geer, 1991). For earlier months of the biological year, threshold values are usually arbitrary and fall between completely discrete modal size ranges. In the later part of the biological year, when the size of early spawned, rapidly growing individuals of the most recent year class may approach that of later spawned, slower growing individuals of the previous year class, threshold values were selected to preserve the numeric proportionality between year classes despite the potential misclassification of some individuals (Table 2). The extent of overlapping lengths and the proportion within that range attributable to each year class was estimated based on the shapes of the modal curve during the months prior to the occurrence of overlap. A length value was then selected, which preserves the proportional separation of year classes. Although this process involved considerable subjectivity and ignored possible interannual variability in average growth rates, the likelihood of significant error is small, because only a small fraction of the total number of young-of-year individuals falls within the zone of overlap, and furthermore, most of the data used to construct juvenile indices were drawn from months when no overlap was present.

After removing non young-of-year individuals, monthly stratum-specific abundances and occurrence rates are calculated for each target species. Numbers of individuals captured are log transformed (ln(n+1)) prior to abundance calculations following Chittenden (1991). Average catch rates (and the approximate 95% confidence intervals as estimated by  $\pm$  2 standard errors of the mean) are then back-transformed to geometric means. The stratum-specific coefficient of variation is expressed as the standard deviation divided by the log-transformed mean catch: STD/ EY<sub>st</sub> (Cochran, 1977). The catch data were examined for area-time combinations that provided the best basis for juvenile index calculations. Criteria applied during the selection process included identification of maximal abundance levels, uniformity of distribution, minimization of overall variance, and

avoidance of periods in which distribution patterns indicated migratory behavior. Although identification of areas most suitable for index calculations (primary nursery zones) was generally clear, selection of appropriate time windows was more complex. Surveys are timed on regular monthly intervals that may or may not coincide with periods of maximal recruitment to the nursery areas. The use of a single (maximal) month's survey result is therefore inappropriate and would decrease sample size, increase confidence intervals, and increase the risk of sampling artifacts. Conversely, the temporal series of data incorporated into index calculations should not be longer than necessary to capture the period of maximal juvenile use of the nursery area. With this approach, three- or four-month periods (6 months for bay anchovies) that provided reasonable abundance data for each species were identified (Table 2).

Using these catch data from area-time combinations, an annual juvenile index is currently calculated as the weighted geometric mean catch per tow. This is accomplished by calculating stratum-specific means and variances and combining the stratum-specific estimates using weights based on stratum area (Cochran, 1977). Because stratum areas are not uniform, a weighted mean provides an index that more closely approximates actual population abundance.

The following three indices are produced for each species for 1988 to the present: an index based on the current Bay strata plus the fixed mid-channel tributary stations (Bay & River Index – BRI), an index based on the fixed mid-channel tributary stations only (River Only – RO), and a random stratified index using all spatially appropriate data (Random Stratified Index – RSI; in previous reports, this index was called the Random Stratified Converted Index, RSCI). Data collected prior to 1988 are excluded from this report because results from the longer time series are considered provisional (i.e., indices prior to 1988 require both gear and vessel conversion factors, and

concerns about conversion factors for this period are being addressed). Multiple indices are presented in this report for completeness, but usually only the RSI will be described in detail. <u>Mobjack Bay</u>

Routine sampling in Mobjack Bay, located on the western side of Chesapeake Bay and north of the York River entrance, was initiated in March 2010 (Figure 1). Extensive seagrass beds occur in Mobjack Bay and likely serve as important nursery habitat for juvenile fishes. Seventeen stations were sampled each month (seven fixed stations and ten random stations) following the depth stratification scheme decribed previously for the trawl survey. Three fixed stations occur in the main portion of Mobjack Bay, one fixed station occurs in each river (Ware, East, Severn, and North rivers), and the ten depth-stratified random stations are selected from sites located throughout the bay and river system.

### RESULTS

A summary of tows completed from 1988 through May 2010 (Table 3) provides a comprehensive synopsis of the sampling completed for this report. For the 2009-2010 project year (June through May), 1224 sites were sampled resulting in 562,635 fishes identified and enumerated from 105 different species (Table 4). Bay anchovy, hogchoker, and Atlantic croaker accounted for 84.4% of the catch by numbers. Ignoring bay anchovy and hogchoker, five species – Atlantic croaker, spot, weakfish, white perch, and blue catfish – represented 77.8% of the catch numerically (Table 4).

Indices were calculated and described for the following species: American eel, Atlantic croaker, bay anchovy, black sea bass, blue catfish, channel catfish, scup, silver perch, striped bass, spot, summer flounder, weakfish, white catfish, and white perch. Length frequency distributions for

each species are shown in Appendix Figure 1 with index-sized fish indicated in red. Actual relative abundance indices are calculated on a subset of the data as described below.

American eel (Anguilla rostrata) – American eel are present along the Atlantic and Gulf coasts of North America and inland in the St. Lawrence Seaway and Great Lakes (Murdy et al., 1997). This catadromous species is panmictic and supported throughout its range by a single spawning population (Haro et al., 2000). Spawning takes place during winter to early spring in the Sargasso Sea. The eggs hatch into leaf-shaped, ribbon-like larvae called leptocephali, which are transported by ocean currents (over 9-12 months) in a generally northwesterly direction. Within a year, metamorphosis into the next life stage (glass eel) occurs in the Western Atlantic near the east coast of North America. Coastal currents and active migration transport the glass eels into rivers and estuaries from February to June in Virginia and Maryland. As growth continues, eels become pigmented (elver stage) and within 12-14 months, eels acquire a dark color with underlying yellow (yellow eel stage). Many eels migrate upriver into freshwater rivers, streams, lakes, and ponds, while others remain in estuaries. Most of the eel's life is spent in these habitats as a yellow eel. Age at maturity varies greatly with location and latitude, and in Chesapeake Bay may range from 8 to 24 years, with most eels in the Bay area less than 7 years old (Owens and Geer, 2003). Eels from Chesapeake Bay mature and migrate at an earlier age than eels from northern areas (Hedgepeth, 1983). Metamorphosis into the silver eel stage occurs during the seaward migration that occurs from late summer through autumn, as mature eels migrate back to the Sargasso Sea to spawn and die (Haro et al., 2000).

The current American eel index includes all eels (> 152 mm TL) collected in the upper half of each of the major tributaries (Figure 1, JA 3 and 4, YK 3 and 4, and RA 3 and 4) during April, May, and June. American eel indices exhibited above average recruitment (mean RSI = 2.13) in the late

1980's and early 1990's and below average recruitment thereafter including 2009 (Table 5; Figure 2, top). American eel are more abundant in the Rappahannock River compared with the James and York rivers (Figure 2, bottom).

Atlantic croaker (*Micropogonias undulatus*) – Atlantic croaker are typically captured in high abundance and are widely distributed throughout the survey area (Figure 3, bottom). Spawning takes place over a protracted period, such that small juveniles (<30 mm TL) can be present in catches year-round (Norcross, 1983; Colvocoresses and Geer, 1991; Colvocoresses et al., 1992; Geer et al., 1994). For some year classes, peak abundance occurs in the fall at lengths less than 100 mm TL, but for other year classes, the peak occurs the following spring. Previously, we provided two estimates of the index: a juvenile fall index (October - December) based on catches in the tributaries, and a spring recruit index (May - August) based on catches in the Bay and tributaries combined. Because the fall index does not reflect over-winter mortality, only the spring index is presented. The Atlantic croaker spring RSI has remained stable throughout the time series with recruitment in 2009 above the historic average (Table 6 and Figure 3, top; mean RSI = 1.26).

**Bay anchovy** (*Anchoa mitchilli*) – Bay anchovy are the most abundant finfish throughout Chesapeake Bay and its tributaries, and are found in salinities ranging from 1-33 ‰ (Murdy et al., 1997). Bay anchovy feed mostly on zooplankton and are an important prey of other Bay fishes (Murdy et al., 1997). In years of average freshwater inflow (e.g.,1997-2000), Atlantic menhaden, bay anchovy, and Atlantic croaker often dominate fish biomass in Chesapeake Bay (Jung, 2002). Bay anchovy abundance has increased in recent years from a period of low recruitment observed during 2001 - 2002 (Table 7; Figure 4, top). The 2009 index was above the time-series average (mean RSI = 24.32). As expected, bay anchovy are ubiquitous in trawl survey catches (Figure 4, bottom). **Black sea bass** (*Centropristis striata*) – Black sea bass are seldom taken in large numbers but regularly occur in survey catches. Young-of-year black sea bass occur throughout the Bay and appear occasionally in the lower portions of the tributaries (Figure 5, bottom). Index calculations are based on all Bay strata and the lower James stratum from May through July. Although some early juveniles appear in the Bay during their first summer and fall, more young-of-year enter the estuary during the following spring. Black sea bass spawn in the summer in the Mid-Atlantic Bight (Musick and Mercer, 1977). Thus, the index is calculated for the year class spawned the previous calendar year (i.e., the index for the 2008 year class is based on catches from May to July 2009). The black sea bass RSI was generally above average (mean RSI = 0.72) prior to 1995, but fell below average the following years with the exception of 2001 and 2007 (Table 8; Figure 5, top).

**Blue catfish** (*Ictalurus furcatus*) – The blue catfish, one of Virginia's largest freshwater fishes (Jenkins and Burkhead, 1993), was introduced to the Chesapeake Bay area as a sportfish in the James, Rappahannock, and Mattaponi rivers between 1974 and 1989 (Virginia Department of Game and Inland Fisheries, 1989 as reported by Connelly, 2001). The blue catfish is a carnivorous bottom feeder that inhabits the main channels and backwaters of rivers (Murdy et al., 1997). Blue catfish are collected from the mesohaline portions of the major tributaries upstream to the limits of the trawl survey (Figures 6 and 7, bottom). The juvenile blue catfish RSI exhibited an increasing trend over time with above average recruitment (mean RSI = 0.74) in 1996, 1997, 2003 – 2006, after which the index declined (Table 9; Figure 6, top). The abundance index for age 1+ blue catfish RSI has been above average (mean RSI = 1.82) since 2004 (Table 10; Figure 7, top).

Blue catfish indices have increased since 1988 and the ecosystem effects of this introduced species are unknown. Diets of small blue catfish are dominated by invertebrates (mostly amphipods, isopods and mud crabs), while larger blue catfish diets include invertebrates, menhaden (*Brevoortia*)

*tyrannus*), and gizzard shad (*Dorosoma cepedianum*; Parthree et al., 2008). Other catfishes (white and channel) have similar diets and may be competing with the introduced blue catfish for the same prey resources.

**Channel catfish** (*Ictalurus punctatus*) and **White catfish** (*Ameiurus catus*) – Channel catfish and white catfish are usually found in the upper portions of the tributaries (Figures 8, 9, 16, and 17, bottom). Although each river system is unique, spawning typically occurs in late May through early July in Virginia (Menzel, 1945); consequently, June was selected as the start of the biological year. The survey typically catches both species up to 600 mm FL with juveniles ( $\leq$  50 mm FL) first recruiting to the gear in June. In most years, juvenile recruitment occurs from January to April for both species in the upriver strata only.

The channel catfish was introduced to Virginia in the late 1800s (Jenkins and Burkhead, 1993). Juvenile channel catfish exhibited low recruitment in most years with a few notable peaks, and in 2009, one juvenile channel catfish was captured by the VIMS trawl survey. The age 1+ channel catfish RSI has exhibited below average (mean RSI = 0.84) values since 1997 (Tables 11 and 12; Figures 8 and 9, top). Similarly, RSIs for juvenile white catfish indicate below average recruitment (mean RSI = 0.38) since 1994 and age 1+ white catfish exhibited below average abundance (mean RSI = 1.25) since 1998 (Tables 19 and 20; Figures 16 and 17, top).

**Scup** (*Stenotomus chrysops*) – Scup are primarily a marine, summer spawning species that use the Bay in a manner similar to black sea bass. The estuary is rarely used as a nursery area by early juveniles, but older juveniles can be found in the Bay during their second summer. Early juvenile scup (25-40 mm FL) occasionally appear in survey catches in June. Older scup first appear in catches in May, and by June, they range from 50 to 215 mm FL. Using the original length threshold for scup that was based on ageing studies (Morse, 1978), trawl survey catches were found

to typically include three age groups (age 0, age 1, and age 2+). Because catches of age 0 and age 2+ scup are highly variable and low, index calculations using trawl survey catches are based on age 1 individuals only. Age 1 fish are present in the Bay and available to the gear for the entire summer and early fall.

Scup are predominantly collected in the lower Bay (Figure 10, bottom). Catch rates for scup usually peak in July, and the index is calculated from catches taken in June to September. Scup indices have increased in recent years with the 2006 index approaching the maximum observed since 1988 (mean RSI = 1.37; Table 13; Figure 10, top).

Silver perch (*Bairdiella chrysoura*) – Silver perch are found in all sampling strata (Figure 11, bottom). Spawning occurs in the deep waters of the Bay and offshore from May to July, and juveniles ( $\leq 100 \text{ mm TL}$ ) enter the Bay by July (Chao and Musick, 1977; Rhodes, 1971). Abundance indices for silver perch from the RSI are consistent and stable (Table 14; Figure 11, top). The time series average RSI index for silver perch is 0.61.

**Spot** (*Leiostomus xanthurus*) – Spot indices are calculated using all strata from July to October. Spot is often one of the most abundant recreational species captured by the survey, however compared with catches in the late 1980s and early 1990s, numbers have declined (Table 15; Figure 12, top). The RSI index in 2009 was below average for the time series (mean RSI = 12.42). Spot are widely distributed throughout the Bay and tributaries (Figure 12, bottom).

**Striped bass** (*Morone saxatilis*) – Striped bass use the upper tributaries of the Bay as spawning and nursery grounds; spawning occurs from early to mid-April through the end of May, in tidal freshwater areas just above the salt wedge. Juvenile striped bass often appear in catches from May to July in size classes less than 50 to 100 mm FL (Table 16) during years of greater abundance, but then diminish in abundance until the following winter. A second, stronger and more consistent

period of juvenile abundance occurs in December and continues through February in the upper regions of the rivers. The trawl survey index for striped bass is based on this winter recruitment period. Collections of juvenile striped bass occurred in the major tributaries only (Figure 13, bottom).

Juvenile striped bass showed strong recruitment peaks for the 1993 and 2000 year classes but recruitment has declined in recent years (Table 16; Figure 13, top). Index values for the past 9 years have been below the time-series average (mean RSI = 0.87).

**Summer flounder** (*Paralichthys dentatus*) – Summer flounder spawn on the continental shelf from September through January with the peak occurring in October and November (Murdy et al., 1997). Flounder larvae enter the Bay and other Virginia estuaries from October through May with juveniles using shallow fine-substrate areas adjacent to seagrass beds (Murdy et al., 1997; Wyanski, 1990). Low water temperatures can have significant effects on growth and survival of individuals that enter the estuary in the winter (Able and Fahay, 1998). Juvenile summer flounder first appear in catches in late March, which is used as the beginning of the biological year. Juvenile summer flounder abundance continues to increase steadily throughout the summer and early fall to a late fall peak, and then trawl catches decline, presumably reflecting emigration of young fish during December. For our trawl survey, September, October, and November usually encompass the months of greatest abundance of juvenile summer flounder. Juvenile flounder are broadly distributed throughout the Bay and lower rivers. Consequently, index calculations are based on catches from the Bay and lower river strata during September, October, and November.

Juvenile summer flounder indices were greater during the early 1990s compared with recent years, but recruitment appears to be consistent since 1995 (Table 17; Figure 14, top). The 2009 index

was below average (mean RSI = 1.03). Juvenile summer flounder were captured throughout the Bay and lower portions of the rivers, particularly the York River (Figure 14, bottom).

**Weakfish** (*Cynoscion regalis*) – Weakfish are one of the dominant species in our catches, and juveniles are found throughout the Bay and tributaries (Figure 15, bottom). Juveniles have occurred in catches in late May and June, with June considered the beginning of the biological year. The weakfish index (RSI) is consistent and indicates steady recruitment since 1988 (mean RSI = 6.48; Table 18; Figure 15, top).

White perch (*Morone americana*) – Spawning of white perch occurs in the upper tributaries from March to July with a peak occurring from late April to early May. Index months include December to February for juveniles and November to February for age 1+. Index stations are from the upper river strata.

Juvenile (age 0) white perch recruitment has been variable with a time series average RSI of 8.22, whereas abundance of age 1+ white perch has been declining (Tables 21 and 22; Figures 18 and 19, top). Abundance of age 1+ white perch has been below average (mean RSI = 14.88) in 15 of the last 18 years. White perch are collected throughout the rivers, but are most abundant in the upper river stations (Figure 18 and 19, bottom).

#### Mobjack Bay

Routine sampling in Mobjack Bay began in March 2010 and a total of 34,593 fishes were collected in 51 tows (Appendix Table 2). Twenty-six species were captured with an average of 678 individuals per tow. Compared with the full trawl survey from the same time period (March – May), Mobjack Bay produced 1.8 times as many fish per tow. Eleven species were captured in greater relative abundance compared with the full survey from March to May: bay anchovy, spot, weakfish,

Atlantic herring, northern pipefish, Atlantic cutlassfish, butterfish, naked goby, lined seahorse, feather blenny, and rough scad.

#### DISCUSSION

Juvenile indices contribute to the assessment and management of important recreational and commercial species in Chesapeake Bay and the mid-Atlantic Bight. For example, the VIMS Trawl Survey was recognized by the Mid-Atlantic Fisheries Management Council (MAFMC) as an important source of the summer flounder recruitment index; the VIMS index was instrumental in shaping more protective harvest regulations in Virginia. Other indices utilized by management agencies include those for Atlantic croaker, spot, and weakfish. Though the trawl is not the preferred gear with which to sample American eel, eel indices from the trawl survey played an important role in the 2006 ASMFC American Eel FMP (ASMFC, 2006) and the U.S. Fish and Wildlife Service American Eel Status Review. In addition to management needs, the VIMS Trawl Survey also fulfills data and specimen requests from a variety of agencies, institutes, and individuals for research and educational purposes (Appendix Table 1).

Efforts to improve recruitment indices continue and include evaluation of the size ranges and months (the index period) used in index calculations. A recent VIMS Master's thesis addressed the distributional assumptions of the catch of YOY weakfish and Atlantic croaker (Woodward, 2009). The results showed that weakfish catch data can be described by a gamma distribution and those for Atlantic croaker appear to follow a lognormal distribution. Such findings indicate that indices of abundance calculated for these species could benefit from further refinements. In addition, the use of different index months for weakfish and Atlantic croaker may improve YOY indices by ensuring fewer age 1+ fish are included in YOY index calculations. However, additional work needs to be

conducted to address potential effects of depth on the distribution and catch of these species before refined indices can be recommended and adopted (Woodward, 2009).

The Trawl Survey monitoring program provides more than relative abundance indices used to tune stock assessments and aid in management activities. The data can also be used to investigate factors that influence species abundance that operate on time periods beyond annual recruitment cycles. For example, using fishery-independent survey data from 1968 – 2004 for estuarine-dependent species, Wood and Austin (2009) found that recruitment of anadromous species was negatively correlated with recruitment of species that spawn on the continental shelf. Furthermore, recruitment patterns favored one group over the other for periods greater than a decade and shifts between recruitment regimes occurred within a short period of time (2-3 years; Wood and Austin, 2009). Understanding that long-term recruitment cycles dominate for decades is an important development that affects management options. It may be difficult to observe the effects of regulations aimed at improving recruitment if regulations are enacted during a cycle of low recruitment.

Information from the Trawl Survey also provides a basis for monitoring species interactions. For example, annual catch rates of channel catfish and white catfish have declined since 1991, while catches of the introduced blue catfish have increased dramatically (Connelly, 2001; this report). Because diets and distributions of these species overlap, the observed trends may be due to competition and thus, species interactions warrant further study. Furthermore, the shift in diet of older blue catfish to include other fishes may affect ecosystem function. A recent collaborative summary of blue catfish data from the Chesapeake Bay region was presented at the 2nd International Catfish Symposium and will be published in a conference proceedings journal next year (Schloesser et al. 2010).

Changes in catches of important recreational species may be associated with degradation of estuarine nursery habitats, overfishing, poor recruitment, or a combination of these factors (Murdy et al., 1997). Although it is not possible to determine the cause of recruitment variability from trawl survey data alone, some general observations are possible. Spot recruitment indices have declined greatly over the past 50 years, but year-class strength of this oceanic spawner appears to be controlled by environmental factors occurring outside the Bay (Homer and Mihursky, 1991; Bodolus, 1994). Atlantic croaker recruitment indices show the greatest interannual variability with fluctuations possibly related to environmental conditions that vary annually. Norcross (1983) found that cold winters increased mortality in overwintering juvenile Atlantic croaker and during some years may "push" the spawning population further south, preventing access to nursery areas in Chesapeake Bay. Weakfish indices have remained low since the mid-1990s, and the decline may be attributed to both habitat degradation (loss of seagrass beds in coastal areas) and overfishing. Declines in summer flounder abundance have been observed and may be due to overfishing or year-class failure (Terceiro, 2006). Striped bass display great recruitment variability and one or two strong year classes may dominate the population at any one time (Richards and Rago, 1999). After closure of the fishery in the mid- to late-1980s due to overfishing, poor recruitment, and low stock abundance (Richards and Rago, 1999), the striped bass recruitment index peaked in 1987. Finally, white catfish and channel catfish indices, while variable, have decreased over the past 19 years, possibly due to competition with the introduced blue catfish.

The VIMS trawl survey program supplies critical data for management of fishery resources that use Chesapeake Bay as a spawning or nursery ground. Because the Bay serves as a nursery area for many coastal migratory fish, annual recruitment data are critical for multi-state management efforts along the Atlantic Coast. Furthermore, the trawl survey serves as a foundation to conduct

research on basic biological characteristics of Bay and tributary fishes, as well as a platform from which emerging issues may be addressed.

### LITERATURE CITED

- Able, K. W. and M. P. Fahay. 1998. The first year in the life of estuarine fishes in the middle Atlantic Bight. Rutgers University Press, New Jersey.
- ASMFC, 2006. Terms of Reference and Advisory Report to the American Eel Stock Assessment Peer Review. ASMFC American Eel Stock Assessment Review Panel. Stock Assessment Report No. 06-01 of the Atlantic States Marine Fisheries Commission.
- Bodolus, D. A. 1994. Mechanisms of larval spot transport and recruitment to the Chesapeake Bay. Ph. D Dissertation. College of William and Mary, Williamsburg, VA.
- Chao, L. N. and J.A. Musick. 1977. Life history, feeding habits, and functional morphology of juvenile sciaenid fishes in the York River estuary, Virginia. Fishery Bulletin 75:657-702.
- Chesapeake Executive Council. 1988. Chesapeake Bay Program Stock Assessment Plan. Agreement Commitment Report. Annapolis, MD.
- Chittenden, M. E., Jr. 1991. Evaluation of spatial/temporal sources of variation in nekton catch and the efficacy of stratified sampling in the Chesapeake Bay. Final report to Chesapeake Bay Stock Assessment Committee & NOAA/NMFS. Virginia Institute of Marine Science, Gloucester Point, VA.

Cochran, W. G. 1977. Sampling techniques. John Wiley & Sons. New York.

- Colvocoresses, J. A. and P. J. Geer. 1991. Estimation of relative juvenile abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R1. July 1990 to June 1991. Virginia Institute of Marine Science, Gloucester Point, VA.
- Colvocoresses, J. A., P. J. Geer and C. F. Bonzek. 1992. Estimation of relative juvenile abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104-2. July 1991 to June 1992. Virginia Institute of Marine Science, Gloucester Point, VA.
- Connelly, W. J. 2001. Growth patterns of three species of catfish (*Ictaluridae*) from three Virginia tributaries of the Chesapeake Bay. Master's Thesis. College of William and Mary, Williamsburg, VA.

- Geer, P. J. and H. M. Austin. 1994. Estimation of relative abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R4. July 1993 to June 1994. Virginia Institute of Marine Science, Gloucester Point, VA.
- Geer, P. J. and H. M. Austin. 1996. Estimation of relative abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R6. July 1995 to June 1996. Virginia Institute of Marine Science, Gloucester Point, VA.
- Geer, P. J. and H. M. Austin. 1997. Estimation of relative abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R7. July 1996 to June 1997. Virginia Institute of Marine Science, Gloucester Point, VA.
- Geer, P. J., C. F. Bonzek, and H. M. Austin. 1994. Juvenile finfish and blue crab stock assessment program bottom trawl survey annual data summary report series. Volume 1993. Virginia Institute of Marine Science Special Scientific Report No. 124. Virginia Institute of Marine Science, Gloucester Point, VA.
- Goodyear, C. P. 1985. Relationship between reported commercial landings and abundance of young striped bass in Chesapeake Bay, Maryland. Transactions of the American Fisheries Society 114: 92-96.
- Haro, A., W. Richkus, K. Whalen, W. D. Busch, S. Lary, T. Brush, and D. Dixon. 2000. Population decline of the American eel: Implications for Research and Management. Fisheries 25(9): 7-16.
- Hedgepeth, M.Y. 1983. Age, growth and reproduction of American eels, *Anguilla rostrata* (Lesueur), from the Chesapeake Bay area. Master's Thesis. College of William and Mary, Williamsburg, VA.
- Homer, M. L. and J. A. Mihursky. 1991. Spot. Pages 11.1-11.19 in S.L. Funderburk, J.A. Mihursky, S.J. Jordan, and D. Reiley, eds., Habitat requirements for Chesapeake Bay Living Resources, 2<sup>nd</sup> Edition. Living Resources Subcommittee, Chesapeake Bay Program. Annapolis, MD.
- Jenkins, R. E. and N. M. Burkhead. 1993. Freshwater fishes of Virginia. American Fisheries Society, Bethesda, MD.
- Jung, S. 2002. Fish community structure and the temporal variability in recruitment and biomass production in Chesapeake Bay. Ph.D. Dissertation. University of Maryland, College Park.
- Lowery, W. A. and P. J. Geer. 2000. Juvenile finfish and blue crab stock assessment program bottom trawl survey annual data summary report series. Volume 1999. Virginia Institute of Marine

Science Special Scientific Report No. 124. Virginia Institute of Marine Science, Gloucester Point, VA.

- Menzel, R.W. 1945. The catfishery of Virginia. Transactions of the American Fisheries Society 73: 364-372.
- Montane, M. M. and H. M. Austin. 2005. Effects of hurricanes on Atlantic croaker (*Micropogonias undulatus*) recruitment to Chesapeake Bay. Pages 185-192 in K. Sellner, ed., Hurricane Isabel in Perspective. Chesapeake Research Consortium, CRC Publication 05-160, Edgewater, MD.
- Morse, W. W. 1978. Biological and fisheries data on scup, *Stenotomus chrysops* (Linnaeus). National Marine Fisheries Service, Sandy Hook Laboratory, Technical Series Report No. 12.
- Murdy, E. O., R. S. Birdsong and J. A. Musick. 1997. Fishes of Chesapeake Bay. Smithsonian Institution Press.
- Musick, J. A. and L. P. Mercer. 1977. Seasonal distribution of black sea bass, *Centropristis striata*, in the Mid-Atlantic Bight with comments on the ecology and fisheries of the species. Transactions of the American Fisheries Society 106:12-25.
- Norcross, B. L. 1983. Climate scale environmental factors affecting year-class fluctuations of Atlantic croaker, *Micropogonias undulatus* in the Chesapeake Bay, VA. Ph.D Dissertation. College of William and Mary, Williamsburg, VA.
- Owens, S. J. and P. J. Geer. 2003. Size and age structure of American eels in tributaries of the Virginia portion of the Chesapeake Bay. Pages 117-124 *in* D. A. Dixon (Editor), Biology, Management and Protection of Catadromous Eels. American Fisheries Society Symposium Series 33, Bethesda, MD.
- Parthree, D. J., C. F. Bonzek and R. J. Latour. 2008. Chesapeake Bay Trophic Interactions Laboratory Services. Project NA06NMF4570299. VIMS, Gloucester Point, VA.
- Rhodes, S. F. 1971. Age and growth of the silver perch *Bairdiella chrysoura*. Master's Thesis. College of William & Mary. Williamsburg, VA.
- Richards, R. A., and P. J. Rago. 1999. A case history of effective fishery management: Chesapeake Bay striped bass. North American Journal of Fisheries Management 19:356-375.
- Schloesser, R. W., M. C. Fabrizio, R. J. Latour, G. C. Garman, B. Greenlee, M. Groves, and J. Gartland. 2010. Ecologial role of blue catfish (*Ictalurus furcatus*) in Chesapeake Bay communities and implications for management. Presented at the 2<sup>nd</sup> International Catfish Symposium St. Louis, MO.
- Terceiro, M. 2006. Summer flounder assessment and biological reference point update for 2006. Northeast Fisheries Science Center Reference Document.

- Wojcik, F. J. and W. A. Van Engel. 1988. A documentation of Virginia trawl surveys, 1955 1984, listing pertinent variables. Volume II – York River. College of William and Mary, VIMS, Gloucester Point, VA. 198 p. (cited in Table 3)
- Wood, R. J. 2000. Synoptic scale climatic forcing of multispecies recruitment patterns in Chesapeake Bay. Ph.D. Dissertation. College of William and Mary, Williamsburg, VA.
- Wood, R. J. and H. M. Austin. 2009. Synchronous multidecadal fish recruitment patterns in Chesapeake Bay, USA. Canadian Journal of Fisheries and Aquatic Sciences 66:496-508.
- Woodward, J. R. 2009. Investigating the relationships between recruitment indices and estimates of adult abundance for striped bass, weakfish, and Atlantic croaker. Master's Thesis. College of William and Mary, Williamsburg, VA.
- Wyanski, D. M. 1990. Patterns of habitat utilization in 0-age summer flounder (*Paralichthys dentatus*). Master's Thesis. College of William and Mary, Williamsburg, VA.

TABLES

	1998 -	1999	1999 -	- 2000	2000 ·	- 2001	2001 -	2002	2002	- 2003	2003 -	2004	2004 -	2005	2005	- 2006	2006 -	2007	2007 -	- 2008	2008 -	2009	2009 -	2010
Substrate Description	%of Stations	Max. Qty.	%of	Max. Qty.	%of	Max. Qty.																		
	4	,								,	4		4				4			,	Stations		Stations	
Artificial	0.68	2.0	3.90	3.0	4.83	2.0	5.68	15.0	5.72	4.0	7.03	7.0	3.27	2.0	1.98	2.0	2.34	3.0	4.17	16.0	4.41	12.0	2.53	12.0
Dead man's fingers (Bryozoan)	7.21	5.0	9.54	4.0	7.42	5.0	8.89	6.0	12.16	16.0	10.29	1.0	3.76	0.5	5.04	24.0	8.30	2.0	9.97	3.0	21.40	23.0	23.28	24.0
Detritus	27.55	6.0	40.53	10.0	37.03	4.0	36.91	6.0	50.09	7.0	61.36	10.0	65.69	8.0	54.91	2.0	59.34	19.0	59.80	10.0	50.93	3.0	49.75	5.0
Hydroids	38.59	5.0	54.19	4.0	43.20	5.0	57.01	10.5	49.36	5.0	59.15	12.0	72.63	5.0	61.77	4.0	59.09	5.0	60.29	5.0	62.21	6.0	57.68	5.0
Sea Squirts (Molgula spp.)	20.35	5.0	29.55	12.0	22.54	14.0	27.86	18.0	15.34	5.0	21.08	9.0	25.57	8.0	19.98	14.0	34.03	7.0	30.88	12.0	29.22	24.0	27.45	4.0
Seaweeds	14.79	4.0	23.84	10.0	25.67	5.0	29.86	30.0	34.30	18.0	41.75	3.0	33.74	4.0	22.79	2.0	24.73	8.0	23.20	1.0	18.10	5.0	27.45	4.0
Shell (oyster, clam, or mussel)	18.84	3.0	26.73	4.0	22.00	5.0	29.06	8.0	32.12	4.0	25.98	3.0	21.08	4.0	23.12	12.0	21.29	2.0	21.98	8.0	21.60	4.0	20.02	3.0
Sponges	7.96	6.0	9.47	5.0	9.84	5.0	12.33	10.0	14.34	18.0	11.03	3.0	9.48	4.0	7.10	4.0	8.21	4.0	11.11	17.0	14.54	25.0	14.46	5.0
Submerged Aquatic Vegetation	4.35	3.0	8.60	1.0	10.47	2.0	5.04	2.0	6.08	1.0	3.02	0.5	6.70	2.0	2.97	0.5	4.53	1.0	5.15	0.1	4.86	0.5	7.11	1.0
WormTubes	5.33	1.0	9.54	1.0	10.47	1.0	9.29	1.0	12.34	1.0	14.05	2.0	10.87	1.0	5.45	1.0	8.05	1.0	11.11	0.5	10.83	4.0	8.82	3.0
Mud <sup>2</sup>	7.36		6.50		7.60		6.73		10.89		8.99		14.30		14.37		6.62		6.86		7.22		5.23	-
Sand <sup>2</sup>	10.21		0.87		1.43		1.04		0.36		0.49		0.57		0.08		1.00		8.33		5.96		3.59	-
Unknown <sup>3</sup>	13.66		5.42	-	4.83	-	2.80		2.72	-	1.06	-	0.74	-	4.38	-	6.71		3.51	-	1.85	-	3.76	-
NUMBER OF TRAWLS:	1,3	32	1,384 1,118		1,2	49	1,1	02	1,2	24	1,2	24	1,211		1,193		1,224		1,224		1,224			

Table 1. Annual comparisons of substrate (habitat) type from June 1998 – May 2010.

1. Based on the number of occurrences of a habitat type divided by the total number of trawls.

2. Sand and Mud are used when verification can be confirmed by direct observation.

3. Unknown is used when none of the categories are found in the trawl.

Abundance is estimated relative to the capacity of a commercial tote (internal dimensions  $25.7" \times 16.6" \times 10"$ , approximately 72 liters). Categories include: 0.5 = < 1/4 bin, 1 = 1/4 bin, 2 = 1/2 bin, 3 = 3/4 bin, 4 = full bin, etc.

Table 2. Spatial, temporal, and length (mm) criteria used to calculate recruitment indices.

								V	IMS	SТ	rawl S	Survey	- Area	/ Time	e / Size	e Valu	es by S	Specie	S				
Species - Age	VIMS			S	tra	ta I	Use	ed						Month									
	SP. CODE		Bay		Jar	mes	Yo	ork	Ra	ıрр		S	Size Cut-	off Valu	es (mm)	- Darker	ed Area	s Repre	sent Inde	ex Month	IS		
		В	L	U	L	U	L	U	L	U													
		0 tt	w	p p	w	p	w	p p	o w	p p													
		0	е	е	е	е	е	е	е	е	lanuaru	February	Marah	Amril	Max	luna	lulu.	August	Contombor	Ostabar	Nevenber	December	
American Fel 1+	0060	m	r	r	r	ŕ	r	r	ſ	ſ	January			>152	>152	>152	July	August	September			December	
Atlantic Croaker Y-O-Y	0005		-								0-100	0-100	0-100	0-110	0-135	0-160	0-180	0-220	0-50	0-80	0-100	0-100	
Atlantic Croaker Recruits	0005	r	1								0-100	0-100	0-100	0-110	0-135	0-160	0-180	0-220	0-50	0-80	0-100	0-100	
Bay Anchovy Y-O-Y	0103										0-77	0-80	0-80	0-80	0-80	0-80	0-44	0-51	0-56	0-61	0-65	0-70	
Black Seabass Y-O-Y	0002										0-110	0-110	0-110	0-110	0-110	0-150	0-175	0-70	0-85	0-100	0-105	0-110	
Blue Catfish Y-O-Y	0314										0-165	0-165	0-165	0-175	0-225	0-250	0-250	0-115	0-125	0-140	0-150	0-165	
Blue Catfish 1+	0314										>165	>165	>165	>175	>225	>250	>250	>115	>125	>140	>150	>165	
Channel Catfish Y-O-Y	0040										0-130	0-130	0-130	0-140	0-150	0-50	0-80	0-105	0-120	0-130	0-130	0-130	
Channel Catfish 1+	0040										>130	>130	>130	>140	>150	>50	>80	>105	>120	>130	>130	>130	
Scup	0001										90-170	90-170	90-170	90-170	35-90	40-100	50-125	60-145	75-160	85-170	90-170	90-170	
Silver Perch Y-O-Y	0213										0-160	0-160	0-160	0-160	0-165	0-170	0-100	0-130	0-150	0-160	0-160	0-160	
Spot Y-O-Y	0033										0-200	0-200	0-50	0-75	0-100	0-135	0-160	0-180	0-200	0-200	0-200	0-200	
Striped Bass Y-O-Y	0031										0-200	0-200	0-200	0-200	0-50	0-80	0-100	0-120	0-135	0-150	0-175	0-190	
Summer Flounder Y-O-Y	0003										0-290	0-290	0-60	0-100	0-140	0-170	0-200	0-225	0-250	0-275	0-290	0-290	
Weakfish Y-O-Y	0007										0-200	0-200	0-200	0-225	0-240	0-90	0-120	0-150	0-180	0-200	0-200	0-200	
White Catfish Y-O-Y	0039										0-110	0-110	0-110	0-110	0-120	0-50	0-65	0-80	0-90	0-100	0-110	0-110	
White Catfish 1+	0039										>110	>110	>110	>110	>120	>50	>65	>80	>90	>100	>110	>110	
White Perch Y-O-Y	0032										0-85	0-85	0-85	0-95	0-35	0-65	0-73	0-80	0-85	0-85	0-85	0-85	
White Perch 1+	0032										86-300	86-300	86-300	96-300	36-300	66-300	74-300	81-300	86-300	85-300	86-300	86-300	

Table 3. Sample collection history of the VIMS Trawl Survey, 1988 – May 2010. Each entry in the table represents the number of completed tows; YR is year, TOT is total, STAT. TYPE is station type. Other codes are below and are based on Wojcik and Van Engel (1988). Appendices A - C

YR	TOT			MONTH											WA	TER S	SYSTE	M		V	essel		Ge	ar	STAT.	TYPE	TOW DUR	ATION/DIS	STANCE
		J	F	М	А	М	J	J	А	S	0	Ν	D	CL	JA	PO	RA	ΥK	ZZ	FH	JS	LN	070	108	F	R	5	OT	DIS
1988	889	69	69	62	48	82	82	82	82	82	82	80	69	576	97	0	105	111	0	0	889	0	889	0	313	576	885	0	4
1989	840	61	61	61	66	76	76	76	76	76	76	76	59	479	108	0	124	129	0	0	840	0	840	0	361	479	840	0	0
1990	827	61	61	61	61	76	76	77	75	76	69	76	58	473	108	0	119	127	0	279	548	0	827	0	354	473	826	0	1
1991	930	61	25	61	61	73	94	95	95	97	97	97	74	411	108	0	120	291	0	930	0	0	0	930	357	573	928	1	1
1992	982	79	47	79	79	97	88	88	88	89	88	88	72	404	110	0	124	344	0	982	0	0	0	982	361	621	975	7	0
1993	915	40	73	40	71	88	89	88	88	88	88	87	75	370	110	0	126	309	0	915	0	0	0	915	365	550	914	1	0
1994	911	40	73	40	73	88	88	88	88	88	88	88	69	368	110	0	124	309	0	911	0	0	0	911	363	548	906	5	0
1995	993	40	73	40	73	92	88	88	88	105	105	407	102	411	96	0	201	285	0	993	0	0	0	993	314	6/9	984	9	0
1996	1176	52	105	66	106	106	107	108	108	107	108	107	105	435	228	0	258	255	0	1176	0	0	0	1176	2/9	010	108	0	2
1997	1220	00 22	105	00 66	90 105	110	111	179	50	120	12/	120	105	420	200	0	204	200	0 80	1220	0	0	0	1220	302	910	1217	3 1	0
1990	1202	70	100	80	100	120	118	120	118	120	124	130	119	300 //02	200	0	200	204	09 187	1202	0	0	0	1202	363	940 1010	1201	2	0
2000	1367	52	122	85	101	158	111	128	125	122	141	111	105	433	250	17	266	265	136	1367	0	0	0	1367	363	1013	1367	0	0
2001	1122	30	30	30	75	112	144	111	112	135	136	111	96	384	230	35	230	230	13	1017	0	105	Ő	1122	277	845	1119	1	2
2002	1090	66	90	66	90	96	106	96	97	.00	.00	96	96	288	264	0	264	264	10	1090	0	0	0	1090	300	790	1089	1	0
2003	1191	66	96	66	96	96	111	111	111	111	111	111	105	399	264	0	264	264	0	1191	0	0	0	1191	300	891	1191	0	0
2004	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0
2005	1211	66	105	66	105	111	111	111	111	113	111	111	90	419	264	0	264	264	0	1211	0	0	0	1211	300	911	1211	0	0
2006	1193	66	105	66	105	111	111	111	111	113	111	78	105	423	242	0	264	264	0	1193	0	0	0	1193	292	901	1193	0	0
2007	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0
2008	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0
2009	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0
2010	504	66	105	83	122	138			0400	0044			0054	123	110	0	110	110	51	504	0	0	0	504	146	358	504	0	0
101	24901	1392	1985	1453	2077	2385	2266	2260	2188	2311	2311	2232	2051	9339	4549	52	4803	5672	486	22519	2277	105	2556	22345	7232	17669	24854	37	10
Syste	em:		CL	I	Lowe	er Ch	esap	eake	Bay	(Vii	ginia	a Poi	tion	)									V	essel	: FH	]	Fish Hawl	ζ.	
			JA		Jame	s Riv	ver																		JS		John Smit	h	
			PO	]	Poto	nac	Rive	r																	LN	]	Langlev II	[	
			RA	1	Rann	ahar	nocl	k Riv	er																				
			VV	-	Vork	Din	or	X IVIV	C1																				
			1 1						0		<b>T</b> )	1071	70	70.20	<b>00</b> .1	n:	1	1 D (		107	0 71	00	00.						
			LL	]	nciu	aes:	Atla	ntic	Ucea	n (A	1)-	19/1	, /8	-79,20	J02; I	Plan	Katar	IK K. (	PK)	- 19/	0-/1	, 98-	00;						
					Mobj	ack	Bay	(MB	) - 1	970-	73, 9	8-01	, 20	10; Po	come	oke S	Sound	d (CP)	) -19	73-81	, 98-	01;							
				(	Grea	t Wie	comi	ico R	. (G'	W) -	1998	8-00.																	
Gear	· Code:								-														St	tatio	n Type	e: ]	F - Fixed		
30' 6	lears		070	1	Line	d tic	kler	chai	n. 60	' brid	lle. 5	4"x2	24" č	loors											~ 1	1	R - Rando	m	
			108	1	[ ine	$\frac{1}{1}$ tio	kler	chai	-, 60	' hrid	lle n	netal	chi	na_v de	ore														
			100	1		<i>a</i> , 110	KIUI	Unall	i, 00	UIIC	ne, n	netal	UIII	ia-v du	015														
Tor	Trung		ОТ :	a ta	. d	otion			for f			at lin	tad																
10W	i ype:			s tow	aur	ation	i in n	ninut	les fo	or the	ose n	ot iis	sted.																
			DIS	is dis	stanc	e.																							

	Number	Day / morie	Percent	Catch	Adjusted	Number	Average	Standard	Minimum	Maximum
Species	of Fish	Frequency	of	Per	Percent of	of Fish	Length	Error	Length	Length
Opecies	(ΔII)	ricqueriey	Catch	Trawl	Catch	YOY	(mm)	(length)	(mm)	(mm)
hav anchow	338 113	1056	60.09	276.24	outen	253 5/0	55	0.06	21	103
bag anchovy	70 161	747	14.07	64.67	•	18 3/1	23	0.00	21	103
Atlantia araakar	57 660	747	10.25	47.11	20.67	45 206	110	0.17	12	193
Aliantic croaker	07,000 16 E 47	797	10.25	47.11	39.07	45,206	110	0.43	13	400
white perch	16,547	284	2.94	13.52	11.38	4,354	129	0.56	19	286
spot	15,341	644	2.73	12.53	10.55	12,099	131	0.35	14	237
weaktish	14,048	554	2.50	11.48	9.66	12,681	107	0.42	19	319
blue catfish	9,526	218	1.69	7.78	6.55	2,722	230	1.01	27	895
blackcheek tonguefish	4,297	402	0.76	3.51	2.96	532	130	0.46	24	193
northern searobin	3,158	178	0.56	2.58	2.17	3,133	100	0.39	21	189
striped anchovy	2,737	152	0.49	2.24	1.88	2,714	83	0.47	34	132
spotted hake	2,618	241	0.47	2.14	1.80	2,584	128	0.81	38	284
silver perch	2,352	268	0.42	1.92	1.62	2,114	123	0.59	24	199
kingfish spp	1,759	238	0.31	1.44	1.21	1,707	75	0.78	23	277
scup	1,678	104	0.30	1.37	1.15	1,663	93	0.33	49	152
gizzard shad	1,474	192	0.26	1.2	1.01	613	208	1.96	83	405
Atlantic menhaden	1,299	236	0.23	1.06	0.89	1,096	85	1.86	27	320
smallmouth flounder	1,097	175	0.19	0.9	0.75	864	83	0.5	38	139
summer flounder	1,095	406	0.19	0.89	0.75	685	239	2.92	25	670
southern kinafish	937	179	0.17	0.77	0.64	803	120	1.69	35	303
blueback herring	923	129	0.16	0.75	0.63	921	76	0.41	29	177
harvestfish	810	178	0.14	0.66	0.56	788	64	0.99	13	172
striped bass	676	170	0.14	0.00	0.00	568	138	3.54	10	623
threadfin shad	508	37	0.12	0.00	0.41	500	88	0.67	58	110
buttorfish	594	134	0.11	0.43	0.41	403	20	1.45	14	102
oveter toodfich	544	154	0.10	0.40	0.40	493	160	2.40	24	192
	541	153	0.10	0.44	0.37		100	3.10	24	3//
alewire	392	75	0.07	0.32	0.27	383	117	1.02	74	255
white catfish	368	106	0.07	0.3	0.25	153	175	5.04	56	428
Atlantic silverside	345	65	0.06	0.28	0.24	345	92	0.62	51	119
northern pipefish	310	157	0.06	0.25	0.21		142	1.99	52	245
black seabass	180	95	0.03	0.15	0.12	127	108	2.76	31	221
seaboard goby	167	39	0.03	0.14	0.11		37	0.71	22	60
striped searobin	156	79	0.03	0.13	0.11	•	73	2.76	29	188
naked goby	133	68	0.02	0.11	0.09	•	40	0.65	19	56
American eel	124	73	0.02	0.1	0.09		298	9.66	53	658
northern puffer	103	71	0.02	0.08	0.07	72	110	3.86	16	205
clearnose skate	103	46	0.02	0.08	0.07		430	5.63	105	528
lined seahorse	98	71	0.02	0.08	0.07		77	2.37	38	126
red hake	93	23	0.02	0.08	0.06		138	2.98	65	200
banded drum	86	28	0.02	0.07	0.06		78	5.97	22	229
Atlantic thread herring	72	27	0.01	0.06	0.05		89	4.61	40	182
inshore lizardfish	68	42	0.01	0.06	0.05	53	151	7.39	53	314
windowpane	57	35	0.01	0.05	0.04	49	141	7.84	48	291
American shad	57	32	0.01	0.05	0.04	56	116	1.87	82	150
skilletfish	55	31	0.01	0.04	0.04		39	1.95	19	67
Atlantic moonfish	55	23	0.01	0.04	0.04		79	2.23	44	126
bluefish	50	31	0.01	0.04	0.03		168	6.03	119	276
Atlantic cutlassfish	46	24	0.01	0.04	0.03		347	18.83	106	607
feather blenny	40	27	0.01	0.03	0.03		69	3.08	31	109
striped mullet	31	12	0.01	0.03	0.02	-	205	7 32	151	326
niafish	30	18	0.01	0.02	0.02	•	146	4 53	106	190
sea lamprev	28	21	0.00	0.02	0.02	•	162	2.84	137	201
Atlantic spadefish	25	20	0.00	0.02	0.02	•	75	6 69	24	155
silver bake	20	12	0.00	0.02	0.02		150	5 00	2-7 Q()	100
stripod cusk-ool	24	12	0.00	0.02	0.02	•	105	5.90	90 94	211
rod drum	23	12	0.00	0.02	0.02	•	67	1 4 2	47	211
	19	13	0.00	0.02	0.01		202	4.42	47	113 E0E
channel callish	19	11	0.00	0.02	0.01	1	303	10.00	00	505
anottod contract	18	14	0.00	0.01	0.01		128	0.00	04	318
sponed seatrout	18	14	0.00	0.01	0.01		1/6	8.92	128	
iongnose gar	17	15	0.00	0.01	0.01	•	743	49.2	353	1045
Spanish mackerel	13	9	0.00	0.01	0.01	•	116	16.82	37	219
green goby	11	10	0.00	0.01	0.01		43	1.62	35	52
bluntnose stingray	11	6	0.00	0.01	0.01		426	54.18	196	780
common carp	10	8	0.00	0.01	0.01		490	40.93	192	660

 Table 4. VIMS trawl survey pooled catch for June 2009 to May 2010 from 1,224 tows.

 Adjusted Percent of Catch Excludes Bay Anchowy and Hogchoker
## Table 4 (continued)

	Number		Percent	Catch	Adjusted	Number	Average	Standard	Minimum	Maximum
Species	of Fish	Frequency	of	Per	Percent of	of Fish	Length	Error	Length	Length
	(All)		Catch	Trawl	Catch	YOY	(mm)	(length)	(mm)	(mm)
spiny butterfly ray	10	) 7	0	0.01	0.01		591	81.26	473	1320
black drum	9	8	0	0.01	0.01		. 207	10.12	151	249
northern stargazer	9	8	0	0.01	0.01		56	22.36	23	234
northern sand lance	8	3	0	0.01	0.01		140	7.18	115	166
Atlantic stingray	7	7	0	0.01	0		278	36.1	172	401
northern kingfish	7	6	0	0.01	0	5	5 105	17.55	45	164
smooth dogfish	6	6	0	0	0		456	38.57	343	580
smooth butterfly ray	6	6	0	0	0		436	47.26	270	630
spottail shiner	6	i 4	0	0	0		87	2.3	78	95
rough silverside	6	i 4	0	0	0		62	6.62	44	92
striped burrfish	5	4	0	0	0		140	34.12	31	232
Atlantic herring	4	4	0	0	0		49	3.99	40	59
brown bullhead	4	4	0	0	0		148	33.48	81	227
conger eel	4	4	0	0	0		519	57.44	447	690
lookdown	4	4	0	0	0		99	8.44	86	124
silver seatrout	4	4	0	0	0		148	15.12	123	191
tessellated darter	4	3	0	0	0		. 72	2.78	66	79
blue runner	4	3	0	0	0		. 149	7.31	128	159
dusky pipefish	3	3	0	0	0		. 101	3.06	97	107
star drum	3	3 3	0	0	0		. 81	1.86	79	85
inland silverside	3	2	0	0	0		66	1.76	63	69
Atlantic bumper	3	2	0	0	0		. 69	2	65	71
gulf kingfish	3	3 1	0	0	0		. 127	13.37	101	146
round herring	2	2 2	0	0	0		. 235	42.5	192	277
eastern silvery minnow	2	2 2	0	0	0		. 91	5.5	85	96
banded killifish	2	2 2	0	0	0		. 44	3	41	47
striped killifish	2	2 2	0	0	0		. 44	8.5	35	52
bluegill	2	2 2	0	0	0		43	3.5	39	46
roughtail stingray	2	2 2	0	0	0		. 447	60	387	507
rough scad	2	2 2	0	0	0		. 57	4.5	52	61
fringed flounder	2	2 2	0	0	0		. 95	6	89	101
chain pipefish	2	2 2	0	0	0		224	48.5	175	272
striped blenny	2	! 1	0	0	0		. 30	0	30	30
winter flounder	1	1	0	0	0		. 86		86	86
tautog	1	1	0	0	0		. 136		136	136
bluespotted cornetfish	1	1	0	0	0		284		284	284
threespine stickleback	1	1	0	0	0		. 67		67	67
pumpkinseed	1	1	0	0	0		. 58		58	58
pinfish	1	1	0	0	0		. 131		131	131
planehead filefish	1	1	0	0	0		40		40	40
bluespotted sunfish	1	1	0	0	0		46		46	46
spotfin mojarra	1	1	0	0	0		61		61	61

Adjusted Percent of Catch Excludes Bay Anchovy and Hogchoker

All species Combined 562,635

Table 5. American eel indices (1988–2009).

					<u></u>	-
	Ran	dom Stratified I	index (RS	51)	Original In	Idex
Year	Geo.	95% C.I.'s	C.V.	N	River Only	Ν
	Mean				(RO)	
1988	1.26	0.48 - 2.46	26.08	18	2.30	18
1989	7.93	4.62 - 13.18	10.57	31	8.82	31
1990	4.85	3.25 - 7.04	9.02	30	6.67	31
1991	2.06	0.70 - 4.50	26.34	31	2.12	31
1992	8.29	6.24 - 10.92	5.60	31	4.01	31
1993	3.56	2.44 - 5.05	9.29	31	3.68	31
1994	2.23	1.06 - 4.08	19.25	31	2.48	31
1995	2.69	2.01 - 3.53	7.83	33	2.44	33
1996	2.57	1.77 - 3.59	9.94	84	2.81	33
1997	2.29	1.11 - 4.13	18.69	90	1.37	39
1998	2.00	1.00 - 3.51	18.49	90	2.30	39
1999	1.25	0.58 - 2.19	21.67	90	1.14	39
2000	1.42	0.75 - 2.35	18.42	90	1.15	38
2001	0.79	0.18 - 1.72	35.92	90	0.46	39
2002	0.80	0.30 - 1.52	28.11	90	0.93	39
2003	0.79	0.22 - 1.61	32.68	90	0.60	39
2004	0.43	0.21 - 0.68	22.95	90	0.50	39
2005	0.35	0.21 - 0.51	18.66	90	0.47	39
2006	0.15	0.00 - 0.32	49.69	90	0.06	39
2007	0.22	0.07 - 0.38	31.92	90	0.14	39
2008	0.25	0.04 - 0.49	40.09	90	0.23	39
2009	0.57	0.23 - 1.01	27.37	90	0.34	39

	Rano	dom Stratified Ir	ndex (RS	l)	Original Index				
	Geo.				Bay & River		River Only		
Year	Mean	95% C.I.'s	C.V.	Ν	(BRI)	Ν	(RO)	Ν	
1988	0.36	0.21 - 0.44	16.05	234	0.38	234	2.22	84	
1989	0.65	0.38 - 0.85	15.63	252	0.78	252	4.63	84	
1990	0.48	0.23 - 0.67	20.56	252	0.52	252	2.98	85	
1991	4.12	2.83 - 5.84	8.87	307	4.35	238	12.87	83	
1992	1.17	0.77 - 1.67	13.17	309	1.34	240	10.26	84	
1993	1.98	1.33 - 2.80	11.20	301	2.21	240	19.40	84	
1994	0.86	0.56 - 1.22	14.33	300	0.95	240	2.98	84	
1995	0.95	0.67 - 1.28	11.55	306	0.93	246	5.55	90	
1996	0.19	0.11 - 0.28	19.63	405	0.16	242	0.36	88	
1997	1.47	1.15 - 1.85	7.78	419	0.87	255	7.78	100	
1998	1.19	0.95 - 1.47	7.51	374	0.48	214	6.21	96	
1999	1.50	1.05 - 2.05	10.83	397	1.28	232	4.08	100	
2000	0.60	0.42 - 0.80	12.68	413	0.44	245	1.39	97	
2001	0.37	0.25 - 0.49	14.38	420	0.32	256	1.18	100	
2002	1.59	1.07 - 2.22	11.59	361	1.11	197	4.80	100	
2003	0.49	0.28 - 0.74	19.19	405	0.52	241	0.28	100	
2004	0.96	0.73 - 1.22	9.34	420	0.70	255	4.42	99	
2005	0.47	0.35 - 0.59	10.46	420	0.31	256	1.85	100	
2006	1.27	1.00 - 1.59	7.90	420	0.77	256	3.92	100	
2007	1.04	0.76 - 1.37	10.34	420	0.76	256	3.05	100	
2008	4.10	3.04 - 5.45	7.16	420	2.84	256	18.62	100	
2009	1.82	1.41 - 2.30	7.65	420	1.48	256	10.06	100	

Table 6. Spring Atlantic croaker indices (1988–2009).

1	Ra	andom Stratified Ir	ndex (RSI	)		Origina	l Index	
-	Geo.				Bay & River		River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(BRI)	Ν	(RO)	Ν
1988	18.25	12.17 - 27.15	6.42	346	18.06	346	32.66	128
1989	52.47	36.27 - 75.71	4.54	374	51.59	374	22.74	128
1990	6.79	4.41 - 10.22	8.89	369	6.65	369	8.78	124
1991	22.51	15.05 - 33.43	6.04	491	22.83	350	33.41	125
1992	40.14	27.17 - 59.09	5.10	448	40.79	355	14.53	128
1993	43.31	28.80 - 64.89	5.23	449	42.71	360	28.93	132
1994	14.67	9.93 - 21.46	6.54	444	14.36	354	19.86	130
1995	18.36	12.84 - 26.07	5.66	540	18.52	362	18.57	138
1996	15.31	11.20 - 20.82	5.21	607	16.91	363	5.11	135
1997	18.96	13.63 - 26.23	5.19	625	17.33	378	12.64	150
1998	30.26	20.75 - 43.93	5.27	579	30.47	336	9.70	146
1999	15.47	11.20 - 21.22	5.35	606	14.38	360	21.26	150
2000	36.58	26.69 - 49.99	4.21	619	40.36	369	16.24	147
2001	9.55	6.93 - 13.04	6.06	627	9.23	377	4.56	150
2002	5.51	3.58 - 8.24	9.36	540	4.09	294	9.30	150
2003	18.03	13.17 - 24.56	5.01	624	20.65	378	3.41	150
2004	23.06	16.71 - 31.70	4.82	624	21.45	377	7.02	149
2005	22.27	16.01 - 30.85	4.98	613	21.26	367	8.43	150
2006	19.31	14.00 - 26.50	5.03	592	16.99	360	10.59	142
2007	23.76	17.33 - 32.44	4.69	624	21.15	378	10.27	150
2008	50.29	36.21 - 69.68	4.07	624	43.11	378	49.06	150
2009	30.12	22.30 - 40.55	4.21	624	25.64	378	25.09	150

Table 7. Bay anchovy indices (1988–2009).

	R	andom Stratified	d Index (RS	SI)	Original Index				
-	Geo.				Bay & River	<u> </u>	River Only		
Year	Mean	95% C.I.'s	C.V.	Ν	(BRI)	Ν	(RO)	Ν	
1988	0.84	0.59 - 1.13	11.89	138	0.83	138	1.04	12	
1989	2.36	1.70 - 3.17	8.93	138	2.36	138	1.52	12	
1990	1.12	0.78 - 1.53	11.63	128	1.12	128	0.50	12	
1991	1.28	0.91 - 1.72	10.76	129	1.29	129	2.35	12	
1992	0.22	0.13 - 0.32	18.86	129	0.22	129	0.19	12	
1993	1.05	0.74 - 1.42	11.46	129	1.04	129	0.76	12	
1994	1.06	0.74 - 1.45	11.85	129	1.06	129	0.60	12	
1995	0.50	0.33 - 0.69	14.47	151	0.54	127	0.62	12	
1996	0.36	0.22 - 0.52	17.99	152	0.35	128	0.38	12	
1997	0.46	0.31 - 0.63	14.63	153	0.47	129	0.23	12	
1998	0.57	0.35 - 0.82	16.40	135	0.59	111	0.32	12	
1999	0.58	0.41 - 0.77	12.22	146	0.60	122	0.48	12	
2000	0.74	0.50 - 1.02	13.39	153	0.78	129	0.93	12	
2001	1.29	0.85 - 1.84	12.89	108	1.33	84	1.31	12	
2002	0.64	0.41 - 0.90	15.16	138	0.69	114	0.57	12	
2003	0.12	0.06 - 0.18	25.11	153	0.11	129	0.12	12	
2004	0.06	0.02 - 0.10	34.69	153	0.05	129	0.06	12	
2005	0.19	0.12 - 0.26	17.66	153	0.20	129	0.06	12	
2006	0.44	0.30 - 0.60	14.14	153	0.48	129	0.06	12	
2007	0.83	0.53 - 1.18	14.68	153	0.90	129	0.49	12	
2008	0.41	0.27 - 0.57	14.90	153	0.45	129	0.43	12	

Table 8. Black sea bass indices (1988–2008).

	Rando	Original	Index			
-	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	Ν
1988	0.00	0.00 - 0.01	100.00	52	0.01	52
1989	0.43	0.00 - 1.24	61.91	51	0.25	52
1990	0.14	0.02 - 0.28	42.14	52	0.29	52
1991	0.39	0.27 - 0.53	14.11	52	0.19	52
1992	0.36	0.17 - 0.59	24.87	52	0.26	52
1993	0.18	0.07 - 0.30	28.51	52	0.45	52
1994	0.17	0.03 - 0.33	40.81	52	0.38	52
1995	0.64	0.34 - 1.00	20.18	109	0.91	52
1996	0.92	0.40 - 1.63	24.21	120	1.24	52
1997	2.40	1.55 - 3.54	11.81	120	2.33	52
1998	0.31	0.14 - 0.52	26.57	120	0.54	52
1999	0.14	0.04 - 0.25	36.47	114	0.30	52
2000	0.22	0.00 - 0.60	66.93	120	0.10	52
2001	0.02	0.00 - 0.04	67.15	120	0.02	52
2002	0.61	0.14 - 1.28	36.48	120	0.50	52
2003	1.33	0.75 - 2.10	16.88	120	2.50	52
2004	1.82	0.83 - 3.35	20.95	120	3.34	52
2005	2.59	1.53 - 4.10	13.75	120	3.88	52
2006	1.52	0.74 - 2.65	19.95	120	3.34	52
2007	0.60	0.18 - 1.17	32.32	120	0.66	52
2008	0.54	0.11 - 1.12	37.77	120	0.71	52
2009	0.88	0.52 - 1.33	16.99	120	1.50	52

Table 9. Blue catfish juvenile indices (1988–2009).

	Rando	Original	Index			
-	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	Ν
1988	0.02	0.00 - 0.05	100.00	52	0.05	52
1989	0.10	0.00 - 0.28	82.29	51	0.10	52
1990	0.26	0.10 - 0.45	29.18	52	0.61	52
1991	0.85	0.51 - 1.28	16.69	52	0.42	52
1992	1.22	0.71 - 1.87	16.17	52	0.84	52
1993	0.47	0.06 - 1.03	42.59	52	0.57	52
1994	0.52	0.16 - 0.99	32.59	52	1.03	52
1995	0.48	0.14 - 0.93	33.56	109	0.62	52
1996	1.38	0.62 - 2.49	22.11	120	2.33	52
1997	3.85	2.41 - 5.89	11.17	120	4.41	52
1998	1.99	0.95 - 3.59	19.57	120	3.34	52
1999	1.06	0.54 - 1.75	19.96	114	1.73	52
2000	0.88	0.33 - 1.65	27.38	120	0.89	52
2001	0.55	0.30 - 0.85	20.39	120	0.98	52
2002	0.96	0.42 - 1.70	23.81	120	0.84	52
2003	1.81	0.94 - 3.08	18.02	120	2.38	52
2004	2.62	1.78 - 3.70	10.23	120	4.99	52
2005	5.96	3.48 - 9.81	11.36	120	13.52	52
2006	4.43	2.57 - 7.24	12.36	120	14.70	52
2007	4.79	2.80 - 7.81	11.97	120	7.00	52
2008	2.89	1.84 - 4.32	11.52	120	5.85	52
2009	3.03	1.61 - 5.23	15.61	120	4.46	52

Table 10. Blue catfish age 1+ indices (1988–2009).

	Rar	ndom Stratified	Index (RSI)	)	Original	Index
-	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	Ν
1988	0.02	0.00 - 0.06	85.43	52	0.03	52
1989	1.92	1.03 - 3.22	17.10	51	1.27	52
1990	0.04	0.00 - 0.01	72.68	52	0.09	52
1991	0.03	0.00 - 0.09	100.00	52	0.02	52
1992	0.00	0		52	0.00	52
1993	0.04	0.00 - 0.12	77.30	52	0.08	52
1994	0.05	0.00 - 0.11	58.60	52	0.09	52
1995	0.22	0.07 - 0.40	33.76	109	0.40	52
1996	0.13	0.02 - 0.26	43.48	120	0.24	52
1997	0.05	0.00 - 0.12	63.47	120	0.03	52
1998	0.06	0.00 - 0.12	49.85	120	0.04	52
1999	0.00	0		114	0.00	52
2000	0.01	0.00 - 0.02	42.25	120	0.04	52
2001	0.00	0.00 - 0.01	100.00	120	0.00	52
2002	0.00	0.00 - 0.01	100.00	120	0.00	52
2003	0.32	0.16 - 0.50	23.67	120	0.83	52
2004	0.19	0.08 - 0.32	28.85	120	0.39	52
2005	0.02	0.00 - 0.05	56.41	120	0.09	52
2006	0.01	0.00 - 0.02	83.74	120	0.00	52
2007	0.00	0.00 - 0.01	100.00	120	0.00	52
2008	0.00	0		120	0.00	52
2009	0.01	0.00 - 0.02	100.00	120	0.02	52

Table 11. Channel catfish juvenile indices (1988–2009).

	Rand	dom Stratified	Index (RS	SI)	Original	Inde
-	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	N
1988	1.41	1.05 - 1.82	9.09	52	0.91	52
1989	1.10	0.52 - 1.91	21.82	51	1.20	52
1990	2.67	1.79 - 3.83	10.56	52	1.52	5
1991	3.70	2.47 - 5.36	9.78	52	1.73	5
1992	2.12	1.46 - 2.96	10.47	52	1.48	5
1993	0.83	0.20 - 1.80	35.01	52	1.15	5
1994	0.84	0.50 - 1.27	17.04	52	1.49	5
1995	0.69	0.39 - 1.05	18.45	109	0.58	5
1996	1.08	0.60 - 1.71	17.84	120	1.17	5
1997	0.84	0.47 - 1.30	18.21	120	1.06	5
1998	0.79	0.46 - 1.19	17.60	120	0.68	5
1999	0.33	0.13 - 0.56	28.23	114	0.77	5
2000	0.25	0.11 - 0.41	26.84	120	0.31	5
2001	0.17	0.04 - 0.33	38.79	120	0.16	5
2002	0.37	0.16 - 0.61	26.31	120	0.36	5
2003	0.28	0.15 - 0.44	22.78	120	0.37	5
2004	0.32	0.14 - 0.53	26.58	120	0.54	5
2005	0.28	0.10 - 0.49	30.49	120	0.31	5
2006	0.21	0.10 - 0.33	25.62	120	0.28	5
2007	0.06	0.02 - 0.11	32.35	120	0.14	5
2008	0.06	0.02 - 0.10	32.99	120	0.10	5
2009	0.06	0.00 - 0.13	57.37	120	0.08	5

Table 12. Channel catfish age 1+ indices (1988–2009).

Table 13. Scup indices (1988–2008).

				D.
-	Rai	ndom Stratified	Index (RS	I)
	Geo.			
Year	Mean	95% C.I.'s	C.V.	N
1988	3.06	2.05 - 4.41	10.20	112
1989	4.92	3.14 - 7.45	10.03	112
1990	1.90	1.11 - 2.99	14.99	103
1991	0.65	0.41 - 0.93	15.67	104
1992	3.36	2.16 - 5.01	10.90	104
1993	0.90	0.53 - 1.35	16.67	104
1994	0.39	0.21 - 0.59	21.36	104
1995	0.54	0.29 - 0.83	20.37	104
1996	0.21	0.09 - 0.35	28.00	104
1997	0.50	0.27 - 0.75	19.83	79
1998	0.27	0.06 - 0.52	37.91	88
1999	0.13	0.02 - 0.25	41.14	105
2000	1.34	0.88 - 1.90	12.80	111
2001	0.24	0.11 - 0.37	24.52	64
2002	0.96	0.58 - 1.42	15.89	104
2003	0.46	0.28 - 0.67	17.38	104
2004	1.11	0.71 - 1.59	13.89	104
2005	1.58	0.99 - 2.36	13.77	104
2006	2.99	2.07 - 4.19	9.47	104
2007	0.20	0.09 - 0.31	25.12	104
2008	2.97	2.07 - 4.13	9.28	104

	Random	Stratified Index	(RSI)			Origina	al Index	
-	Geo.				Bay & River		River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(BRI)	Ν	(RO)	Ν
1988	0.61	0.35 - 0.92	18.30	172	0.65	172	1.02	65
1989	0.53	0.33 - 0.76	16.32	189	0.56	189	1.63	63
1990	0.69	0.49 - 0.92	11.94	185	0.75	185	4.08	59
1991	0.35	0.21 - 0.51	17.33	179	0.40	179	1.47	62
1992	0.81	0.49 - 1.18	15.80	178	0.86	178	1.95	61
1993	0.45	0.29 - 0.63	16.01	180	0.45	180	0.60	63
1994	0.25	0.11 - 0.40	25.42	180	0.26	180	0.37	63
1995	0.58	0.34 - 0.87	15.65	180	0.65	180	1.81	67
1996	0.59	0.38 - 0.84	15.63	304	0.58	183	1.18	66
1997	0.71	0.50 - 0.94	12.07	316	0.79	192	1.43	75
1998	0.24	0.15 - 0.33	16.77	316	0.24	192	0.53	75
1999	0.70	0.49 - 0.94	12.42	309	0.74	186	2.51	75
2000	0.68	0.46 - 0.93	13.56	317	0.76	192	2.12	74
2001	0.70	0.47 - 0.97	13.77	327	0.85	200	3.17	75
2002	0.44	0.24 - 0.67	20.16	269	0.41	146	1.67	75
2003	0.63	0.40 - 0.90	15.49	315	0.66	192	0.71	75
2004	0.34	0.22 - 0.48	16.50	315	0.36	192	0.80	75
2005	0.76	0.52 - 1.03	12.64	315	0.77	192	2.20	75
2006	1.21	0.84 - 1.64	11.31	283	1.22	174	4.45	67
2007	0.75	0.50 - 1.03	13.53	315	0.68	192	2.26	75
2008	0.49	0.34 - 0.66	13.31	315	0.46	192	0.84	75
2009	1.00	0.72 - 1.32	10.83	315	0.92	192	1.74	75

Table 14. Silver perch indices (1988–2009).

	Rano	dom Stratified Index	(RSI)		(	Drigina	l Index	
-	Geo.				Bay & River		<b>River Only</b>	
Year	Mean	95% C.I.'s	C.V.	Ν	(BRI)	Ν	(RO)	Ν
1988	67.01	46.36 - 96.67	4.29	231	67.45	231	50.20	84
1989	31.41	24.51 - 40.18	3.44	252	32.27	252	54.19	84
1990	44.78	32.34 - 61.85	4.14	248	45.28	248	53.06	81
1991	16.83	12.28 - 21.60	4.66	238	16.56	238	21.44	83
1992	1.92	1.45 - 2.49	8.20	238	1.96	238	4.39	82
1993	9.78	7.23 - 13.13	5.68	240	9.74	240	11.85	84
1994	9.23	6.88 - 12.27	5.61	240	9.07	240	8.88	84
1995	1.56	1.15 - 2.05	9.25	248	1.52	248	2.37	92
1996	5.26	4.15 - 6.60	5.30	407	4.52	244	4.84	88
1997	11.50	9.11 - 14.45	4.20	421	8.63	256	19.68	100
1998	2.51	1.92 - 3.23	7.36	374	1.88	214	3.04	96
1999	4.72	3.63 - 6.07	6.07	402	3.98	238	6.61	100
2000	3.32	2.57 - 4.23	6.51	421	2.70	253	4.94	97
2001	3.09	2.45 - 3.85	6.06	432	2.83	264	3.69	100
2002	2.89	2.10 - 3.88	8.38	360	2.09	196	3.12	100
2003	2.85	2.25 - 3.56	6.32	420	2.58	256	2.32	100
2004	3.96	3.14 - 4.95	5.68	420	3.21	255	6.91	99
2005	12.12	9.80 - 14.94	3.78	420	8.91	256	16.58	100
2006	3.37	2.71 - 4.16	5.61	420	2.67	256	3.20	100
2007	9.17	7.38 - 11.35	4.18	420	7.79	256	12.75	100
2008	19.89	15.16 - 26.01	4.22	420	16.83	256	16.77	100
2009	6.08	4.96 - 7.40	4.39	420	4.74	256	9.05	100

Table 15. Spot indices (1988–2009).

	Random Stratified Index (RSI) Original Index					
•	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	Ν
1988	1.24	0.65 - 2.06	19.19	35	1.93	35
1989	1.65	1.12 - 2.32	11.51	37	1.59	37
1990	1.06	0.49 - 1.84	22.33	36	1.14	36
1991	1.09	0.31 - 2.33	31.00	36	1.02	36
1992	1.22	0.76 - 1.81	13.18	39	2.15	39
1993	2.52	1.09 - 4.94	19.32	41	3.30	41
1994	1.31	0.85 - 1.87	12.58	39	1.07	39
1995	0.63	0.34 - 0.99	20.19	61	1.22	39
1996	0.61	0.32 - 0.95	20.56	90	1.19	39
1997	0.55	0.25 - 0.93	24.75	90	0.41	39
1998	0.89	0.44 - 1.47	21.30	90	1.22	39
1999	0.21	0.00 - 0.47	51.55	84	0.26	39
2000	1.54	0.76 - 2.67	19.70	90	2.72	39
2001	0.53	0.27 - 0.85	21.84	90	1.94	39
2002	0.71	0.42 - 1.07	17.34	90	1.68	39
2003	0.63	0.24 - 1.13	27.59	90	1.01	39
2004	0.33	0.17 - 0.52	22.68	90	0.45	39
2005	0.59	0.30 - 0.95	21.79	90	0.53	39
2006	0.27	0.13 - 0.42	23.65	90	0.55	39
2007	0.37	0.21 - 0.55	20.10	90	0.74	39
2008	0.62	0.22 - 1.15	29.31	90	1.58	39
2009	0.48	0.28 - 0.70	17.93	90	1.06	39

Table 16. Striped bass indices (1988–2009).

	Ra	andom Stratifie	d Index (		Origi	nal Index		
-	Geo.				Bay & River		River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(BRI)	Ν	(RO)	Ν
1988	0.54	0.35 - 0.75	14.99	143	0.53	143	0.54	36
1989	1.24	0.94 - 1.58	8.77	162	1.23	162	0.96	36
1990	2.54	2.06 - 3.09	5.73	162	2.54	162	2.61	36
1991	2.79	2.26 - 3.41	5.66	153	2.78	153	1.42	36
1992	0.92	0.70 - 1.17	9.25	153	0.91	153	0.49	36
1993	0.52	0.38 - 0.68	11.87	153	0.53	153	0.49	36
1994	2.54	2.01 - 3.15	6.39	153	2.50	153	1.08	36
1995	0.71	0.52 - 0.92	10.89	149	0.72	149	0.74	36
1996	0.81	0.62 - 1.02	9.32	224	0.86	153	0.62	36
1997	0.89	0.69 - 1.12	8.77	226	0.97	153	0.70	36
1998	0.73	0.55 - 0.93	9.92	226	0.78	153	0.17	36
1999	0.53	0.41 - 0.67	9.94	219	0.58	147	0.36	36
2000	0.57	0.43 - 0.73	10.81	227	0.62	154	0.52	36
2001	0.47	0.34 - 0.61	11.84	236	0.52	161	0.53	36
2002	0.77	0.54 - 1.04	12.21	179	0.80	107	0.43	36
2003	0.44	0.33 - 0.56	10.95	225	0.43	153	0.50	36
2004	1.30	1.03 - 1.60	7.50	225	1.40	153	1.17	36
2005	0.35	0.25 - 0.46	13.18	225	0.36	153	0.29	36
2006	0.80	0.60 - 1.02	10.03	203	0.87	139	0.59	32
2007	1.00	0.78 - 1.24	8.22	225	1.04	153	0.53	36
2008	1.35	1.10 - 1.63	6.68	225	1.49	153	1.09	36
2009	0.75	0.58 - 0.92	8.76	225	0.82	153	0.84	36

Table 17. Summer flounder indices (1988–2009).

	Rand	om Stratified Inde	(	Drigina	l Index			
	Geo.				Bay & River	-	River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(BRI)	Ν	(RO)	Ν
1988	8.13	5.37 - 12.07	8.12	173	8.89	173	21.72	63
1989	11.74	8.18 - 16.88	6.44	189	12.22	189	21.27	63
1990	4.46	3.10 - 6.26	8.44	184	4.87	184	30.01	59
1991	3.16	2.32 - 4.21	7.92	179	3.56	179	15.32	62
1992	6.78	4.74 - 9.53	7.39	178	6.93	178	15.91	61
1993	5.81	4.06 - 8.17	7.76	180	6.12	180	15.42	63
1994	2.51	1.76 - 3.47	9.59	180	2.67	180	7.04	63
1995	5.95	4.26 - 8.18	7.20	186	6.07	186	11.00	69
1996	7.26	5.33 - 9.78	6.31	305	7.85	183	7.42	66
1997	6.81	5.26 - 8.74	5.38	316	7.15	192	14.82	75
1998	7.60	5.46 - 10.45	6.65	269	8.18	150	9.95	71
1999	6.78	5.01 - 9.06	6.28	303	7.38	180	16.25	75
2000	8.35	6.34 - 10.92	5.42	316	9.39	191	11.09	74
2001	5.09	3.74 - 6.82	6.93	327	5.14	200	11.52	75
2002	6.93	4.27 - 10.94	9.89	270	6.30	147	8.59	75
2003	9.23	6.72 - 12.54	6.04	315	9.34	192	5.42	75
2004	6.66	4.94 - 8.88	6.24	315	7.27	192	10.47	75
2005	5.69	4.26 - 7.50	6.31	315	5.93	192	7.10	75
2006	6.34	4.83 - 8.25	5.80	315	6.21	192	6.20	75
2007	5.35	3.99 - 7.08	6.51	315	5.30	192	14.37	75
2008	5.77	4.33 - 7.60	6.26	315	5.51	192	25.87	75
2009	6.18	4.75 - 7.96	5.63	315	6.25	192	11.44	75

Table 18. Weakfish indices (1988–2009).

	Rand	om Stratified	Original	Index		
-	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	N
1988	0.25	0.11 - 0.41	26.68	52	0.61	52
1989	3.63	2.01 - 6.12	14.03	51	3.33	52
1990	0.76	0.57 - 0.97	9.89	52	0.82	52
1991	0.07	0.02 - 0.12	34.21	52	0.19	52
1992	0.81	0.62 - 1.01	9.08	52	0.50	52
1993	0.76	0.39 - 1.24	21.11	52	1.14	52
1994	0.26	0.19 - 0.32	11.29	52	0.34	52
1995	0.21	0.08 - 0.35	29.33	109	0.46	52
1996	0.36	0.18 - 0.55	22.23	120	1.18	53
1997	0.37	0.23 - 0.53	17.47	120	0.94	52
1998	0.07	0.04 - 0.10	22.96	120	0.34	52
1999	0.00	0.00 - 0.01	100.00	114	0.00	52
2000	0.05	0.00 - 0.12	58.53	120	0.09	52
2001	0.02	0.00 - 0.04	73.60	120	0.03	52
2002	0.00	0		120	0.00	52
2003	0.29	0.17 - 0.42	19.28	120	0.99	52
2004	0.12	0.04 - 0.20	33.23	120	0.19	52
2005	0.04	0.00 - 0.09	44.35	120	0.18	52
2006	0.08	0.03 - 0.14	32.32	120	0.18	52
2007	0.02	0.00 - 0.04	46.28	120	0.07	52
2008	0.03	0.00 - 0.08	80.25	120	0.02	52
2009	0.13	0.07 - 0.20	23.87	120	0.36	52

Table 19. White catfish juvenile indices (1988–2009).

	Ran	dom Stratified	Original	Index		
-	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	Ν
1988	1.88	1.29 - 2.62	10.81	52	3.16	52
1989	3.23	1.68 - 5.67	15.78	51	4.35	52
1990	3.46	2.13 - 5.34	11.82	52	6.75	52
1991	2.12	0.90 - 4.11	21.68	52	2.31	52
1992	4.27	3.41 - 5.31	5.38	52	3.97	52
1993	2.13	1.12 - 3.64	17.19	52	1.66	52
1994	1.79	1.23 - 2.48	10.86	52	2.72	52
1995	0.94	0.45 - 1.61	22.21	109	1.77	52
1996	1.05	0.76 - 1.40	10.78	120	3.11	53
1997	1.85	1.32 - 2.49	9.82	120	3.45	52
1998	1.21	0.76 - 1.77	14.40	120	2.45	52
1999	0.56	0.36 - 0.79	15.31	114	1.51	52
2000	0.29	0.15 - 0.45	22.91	120	0.66	52
2001	0.29	0.14 - 0.47	24.65	120	0.54	52
2002	0.36	0.11 - 0.66	33.57	120	0.52	52
2003	0.48	0.26 - 0.74	20.34	120	1.13	52
2004	0.28	0.15 - 0.42	21.66	120	0.66	52
2005	0.41	0.23 - 0.62	19.83	120	0.98	52
2006	0.34	0.22 - 0.11	16.10	120	0.73	52
2007	0.23	0.11 - 0.35	24.29	120	0.62	52
2008	0.20	0.10 - 0.30	23.09	120	0.55	52
2009	0.19	0.09 - 0.29	24.47	120	0.46	52

Table 20. White catfish age 1+ indices (1988–2009).

	Random Stratified Index (RSI)						
Year	Geo.	95% C.I.'s	C.V.	Ν	River Only	Ν	
	Mean				(RO)		
1988	6.15	3.68 - 9.91	10.75	35	5.29	35	
1989	12.93	6.69 - 24.25	11.29	37	13.33	37	
1990	3.24	1.84 - 5.32	13.89	36	3.31	36	
1991	4.15	1.35 - 10.30	23.95	36	2.30	36	
1992	1.71	0.90 - 2.86	17.70	39	1.21	39	
1993	19.24	5.55 - 61.55	18.75	41	17.91	41	
1994	14.28	7.85 - 25.39	10.02	39	8.43	39	
1995	2.58	1.34 - 4.48	16.65	61	4.61	39	
1996	24.41	12.94 - 45.29	9.27	90	22.69	39	
1997	9.34	6.04 - 14.19	8.22	90	10.00	39	
1998	3.84	1.98 - 6.86	15.38	90	7.13	39	
1999	0.74	0.39 - 1.19	20.54	84	2.38	39	
2000	8.23	4.01 - 15.99	13.74	90	16.90	39	
2001	1.93	0.95 - 3.39	18.83	90	5.99	39	
2002	4.66	3.47 - 6.16	6.77	90	9.48	39	
2003	21.98	9.91 - 47.40	11.89	90	15.70	39	
2004	6.52	3.27 - 12.26	14.05	90	4.32	39	
2005	11.75	6.70 - 20.11	9.91	90	8.39	39	
2006	4.35	2.52 - 7.14	12.48	90	5.00	39	
2007	3.10	1.43 - 5.93	18.56	90	4.62	39	
2008	8.39	3.40 - 19.04	16.94	90	9.74	39	
2009	7.22	3.95 - 12.64	12.02	90	9.03	39	

Table 21. White perch juvenile indices (1988–2009).

	Ra	Original	Index			
•	Geo.				River Only	
Year	Mean	95% C.I.'s	C.V.	Ν	(RO)	Ν
1988	39.57	26.69 - 58.42	5.15	46	35.10	46
1989	22.78	16.00 - 32.25	5.29	46	25.86	46
1990	35.39	21.9 - 56.83	6.44	45	31.97	45
1991	39.52	28.45 - 54.75	4.31	44	29.49	44
1992	11.94	7.73 - 18.17	7.68	48	15.77	48
1993	10.74	4.89 - 22.37	13.98	50	15.04	50
1994	24.61	15.49 - 38.78	6.79	48	18.77	48
1995	13.69	6.17 - 29.12	13.35	70	40.82	48
1996	9.03	5.29 - 15.00	10.13	116	13.27	49
1997	19.37	10.56 - 34.90	9.40	120	20.25	52
1998	10.89	6.70 - 17.36	8.78	120	27.44	52
1999	10.34	5.97 - 17.46	10.03	114	23.48	52
2000	7.65	3.79 - 14.63	13.72	120	17.31	52
2001	4.62	2.54 - 7.92	13.36	120	17.09	52
2002	7.22	4.99 - 10.28	7.51	120	20.61	52
2003	19.13	9.95 - 36.00	10.14	120	27.35	52
2004	6.84	3.83 - 11.72	11.76	120	8.71	52
2005	8.40	5.30 - 13.04	8.95	120	9.34	52
2006	6.45	4.09 - 9.89	9.48	120	13.11	52
2007	6.48	3.88 - 10.46	10.61	120	19.58	52
2008	5.08	2.51 - 9.53	15.20	120	17.27	52
2009	7.61	4.63 - 12.17	9.87	120	20.68	52

Table 22. White perch age 1+ indices (1988–2009).

FIGURES

Figure 1. The VIMS trawl survey random stratified design in the Chesapeake Bay. Transect lines indicate geographic regions as designated below.

Chesapeake Bay	B1	Bottom Bay
	B2	Lower Bay
	B3	Upper Bay
James River	J1	Bottom James
	J2	Lower James
	J3	Upper James
	J4	Top James
York River	Y1	Bottom York
	Y2	Lower York
	Y3	Upper York
	Y4	Top York (lower Pamunkey River)
Rappahannock River	R1	Bottom Rappahannock
	R2	Lower Rappahannock
	R3	Upper Rappahannock
	R4	Top Rappahannock
Mobjack Bay	MB	Routine monitoring established March 2010



Figure 1 (continued)



Figure 2. American eel random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices and the time series average based on the RSI (Top), and distribution of American eel (all age classes combined) from index strata and months (Bottom).



Figure 3. Spring juvenile Atlantic croaker random stratified (RSI, 95% C.I.), fixed transect (Rivers only - RO), and Bay and fixed river station (BRI) indices and the time series average based on the RSI (Top), and distribution of Atlantic croaker juveniles from index strata and months (Bottom).



Figure 4. Bay anchovy random stratified (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices and the time series average based on the RSI (Top), and distribution of bay anchovy from index strata and months (Bottom).



Figure 5. Black sea bass random stratified index (RSI, 95% C.I.) and fixed transect (Rivers only - RO), and bay and fixed river station (BRI) indices and the time series average based on the RSI (Top), and distribution of juvenile black sea bass from index strata and months (Bottom).



Figure 6. Juvenile blue catfish random stratified index (RSI, 95% C.I.) and fixed transect (Rivers only -RO) indices and the time series average based on the RSI (Top), and distribution of juvenile blue catfish from index strata and months (Bottom).



Figure 7. Age 1+ blue catfish random stratified index (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices and the time series average based on the RSI (Top), and distribution of age 1+ blue catfish from index strata and months (Bottom).



Figure 8. Juvenile channel catfish random stratified index (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices and the time series average based on the RSI (Top), and distribution of juvenile channel catfish from index strata and months (Bottom). One juvenile channel catfish was captured in this time period.



Figure 9. Age 1+ channel catfish random stratified index (RSI, 95% C.I.) and fixed transect (Rivers only - RO) indices and the time series average based on the RSI (Top), and distribution of age 1+ channel catfish from index strata and months (Bottom).



Figure 10. Juvenile scup random stratified index (RSI, 95% C.I.) and the time series average (Top), and distribution of scup from index strata and months (Bottom).



Figure 11. Juvenile silver perch random stratified (RSI, 95% C.I.), fixed transect (Rivers only – RO), and Bay and fixed river station (BRI) indices and the time series average based on the RSI (Top), and distribution of juvenile silver perch from index strata and months (Bottom).



Figure 12. Juvenile spot random stratified (RSI, 95% C.I.) fixed transect (Rivers only – RO), and Bay and fixed river station (BRI) indices and the time series average based on the RSI (Top), and distribution of juvenile spot from index strata and months (Bottom).



Figure 13. Juvenile striped bass random stratified (RSI, 95% C.I.) and fixed transect (Rivers only – RO) indices and the time series average based on the RSI (Top), and distribution of juvenile striped bass from index strata and months (Bottom).



Figure 14. Juvenile summer flounder random stratified (RSI, 95% C.I.), fixed transect (Rivers only – RO), and Bay and fixed river station (BRI) indices and the time series average based on the RSI (Top), and distribution of juvenile summer flounder from index strata and months (Bottom).



Figure 15. Juvenile weakfish random stratified (RSI, 95% C.I.), fixed transect (Rivers only – RO), and Bay and fixed river station (BRI) indices and the time series average based on the RSI (Top), and distribution of juvenile weakfish from index strata and months (Bottom).


Figure 16. Juvenile white catfish random stratified (RSI, 95% C.I.) and fixed transect (Rivers only – RO) indices and the times series average based on the RSI (Top), and distribution of juvenile white catfish from index strata and months (Bottom).



Figure 17. White catfish age 1+ random stratified (RSI, 95% C.I.) and fixed transect (Rivers only – RO) indices and the time series average based on the RSI (Top), and distribution of age 1+ white catfish from index strata and months (Bottom).



Figure 18. Juvenile white perch random stratified (RSI, 95% C.I.) and fixed transect (Rivers only – RO) indices and the time series average based on the RSI (Top), and distribution of juvenile white perch from index strata and months (Bottom).



Figure 19. White perch age 1+ random stratified (RSI, 95% C.I.) and fixed transect (Rivers only – RO) indices and the time series average based on the RSI (Top), and distribution of age 1+ white perch from index strata and months (Bottom).

Appendix Table 1. Trawl survey advisory requests, data requests, and specimen requests from June 2009 to May 2010.

Contact	Agency	Nature of Request			
Pat McGrath	VIMS	Gar specimens			
Karin Limburg	SUNY	Alosine specimens			
Edward McGinley	WVU	Bay anchowy specimens			
Edward McGinley	WVU	Atlantic croaker specimens			
Edward McGinley	WVU	Atlantic menhaden specimens			
Joel Boehm	River Project	Dusky pipefish specimens			
Joel Boehm	River Project	Northern pipefish specimens			
Joel Boehm	River Project	Lined seahorse specimens			
Paul Gerdes	VIMS	Naked goby specimens			
Paul Gerdes	VIMS	Bay anchowy specimens			
Paul Gerdes	VIMS	Striped anchowy specimens			
Paul Gerdes	VIMS	Scup specimens			
Alicia Nelson	VMRC	2009 Horseshoe Crab data			
Sally Upton	VIMS	Fish specimens for education dept			
Laura Lee	VMRC	Eel data			
Jack Travelstead	VMRC	Blue catfish data summary			
Mike Newman	VIMS	Fish for mercury study			
Laura Lee	VMRC	Spot indices			
Rob O'Reilly	VMRC	Atlantic Croaker Indices, fall and spring 2008			
Eric Hilton	VIMS	Feather blenny specimens			
Eric Hilton	VIMS	Striped blenny specimens			
Eric Hilton	VIMS	Crested blenny specimens			
Troy Tuckey	VIMS	White perch juveniles			
Lyle Varnell	VIMS	Fish distribution data			
Lyle Varnell	VIMS	James River water temperature data			
VIMS Aquarium	VIMS	Live specimens for display and education			
Tracey Sutton	VIMS	Spiny dogfish data			
Chris Hager	VA Sea Grant	Atlantic sturgeon data			
Tom Miller	CBL	Blue crab indices 1968-2009, age 0 spring/fall, age 1+, adult female			

	Number		Percent	Catch	Adjusted	Number	Average	Standard	Minimum	Maximum
Species	of Fish	Frequency	of	Per	Percent of	of Fish	Length	Error	Length	Length
	(All)		Catch	Trawl	Catch	YOY	(mm)	(length)	(mm)	(mm)
bay anchovy	31,806	44	91.94	623.65	•	31,606	59	0.19	33	88
spot	2,065	28	5.97	40.49	74.09	1,641	81	2.12	19	194
weakfish	246	15	0.71	4.82	8.83	245	166	1.25	117	245
Atlantic croaker	219	17	0.63	4.29	7.86	5	202	1.85	97	343
Atlantic menhaden	86	15	0.25	1.69	3.09	16	114	4.76	29	257
spotted hake	62	18	0.18	1.22	2.22	62	142	2.87	97	180
Atlantic herring	27	7	0.08	0.53	0.97		51	2.35	31	59
northern pipefish	12	8	0.03	0.24	0.43		143	7.2	103	169
alewife	12	1	0.03	0.24	0.43	12	120	2.59	105	134
hogchoker	11	6	0.03	0.22	0.39	0	105	1.96	92	114
summer flounder	10	8	0.03	0.2	0.36	1	398	42.12	89	497
silver perch	6	4	0.02	0.12	0.22	5	137	6.35	124	166
oyster toadfish	6	1	0.02	0.12	0.22		67	9.28	49	97
Atlantic cutlassfish	5	3	0.01	0.1	0.18		365	22.3	329	452
butterfish	4	3	0.01	0.08	0.14	4	63	12.61	43	99
naked goby	3	3	0.01	0.06	0.11		54	3.71	47	59
kingfish spp	2	2	0.01	0.04	0.07	1	202	62.5	139	264
lined seahorse	2	2	0.01	0.04	0.07		77	8	69	85
American shad	2	1	0.01	0.04	0.07	2	126	4	122	130
blueback herring	1	1	0.00	0.02	0.04	0	233		233	233
striped bass	1	1	0.00	0.02	0.04	0	486		486	486
northern puffer	1	1	0.00	0.02	0.04	0	199		199	199
northern searobin	1	1	0.00	0.02	0.04	1	51		51	51
striped blenny	1	1	0.00	0.02	0.04		65		65	65
feather blenny	1	1	0.00	0.02	0.04		55		55	55
rough scad	1	1	0.00	0.02	0.03		50	<u>.</u>	50	50
	04 500									

Appendix Table 2. Mobjack Bay pooled catch for March to May 2010 from 51 tows.

All Species Combined 34,593

Appendix Figure 1. Length frequency distributions by species from June 2009 to May 2010. Red bars are index-sized fish. (Note that actual indices are calculated using a subset of months and strata. Therefore, not all index-sized fish are included.)



Appendix Figure 1. (continued)











