
Reports

2-2-2020

Evaluation of Striped Bass Stocks in Virginia: Monitoring and Tagging Studies, 2020 Progress Report - 1 December 2019 - 30 November 2020

Robert J. Latour
Virginia Institute of Marine Science

Christopher F. Bonzek
Virginia Institute of Marine Science

James Gartland
Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Latour, R. J., Bonzek, C. F., & Gartland, J. (2020) Evaluation of Striped Bass Stocks in Virginia: Monitoring and Tagging Studies, 2020 Progress Report - 1 December 2019 - 30 November 2020. Virginia Institute of Marine Science, William & Mary. <https://www.doi.org/10.25773/664n-5725>

This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

Evaluation of Striped Bass Stocks in Virginia: Monitoring and Tagging Studies, 2020

Progress Report

Contract Number: F-77-R-33

Project Period: 1 December 2019 - 30 November 2020

Principal Investigator: Robert J. Latour

Prepared by:

Robert J. Latour, Christopher F. Bonzek, James Gartland

**Department of Fisheries Science
School of Marine Science
Virginia Institute of Marine Science
William & Mary
Gloucester Point, VA 23062-1346**

Submitted To:

**Virginia Marine Resources Commission
P.O. Box 756
Newport News, VA 23607-0756**

2 February 2020

Preface

This report presents the results of striped bass (*Morone saxatilis*) tagging and monitoring activities in Virginia during the period 1 December 2019 through 30 November 2020. It includes an assessment of the biological characteristics of striped bass taken from the 2020 spring spawning run and estimates of annual survival and fishing mortality based on annual spring tagging. The information contained in this report is required by the Atlantic States Marine Fisheries Commission and is used to implement a coordinated management plan for striped bass in Virginia, and along the eastern seaboard.

Striped bass have historically supported one of the most important recreational and commercial fisheries along the Atlantic coast. In colonial times, striped bass were abundant in most coastal rivers from New Brunswick to Georgia, but overfishing, pollution and reduction of spawning habitat have resulted in periodic declines in stocks and an overall reduction of biomass (Merriman 1941, Pearson 1938). Striped bass populations at the northern and southern extremes of the Atlantic are apparently non-migratory (Raney 1957). Presently, important sources of striped bass in their native range are found in the Roanoke, Delaware and Hudson rivers and the major tributaries of Chesapeake Bay (Lewis 1957) with the Chesapeake Bay and Hudson River being the primary sources of the coastal migratory population (Dorazio *et al.* 1994).

Examination of meristic characteristics indicate that the coastal migratory population consists of distinct sub-populations from the Hudson River, James River, Rappahannock - York rivers, and upper Chesapeake Bay (Raney 1957). The Roanoke River striped bass may represent another distinct sub-population (Raney 1957). The relative contribution of each area to the coastal population varies. Berggren and Lieberman (1978) concluded from a morphological study that Chesapeake Bay striped bass were the major contributor (90.8%) to the Atlantic coast fisheries, and the Hudson River and Roanoke River stocks were minor contributors. However, they estimated that the exceptionally strong 1970 year class constituted 40% of their total sample. Van Winkle *et al.* (1988) estimated that the Hudson River stock constituted 40% - 50% of the striped bass caught in the Atlantic coastal fishery in 1965. Regardless of the exact proportion, management of striped bass is a multi-jurisdictional concern as spawning success in one area probably influences fishing success in many areas. Furthermore, recent evidence suggests the presence of divergent migratory behavior at intra-population levels (Secor 1999). The extent to which these levels of behavioral complexity impact management strategies in Chesapeake Bay and other stocks is unknown.

Concern about the decline in striped bass landings along the Atlantic coast since the mid-1970s prompted the development of an interstate fisheries management plan (FMP) under the auspices of the Atlantic States Marine Fisheries Management Program (ASMFC). Federal legislation was enacted in 1984 (Public Law 98-613, the Atlantic Striped Bass Conservation Act) which enables Federal imposition of a moratorium for an indefinite period in those states that fail to comply with the coast-wide plan. To be in compliance with the plan, coastal states have

imposed restrictions on their commercial and recreational striped bass fisheries ranging from combinations of catch quotas, size limits, closed periods and year-round moratoriums. Due to an improvement in spawning success, as judged by increases in annual values of the Maryland juvenile index, a limited fishery was established in fall, 1990. This transitional fishery existed until 1995 when spawning stock biomass reached sufficiently healthy levels (Field 1997). ASMFC subsequently declared Chesapeake Bay stocks to have reached benchmark levels and adopted Amendment 5 to the original FMP that allowed expanded state fisheries.

To document continued compliance with Federal law, the Virginia Institute of Marine Science (VIMS) has monitored the size and age composition, sex ratio and maturity schedules of the spawning striped bass stock in the Rappahannock River since December 1981 utilizing commercial pound nets and, from 1991-2014, variable-mesh experimental gill nets. Spawning stock assessment was expanded to include the James River in 1994, utilizing commercial fyke nets and variable-mesh experimental gill nets. An experimental fyke net was established in the James River to assess its potential as a source for tagging striped bass. The use of fyke nets was discontinued after 1997. In conjunction with the monitoring studies, tagging programs have been conducted in the James and Rappahannock rivers since 1987. These studies were established to document the migration and relative contribution of these Chesapeake Bay stocks to the coastal population and to provide a means to estimate annual survival rates (S). With the re-establishment of fall recreational fisheries in 1993, the tagging studies were expanded to include the York River and western Chesapeake Bay to provide a direct estimation of the resultant fishing mortality (F). Commencing in 2005, these estimates of F were estimated from the striped bass tagged during the spring in the Rappahannock River. In 2015, tagging and monitoring activities were expanded to encompass three rivers – the James, York and Rappahannock Rivers. In a meeting in September 2017 the ASMFC Striped Bass Technical Committee concluded that the Virginia pound net spawning stock monitoring program had inherent shortcomings which rendered the resulting CPUE indices unsuitable for inclusion in future stock assessments. That action, combined with budget cuts necessitated by VMRC resulted in a major change in methodology to both the spawning stock monitoring and tagging portions of the program. Beginning in 2018 monitoring has been accomplished using multi-panel anchor gill nets and tagging was conducted via electrofishing. Methodologies are fully described later in this report.

List of Tables

I. Spawning Stock Monitoring

1	Number of striped bass captured, water temperature, and salinity by week in the Rappahannock and James Rivers, spring 2020.	8
2	Catch rates by week and sex for the Rappahannock and James Rivers, spring 2020. ...	8
3	Otolith age frequencies by river and sex, spring 2020.	9
4	Weekly total biomass, by sex, of striped bass captured in the Rappahannock and James Rivers, spring 2020.	9
5	Average length (mm), weight (g) and age of striped bass by year, river and sex, spring 2020.	10
6	Length frequencies (mm TL) of striped bass sampled from the gill nets, spring 2020. .	11
7	Average length (mm) and weight (g), with standard deviations (Std Dev) of striped bass by year class, spring 2020.	12
8	Average catch per day by river and sex, in numbers and biomass with lower (LCL) and upper (UCL) confidence limits, spring 2020.	12
9	Average catch per day by river and mesh size, in numbers and biomass with lower (LCL) and upper (UCL) confidence limits, spring 2020.	13
10	Average catch per day by river, sex and mesh size, in biomass with lower (LCL) and upper (UCL) confidence limits, spring 2020.	14
11	Average, minimum and maximum scale ages for each otolith age class from ages derived from the same specimen, spring 2020.	15

II. Tagging

1	Summary data of striped bass tagged and released in the Rappahannock River, spring 2020.	36
2	Summary data of striped bass tagged and released in the James River, spring 2020.	36
3	Summary data of striped bass tagged and released York River, spring 2020.	36
4	Location of striped bass (≥ 457 mm TL), recaptured in 2020, that were originally tagged and released in the Rappahannock River during springs 1990-2019.	37
5	Location of migratory striped bass (≥ 710 mm TL), recaptured in 2020, that were originally tagged and released in Virginia during springs 1990-2019.	37
6	Description of IRCR models with varying time periods for fishing mortality, tagging mortality and natural mortality.	38
7a	Input recapture matrix for IRCR analysis: from striped bass (>457 mm TL) tagged and released in the springs of 1990-2019 (Rappahannock River only 1990-2017, all Virginia waters 2018-2019). Harvested recaptures only.	39
7b	Input recapture matrix for IRCR analysis: from striped bass (>457 mm TL) that were tagged and released in the springs of 1990-2019 (Rappahannock River only 1990-2017, all Virginia waters 2018-2019). Recaptures released with streamers cut off only.	39
8a	Input recapture matrix for IRCR analysis: from striped bass (>710 mm TL) that were tagged and released in the Rappahannock River, springs of 1990-2019 (Rappahannock River only 1990-2017, all Virginia waters 2018-2019). Harvested recaptures only.	40
8b	Input recapture matrix for IRCR analysis: from striped bass (>710 mm TL) that were tagged released in the springs of 1990-2019 (Rappahannock River only 1990-2017, all Virginia waters 2018-2019). Recaptures released with streamers cut off only. ...	40

List of Figures

I. Spawning Stock Monitoring

1	Locations of gill nets sampled in spring spawning stock assessments of striped bass in the Rappahannock and James Rivers, spring 2020.....	16
2	Number of striped bass captured and water temperature by week in the Rappahannock and James Rivers, spring 2020.	17
3	Catch by week and sex in the Rappahannock and James rivers, spring 2020.	17
4a	Otolith age frequencies by river, spring 2020 females.	18
4b	Otolith age frequencies by river, spring 2020 males.	18
5	Weekly total biomass, by sex, of striped bass captured in the Rappahannock and James Rivers, spring 2020.	19
6	Comparison of catch rates, in biomass and numbers, between two nets in each river, 2018 through 2020.	20
7	Average length (mm, a), weight (g, b) and age of striped bass by year, river and sex, 1990 - 2019.	21
8	Length frequencies of striped bass captured in gill nets, spring 2020.	22
9	Average length (mm, a) and weight (g, b), by sex of striped bass by year class, spring 2020, Rappahannock and James Rivers combined.	23
10	Average catch per day by river and mesh size, in biomass and numbers, spring 2020.	24
11	Average catch per day by river, sex and mesh size, in biomass, spring 2020.	25
12	Comparison of otolith age to scale age for samples derived from the same specimen, spring 2020.	26

II. Tagging

1a	IRCR generated estimates of annual survival (S) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.	41
1b	IRCR generated estimates of annual natural mortality (M) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.	41
1c	IRCR generated estimates of annual fishing mortality (F) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.	42
1d	IRCR generated estimates of annual tagging mortality (F-tag) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.	42
2a	IRCR generated estimates of annual survival (S) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.	43
2b	IRCR generated estimates of annual natural mortality (M) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.	43
2c	IRCR generated estimates of annual fishing mortality (F) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.	44
2d	IRCR generated estimates of annual tagging mortality (F-tag) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.	44

Spawning Stock Monitoring

Spring 2020

Introduction

Every year, striped bass migrate along the continental shelf waters of the US east coast and then enter brackish or fresh water to spawn. Historically, the principal spawning areas in the northeastern US have been the Hudson, Delaware and Chesapeake estuarine systems (Hardy 1998). The importance of the Chesapeake Bay spawning grounds to maintaining the productivity of these stocks has long been recognized (Merriman 1941, Raney 1952). In the Virginia tributaries of Chesapeake Bay, peak spawning activity is usually observed in April and is associated with rapidly rising water temperatures in the range of 13-19° C (Grant and Olney 1991). Spawning is often completed by mid-May, but may continue until June (Chapoton and Sykes 1961). Spawning grounds have been associated with rock-strewn coastal rivers characterized by rapids and strong currents on the Roanoke and the Susquehanna Rivers (Pearson 1938). In Virginia, spawning occurs over the upper 40 km of the tidal freshwater portions of the James, Rappahannock, Pamunkey and Mattaponi rivers (Grant and Olney 1991; Olney et al. 1991; McGovern and Olney 1996).

The Atlantic States Marine Fisheries Commission (ASMFC) declared that the Chesapeake Bay spawning stocks were fully recovered in 1995 after a period of very low stock abundance in the 1980's. This statement of recovered status was based on estimated levels of spawning stock biomass that were found in 1995 to be equal or greater than the average levels of the 1960-72 period (Rugulo et al. 1994). Thus, continued assessment of spawning stock abundance is an important component of ASMFC mandated monitoring programs. To this end, the Virginia Institute of Marine Science (VIMS) began development of spawning indexes that depict annual changes in catch rates of striped bass on the spawning grounds of the James and Rappahannock Rivers. These rivers represent the major contributors to the Chesapeake Bay stocks that originate from Virginia waters.

Materials and Methods

Samples of striped bass for biological characterization of the spring spawning stocks were obtained from the Rappahannock and James Rivers between 18 February and 31 March, 2020 by the VIMS Multispecies Research Group (MRG). Sampling had to be curtailed approximately six weeks early and prior to the anticipated maximum spawning period due to physical distancing requirements imposed in response to the COVID-19 virus. MRG strategized extensively on how to continue sampling while assuring the safety of personnel but could not guarantee that required distancing could be maintained under all sampling scenarios. Note that restrictions have since been modified such that sampling is expected to resume in 2021.

All samples were obtained using two 300-foot multi-panel anchor gill nets consisting of 10 panels each set once per week in approximately the same locations each time. Each panel was 30ft. (18.28m) in length and 10ft. (3.05m) in depth. The ten stretched mesh sizes (in inches) are 3, 3.75, 4.5, 5.25, 6, 6.5, 7, 8, 9, and 10. These mesh sizes correspond to those used by the Maryland Department of Natural Resources spawning stock monitoring program. The order of the panels in each net was determined randomly prior to net construction. Two nets were set in

close proximity (~1/4 mile apart) for each sampling event. The relative locations of the two nets were randomly assigned each week. The gear was set by MRG scientists and retrieved approximately 24 hours later. All specimens were brought back to VIMS for processing (lengths, weight, sex and maturity, with scales, otoliths and spleens preserved for later analysis) and disposal. The fishing locations were within the striped bass spawning areas as defined by VMRC (Figure 1).

Striped bass collected from the monitoring sites were measured on an MRG-made “Ichthystick” electronic fish measuring board and weighed using a Mettler PM 30000-K electronic balance. The board records lengths (FL and TL) to the nearest 1mm, and the scale to the nearest 10g. Both are integrated into the FEED (Fisheries Environment for Electronic Data) software system which allows manual input of sex and gonad maturity into a data base file for subsequent analysis. Scales were collected from between the spinous and soft dorsal fins above the lateral line for subsequent aging, using the method established by Merriman (1941), except that impressions made in acetate sheets replaced the glass slide and acetone. Otoliths were extracted from the striped bass, processed for aging, and compared to scale-derived ages.

Otoliths were cleansed of external tissue material by successive rinses in water immediately after extraction. The right sagitta were prepared for ageing by being placed on melted crystal bond and sectioned to a one millimeter thickness on a Buehler Isomet saw. These transverse sections were then polished using 320-400 grit wet sandpaper. The polished sections were then covered over with a thin layer of crystal bond. The sections are read using a Motic stereo dissecting microscope under 25x magnification.

Each otolith was aged by each of three readers using the methods described by Wischniowski and Bobko (1998). All readable scales from the monitoring specimens were aged by a single reader then a subsample was verified by a second readers. For otoliths, final ages assigned if at least two readers agreed. In cases in which all three readers disagreed the structure was re-analyzed by each reader and if agreement was still not found then the readers would conference together until consensus was reached. As scales are considered a secondary source for this part of the program, ages were assigned by just a single reader, with a selection of samples then also read by a senior scientist. Agreement between the two readers was greater than 80%. The annual birthdate is assumed to be 1 January of each year.

River-specific spawning stock catch-per-unit-effort (CPUE) indices were calculated for all mature specimens captured and for several subgroups of fish (e.g. by sex). The unit of effort used was a standardized 24-hour set (the actual number of hours fished divided by 24). Data from the two nets in each location were treated as independent samples. The assumption of independence will be tested as more data are collected in future years. As each net contained the same selection of mesh sizes and equal panel dimensions, the net measurements were dropped from the calculations. This may also have to be amended in future years if individual nets or net panels are rendered inoperable during a sampling day. Careful interpretation of 2020 data is recommended due to the shortened sampling season.

Results

Catch Summary

Striped bass (n= 837) were collected from gill nets in the Rappahannock River while 247 fish were sampled from the James between 18 February and 31 March, 2020. Considering the shortened sampling season, the numbers caught in the both rivers are larger than might be expected compared to 2019 (Rapp: n=379, James: n=77) though the numbers captured in 2019 were very low compared to historical averages.

In the Rappahannock, total daily catches varied from 12-429 striped bass, with the peak catch on 31 March, the last day of sampling. In the James, daily catches were between 8 and 87 fish with the maximum falling on 12 March. Surface water temperatures in the James hovered between 8-9°C for the first three weeks of sampling (through 1 March) then rose quickly to 12.4°C by 12 March, rose again to about 14°C by 18 March and jumped again to 16°C by the termination of sampling on 31 March. Water temperatures in the Rappahannock followed a very similar pattern but were generally 1°C-2°C cooler than those in the James. Typically, peak spawning in the Rappahannock occurs once the temperature rises above 16°C so it is likely that sampling was curtailed prior to peak abundance. In both rivers, catch rates started rising during the week of 1 March while surface temperature was still near its low values in previous weeks. Due to the curtailed sampling it is difficult to determine if catch rates were significantly related to water temperatures (Table 1, Figure 2). Salinities in both rivers were very low (0.03 – 0.45) during the entire sampling period though beginning on 12 March salinity in the Rappahannock was somewhat higher than that in the James (Table 1, Figure 2).

Males dominated the catch in both rivers (Rapp-826M:9F, James-221M:25F) though the proportions varied between rivers (Rapp - 98.9%, James – 89.8% / Table 2, Figure 3). However, sampling was curtailed prior to the times at which abundance of female striped bass in both rivers would have peaked.

In the Rappahannock, males ranged in age from 2 to 9, the same age range as in 2019. Rappahannock females were between ages 2 to 17. Males captured in the Rappahannock were predominantly ages 2-5 (2015-2018 year classes) with the 2015 (age-5) fish being predominant followed by fish from the 2018 year class (age-2). In the James the age range of male fish was the same as in the Rappahannock (ages 2 – 9). Similarly to 2019, males 3 to 5 (2015-2017 yc) were the most abundant in the James.

Females in the Rappahannock ranged between 1 and 17 years (2003-2018 yc) with fish from the 2011 yc (age-9) being somewhat more abundant than others. James females ranged between age-2 and age-23 (1997 – 2018 yc). As stated previously, sampling was curtailed prior to the weeks when female abundance would be expected to reach its peaks (Table 3, Figures 4a,4b).

Biomass catch rate (kg/set) followed a similar weekly pattern as catch in numbers, beginning at low values in February and beginning to climb by mid-March. In the Rappahannock abundance, especially of males, rose swiftly though with one lower value. Biomass in the James basically plateaued for the last four weeks of sampling prior to that

sampling ending prematurely (Table 4, Figure 5).

The ratio of males:females varied weekly, between 4.3:1 – 12.1:1 in the Rappahannock and between 0.7:1 – 3.6:1 in the James, excluding several weeks in which no females were captured. Over the entire season the sex ratio in the Rappahannock was 9.6:1 and in the James was 1.8:1 during an abbreviated sampling season (Table 4).

For all three years of the current monitoring protocol (2018 through 2020), the upstream or downstream position of the two nets fished at each location did not appear to have an effect on the catch rate, though sample sizes are still relatively small (Figure 6). Over that three-year period in the Rappahannock, the average catch of whichever net was set in the downstream position was 50.8 kg (35.7 fish) compared to the upstream net which averaged 66.0 kg (45.4 fish). In the James however, the downstream net averaged slightly more biomass and more fish (22.3 kg / 10.3 fish) than the upstream net (35.7 kg / 45.4 fish). This pattern has held true for each year of the current study. A paired t-test with all data combined found no significant difference due to the relative net position ($p = 0.69$ – biomass / $p = 0.65$ – count / $df = 54$). Similarly, considering only those sets in the Rappahannock river a t-test found no difference between upper and lower net position either in count ($p = 0.45$ – biomass / $p = 0.57$ – count / $df = 27$) as well as in the James ($p = 0.12$ – biomass / $p = 0.25$ – Count / $df = 26$).

In the Rappahannock, 826 male striped bass sampled in 2020 averaged 458.6mm (TL), 1.229kg and 4.3 years which were all similar to the respective values from 2019 (475.5 mm, 1.218 kg, 4.4 yr). Rappahannock females (9 specimens) had a mean length of 930.0mm, a mean weight of 11.689kg and on average were 10.3 years old which generally were moderate decreases compared to the spring of 2019. In the James River, 221 male specimens averaged 475.2mm in length, 1.555kg and 4.0 years (somewhat larger but younger than 2019 fish) while averages for the 25 sampled females were 748.6mm, 7.580kg and 8.6 years (both larger and older than averaged fish from 2019 – Table 5, Figure 7).

Rappahannock male fish displayed a peak in abundance between 300-380 mm with a much more abundant group at 390-600 mm and a scattering of individuals up to 870mm total lengths. With the truncated sampling season there were just 8 Rappahannock females with two fish smaller than 570mm and the rest between 960mm and 1070mm total length. While fewer in number and with a narrower size range, males in the James River showed peak abundance somewhat smaller than their Rappahannock counterparts with the largest cohort ranging between 400mm and 560mm. Females in the James were actually more abundant in the abbreviated season with a broad size group ranging between 450 to 640mm TL and other between 980 and 1220mm TL (Table 6, Figure 8). Within each year class, females on average were slightly larger and heavier than males (Table 7, Figure 9).

Spawning Stock Biomass Indexes

The overall (all data pooled) mean biomass index for the Rappahannock and James rivers were 92.2 kg/day (confidence interval (CI) = 18.8-165.6) and 44.1 kg/day (CI = 2372-64.4) respectively, representing 68.2 (Rappahannock, CI = 11.6-124.8) and 20.2 (James, CI = 11.8-28.6) fish per day on average. For females the average catch was 27.4 kg/day (CI = 0.0 – 64.8) in the Rappahannock and 18.9 (CI = 6.8-31.0) in the James. The average daily catch for

males Rappahannock was 72.7 kg/day (CI = 22.9-122.5) and in the James was 25.1 kg/day (CI = 13.5 – 36.7 / Table 8). Again, comparing these estimates with those from other years must be done with extreme caution due to the pandemic-truncated sampling season.

In both rivers, most fish were captured in the smaller mesh panels ($\leq 4.5''$) and in the largest meshes (9'' and 10'' - Table 9, Figure 10). Female fish in the Rappahannock were primarily captured in the largest mesh panels (8'', 9'', 10'') while males were captured across multiple mesh sizes with peaks at the smallest and largest meshes. In the James, the pattern of catch rates across mesh sizes was similar for both sexes with maximum captures between 3.75'' to 6'' and at the larger mesh panels, 8'', 9'', 10'' (Table 10, Figure 11).

Age Determinations using Scales and Otoliths

With one exception, each of the 1082 striped bass captured in the gill net monitoring project were subjected to age determination via otolith extraction and of these 651 were also aged using scales. In past years scale-otolith comparison were performed on virtually all fish. MRG determined that after many years of scale-otolith comparison data the relationship was well established and the paired sample rate could be decreased. Still, roughly 60% of the fish were aged via both methods. Compared to otolith ages (considered to be the more accurate and preferred method), between ages 2 and 10, mean scale ages were very close with relatively narrow ranges at each otolith age class. With the shorted sampling season very few fish older than age-9 were captured but as has been demonstrated many times, scale ages at the older age classes tend underestimated compared to the otolith age (Table 11, Figure 12).

Literature Cited

- Berggren, T.J. and J.T. Lieberman. 1978. Relative contribution of Hudson, Chesapeake and Roanoke striped bass, *Morone saxatilis*, stocks to the Atlantic coast fishery. U. S. Fish. Bull. 76(2): 335-345.
- Chapoton, R.B. and J.E. Sykes. 1961. Atlantic coast migration of large striped bass as evidenced by fisheries and tagging. Trans. Amer. Fish. Soc. 90(1):13-20.
- Dorazio, R.M., K.A. Hattala, C.B. McCollough and J.E. Skjeveland. 1994. Tag recovery estimates of migration of striped bass from spawning areas of the Chesapeake Bay. Trans. Amer. Fish. Soc. 123: 950-963.
- Field, J.D. 1997. Atlantic striped bass management: where did we go right? Fisheries 22(7): 6-8.
- Grant, G.C. and J.E. Olney. 1991. Distribution of striped bass *Morone saxatilis* (Walbaum) eggs and larvae in major Virginia rivers. U. S. Fish. Bull. 89:187-193.
- Hardy, J.D. Jr. 1978. Development of fishes of the mid-Atlantic bight. Vol. III, Aphrederidae through Rachycentridae. U. S. Fish Wildl. Serv. FWS/OBS-78/12.
- Lewis, R.M. 1957. Comparative study of populations of the striped bass. U. S. Fish and Wildlife Service Spec. Rep. Fisheries 204:1-54.
- McGovern, J.C. and J.E. Olney. 1996. Factors affecting survival of early life stages and subsequent recruitment of striped bass on the Pamunkey River, Virginia. Can. J. Fish. Aquat. Sci. 53: 1713-1726.
- Merriman, D. 1941. Studies on the striped bass (*Roccus saxatilis*) of the Atlantic Coast. Fish. Bull. U.S. Fish Wildl. Serv. 50(35):1-77.
- Olney, J.E., J.D. Field, and J.C. McGovern. 1991. Striped bass egg mortality, production and female biomass in Virginia rivers, 1980-1989. Trans. Amer. Fish. Soc. 120: 354-367.
- Pearson, J.C. 1938. The life history of the striped bass, or rockfish, *Roccus saxatilis* (Walbaum). U. S. Fish. Bull. 49: 825-851.
- Raney, E.C. 1957. Subpopulations of the striped bass *Roccus saxatilis* (Walbaum), in tributaries in Chesapeake Bay. U. S. Fish Wildl. Serv., Spec. Sci. Fish. 208: 85-107.

Literature Cited (cont.)

- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Fish. Res. Bd. Can. Bull. 191: 382 p.
- Rugolo, L.J., P.W. Jones, R.K. Schaefer, K.S. Knotts, H.T. Hornick and J.L. Markham. 1994. Estimation of Chesapeake Bay-wide exploitation rate and population abundance for the 1993 striped bass stock. Manuscript, Maryland Department of Natural Resources, Annapolis, Md.
- Secor, D.H. 1999. Specifying divergent migrations in the concept of stock: the contingent hypothesis. Fisheries research 43: 13-34.
- Secor, D.H., T.M. Trice and H.T. Hornick. 1995. Validation of otolith-based ageing and a comparison of otolith and scale-based ageing in mark-recaptured Chesapeake Bay striped bass, *Morone saxatilis*. Fish. Bull. 93:186-190.
- Van Winkle, W., K.D. Kumar, and D.S. Vaughan. 1988. Relative contributions of Hudson River and Chesapeake Bay striped bass stocks to the Atlantic Coast population. Amer. Fish. Soc. Mono. 4: 255-266.
- Wischniowski, W. and S. Bobko. 1998. Age and growth laboratory manual. Final report Old Dominion Univ. Center for Quantitative Fisheries Ecology.

Table 1. Number and biomass of striped bass captured, water temperature, and salinity by week in the Rappahannock and James Rivers, spring 2020.

River	Week-of-Year	Sample Date	Number Caught	Biomass Caught (kg)	Water Temp (C)	Salinity (ppt)
Rappahannock	8	18-Feb-20	12	14.38	8.1	0.03
	9	25-Feb-20	26	24.79	7.9	0.03
	10	01-Mar-20	12	14.2	7.9	0.03
	11	12-Mar-20	40	58.72	10.7	0.19
	12	19-Mar-20	278	351.92	12.5	0.23
	13	26-Mar-20	38	69.3	13.2	0.45
	14	31-Mar-20	429	587.09	14.7	0.17
James	8	18-Feb-20	10	14.8	9.0	0.05
	9	25-Feb-20	8	9.3	8.7	0.05
	10	01-Mar-20	9	13.31	7.4	0.06
	11	12-Mar-20	87	174.33	12.4	0.08
	12	18-Mar-20	43	81.02	13.9	0.09
	13	26-Mar-20	48	88.93	13.9	0.09
	14	31-Mar-20	42	153.14	15.9	0.08

Table 2. Catch rates by week and sex for the Rappahannock and James Rivers, spring 2020.

River	Week-of-Year	Sample Date	Males	Females	Percent Males
Rappahannock	8	18-Feb-20	12	0	100%
	9	01-Mar-20	12	0	100%
	10	25-Feb-20	24	2	92%
	11	12-Mar-20	40	0	100%
	12	19-Mar-20	278	0	100%
	13	26-Mar-20	37	1	97%
	14	31-Mar-20	423	6	99%
Total			826	9	99%
James	8	18-Feb-20	8	2	80%
	9	01-Mar-20	6	2	75%
	10	25-Feb-20	8	0	100%
	11	12-Mar-20	81	6	93%
	12	18-Mar-20	38	5	88%
	13	26-Mar-20	45	3	94%
	14	31-Mar-20	35	7	83%
Total			221	25	90%

Table 3. Otolith age frequencies by river and sex, spring 2020.

Otolith Age	Rappahannock		James	
	Females	Males	Females	Males
1	0	0	0	0
2	1	142	1	13
3	0	93	3	84
4	0	51	3	54
5	0	490	2	37
6	1	36	4	25
7	0	7	1	5
8	0	4	1	1
9	3	3	3	2
10	1	0	1	0
11	0	0	0	0
12	0	0	0	0
13	0	0	0	0
14	0	0	0	0
15	1	0	1	0
16	1	0	1	0
17	1	0	3	0
18	0	0	0	0
19	0	0	0	0
20	0	0	0	0
21	0	0	0	0
22	0	0	0	0
23	0	0	1	0
24	0	0	0	0
25	0	0	0	0

Table 4. Weekly total biomass, by sex, of striped bass captured in the Rappahannock and James Rivers, spring 2020.

River	Week-of-Year	Sample Date	Males	Females	Ratio Males:Females
Rappahannock	8	18-Feb-20	14.38	0.00	n/a
	9	01-Mar-20	14.2	0.00	n/a
	10	25-Feb-20	22.89	1.90	12:1
	11	12-Mar-20	58.72	0.00	n/a
	12	19-Mar-20	351.92	0.00	n/a
	13	26-Mar-20	56.21	13.09	4.3:1
	14	31-Mar-20	496.88	90.21	5.5:1
	Overall		1015.2	105.20	9.7:1
James	8	18-Feb-20	10.99	3.81	2.9:1
	9	01-Mar-20	8.37	3.25	2.6:1
	10	25-Feb-20	9.3	0.00	n/a
	11	12-Mar-20	127.11	47.22	2.7:1
	12	18-Mar-20	55.47	25.55	2.2:1
	13	26-Mar-20	69.56	19.37	3.6:1
	14	31-Mar-20	62.83	90.31	0.7:1
	Overall		343.63	189.51	1.8:1

Table 5. Average length (mm), weight (g) and age of striped bass by year, river and sex, spring 2020. Double lines indicate changes in sampling methods and/or locations.

Year	Rappahannock River								James River								Overall							
	Males				Females				Males				Females				Males				Females			
	n	Mean TL (mm)	Mean Wt (g)	Mean Age	n	Mean TL (mm)	Mean Wt (g)	Mean Age	n	Mean TL (mm)	Mean Wt (g)	Mean Age	n	Mean TL (mm)	Mean Wt (g)	Mean Age	n	Mean TL (mm)	Mean Wt (g)	Mean Age	n	Mean TL (mm)	Mean Wt (g)	Mean Age
1990	595	476.5	1352	4.3	176	528.1	2330	5.1	15	493.9	1595		27	549.9	2157		625	478.7	1373	4.3	210	532.2	2297	5.1
1991	1549	524.5	1695	3.6	569	549.6	2059	3.7	0		1070		1	846.0	3404		1687	527.3	1728	3.6	602	550.0	2058	3.7
1992	694	521.8	1620	3.6	332	595.8	2865	4.8									709	523.0	1639	3.6	341	600.8	2944	4.8
1993	1229	515.8	1762	3.8	561	610.4	3425	4.9									1229	515.8	1762	3.8	561	610.4	3425	4.9
1994	936	536.7	1864	4.0	342	702.9	4749	6.0	171	537.1	1868	4.2	66	709.7	4822	6.4	1107	536.8	1864	4.0	408	704.0	4760	6.1
1995	1327	445.3	1224	3.3	405	589.6	3583	5.1	1143	512.7	1743	3.9	423	687.4	4600	6.1	2470	476.5	1464	3.6	828	639.6	4102	5.6
1996	647	467.3	1221	3.5	136	578.6	2715	4.8									647	467.3	1221	3.5	136	578.6	2715	4.8
1997	522	482.1	1401	3.9	133	862.3	8408	9.1									522	482.1	1401	3.9	133	862.3	8408	9.1
1998	697	471.6	1192	3.7	119	897.0	8498	9.3									697	471.6	1192	3.7	119	897.0	8498	9.3
1999	1103	447.3	1029	3.5	51	825.6	7186	8.6									1103	447.3	1029	3.5	51	825.6	7186	8.6
2000	1937	471.0	1113	2.7	80	853.5	7408	7.7									1937	471.0	1113	2.7	80	853.5	7408	7.7
2001	472	511.9	1571	3.3	105	875.4	7339	8.1									472	511.9	1571	3.3	105	875.4	7339	8.1
2002	111	534.3	1938	3.5	41	781.2	5833	6.5									111	534.3	1938	3.5	41	781.2	5833	6.5
2003	283	585.0	2413	4.2	186	893.2	8306	8.5									283	585.0	2413	4.2	186	893.2	8306	8.5
2004	631	585.8	2529	4.3	236	877.8	8014	8.4									631	585.8	2529	4.3	236	877.8	8014	8.4
2005	446	550.7	2104	3.9	171	875.3	7996	8.6									446	550.7	2104	3.9	171	875.3	7996	8.6
2006	623	479.8	1370	3.5	119	828.0	6876	5.6									623	479.8	1370	3.5	119	828.0	6876	5.6
2007	748	561.5	2233	5.0	356	896.9	8427	10.5									748	561.5	2233	5.0	356	896.9	8427	10.5
2008	413	504.5	1594	4.3	74	801.7	6632	8.6									413	504.5	1594	4.3	74	801.7	6632	8.6
2009	437	573.5	1821	5.1	183	786.0	5037	8.3									437	573.5	1821	5.1	183	786.0	5037	8.3
2010	828	568.4	1040	5.2	219	871.3	5481	10.1									828	568.4	1040	5.2	219	871.3	5481	10.1
2011	131	625.9	1140	6.1	84	851.3	7123	9.5									131	625.9	1140	6.1	84	851.3	7123	9.5
2012	321	577.1	2390	5.5	117	859.5	8405	9.8									321	577.1	2390	5.5	117	859.5	8405	9.8
2013	152	556.6	745	5.3	94	855.4	5514	10.1									152	556.6	745	5.3	94	855.4	5514	10.1
2014	126	507.9	1958	4.8	95	925.0	9910	11.1									126	507.9	1958	4.8	95	925.0	9910	11.1
2015	108	508.9	1565	4.5	44	917.2	9795	11.4									108	508.9	1565	4.5	44	917.2	9795	11.4
2016	305	480.3	1260	4.1	57	906.3	9830	10.8									305	480.3	1260	4.1	57	906.3	9830	10.8
2017	204	453.8	1138	4.0	17	746.6	6475	8.4									204	453.8	1138	4.0	17	746.6	6475	8.4
2018	916	434.1	986	3.9	25	912.0	10997	10.4	179	483.7	1347	3.9	36	669.7	4664	5.2	1095	442.2	1045	3.9	61	769.0	7259	7.3
2019	356	471.5	1218	4.4	23	926.9	14110	11.7	62	455.6	1205	4.2	15	725.7	6252	7.2	418	469.1	1216	4.4	38	847.4	11008	9.9
2020	826	458.6	1229	4.3	9	930.0	11689	10.3	221	475.2	1555	4.0	25	748.6	7580	8.6	1047	462.1	1298	4.2	34	796.6	8668	9.0

Table 6. Length frequencies (mm TL) of striped bass sampled from the gill nets, spring 2020.

Length	Rapp- males	Rapp- females	James- males	James- females	Length	Rapp- males	Rapp- females	James- males	James- females
300	8	0	1	0	800	0	0	0	0
310	18	0	0	0	810	0	0	0	0
320	35	1	2	0	820	1	0	0	0
330	27	0	0	0	830	1	0	0	0
340	27	0	1	1	840	0	0	0	0
350	17	0	4	0	850	0	0	0	0
360	19	0	1	0	860	0	0	0	0
370	8	0	1	0	870	1	0	0	0
380	14	0	4	0	880	0	0	0	0
390	10	0	2	0	890	0	0	0	0
400	14	0	3	0	900	0	0	0	0
410	20	0	9	0	910	0	0	0	0
420	33	0	14	0	920	0	0	0	0
430	30	0	5	0	930	0	0	0	0
440	32	0	16	0	940	0	0	0	0
450	28	0	11	1	950	0	0	0	0
460	40	0	29	0	960	0	1	0	0
470	38	0	10	0	970	0	1	0	0
480	48	0	13	2	980	0	1	0	2
490	39	0	15	0	990	0	0	0	0
500	68	0	20	1	1000	0	0	0	0
510	45	0	15	1	1010	0	0	0	0
520	65	0	9	0	1020	0	0	0	0
530	33	0	7	0	1030	0	1	0	0
540	32	0	5	1	1040	0	0	0	0
550	21	0	6	0	1050	0	0	0	0
560	13	0	2	0	1060	0	0	0	0
570	11	1	4	1	1070	0	2	0	1
580	8	0	4	2	1080	0	0	0	0
590	7	0	3	0	1090	0	0	0	0
600	3	0	1	0	1100	0	0	0	0
610	0	0	0	0	1110	0	0	0	0
620	1	0	0	1	1120	0	0	0	0
630	0	0	1	1	1130	0	0	0	1
640	2	0	0	2	1140	0	0	0	0
650	0	0	0	0	1150	0	0	0	0
660	1	0	1	0	1160	0	0	0	1
670	1	0	1	0	1170	0	0	0	0
680	2	0	1	0	1180	0	0	0	0
690	1	0	0	0	1190	0	0	0	1
700	1	0	0	1	1200	0	0	0	1
710	0	0	0	0	1210	0	0	0	0
720	0	0	0	0	1220	0	0	0	1
730	0	0	0	0	1230	0	0	0	0
740	1	0	0	1	1240	0	0	0	0
750	0	0	0	0	1250	0	0	0	0
760	0	0	0	0	1260	0	0	0	0
770	0	0	0	0	1270	0	0	0	0
780	0	0	0	1	1280	0	0	0	0
790	0	0	0	0	1290	0	0	0	0
					1300	0	0	0	0
					Total	824	8	221	25

Table 7. Average length (mm) and weight (g), with standard deviations (Std Dev) of striped bass by year class, spring 2020.

Year Class	Females					Males				
	n	Mean TL (mm)	Std Dev	Mean Wt. (g)	Std Dev	n	Mean TL (mm)	Std Dev	Mean Wt. (g)	Std Dev
2018	2	329.0	8.5	415	49.5	155	336.9	27.4	429	126.2
2017	3	470.3	15.9	1,400	168.2	177	419.2	47.5	945	380.1
2016	3	533.0	43.4	2,100	437.1	105	479.8	39.7	1,444	413.0
2015	2	581.0	56.6	2,845	1,011.2	527	494.7	42.7	1,469	457.8
2014	5	598.4	38.7	2,854	1,026.5	61	527.5	51.0	1,977	678.8
2013	1	627.0		2,550		12	605.8	93.0	2,929	1,443.1
2012	1	984.0		9,600		5	721.0	119.1	4,886	2,133.7
2011	6	856.3	130.2	8,250	3,505.6	5	654.8	69.2	3,784	1,309.2
2010	2	1,002.0	36.8	12,295	1,124.3					
2005	2	1,071.5	2.1	16,060	226.3					
2004	2	1,306.0	121.6	19,715	6,385.2					
2003	4	1,155.8	59.3	19,213	4,105.9					
2002	1	1,133.0		28,780						
2001										
2000										
1999										
1998										
1997	1	1,133.0		28,780						

Table 8. Average catch per day by river and sex, in numbers and biomass with lower (LCL) and upper (UCL) confidence limits, spring 2020.

River	Sex	LCL	Number		LCL	KG per day	
			per Day	UCL		UCL	UCL
Rapp	Combined	11.6	68.2	124.8	18.8	92.2	165.6
	F	0.3	2.3	4.3	0.0	27.4	64.8
	M	17.0	58.0	99.1	22.9	72.7	122.5
James	Combined	11.8	20.2	28.6	23.7	44.1	64.4
	F	1.6	2.4	3.2	6.8	18.9	31.0
	M	9.0	16.2	23.4	13.5	25.1	36.7

Table 9. Average catch per day by river and mesh size, in numbers and biomass with lower (LCL) and upper (UCL) confidence limits, spring 2020.

River	Mesh (in)	Number			KG per Day		
		LCL	per Day	UCL	LCL	Day	UCL
Rapp	3.00	0.00	14.14	30.71	0.00	7.68	16.82
	3.75	0.64	21.39	42.14	0.60	24.31	48.03
	4.50	0.48	18.29	36.11	0.11	26.93	53.75
	5.25	0.95	4.97	8.99	1.86	9.36	16.85
	6.00	0.14	0.87	1.59	0.04	2.35	4.67
	6.50	0.08	0.42	0.75	0.17	1.08	1.99
	7.00	0.00	0.28	0.59	0.00	1.18	2.69
	8.00	0.00	0.15	0.35	0.00	1.06	2.50
	9.00	0.00	0.37	0.84	0.00	4.71	10.82
	10.00	0.00	0.15	0.45	0.00	2.99	8.96
James	3.00	0.27	1.15	2.02	0.15	0.93	1.71
	3.75	1.41	3.24	5.07	1.09	3.74	6.40
	4.50	3.36	6.98	10.59	5.02	10.17	15.33
	5.25	3.30	5.28	7.26	6.01	9.71	13.42
	6.00	0.29	0.86	1.43	0.08	3.32	6.55
	6.50	0.09	0.49	0.90	0.41	2.02	3.62
	7.00	0.00	0.07	0.21	0.00	0.25	0.76
	8.00	0.00	0.14	0.32	0.00	1.73	4.16
	9.00	0.00	0.07	0.21	0.00	2.03	6.09
	10.00	0.00	0.29	0.63	0.00	5.56	12.09

Table 10. Average catch per day by river, sex and mesh size, in biomass with lower (LCL) and upper (UCL) confidence limits, spring 2020.

River	Sex	Mesh (in)	LCL	Number per Day	UCL	LCL	KG per Day	UCL
Rapp	Female	3.00	0.00	1.45	3.72	0.00	0.88	2.39
		3.75	0.00	2.41	5.23	0.81	2.25	3.68
		4.50	0.00	4.04	8.85	0.00	4.82	10.38
		5.25	0.00	1.24	2.87	0.00	2.11	4.76
		6.00	0.00	0.14	0.41	0.00	0.37	1.10
		6.50	0.00	0.10	0.29	0.00	0.16	0.47
		7.00	0.00	0.00	0.00	0.00	0.00	0.00
		8.00	0.00	0.00	0.00	0.00	0.00	0.00
		9.00	0.00	0.37	0.84	0.00	4.71	10.82
		10.00	0.00	0.15	0.45	0.00	2.99	8.96
	Male	3.00	2.63	15.93	29.22	1.35	8.69	16.03
		3.75	7.20	22.63	38.05	8.14	25.76	43.37
		4.50	6.30	20.00	33.71	8.96	29.58	50.20
		5.25	2.34	5.42	8.50	4.50	10.23	15.95
		6.00	0.48	1.11	1.74	0.95	3.01	5.07
		6.50	0.28	0.59	0.91	0.71	1.59	2.48
		7.00	0.12	0.45	0.79	0.29	1.94	3.58
		8.00	0.03	0.26	0.49	0.19	1.85	3.51
		9.00	0.00	0.37	0.84	0.00	4.71	10.82
10.00		0.00	0.15	0.45	0.00	2.99	8.96	
James	Female	3.00	0.00	0.30	0.68	0.00	0.18	0.42
		3.75	0.00	0.69	1.38	0.00	0.50	1.05
		4.50	0.34	1.02	1.70	0.55	1.50	2.45
		5.25	0.43	1.53	2.62	1.05	2.80	4.55
		6.00	0.09	0.71	1.34	0.00	3.40	7.37
		6.50	0.00	0.18	0.41	0.00	0.77	1.82
		7.00	0.00	0.07	0.21	0.00	0.25	0.76
		8.00	0.00	0.14	0.32	0.00	1.73	4.16
		9.00	0.00	0.07	0.21	0.00	2.03	6.09
		10.00	0.00	0.29	0.63	0.00	5.56	12.09
	Males	3.00	0.66	1.43	2.19	0.49	1.17	1.86
		3.75	2.17	3.55	4.92	2.12	4.13	6.14
		4.50	4.44	7.44	10.44	6.53	10.79	15.06
		5.25	3.23	4.60	5.97	5.85	8.44	11.04
		6.00	0.50	0.71	0.93	1.25	1.96	2.67
		6.50	0.23	0.59	0.94	1.04	2.32	3.59
		7.00	0.00	0.07	0.21	0.00	0.25	0.76
		8.00	0.00	0.14	0.32	0.00	1.73	4.16
		9.00	0.00	0.07	0.21	0.00	2.03	6.09
10.00		0.00	0.29	0.63	0.00	5.56	12.09	

Table 11. Average, minimum and maximum scale ages for each otolith age class from ages derived from the same specimen, spring 2020.

Otolith Age	n	Minimum Scale Age	Mean Scale Age	Maximum Scale Age
2	75	2	2.05	3
3	131	3	3.12	5
4	81	2	4.02	5
5	292	3	4.85	6
6	45	4	5.78	8
7	9	5	7.00	10
8	4	7	7.75	8
9	5	9	9.00	9
10	2	9	9.50	10
11	0			
12	0			
13	0			
14	0			
15	1	12	12.00	12
16	1	10	10.00	10
17	3	10	12.67	14
18	0			
19	0			
20	0			
21	0			
22	0			
23	1	16	16.00	16
24				
25				

Figure 1. Locations of gill nets sampled in spring spawning stock assessments of striped bass in the Rappahannock and James Rivers, spring 2020.




Figure 2. Number of striped bass captured and water temperature by week in the Rappahannock and James Rivers, spring 2020.





Figure 3. Catch by week and sex in the Rappahannock and James rivers, spring 2020.



Rapp - Males	12	12	24	40	278	37	423											
Rapp - Females	0	0	2	0	0	1	6											
James - Males	8	6	8	81	38	45	35											
James - Females	2	2	0	6	5	3	7											

Figure 4a. Otolith age frequencies by river, spring 2020 females.




Figure 4b. Otolith age frequencies by river, spring 2020 males.




Figure 5. Weekly total biomass, by sex, of striped bass captured in the Rappahannock and James Rivers, spring 2020.




Figure 6. Comparison of catch rates, in biomass and numbers, between upstream and downstream paired nets in each river, 2018 through 2020.





Figure 7. Average length (mm, a), weight (g, b) and age of striped bass by year and sex, 1990 - 2020. Double lines indicate changes in sampling methods and/or locations.




Figure 8. Length frequencies of striped bass captured in gill nets, spring 2020.




Figure 9. Average length (mm) and weight (g), by sex of striped bass by year class, spring 2020, Rappahannock and James Rivers combined.





Figure 10. Average catch per day by river and mesh size, in biomass and numbers, spring 2020.




Figure 11. Average catch per day by river, sex and mesh size, in biomass, spring 2020.





Figure 12. Comparison of otolith age to scale age for samples derived from the same specimen, spring 2020. Boxes represent 25% and 75% quartiles, diamond symbols are the mean, horizontal lines are the median, circles are outliers, the solid black line is the 1:1 line.



Tagging
Spring, 2020

Introduction

The Striped Bass Program, a component of the Multispecies Research Group at the Virginia Institute of Marine Science (VIMS), has monitored the size and age composition, sex ratio and maturity schedules of the spawning striped bass stock in the Rappahannock River since 1981. In conjunction with the monitoring studies, VIMS established a tagging program in 1988 to provide information on the migration, relative contribution to the coastal population, and annual survival of striped bass that spawn in the Rappahannock River. This program is part of an active cooperative tagging study that currently involves 15 state and federal agencies along the Atlantic coast. The U.S. Fish and Wildlife Service manages the coast-wide tagging database. Hence, commercial and recreational anglers that target striped bass are encouraged to report all recovered tags to that agency.

Although the initial purpose of the coast-wide tagging study was to evaluate efforts to restore Atlantic striped bass stocks (Wooley *et al.* 1990), tagging data are now being collected to monitor striped bass mortality rates in a recovered fishery.

Multi-year Tagging Models

Tag return data are generally represented by constructing an upper triangular matrix of tag recoveries, where each cell of the matrix contains the number of tag returns from a particular year of tagging and recovery. For example, a study with I years of tagging and J years of recovery would yield the following data matrix

$$R = \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1J} \\ - & r_{22} & \cdots & r_{2J} \\ \vdots & \vdots & \ddots & \vdots \\ - & - & - & r_{IJ} \end{bmatrix}, \quad (1)$$

where r_{ij} is the number of tags recovered in year j that were released in year i (note, $J \geq I$). Tagging periods do not necessarily have to be yearly intervals; however, data analysis is easiest if all periods are the same length and all tagging events are conducted at the beginning of each period.

Application of tagging models involves constructing an upper triangular matrix of expected values and comparing them to the observed data. Since the recovery data over time for each year's batch of tagged fish can be assumed to follow a multinomial distribution, the method of maximum likelihood can be used to obtain parameter estimates. Analytical solutions for the maximum likelihood parameter estimates are generally not available. Hence, several software packages that numerically maximize a product multinomial likelihood function have been developed for application of tagging models. They include programs

SURVIV (White 1983) and MARK (White and Burnham 1999).

Seber models: White and Burnham (1999) reformulated the original Brownie et al. (1985) models in the way originally suggested by Seber (1970) to create a consistent framework for modeling mark-recapture data (Smith et al. 2000). This framework served as the foundation for program MARK, which is a comprehensive software package for the application of capture-recapture models. For time-specific parameterization of the Seber models, the matrix of expected values associated with equation (1) would be

$$E(R) = \begin{bmatrix} N_1(1-S_1)r_1 & N_1S_1(1-S_2)r_2 & \cdots & N_1S_1\cdots S_{J-1}(1-S_J)r_J \\ - & N_2(1-S_2)r_2 & \cdots & N_2S_2\cdots S_{J-1}(1-S_J)r_J \\ \vdots & \vdots & \ddots & \vdots \\ - & - & - & N_I(1-S_I)r_I \end{bmatrix} \quad (2)$$

Where N_i is the number tagged in year i , S_i is the survival rate in year i and r_i is the probability that a tag is recovered from a killed fish regardless of the source of mortality. For the 2006 estimates the updated version of MARK (version 4.3) replaced the version used in previous years (version 4.2).

The Seber models are simple and robust, but they do not yield direct information about exploitation (u) or instantaneous rates of fishing and natural mortality, which are often of interest to fisheries managers. Estimates of S can be converted to the instantaneous total mortality rate via the equation (Ricker 1975)

$$Z = -\log_e(S) \quad (3)$$

and, if information about the instantaneous natural mortality rate is available, estimates of the instantaneous fishing mortality can be recovered. Given estimates of the instantaneous rates, it is possible to recover estimates of u if the timing of the fishery (Type I or Type II) is known (Ricker 1975).

Instantaneous rate models: Hoenig et al. (1998a) modified the Brownie et al. (1985) models to allow for the estimation of instantaneous rates of fishing and natural mortality. This extension showed how information on fishing effort could be used as an auxiliary variable and also discussed generalizing the pattern of fishing within the year. The matrix of expected values corresponding to equation (1) for a model that assumes time-specific fishing mortality rates and a constant natural mortality rate would be

$$E(R) = \begin{bmatrix} N_1\phi\lambda u_1(F_1, M) & N_1\phi\lambda u_2(F_2, M)e^{-(F_1+M)} & \dots & N_1\phi\lambda u_J(F_J, M)e^{-\sum_{k=1}^{J-1} F_k+(J-1)M} \\ - & N_2\phi\lambda u_2(F_2, M) & \dots & N_2\phi\lambda u_J(F_J, M)e^{-\sum_{k=2}^{J-1} F_k+(J-2)M} \\ \vdots & \vdots & \ddots & \vdots \\ - & - & - & N_1\phi\lambda u_J(F_J, M) \end{bmatrix}$$

(4)

where ϕ is the probability of surviving being tagged and retaining the tag in the short-term, λ is the tag-reporting rate, and $u_k(F_k, M)$ is the exploitation rate in year k which, as mentioned above, depends on whether the fishery is Type I or Type II. For striped bass, a Type II (continuous) fishery is assumed. Note that ϕ and λ are considered constant over time.

Materials and Methods

Capture and Tagging Protocol

1991 – 2017

Each year in the Rappahannock River during the months of March, April and May, VIMS scientists obtained samples of mature striped bass on the spawning grounds of the Rappahannock River. Samples were taken twice-weekly from pound nets owned and operated by cooperating commercial fishermen. The pound net is a fixed trap that is presumed to be non-size selective in its catch of striped bass, and has been historically used by commercial fishermen in the Rappahannock River. These pound nets were located between river miles 45 – 56. All captured striped bass were removed from each pound net and placed into a floating holding pocket (1.2m x 2.4m x 1.2m deep, with 25.4mm mesh and a capacity of approximately 200 fish) anchored adjacent to the pound net. Fish were dip-netted from the holding pocket and examined for tagging.

In order to diversify the tagging locations of striped bass and to increase the number of fish tagged each year, in some years specimens from the James and York River systems were captured in multi-mesh gill nets, then tagged and released similarly as described above. Full descriptions of the gear and methods are described in earlier project reports.

2018 - present

In an effort to increase sampling efficiency and decrease costs, in 2018 MRG commenced capturing striped bass to be tagged using electrofishing gear rather than the pound nets and gill nets used in earlier years. In 2018 this was accomplished in cooperation with the Virginia Department of Game and Inland Fisheries (VDGIF) which possessed the requisite vessels, equipment and expertise and which regularly conducts such investigations at approximately the same locations and time of year. Subsequent to the 2018 tagging season, having demonstrated that this gear could be an effective method for this program in Virginia waters, MRG acquired its own specialized vessel and electrofishing rig, sent personnel to training, and in 2019 we performed all sampling using only VIMS equipment and personnel.

During most sampling events, all operations were performed on the single vessel described above. Trained VIMS personnel piloted the vessel and operated the apparatus while other VIMSbiologists would scoop specimens from the water using dip nets and perform the tagging operation described below. Depending upon the sampling schedule on any given day, during some tagging events the specimen processing could be done on a second, following vessel. Tagging was done at several locations in the Rappahannock River, in the James River main stem as well as in the James River tributaries as well as in the York River tributaries the Pamunkey River and the Mattaponi River.

Once onboard, fork length (FL) and total length (TL) measurements were taken and

whenever possible the sex of each fish was determined. Striped bass not previously marked and larger than 457 mm TL were tagged with sequentially numbered internal anchor tags (Floy Tag and Manufacturing, Inc.). Each internal anchor tag was applied through a small incision in the abdominal cavity of the fish. A small sample of scales from between the dorsal fins and above the lateral line on the left side was removed and used to estimate age. Each fish was released at the site of capture immediately after receiving a tag or after a short recovery period spent in an onboard holding tank which was supplied with fresh aerated water.

Analysis Protocol

For each striped bass assessment through 2016 several different approaches were used to analyze the tagging data. These were, the program MARK, the exploitation rate (R/M) method, the catch equation method, and the instantaneous rates method. Each is fully described in earlier project annual reports (e.g. Sadler, 2016).

For the 2018 Benchmark Assessment only the instantaneous rates method was used. This method allows the estimate of natural mortality to be constant, or to vary by periods and allows for varying fishing mortality under different regulatory periods as well changes in tagging mortality. Virginia data were included under 11 sets of test assumptions regarding changes in fishing mortality, tagging mortality, and natural mortality (Table 1).

All analytical approaches were applied to striped bass greater than 457 mm total length (minimum legal size) and to striped bass greater than 710 mm TL (coastal migrants). Coast wide model results and selection are published in the 2019 benchmark assessment (National Marine Fisheries Service, 2019). Model fit was evaluated using Akaike's Information Criterion (AOC) (Akaike 1973; Burnham and Anderson 1992), quasi-likelihood AIC (QAIC) (Akaike 1985), and goodness-of-fit (GOF) diagnostics are used to evaluate their fit (Burnham et al. 1995).

Results

Spring 2020 Tag Release summary

Electrofishing tagging events (24 of them) occurred between 5 March and 5 May 2020 in the Rappahannock (13), James (7), Mattaponi (4). During each event fishing occurred nearly continuously, generally in a grid pattern covering the location thoroughly. Between 0 (3 events in early March) and 192 fish were tagged on a given day.

A total of 1,023 fish were tagged and released, slightly exceeded the target of 1,000, compared to 859 fish in 2018 and just 119 in 2019. The median date of release for both rivers combined was 19 April 2019.

In the Rappahannock River, 11 electrofishing events were conducted between 5 March and 5 May, 2020 which resulted in 789 fish being tagged and released (Table 1). There were 746 resident striped bass (457-710 mm TL) tagged and released. Coastal migrant fish (>710mm) tagged totaled 46 specimens with the largest measured at 1,232mm.

In the James River, resident fish (457-710mm TL) accounted for 191 out of the total of 212 striped bass which were tagged and released between 17 March and 24 April, 2020 (6 sampling dates). An additional 21 coastal migrants (457-710 mm TL) were also tagged and released (Table 2).

In the Mattaponi River (a tributary to the York River), an additional 23 fish were tagged and released during three sampling events between 9 April and 1 May 2020 (Table 3). Of these 16 were residents (457-710mm TL) and 7 were coastal migrants (>710mm TL).

Mortality Estimates

Tag recapture summary: A total of 48 striped bass >457 mm TL were recaptured between 1 January and 31 December 2020. The largest source of recaptures (41 / 85.4%) was from Chesapeake Bay (20 / 41.7% in Virginia, and 21 / 43.8% in Maryland, Table 4). Other recaptures occurred in Rhode Island (2), New York (2) and New Jersey (3). The peak months for recaptures were May (10) and June (12) with relatively equal numbers (≤ 6) during each succeeding month. No recaptured fish were reported in January, February or April and just one tag was returned in March.

From the 48 total recoveries, 5 were migratory striped bass (>710 mm TL at time of tagging) recaptured between 1 January and 31 December, 2020 (Table 5). These fish were recaptured in Rhode Island (1), New York (1), New Jersey (2), and Virginia (1). Recapture events for the coastal migrants occurred in May (1), June (1), August (1) and December (2).

The small number of recaptured fish may be due to decreased recreational fishing effort caused by the combination of lower bag limits imposed by ASMFC and Covid-19 restrictions.

Instantaneous rates model estimates of survival, fishing and natural mortality

Consistent with the analyses performed as part of the 2019 benchmark assessment, eleven models with varying time periods (corresponding to past management actions) for estimates of fishing mortality (F) and tagging mortality (F_{tag}) were included. All models included two natural mortality periods 1990-1997 and 1998-2019 (Table 6). These models were not updated for management actions initiated in 2019.

Virginia releases: From the 849 resident striped bass (≥ 457 mm TL) tagged in spring 2018, 21 were recaptured and harvested in 2018 and 7 more in 2019. Five fish tagged in previous springs were harvested in 2018. Just 2 of the 101 fish tagged in 2019 have been harvested to date (Table 7a). There were 22 2018-tagged fish which were recaptured and re-released in 2018 and 6 more were captured in 2019. Two fish tagged in 2019 have been subsequently recaptured and returned to the water (Table 7b). These were added to their respective input matrixes for estimating survival and mortality parameters using the instantaneous rates model.

Likewise, there were 4 harvested and 3 re-released migratory striped bass (≥ 711 mm TL) tagged in spring 2018 and recaptured during the 2018-2019. No migratory fish tagged in 2019 have yet been recaptured. These data were also used to complete their respective instantaneous rate model input matrixes (Tables 8a,b).

For striped bass ≥ 457 mm TL, Model 9 received the most support, with Models 10, 14, 4 and 5 also receiving a measure of support. All models estimated similar values of annual survival, averaging about 0.62 during the period 1990-1997 and 0.48-0.51 during 1998-2019 (Figure 1a). Similarly, all models resulted in natural mortality (M) estimates averaging 0.36 during 1990-1997 and 0.61 during 1998-2019 (Figure 1b). Estimates of fishing mortality (F) were more variable, with those models which allow year-specific estimates of F differing from those allowing only periodic changes. Considering only Model 9, F estimates ranged between 0.07 and 0.11, with recent years estimated at 0.04-0.05 (Figure 1c). F-tag estimates followed a general downward trend for nearly all models, with very low 0.01-0.02 values in recent years (Figure 1d).

For migratory striped bass (≥ 711 mm TL), Model 8 received the most support, with models 9 and 3 also receiving a measure of support with several additional models having lower AIC values. All models except Models 1 and 8 estimated similar values of annual survival, averaging about 0.68 during the period 1990-1997 and 0.60-0.63 during 1998-2019 (Figure 2a). All models resulted in natural mortality (M) estimates averaging 0.21-0.22 during 1990-1997 and 0.39-1.40 during 1998-2019 (Figure 2b). Estimates of fishing mortality (F) were somewhat more variable, with those models which allow year-specific estimates of F differing from those allowing only periodic changes. Considering only Model 8, F estimates generally increased between 1990 (0.09) and 1999 (0.26) then ranged between 0.08 and 0.11 until 2008 and have remained near 0.06 since (Figure 2c). F-tag estimates followed a general downward trend for all models, with very low 0.01-0.02 values in recent years (Figure 2d).

Literature Cited

- Akaike, H. 1973. Information theory as an extension of the maximum likelihood principle. In Second International Symposium on Information Theory. Edited by B. N. Petrov and F. Csaki. Budapest. Akademiai Kiado.
- Akaike, H. 1985. Prediction and entropy. In A Celebration of Statistics. Edited by A.C. Atkinson and S.E. Fienberg. New York: Springer.
- Brownie, C., D.R. Anderson, K.P. Burnham, and D.R. Robson. 1985. Statistical inference from band recovery data: a handbook, 2nd ed., U.S. Fish and Wildl. Serv. Resour. Publ. No. 156.
- Burnham, K.P., G.C. White, and D.R. Anderson. 1995. Model selection strategy in the analysis of capture-recapture data. *Biometrics* 51:888-898.
- Hoenig, J.M., N.J. Barrowman, W.S. Hearn, and K.H. Pollock. 1998a. Multiyear tagging studies incorporating fishing effort data. *Can. J. Fish. Aquat. Sci.* 55:1466-1476.
- National Marine Fisheries Service, Northeast Fisheries Science Center. 2019. 66th Northeast Regional Stock Assessment Workshop (66th SAW) Assessment Report. Woods Hole, MA. <https://doi.org/10.25923/nhqe-jd35> (accessed 1 February 2021).
- Ricker, W.E. 1975. Computation and Interpretation of Biological Statistics of Fish Populations. *Bull. Fish. Res. Board Can.* No 191.
- Seber, G.A.F. 1970. Estimating time-specific survival and reporting rates for adult birds from band returns. *Biometrika*, 57: 313-318.
- Sadler, P.W., Hoenig, J.M., Michaelson, S., Goins, L. M. and R.E. Harris. 2016. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 2015-2019. Annual report, Virginia Institute of Marine Science: 141pp.
- Smith, D.R., K.P. Burnham, D.M. Kahn, X. He, C.J. Goshorn, K.A. Hattala, and A.W. Kahnle. 2000. Bias in survival estimates from tag-recovery models where catch-and-release is common, with an example from Atlantic striped bass (*Morone saxatilis*). *Can. J. Fish. Aquat. Sci.* 57:886-897.
- White, G.C. 1983. Numerical estimation of survival rates from band-recovery and biotelemetry data. *J. Wildl. Manage.* 47:716-728.
- White, G.C. and K. P. Burnham. 1999. Program MARKB survival estimation from populations of marked animals. *Bird Study* 46:120-138.
- Wooley, C.M., N.C. Parker, B.M. Florence and R.W. Miller. 1990. Striped bass restoration along the Atlantic Coast: a multistate and federal cooperative hatchery and tagging program.

Table 1. Summary data of striped bass tagged and released in the Rappahannock River, spring 2020.

Date	<457mm TL			457mm - 710mm TL						> 710mm TL			Total		
	Unknown		Males	Unknown		Males		Females	Unknown		Males	Females			
	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n
3/5/2020							1	470.0							1
3/27/2020							5	503.0			3	1165.0			8
3/30/2020				1	481.0	54	509.1			2	1104.5	2	895.0		59
4/8/2020						89	511.7			2	1147.5	1	723.0		92
4/14/2020				4	581.5	53	537.3			7	1041.0	3	864.7		67
4/16/2020				1	500.0	121	518.6			1	973.0	2	952.0		125
4/21/2020				4	594.0	148	524.9					8	933.6		160
4/22/2020				8	558.0	181	516.4					3	890.7		192
4/28/2020				1	572.0	11	553.6			3	979.7	1	886.0		16
5/4/2020				2	500.5	16	532.8			1	782.0	1	907.0		20
5/5/2020				2	569.0	44	542.3			1	1152.0	2	1005.5		49
Total	0	0	0	23		723		0		20	23	0			789

Table 2. Summary data of striped bass tagged and released James River, spring 2020.

Date	<457mm TL			457mm - 710mm TL						> 710mm TL			Total		
	Unknown		Males	Unknown		Males		Females	Unknown		Males	Females			
	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n
3/17/2020				1	485.0	11	510.5								12
3/24/2020						13	486.9								13
3/26/2020				8	596.1	40	512.2			1	1100.0	1	757.0	1	1009.0
4/7/2020				12	575.8	33	521.2			7	975.0				52
4/15/2020				10	554.1	35	494.2			2	1039.5	1	775.0		48
4/24/2020				9	607.4	19	529.7			7	1057.7	1	1010.0		36
Total	0	0	0	40		151		0		17	3	1			212

Table 3. Summary data of striped bass tagged and released Mattaponi River, spring 2020.

Date	<457mm TL			457mm - 710mm TL						> 710mm TL			Total		
	Unknown		Males	Unknown		Males		Females	Unknown		Males	Females			
	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n	Avg TL	n
4/9/2020						9	527.8			2	1093.5				11
4/17/2020				2	613.5	4	563.8			2	1109.0	2	892.5		10
5/1/2020						1	511.0					1	933.0		2
Total	0	0	0	2		14		0		4	3	0			23

Table 4. Location of striped bass (≥ 457 mm TL), recaptured in 2020, that were originally tagged and released in Virginia during springs 1990-2019.

State	Month												Total	
	1	2	3	4	5	6	7	8	9	10	11	12		
Maine	0	0	0	0	0	0	0	0	0	0	0	0	0	0
New Hampshire	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Massachusetts	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Rhode Island	0	0	0	0	0	0	0	1	1	0	0	0	0	2
Connecticut	0	0	0	0	0	0	0	0	0	0	0	0	0	0
New York	0	0	0	0	0	2	0	0	0	0	0	0	0	2
New Jersey	0	0	0	0	1	1	0	0	0	0	0	0	1	3
Pennsylvania	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Delaware	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Maryland	0	0	1	0	4	5	2	3	2	4	0	0	0	21
District of Columbia	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Virginia	0	0	0	0	5	4	2	2	0	2	2	3	0	20
North Carolina	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	0	0	1	0	10	12	4	6	3	6	2	4	0	48

Table 5. Location of migratory striped bass (≥ 710 mm TL), recaptured in 2020, that were originally tagged and released in Virginia during springs 1990-2019.

State	Month												Total	
	1	2	3	4	5	6	7	8	9	10	11	12		
Maine	0	0	0	0	0	0	0	0	0	0	0	0	0	0
New Hampshire	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Massachusetts	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Rhode Island	0	0	0	0	0	0	0	1	0	0	0	0	0	1
Connecticut	0	0	0	0	0	0	0	0	0	0	0	0	0	0
New York	0	0	0	0	0	1	0	0	0	0	0	0	0	1
New Jersey	0	0	0	0	1	0	0	0	0	0	0	0	1	2
Pennsylvania	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Delaware	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Maryland	0	0	0	0	0	0	0	0	0	0	0	0	0	0
District of Columbia	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Virginia	0	0	0	0	0	0	0	0	0	0	0	0	1	1
North Carolina	0	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	0	0	0	0	1	1	0	1	0	0	0	0	2	5

Table 6. Description of IRCR models with varying time periods for fishing mortality, tagging mortality and natural mortality.

	Fishing Mortality	Tagging Mortality	Natural Mortality
Model 1:	Year-specific	Year-specific	2 periods: 1990 - 1997 / 1998 - 2019
Model 2:	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 - 2019	Year-specific	2 periods: 1990 - 1997 / 1998 - 2019
Model 3:	Year-specific	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 - 2019	2 periods: 1990 - 1997 / 1998 - 2019
Model 4:	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 - 2019	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 - 2019	2 periods: 1990 - 1997 / 1998 - 2019
Model 5:	7 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 - 2016 / 2019	7 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 - 2016 / 2017	2 periods: 1990 - 1997 / 1998 - 2017
Model 6:	7 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 / 2016 - 2019	7 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2014 / 2015 / 2016 - 2019	2 periods: 1990 - 1997 / 1998 - 2019
Model 7:	5 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2019	Year-specific	2 periods: 1990 - 1997 / 1998 - 2019
Model 8:	Year-specific	5 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2019	2 periods: 1990 - 1997 / 1998 - 2019
Model 9:	5 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2019	5 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2019	2 periods: 1990 - 1997 / 1998 - 2019
Model 10:	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2016 / 2017-2019	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2016 / 2017-2019	2 periods: 1990 - 1997 / 1998 - 2019
Model 11:	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2015 / 2016 - 2019	6 periods: 1990 - 1994 / 1995 - 1999 / 2000 - 2002 / 2003 - 2006 / 2007 - 2015 / 2016 - 2019	2 periods: 1990 - 1997 / 1998 - 2019

Figure 1a. IRCR generated estimates of annual survival (S) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.




Figure 1b. IRCR generated estimates of annual natural mortality (M) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.




Figure 1c. IRCR generated estimates of annual fishing mortality (F) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.




Figure 1d. IRCR generated estimates of annual tagging mortality (F-tag) for striped bass ≥ 457 mm TL tagged in Virginia, 1990-2019.




Figure 2a. IRCR generated estimates of annual survival (S) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.




Figure 2b. IRCR generated estimates of annual natural mortality (M) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.




Figure 2c. IRCR generated estimates of annual fishing mortality (F) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.




Figure 2d. IRCR generated estimates of annual tagging mortality (F-tag) for striped bass ≥ 711 mm TL tagged in Virginia, 1990-2019.

