
Reports

9-2024

Estimating Relative Juvenile Abundance of Ecologically Important Finfish in the Virginia Portion of Chesapeake Bay: 2024 Annual Report (1 July 2023 – 30 June 2024)

Troy D. Tuckey
Virginia Institute of Marine Science

Mary C. Fabrizio
Virginia Institute of Marine Science

Follow this and additional works at: <https://scholarworks.wm.edu/reports>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Tuckey, T. D., & Fabrizio, M. C. (2024) Estimating Relative Juvenile Abundance of Ecologically Important Finfish in the Virginia Portion of Chesapeake Bay: 2024 Annual Report (1 July 2023 – 30 June 2024). Project Number: F-104-R-28. Virginia Institute of Marine Science, William & Mary. <https://www.doi.org/10.25773/zs7v-t423>


This Report is brought to you for free and open access by W&M ScholarWorks. It has been accepted for inclusion in Reports by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.



2024 ANNUAL REPORT



Estimating Relative Juvenile Abundance of Ecologically Important Finfish in the Virginia Portion of Chesapeake Bay



Prepared by:
Troy D. Tuckey
Mary C. Fabrizio

VIMS | **WILLIAM & MARY**
VIRGINIA INSTITUTE OF MARINE SCIENCE

doi: 10.25773/zs7v-t423

2024 ANNUAL REPORT

Estimating Relative Juvenile Abundance of Ecologically Important Finfish in the Virginia Portion of Chesapeake Bay (1 July 2023 – 30 June 2024)

Project Number: F-104-R-28

Submitted to:

Virginia Marine Resources Commission
380 Fenwick Road, Bldg. 96
Fort Monroe, VA 23651



Prepared by:

Troy D. Tuckey and Mary C. Fabrizio

Virginia Institute of Marine Science
William & Mary, Gloucester Point, Virginia 23062

September 2024

Cover image

The lowest abundance index for Scup was observed in 2023 with only four individuals captured by the survey, three of which were young-of-the-year fish.

TABLE OF CONTENTS

ACKNOWLEDGMENTS	4
EXECUTIVE SUMMARY	5
INTRODUCTION.....	6
METHODS.....	8
RESULTS	12
DISCUSSION	22
LITERATURE CITED	25
TABLES	31
FIGURES	54
APPENDICES	150

ACKNOWLEDGMENTS

Thanks to the many individuals who participated in the field collections, often under difficult circumstances. We thank the crew - Wendy Lowery, Aimee Comer, Jack Buchanan, Katie Nickerson, Daniel Royster, Samantha Dowiarz, and Anna DeMotte. We thank Captains Dale Fitchett and Voight “Bubba” Hogge for their dedication to ensuring safe operations. Appreciation is also expressed to Aimee Comer for data management assistance. We thank Kingsmill Marina on the James River for providing monthly mooring facilities for the *R/V Tidewater*. This study was supported by the Virginia Marine Resources Commission, Project No. F-104-R-28 and VIMS.

EXECUTIVE SUMMARY

1 July 2023 – 30 June 2024

The VIMS Juvenile Finfish Trawl Survey provides crucial data to state, regional, and national fisheries management agencies, including the Virginia Marine Resources Commission (VMRC), the Atlantic States Marine Fisheries Commission (ASMFC), the Mid-Atlantic Fisheries Management Council (MAFMC), and the National Marine Fisheries Service (NMFS). Annual indices of juvenile abundance have been generated from trawl survey data for species of key recreational and ecological importance in the Virginia portion of Chesapeake Bay. These include Spot, Atlantic Croaker, Weakfish, Summer Flounder, Black Sea Bass, Scup, Striped Bass, White Perch, White Catfish, Channel Catfish, Blue Catfish, Silver Perch, American Eel, and Bay Anchovy.

We completed 1,202 tows from July 2023 to June 2024 and collected 645,919 fishes. We were unable to complete sampling in the James River in Dec 2023 due to mechanical issues with the *R/V Tidewater*. Bay Anchovy continue to be the most abundant species observed in the survey, accounting for 61% of all fishes collected, however the Bay Anchovy index for 2023 was below average for the third year in a row. Of the 36 target species categories for which we provide indices of relative abundance, 26 species categories (considering YOY and age 1+ as distinct categories) exhibited below-average abundance in 2023. Scup was particularly low and exhibited the lowest index of the time series with only four individuals captured during the index period; three were of appropriate size to be included in the index. Five species categories exhibited abundances that were similar to their long-term average abundance including Black Sea Bass, Spot, Striped Bass, Summer Flounder, and age 1+ White Perch (York River). Atlantic Croaker, age 1+ Blue Catfish (York River), Silver Perch, and age 1+ White Perch (James and Rappahannock rivers) exhibited above-average abundances in 2023.

INTRODUCTION

Relative abundance estimates of early juvenile (age-0) fishes and invertebrates generated from fishery-independent survey programs provide a reliable and early indicator of year-class strength (Goodyear, 1985) and may be used to evaluate the efficacy of management actions. The Chesapeake Bay Stock Assessment Committee (CBSAC) reviewed available indices of juvenile abundance for important fishery resources in Chesapeake Bay (hereafter referred to as the "Bay") and recommended that "a unified, consistent trawl program should be one of the primary monitoring tools for finfish and crab stock assessment" (Chesapeake Bay Program Stock Assessment Plan, Chesapeake Executive Council, 1988). Subsequently, pilot studies directed at developing a comprehensive trawl survey for the Bay began at the Virginia Institute of Marine Science (VIMS) with monthly trawl sampling in the main stem of the lower Bay. This effort complemented and expanded the monthly trawl sampling conducted in major Virginia tributaries (James, York, and Rappahannock rivers) by VIMS.

The present sampling program, which includes the Bay and its tributaries, ensures that data are of sufficient geographic coverage to generate relative abundance indices for recreationally and ecologically important finfishes. The National Marine Fisheries Service Marine Recreational Information Program shows that recent recreational catches in Virginia are dominated by Atlantic Croaker (*Micropogonias undulatus*), Summer Flounder (*Paralichthys dentatus*), Spot (*Leiostomus xanthurus*), Striped Bass (*Morone saxatilis*), Black Sea Bass (*Centropristis striata*), Bluefish (*Pomatomus saltatrix*), Pigfish (*Orthopristis chrysoptera*), Weakfish (*Cynoscion regalis*), and Kingfishes (*Menticirrhus* spp.). These species depend on the lower Bay and its tributaries as nursery areas and, with the exception of Bluefish, are highly vulnerable to bottom trawls. Additional species of recreational interest, such as Scup (*Stenotomus chrysops*), White Perch (*Morone americana*), Silver Perch (*Bairdiella chrysoura*), White Catfish (*Ameiurus catus*), Channel Catfish (*Ictalurus punctatus*), and Blue Catfish (*I. furcatus*), are also taken with sufficient regularity during trawling operations to provide information suitable for the generation of abundance indices or document changes in relative abundance. Several species of interest are captured in significant numbers across several year classes. As a result, both juvenile and age 1+ (i.e., all fish older than age-0) indices are reported for White Perch, White Catfish, Channel Catfish, and Blue Catfish.

Although annual juvenile indices are the primary focus of this project, data from the survey can be used to address other aspects of finfish population biology, such as habitat utilization, early growth and survival, environmental effects on recruitment, or disease prevalence. For example, episodic climatic events, such as hurricanes, were shown to affect recruitment of shelf-spawning species such as Atlantic Croaker (Montane and Austin, 2005). Tuckey and Fabrizio (2013) used trawl survey data to examine the influence of survey design on fish assemblages comparing fishes captured from fixed sites with those captured at random sites. For some species, the fixed survey

design provided adequate sampling for determination of temporal trends, thus extending the utility of historical (pre-1988) survey catches. Additionally, Tuckey and Fabrizio (2016) examined the indirect effects of hypoxia on fish tissue proximate composition in Bay tributaries. Severe hypoxia, such as that observed in the lower Rappahannock River, reduced the ability of Atlantic Croaker to accumulate lipids in their ovaries, likely affecting their reproductive output (Tuckey and Fabrizio, 2016). Stock assessment models that ignore the effects of hypoxia may yield overly optimistic production estimates for hypoxia-exposed populations, particularly if environmentally invariant fecundity and growth are assumed (Tuckey and Fabrizio, 2016). Using models to describe multi-decadal changes in size of juvenile Summer Flounder, Nys et al. (2015) concluded that variable conditions in nursery habitats may influence production of Summer Flounder year classes through effects on maturation and survival. Additionally, Schloesser and Fabrizio (2015) compared methods to estimate energy density of juvenile Summer Flounder, Striped Bass, and Atlantic Croaker, and demonstrated that the lipid-to-energy conversion factors commonly used in bioenergetics models overestimate energy density of juvenile fishes. Such information can be used to better inform models for ecosystem-based fisheries management because these models rely on inputs from bioenergetics models and juvenile fishes are also an important component of the forage base in the Bay. Schloesser and Fabrizio (2016) described the temporal dynamics of juvenile fish condition using multiple condition metrics; they found that juveniles that remain in the estuary during winter used the liver to store energy and that this energy-storage strategy was common among the species examined. In addition, Striped Bass and Summer Flounder juveniles exhibited density-dependent effects on mean condition (Schloesser and Fabrizio, 2016), suggesting the need to consider relative local abundance when examining condition indices for juvenile fishes. This study provided a more complete understanding of energy-storage strategies for juvenile fishes and highlighted the need to consider temporal dynamics of individual fish condition. In a follow-up study, Schloesser and Fabrizio (2017) described appropriate condition metrics for juvenile fishes based on energy-storage strategies and life history. Together, these studies provide insight on the application of juvenile fish condition metrics as measures of nursery habitat health; additional research on this topic is currently underway. Recently, Nepal and Fabrizio (2020) and Nepal et al. (2020) used trawl survey data to assess population dynamics of invasive Blue Catfish by examining density-dependent growth of juvenile Blue Catfish and phenotypic plasticity of Blue Catfish populations in Virginia. Trawl survey observations were also used to characterize the salinity of tidal habitats used by Blue Catfish and determined that larger fish are more likely to use high salinity (> 10 psu) habitats than smaller fish (Nepal and Fabrizio, 2019). Reproductive characteristics of Blue Catfish were found to differ in the James and York rivers, further supporting the notion of unique stocks in each tributary (Nepal et al., 2021). Fabrizio et al. (2021a) summarized the management objectives for Blue Catfish in Chesapeake Bay and synthesized existing information on this invasive species. The suitability of

Chesapeake Bay habitats for four forage fishes (Bay Anchovy, juvenile Spotted Hake, juvenile Spot, and juvenile Weakfish) was assessed using data from the VIMS trawl survey (Fabrizio et al. 2021b). Extents of suitable habitats for Bay Anchovy in winter and juvenile Spot in summer were reliable predictors of the relative baywide abundance for these two species, suggesting that habitat suitability may limit production of Bay Anchovy and juvenile Spot (Fabrizio et al. 2021b). Tuckey et al. (2021) documented the growth of penaeid shrimp populations in Chesapeake Bay and examined the incidence of black gill disease syndrome. Finally, Marcek et al. (2024) used metabolic scope and hypoxia avoidance to model the distribution of Atlantic Croaker and Spot in the dynamic seascape of Chesapeake Bay. The utility and value of the VIMS trawl survey data grow each year, in step with the questions being asked of the data.

The development of juvenile indices requires a continuous time series of catch data to determine the most appropriate area-time sequences for index calculations. Provisional annual juvenile abundance indices were developed for Spot, Weakfish, Atlantic Croaker, Summer Flounder, and Black Sea Bass (Colvocoresses and Geer, 1991), followed by Scup (Colvocoresses et al., 1992), White Perch, Striped Bass (Geer et al., 1994), White Catfish, Channel Catfish, and Silver Perch (Geer and Austin, 1994). More recently, Blue Catfish, American Eel (*Anguilla rostrata*), and Bay Anchovy (*Anchoa mitchilli*) indices were developed. Through the use of gear conversion factors and post stratification, a time series of index values can be produced back to 1955 for many species (Geer and Austin, 1997).

This report summarizes the activity of the VIMS Juvenile Finfish Trawl Survey (hereafter referred to as the “VIMS Trawl Survey”) from July 2023 through June 2024. We provide abundance indices from 1988 to the present, along with the mean value estimated across the time series; indices for years prior to 1988 are available in previous reports, though sampling design and survey gear differed, and results are not necessarily directly comparable. This year’s indices have been corrected for the change in vessel/gear that occurred in 2015 and values are reported as catches comparable to the R/V *Fish Hawk* for continuity of the time series.

METHODS

Field Sampling

A trawl net with a 5.8-m head line, 40 mm stretch-mesh body, and a 6.4-mm liner was towed along the bottom for 5 minutes during daylight hours. Sampling in the Bay occurred monthly except January and March, when few target species are available. Sampling in the tributaries occurred monthly, at both the random stratified and historical fixed (mid-channel) stations. The stratification system is based on depth and latitudinal regions in the Bay, or depth and longitudinal regions in the rivers. Each Bay region spanned 15 latitudinal minutes and consisted of six strata: western and eastern shore shallow (4-12 ft), western and eastern shoal (12-30 ft), central plain (30-42 ft), and

deep channel (≥ 42 ft). Each tributary was partitioned into four regions of approximately ten longitudinal minutes, with four depth strata in each (4-12 ft, 12-30 ft, 30-42 ft, and ≥ 42 ft; Figure 1). Strata were collapsed in areas where certain depths are limited. Fixed stations were assigned to a stratum according to their location and depth. Additional details of the field sampling protocol are described in Lowery and Geer (2000).

With the exception of the fixed stations in the rivers, trawling sites within strata were selected randomly from the National Ocean Service's Chesapeake Bay bathymetric grid, a database of depth records measured or calculated at 15-cartographic-second intervals. Between two and four trawling sites were randomly selected for each Bay stratum each month, and the number varied seasonally. Exceptions included the shallow-water strata where only a single station was sampled each month. For most river strata, one to two random stations were selected per month. Sampling in the York River was altered slightly as of 1991 to make the deeper depth strata (≥ 30 ft) more similar to those in the James and Rappahannock rivers and main stem Bay. The stratification scheme for the tributaries was modified in January 1996 to further partition the deep strata into two strata of 30-42 ft and ≥ 42 ft (Geer and Austin, 1996). Because tributary sampling occurred at these depths prior to 1996, samples collected previously were reassigned to the strata established in 1996.

Fixed stations were sampled monthly (nearly continuously) since 1980 with sites in each tributary spaced at approximately 5-mile intervals from the river mouth up to the freshwater interface. From the mid-1950s (York River) and early-1960s (James and Rappahannock rivers) to 1972, fixed stations were sampled monthly using an unlined 30' trawl (gear code 010). During 1973-79, semi-annual random stratified sampling was performed by the VIMS Ichthyology Department, while the VIMS Crustaceology Department continued monitoring the fixed tributary stations on a limited monthly basis (May-November). Area-based weightings for the tributaries were previously assigned by dividing each river into two approximately equal length 'strata' by assuming that the stations in each stratum were representative of the channel areas in those reaches (see Lowery and Geer, 2000). As of 1996, all three tributaries were sampled with a random stratified design; the fixed stations were assigned to a stratum based on location and depth. The current design (combined fixed and random stations) provides greater spatial coverage and a long-term historical reference (Tuckey and Fabrizio, 2013).

At the completion of each tow, all fishes were identified to species, counted, and measured to the nearest millimeter fork length (FL), total length (TL), or total length centerline (TLC, Black Sea Bass only). Species that have varying size ranges were measured and counted by size class and large catches of a particular species were randomly subsampled, measured, and the remaining unmeasured catch was counted. In instances of extremely large catches (e.g., Bay Anchovy), subsampling was performed volumetrically.

Juvenile Index Computations

Many of the target species of this study are migratory, so abundance estimation presents special difficulties, particularly if the timing and duration of migration varies annually. Juvenile fishes that use estuarine nursery areas are especially vulnerable to the vagaries of the environment, as many rely on wind-driven and tidal circulation patterns for transport into the estuaries as larvae and early juveniles (Norcross, 1983; Bodolus, 1994; Wood, 2000). In addition, the outward migration of some species from the nursery area may follow annually variable environmental cues (e.g., temperature). Ideally, juvenile abundance should be measured when young fish are fully recruited to the nursery area under study. In practice, however, this can only be accomplished if the time of maximal abundance and size at recruitment to the gear can be predicted (and surveys timed accordingly), or if surveys can be conducted with high frequency over the season of potential maximal abundance. Neither of these two approaches are practical. The period of maximal abundance and the spatial extent of the area occupied by juvenile fish have proven to be variable among years and among species. This observation, coupled with multispecies monitoring objectives, precludes temporally intense surveys. Consequently, the survey is operated with a regular periodicity (monthly) and sample-site selection was performed using a standard sampling design for multispecies resource surveys.

A monthly size threshold value was applied to the length-frequency information collected for each target species to partition the catch data into young-of-year and older components for index calculation (Table 1). Threshold values varied among months for each species and were based on modal analyses of historical, composite length-frequency data and on reviews of ageing studies (Colvocoresses and Geer, 1991). For earlier months of the biological year, threshold values were somewhat arbitrary and fell between completely discrete modal size ranges. In the later part of the biological year, when the size of early spawned, rapidly growing individuals of the most recent year class may approach that of later spawned, slower growing individuals of the previous year class, threshold values were selected to preserve the numeric proportionality between year classes despite the potential misclassification of some individuals (Table 1). The extent of overlapping lengths and the proportion within that range attributable to each year class were estimated based on the shapes of the modal curve during the months prior to the occurrence of overlap. A length value was then selected, which preserves the proportional separation of year classes. Although this process involved considerable subjectivity and ignored possible interannual variability in average growth rates, the likelihood of significant error is small, because only a small fraction of the total number of young-of-year individuals fell within the zone of overlap, and furthermore, with a few exceptions, most of the data used to construct juvenile indices were drawn from months when no overlap was present.

After removing non-target individuals from consideration, monthly stratum-specific abundances and occurrence rates were calculated for each target species. Numbers of individuals captured were natural-log transformed ($\ln(n+1)$) prior to abundance calculations following Chittenden (1991). Average catch rates (and the approximate 95% confidence intervals as estimated by ± 2 standard errors of the mean) were then back-transformed to geometric means. The stratum-specific coefficient of variation was expressed as the standard deviation divided by the log-transformed mean catch: $STD/ E[Y_{st}]$ (Cochran, 1977). The catch data were examined for area-time combinations that provided the best basis for juvenile index calculations. Criteria applied during the selection process included identification of maximal abundance levels, uniformity of distribution, minimization of overall variance, and avoidance of periods during which distribution patterns indicated migratory behavior. Although identification of areas most suitable for index calculations (primary nursery zones) was generally clear, selection of appropriate time windows was more complex. Surveys that are timed on regular monthly intervals may or may not coincide with periods of maximal recruitment to the nursery areas. The use of a single (maximal) month's survey result was therefore inappropriate and would decrease sample size, increase confidence intervals, and increase the risk of sampling artifacts. Conversely, the temporal series of data incorporated into index calculations should not be longer than that necessary to capture the period of maximal juvenile use of the nursery area. With this approach, three- or four-month periods (but six months for Bay Anchovy and American Eel) that provided reasonable abundance data for each species were identified (Table 1).

Using catch data from area-time combinations, an annual juvenile index was calculated as the weighted geometric mean catch per tow (Random Stratified Index, RSI_{GM}) for all species except American Eel, Atlantic Croaker, and Blue Catfish. The RSI_{GM} was obtained by calculating stratum-specific means and variances and combining the stratum-specific estimates using weights based on stratum area (Cochran, 1977). Because stratum areas are not uniform, a weighted mean provides an index that more closely approximates actual population abundance. For American Eel, Atlantic Croaker, and Blue Catfish, whose catches follow a lognormal distribution, an index is calculated assuming a delta lognormal approach (RSI_{Delta}). The delta lognormal index method calculates stratum-specific means on log-transformed positive values only and adjusts the stratum means by the proportion of positive tows within each stratum. The stratified mean is then calculated in the usual manner. We are currently developing confidence interval estimates for the annual relative abundance index, but these are not available for this report.

Indices (Random Stratified Index – RSI_{GM} or RSI_{Delta}) were produced for each species from 1988 to the present using all spatially appropriate data. Data collected prior to 1988 were excluded from this report because results from the longer time series are considered provisional (i.e., indices

prior to 1988 require both gear and vessel conversion factors, and concerns about conversion factors for this period have not yet been addressed).

Data collected after 2015 were adjusted by a species-specific calibration factor to express catches from the R/V *Tidewater* in units comparable to the R/V *Fish Hawk* and thus maintain the continuity of the time series (Appendix Table 2). To accomplish the estimation of calibration factors, we performed side-by-side comparison tows in 2014 and 2015 (40 side-by-side tows were completed in 2016 to target Scup, which were undersampled in 2014-2015 due to low abundances). We compared the catch of the old research vessel and net (R/V *Fish Hawk*) with those of the new research vessel and net (R/V *Tidewater*) based upon completion of 1,141 paired tows. The calibration factor is the ratio of R/V *Fish Hawk* catches to R/V *Tidewater* catches and represents the relative catch efficiency of the *Fish Hawk* to the *Tidewater*. These factors are required for each species-age group because availability, selectivity, and efficiency of the net varies by species and relative size of the individuals captured. Calibration factors (or relative catch efficiency) can be estimated using a number of models, but one of the fundamental characteristics of catch data is that they follow a binomial distribution – either the species is captured by the paired tow or not. The binomial distribution does not account for the additional variation (overdispersion) that is typically observed (McCullagh and Nelder, 1989), so in addition to the binomial model, we considered models that specifically address overdispersion. We considered three additional models to account for the overdispersed catch data: the beta-binomial model, a random-clumped binomial distribution model, and a generalized linear overdispersion model based on the beta-binomial distribution. The best model was selected using information-theoretic approaches, namely Akaike's Information Criterion corrected for small sample sizes (AIC_c). The model selected to calculate the calibration factor varied by species and is discussed under the description for each species. The calibration factor was applied at the individual tow-level and equated catches of fish from the R/V *Tidewater* with those from the R/V *Fish Hawk* (historic data). We included tow direction, current speed, Secchi depth, and an offset for the ratio of the distance towed by each vessel as explanatory factors in the probability of success (i.e., the probability of catching the target group). We used salinity as an explanatory factor for the variation in the overdispersion parameter in models that included an overdispersion parameter.

RESULTS

A summary of tows completed from 1988 through June 2024 (Table 2) provides a synopsis of the sampling completed for this report. For the 2023-2024 project year (1 July 2023 through 30 June 2024), 1,202 sites were sampled resulting in 645,919 fishes identified and enumerated, representing 106 species (Table 3). Bay Anchovy and Hogchoker accounted for 68% of the total catch by numbers

(Table 3). Ignoring Bay Anchovy and Hogchoker, five species – Spot, Atlantic Croaker, Weakfish, Atlantic Menhaden, and Silver Perch – represented 86% of the catch numerically (Table 3).

Water temperature, salinity, and dissolved oxygen were measured at the end of each trawl tow and are presented here as weighted means (with 95% confidence intervals; Figures 2 - 4). Mean annual bottom-water temperature has increased about 1° C in the Rappahannock and York rivers, and about 3-4° C in the James River since 1988 (Figure 2). Mean annual bottom-water temperature exhibited an upward trend in the Virginia portion of the Bay from less than 15° C in 1988 to 18 - 19° C through 2020 and decreased to a mean of 17° C in recent years (Figure 2). Mean annual salinity in the region exhibited similar interannual patterns with highest salinity found in the Bay and lower salinities in the subestuaries (Figure 3). Mean salinity increased slightly in all regions in 2023-2024, returning to levels observed previously in the time series (Figure 3). Annual bottom dissolved oxygen averaged between 6 and 9 mg/L in all regions sampled, with greater variability (wider confidence intervals) observed in the Rappahannock River subestuary compared with other regions (Figure 4).

Relative abundance indices are described below for the following species: American Eel, Atlantic Croaker, Bay Anchovy, Black Sea Bass, Blue Catfish, Channel Catfish, Scup, Silver Perch, Striped Bass, Spot, Summer Flounder, Weakfish, White Catfish, and White Perch. In addition, the distribution of each species, number captured, mean length and size range of fish during index months and from index strata are also reported. Furthermore, mean bottom-water temperature, salinity, and dissolved oxygen at sites where positive catches of each species occurred (during the index period) were compared with water quality observations from all sites and strata sampled during the index period. Length-frequency distributions for each species are shown in Appendix Figure 1 with index-sized fish indicated in black. Relative abundance indices were calculated on a subset of the data as described below and time-series averages included all years except the most recent year.

American Eel (*Anguilla rostrata*) – American Eel are present along the Atlantic and Gulf coasts of North America and inland in the St. Lawrence Seaway and Great Lakes (Murdy et al., 1997). This catadromous species is panmictic and is supported throughout its range by a single spawning population (Haro et al., 2000). Spawning takes place during winter to early spring in the Sargasso Sea. The eggs hatch into leaf-shaped, ribbon-like larvae called leptocephali, which are transported by ocean currents (over 9-12 months) in a generally northwesterly direction. Within a year, metamorphosis into the next life stage (glass eel) occurs in the Western Atlantic near the east coast of North America. Coastal currents and active migration transport the glass eels into rivers and estuaries from February to June in Virginia and Maryland. As growth continues, eels become pigmented (elver stage) and within 12 –14 months they acquire a dark color with underlying yellow (yellow eel stage). Many eels migrate upriver into freshwater rivers, streams, lakes, and ponds,

while others remain in estuaries. Most of the eel's life is spent in these habitats as yellow eel. Age at maturity varies greatly with location and latitude, and in the Bay may range from 8 to 24 years; most eels in the Bay area are immature and less than 7 years old (Owens and Geer, 2003). Eels from the Bay mature and migrate at an earlier age than eels from northern areas (Hedgepeth, 1983). Metamorphosis into the silver eel stage occurs during the seaward migration that occurs from late summer through autumn, as mature eels migrate back to the Sargasso Sea to spawn and die (Haro et al., 2000).

The current American Eel index includes all eels > 152 mm TL collected in each of the major tributaries (Figure 6) during April through September 2023 (Fenske et al., 2011). The best model supported by the data (n=27 paired tows) to estimate the calibration factor was the binomial model resulting in a calibration factor equal to 0.8731.

American Eel indices exhibited above-average abundance in the Rappahannock River (mean $RSI_{Delta} = 1.78$) and the James River (mean $RSI_{Delta} = 1.57$) in the late 1980's and 1990's and below-average abundance thereafter (Table 4; Figure 5). In the York River, below-average abundance (mean $RSI_{Delta} = 0.47$) has been observed since 1999. Indices for American eel were below-average in 2023 in all three subestuaries. American Eel are typically more abundant in the Rappahannock River compared with the James and York rivers, however a similar number of eels were captured in the James River in 2023 as were found in the Rappahannock River (Figure 6). Positive catches of American Eel are associated with average water temperature (Figure 7), average to below-average salinity (Figure 8), and average to above-average dissolved oxygen content compared with conditions at all sites sampled from index months and strata (Figure 9).

Atlantic Croaker (*Micropogonias undulatus*) – Atlantic Croaker are typically captured in high abundance and are widely distributed throughout the survey area (Figure 11). Spawning takes place over a protracted period, such that small juveniles (<30 mm TL) can be present in catches year-round (Norcross, 1983; Colvocoresses and Geer, 1991; Colvocoresses et al., 1992; Geer et al., 1994). For some year classes, peak abundance occurs in the fall at lengths less than 100 mm TL, but for other year classes, the peak occurs the following spring. Previously, we provided two estimates of the index: a juvenile fall index (October-December) based on catches in the tributaries, and a spring recruit index (May-August) based on catches in the Bay and tributaries combined. Because the fall index does not reflect over-winter mortality, only the modified spring index, after Woodward (2009), is presented. The spring index is calculated using a delta-lognormal model with juvenile Atlantic Croaker collected from all regions of the Bay and tributaries from April to June. The beta-binomial model was the best model (n=283) and resulted in a calibration factor of 1.4770.

The 2023 year class of Atlantic Croaker exhibited the highest level of recruitment (Index = 93.77, mean = 15.28) for the time series and the second above-average recruitment event in a row (Figure 10). Atlantic Croaker were found throughout the Bay with most captured in the three

tributaries, particularly the York River (Figure 11). Positive catches of juvenile Atlantic Croaker are found in warm waters, with average to below-average salinity and DO (Figures 12 – 14).

Bay Anchovy (*Anchoa mitchilli*) – Bay Anchovy are the most abundant finfish in the Bay and its tributaries and are found in salinities ranging from 1-33 psu (Murphy et al., 1997). Bay Anchovy feed mostly on zooplankton and are an important prey of other Bay fishes (Murphy et al., 1997). In years of average freshwater inflow (e.g., 1997-2000), Atlantic Menhaden, Bay Anchovy, and Atlantic Croaker often dominate fish biomass in the Bay (Jung, 2002). Only two models converged for estimating the calibration factor for Bay Anchovy and the beta-binomial model was best supported by the data (n=504 paired tows) with a calibration factor of 0.8014.

Bay Anchovy abundance is variable from year to year with highest recruitment observed between 2008 and 2011 (Table 6; Figure 15). However, since 2013, Bay Anchovy have exhibited average to below-average recruitment. The 2023 index ($RSI_{GM} = 9.39$) was below-average (Average $RSI_{GM} = 24.08$). As expected, Bay Anchovy were ubiquitous in trawl survey catches with many stations yielding thousands of individuals in a single tow (Figure 16). Positive catches of Bay Anchovy occurred in cooler waters (Figure 17), with average salinity (Figure 18), and in areas with average to higher dissolved oxygen (Figure 19) compared with all sites sampled from index months and strata.

Black Sea Bass (*Centropristis striata*) – Black Sea Bass are seldom taken in large numbers but regularly occur in survey catches. Young-of-year Black Sea Bass occur throughout the Bay and appear occasionally in the lower portions of the tributaries. Index calculations are based on all Bay strata and the lower James stratum from May through July only (Figure 21). Although some early juveniles appear in the Bay during their first summer and fall, more young-of-year enter the estuary during the following spring. Black Sea Bass spawn in the summer in the Mid-Atlantic Bight (Musick and Mercer, 1977). Thus, the index is calculated for the year class spawned the previous calendar year (i.e., the index for the 2019 year class is based on catches from May to July 2020). The best model for estimating the calibration factor for Black Sea Bass (n=25 paired tows) was the binomial model resulting in a calibration factor equal to 0.6336.

The Black Sea Bass index in 2023 (2022 year class) was average (mean $RSI_{GM} = 0.74$; Table 7). The Black Sea Bass index has been average or below-average for the last ten years (Table 7; Figure 20). Black Sea Bass are found mostly in the Bay with some occurring on the lower James River (Figure 21). Positive catches of Black Sea Bass were variable with respect to water temperature (Figure 22), but were associated with higher salinity (Figure 23) and normoxic conditions (Figure 24) compared with all sites sampled during index months and strata.

Blue Catfish (*Ictalurus furcatus*) – The Blue Catfish, one of Virginia's largest freshwater fishes (Jenkins and Burkhead, 1993), was introduced to the Bay area as a sportfish in the James, Rappahannock, and Mattaponi rivers between 1974 and 1989 (Virginia Department of Game and

Inland Fisheries, 1989 as reported by Connelly, 2001). The Blue Catfish is an omnivorous bottom feeder that inhabits the main channels and backwaters of rivers (Schmitt et al. 2019). Blue Catfish are distributed from the mesohaline portions of the major tributaries upstream into freshwater habitats, beyond the sampling limits of the trawl survey (Figures 26 and 31; Schloesser et al., 2011). Because Blue Catfish are restricted in their distribution, an index of abundance is calculated for each tributary incorporating strata from portions of the lower river and all upper river strata. For estimating calibration factors for juvenile (< 130 mm FL) and age 1+ Blue Catfish (>130 mm FL), the beta-binomial model was best supported by the data ($n_{\text{juvenile}}=78$ paired tows; $n_{\text{age-1+}}=123$ paired tows) and resulted in a calibration factor equal to 1.0023 for juvenile Blue Catfish and 1.4289 for age 1+ Blue Catfish.

The James River typically has the highest juvenile Blue Catfish index with above-average recruitment (mean $RSI_{\text{Delta}} = 2.89$) in 1996, 2003, 2004, 2013, 2016, 2020 and the highest recruitment ever in 2018 (Table 8; Figure 25). The 2023 juvenile index for Blue Catfish in the James River was well below-average ($RSI_{\text{Delta}} = 0.00$), with no individuals captured for the third year in a row. Recruitment of Blue Catfish juveniles in the Rappahannock River was highest in 2003 ($RSI_{\text{Delta}} = 10.40$) followed by 2011 and 2018. The 2023 index for the Rappahannock River was also well below-average ($RSI_{\text{Delta}} = 0.00$) with no individuals captured during the index period. In the York River, Blue Catfish juveniles in 2004 ($RSI_{\text{Delta}} = 18.56$) were 6 times more abundant than the previous record ($RSI_{\text{Delta}} = 2.97$ in 2003; Figure 25). The 2023 index ($RSI_{\text{Delta}} = 0.00$) for the York River population was below-average (mean $RSI_{\text{Delta}} = 1.80$) with no individuals captured. Historically, positive catches of juvenile Blue Catfish occurred in average water temperatures (Figure 27) with lower salinity (Figure 28) and similar dissolved oxygen (Figure 29) compared with conditions at all sites sampled during index months and strata.

Age 1+ Blue Catfish resided in mesohaline portions of the Rappahannock, York, and James rivers (as well as freshwater habitats not sampled by the survey; Figure 31). The 2023 abundance index for age 1+ Blue Catfish in the Rappahannock River ($RSI_{\text{Delta}} = 9.70$) was below average (mean long-term $RSI_{\text{Delta}} = 24.29$; Table 9; Figure 30). In the York River, abundance indices for age 1+ Blue Catfish have been average or above average since 2014 (mean long-term $RSI_{\text{Delta}} = 1.74$; Table 9; Figure 30). Recent age 1+ Blue Catfish abundance estimates from the York River have been among the highest observed in this system; the 2023 abundance index was above-average at 2.46. In the James River, above-average abundance estimates for age 1+ Blue Catfish have been observed for the previous eight years (mean long-term $RSI_{\text{Delta}} = 34.78$; Table 9; Figure 30). The 2023 index for the James River was 16.20 and below the average. Positive catches of Age 1+ Blue Catfish occurred in average water temperature (Figure 32), with lower salinity (Figure 33), and in a wide range of dissolved oxygen conditions (Figure 34) compared with conditions at all sites sampled during index months and strata. We note that positive catches of age 1+ Blue Catfish have been occurring in

higher salinity waters in recent years compared with the early portion of the times series (Figure 33), suggesting a shift in the spatial distribution of the population reflecting adaptation to more saline environments, or a shift in the size structure of the population (larger fish are more tolerant of saline conditions than smaller fish; Nepal and Fabrizio 2019)

Blue Catfish indices have been increasing since the mid-1990's and the ecosystem effects of this invasive species are not fully known (Schloesser et al., 2011). Diets of small Blue Catfish are dominated by invertebrates (mostly amphipods, isopods and mud crabs), whereas larger Blue Catfish consume invertebrates, Atlantic Menhaden (*Brevoortia tyrannus*), and Gizzard Shad (*Dorosoma cepedianum*; Parthree et al., 2008; Schmitt et al., 2017; Schmitt et al., 2019). Other catfishes (White and Channel) have similar diets and may compete with Blue Catfish for the same prey resources (Schloesser et al., 2011). A recently completed study investigating the trophic interactions between Blue Catfish and Blue Crab (*Callinectes sapidus*) in the mesohaline portion of the James River indicated that consumption of Blue Crabs was about 2.3 million crabs annually in the study area between Hog Island and Burwell Bay (Fabrizio et al. 2021c).

Channel Catfish (*Ictalurus punctatus*) and **White Catfish** (*Ameiurus catus*) – Channel Catfish and White Catfish are usually found in the upper portions of the tributaries (e.g., Figure 81). Although each river system is unique, spawning typically occurs from late May to early July in Virginia (Menzel, 1945); consequently, June was selected as the start of the biological year. The survey typically catches both species up to 600 mm FL with young-of-the-year fish (≤ 50 mm FL) first recruiting to the gear in June. In most years, juvenile recruitment occurs from September to November for both species in the upriver strata only. Because of the low number of paired tows that contained juvenile and age 1+ Channel Catfish and juvenile White Catfish, we used the model-based calibration factor for Blue Catfish for these species. The beta-binomial model was the best model for Blue Catfish and resulted in a calibration factor equal to 1.0023 for YOY catfishes and 1.4289 for age 1+ catfishes. In 2018, the recruitment window for juvenile and age 1+ Channel Catfish and White Catfish was changed from January to April to the new period that occurs from September to November. Annual indices were recalculated to reflect the new recruitment period.

The Channel Catfish was introduced to Virginia in the late 1800s (Jenkins and Burkhead, 1993). Juvenile Channel Catfish exhibited low or failed recruitment in most years with a few notable peaks. In the past 15 years, only ten juvenile Channel Catfish were captured by the trawl survey and none were captured in 2023 during index months (Table 10, Figure 35). In the past, positive catches of juvenile Channel Catfish were associated with average water temperature (Figure 37), below-average salinity (Figure 38), and average dissolved oxygen (Figure 39) compared with all sites sampled during index months and strata.

The age 1+ Channel Catfish RSI_{GM} exhibited below-average values since the early 1990's in the Rappahannock River with only three fish captured since 2018 and none in 2023. In the York

River, no age 1+ Channel Catfish have been captured since 2004. In the James River, no age 1+ Channel Catfish have been captured since 2014 (Table 11; Figure 40). When captured, age 1+ Channel Catfish were found in below-average water temperature (Figure 42), below-average salinity (Figure 43), and average dissolved oxygen conditions (Figure 44) compared with all sites sampled during index months and strata.

Juvenile White Catfish reappeared in the York River in 2018 after a succession of years (2003-2017) when no recruitment was observed (Table 18; Figure 75). In 2023, the index of abundance for juvenile White Catfish in the York River was below average with no juvenile White Catfish being captured. Juvenile White Catfish were most abundant in the James River, however abundance in 2023 ($RSI_{\text{Delta}} = 0.0$) was below-average ($RSI_{\text{mean}} = 0.05$; Figure 75). Juvenile White Catfish have not been observed during index months in the Rappahannock River since 2003. Positive catches of juvenile White Catfish occurred in average to below-average water temperature (Figure 77), below-average salinity (Figure 78), and average dissolved oxygen conditions (Figure 79) compared with all sites sampled during index months and strata.

We had a sufficient number of comparison tows ($n=32$ paired tows) to estimate a calibration factor for age 1+ White Catfish and the best model was the binomial model resulting in a calibration factor equal to 0.6918. Abundance indices for age 1+ White Catfish were below-average since 2010 in the James River (average = 0.35), below-average since 2004 in the York River (average = 0.10), and below-average since 1997 in the Rappahannock River (average = 0.15; Table 19; Figure 80). Age 1+ White Catfish in 2023 were found only in the James and York rivers (Figure 81). Positive catches of age 1+ White Catfish tended to occur in cooler water in the Rappahannock and York rivers, compared with all sites sampled in these two systems, whereas age 1+ White Catfish were captured in average to above-average water temperatures in the James River (Figure 82). Low salinity waters (Figure 83) with average dissolved oxygen conditions (Figure 84) were preferred by age 1+ White Catfish compared with all sites sampled from index months and strata.

Scup (*Stenotomus chrysops*) – Scup are primarily a marine, summer spawning species that use the Bay in a manner similar to Black Sea Bass. The estuary is rarely used as a nursery area by early juveniles (25-40 mm FL), which occasionally appear in survey catches in June. Older juveniles can be found in the Bay during their second summer and first appear in catches in May; by June, they range from 45 to 215 mm FL. Using the original length threshold for Scup that was based on ageing studies (Morse, 1978), trawl survey catches were found to typically include three age groups (age 0, age 1, and age 2+). Because catches of age 1 and age 2+ Scup are highly variable and low, index calculations using trawl survey catches are based on age-0 individuals only. Age-0 fish are present in the Bay and available to the gear for the entire summer and early fall (June to September).

During index months, Scup are predominantly collected in the lower Bay (Figure 46). Catch rates for Scup usually peak in July, and the index was calculated from catches taken in June to September. The generalized linear overdispersion model based on the beta-binomial distribution was the model most supported by the data (n=27 paired tows) resulting in a calibration factor equal to 1.3471.

Scup indices were above-average in 2000, 2005, 2006, 2008, 2009, and 2014 and below-average most other years since 1993 (mean long-term $RSI_{GM} = 1.14$; Table 12; Figure 45). Age-0 Scup abundance in 2023 (2022 year-class) was the lowest ever observed with only three individuals captured ($RSI_{GM} = 0.0$; Table 12). Positive catches of Scup occurred in a variety of water temperatures from below-average to above-average conditions (Figure 47), typically with higher salinity (Figure 48), and wide-ranging dissolved oxygen conditions (Figure 49) compared with all sites sampled from index months and strata.

Silver Perch (*Bairdiella chrysoura*) – Silver Perch are found in all sampling strata (Figure 51). Spawning occurs in the deep waters of the Bay and offshore from May to July, and juveniles (≤ 100 mm TL) enter the Bay by July (Chao and Musick, 1977; Rhodes, 1971). The beta-binomial model was best supported by the data (n=118 paired tows) resulting in a calibration factor for juvenile Silver Perch equal to 0.7524.

Abundance indices for Silver Perch were consistent and stable through 2003 and increased to higher levels thereafter until 2018 (Table 13; Figure 50). The time series average RSI_{GM} index for Silver Perch was 0.72 and the 2023 index ($RSI_{GM} = 1.29$) was above-average. Silver Perch were found in all regions sampled but were most abundant in the York and Rappahannock River systems (Figure 51). Positive catches of juvenile Silver Perch were associated with a wide range of water temperatures (Figure 52), salinities (Figure 53), and dissolved oxygen conditions (Figure 54) that varied relative to all sites sampled from index months and strata.

Spot (*Leiostomus xanthurus*) – Spot were widely distributed throughout the Bay and tributaries (Figure 56); abundance indices for this species were calculated using all strata from July to October. Spot is often one of the most abundant recreational species captured by the survey. Two models were equally supported by the data (n=187 paired tows) and we chose the simpler model (Beta-binomial model) resulting in a calibration factor equal to 1.1654.

Below-average abundance for Spot was observed between 1992 and 2007, with average or above-average abundance observed during 2008, 2010, 2012, 2013, and 2018 (Table 14; Figure 55). The RSI_{GM} index in 2023 (15.03) was average (mean long-term $RSI_{GM} = 12.62$). Spot were found throughout the Bay and river systems (Figure 56). Positive catches of Spot were associated with average water temperature (Figure 57), salinity (Figure 58), and dissolved oxygen conditions (Figure 59) compared with all sites sampled from the same index months and strata.

Striped Bass (*Morone saxatilis*) – Striped Bass use the upper tributaries of the Bay as spawning and nursery grounds; spawning occurs from early to mid-April through the end of May, in tidal freshwater areas just above the salt wedge. Juvenile Striped Bass often appear in catches from May to July in size classes less than 50 mm and up to 100 mm FL during years of high abundance, and we used August to November as the index period. Index calculations were from those juvenile Striped Bass captured in the major tributaries only (Figure 61), although Striped Bass were encountered in other areas throughout the year. The beta-binomial model was best supported by the data (n=90 paired tows) resulting in a calibration factor equal to 1.2009. In 2018, the recruitment window for juvenile Striped Bass was changed from the period December to February to the new period between August and November. Annual indices were recalculated to reflect the new recruitment period.

Juvenile Striped Bass showed periods of stronger recruitment after the 1995 year class (Table 15; Figure 60). The index value for the 2023 year class ($RSI_{GM} = 0.24$, $CI = 0.0 - 0.59$) was average for the time-series (mean $RSI_{GM} = 0.38$). Juvenile Striped Bass were more abundant in the James River (Figure 61) than in the York or Rappahannock rivers. Positive catches of juvenile Striped Bass varied with respect to water temperatures (Figure 62) but tended to occur in lower salinity (Figure 63) and higher dissolved oxygen conditions (Figure 64) compared with all sites sampled from index months and strata.

Summer Flounder (*Paralichthys dentatus*) – Summer Flounder spawn on the continental shelf from September through January with a peak occurring in October and November (Murdy et al., 1997). Summer Flounder larvae enter the Bay and other Virginia estuaries from October through May with juveniles using shallow fine-substrate areas adjacent to seagrass beds (Murdy et al., 1997; Wyanski, 1990). Whereas low water temperatures can have significant effects on growth and survival of individuals that enter the estuary in the winter (Able and Fahay, 1998), high water temperatures during summer can also have negative effects on the size of individual juvenile Summer Flounder (Nys et al., 2015). Juvenile Summer Flounder first appear in catches in late March, which was used as the beginning of the biological year. Juvenile Summer Flounder abundance increases steadily throughout the summer and early fall to a late fall peak, and then trawl catches decline, presumably reflecting emigration of young fish during December. For our trawl survey, September, October, and November usually encompass the months of greatest abundance of juvenile Summer Flounder. Juvenile Summer Flounder are broadly distributed throughout the Bay and lower rivers. Consequently, index calculations are based on catches from the Bay and lower river strata during September, October, and November (Figure 66). The beta-binomial model was the best model (n=146 paired tows) resulting in a calibration factor equal to 1.0308

Juvenile Summer Flounder indices were greater during the early 1990s compared with recent years with notable peaks in 2004, 2008, and 2012 (Table 16; Figure 65). The 2023 Summer

Flounder index ($RSI_{GM} = 0.81$, $CI = 0.60 - 1.05$) was average for the time-series (mean $RSI_{GM} = 0.87$). Positive catches of juvenile Summer Flounder occurred in average to cooler water temperatures (Figure 67) with above average salinity (Figure 68) and variable dissolved oxygen conditions (Figure 69) compared with all sites sampled from index months and strata.

Weakfish (*Cynoscion regalis*) – Historically, weakfish were one of the dominant species in trawl survey catches, and juveniles are found throughout the Bay and tributaries (Figure 71). Juvenile catches occur in late May and June, with June considered the beginning of the biological year. The best model for estimating the calibration factor was the beta-binomial model ($n=220$ paired tows) resulting in a calibration factor equal to 1.1924.

Overall, the Weakfish index was consistent through time indicating steady recruitment between 1988 and 2013 (mean $RSI_{GM} = 5.60$; Table 17; Figure 70). However, we observed below-average recruitment since 2014, with the 2023 index ($RSI_{GM} = 2.35$) below average for the tenth year in a row. Juvenile Weakfish are found throughout the Bay and rivers with most being found in the York and Rappahannock rivers (Figure 71). Positive catches of juvenile Weakfish occurred in a wide range of water temperatures (Figure 72) with average salinity (Figure 73) and normoxic dissolved oxygen conditions (Figure 74) compared with all sites sampled from index months and strata.

White Perch (*Morone americana*) – Spawning of White Perch occurs in the upper tributaries (Figure 86) from March to July with a peak occurring from late April to early May. Index months include August to October for juveniles (< 85 mm TL) and age 1+ (>81 mm TL). Index stations are from the upper river strata and a separate index is calculated for each river. The beta-binomial model was best supported by the data for each size group ($n_{juvenile}=161$ paired tows; $n_{age-1+}=210$ paired tows) resulting in a calibration factor equal to 1.0679 for YOY White Perch and 1.1676 for age-1+ White Perch. In 2018, the recruitment window for juvenile White Perch was changed from the period December to February to the new period between August and October. Similarly, the recruitment window for White Perch age 1+ was changed from November to February to the new period between August and October. Annual indices were recalculated to reflect the new recruitment period.

Juvenile White Perch recruitment has been stable in each of the rivers, with high and low periods of recruitment (Table 20; Figure 85). The three rivers exhibited different time-series averages with greater numbers of juvenile White Perch occurring in the James River (mean $RSI_{GM} = 1.13$) followed by the Rappahannock River (mean $RSI_{GM} = 0.13$) and the York River (mean $RSI_{GM} = 0.02$). No juvenile White Perch were captured in the York or Rappahannock rivers in 2023, and the index was below-average in the James River in 2023 with only one individual captured (Figure 85). Positive catches of juvenile White Perch were associated with average water temperature (Figure

87), average to below-average salinity (Figure 88), and average dissolved oxygen (Figure 89) compared with all sites sampled from index months and strata.

Abundance of age-1+ White Perch was average in the Rappahannock River (mean $RSI_{GM} = 0.68$) for most of the time series, however the index was above-average in 2023 tying the previous record observed in 2003 ($RSI_{GM} = 2.42$; Table 21; Figure 90). In the York River, abundance of age-1+ White Perch was average or above-average (mean $RSI_{GM} = 0.05$) prior to 1997 and average to below-average values have been observed since 1998, except in 2003 and 2004 (Table 21; Figure 90). In 2023, abundance of age-1+ White Perch was average in the York River ($RSI_{GM} = 0.05$). Abundance of age-1+ White Perch has been average in the James River (mean $RSI_{GM} = 1.32$) with above-average abundance observed between 1995 and 2004 (Table 21; Figure 90). Abundance of age-1+ White Perch was above average in 2023 in the James River ($RSI_{GM} = 3.06$). Age-1+ White Perch were collected throughout the upper rivers with most occurring in the James and Rappahannock rivers (Figure 91). Positive catches of age-1+ White Perch occurred in average water temperature (Figure 92), average salinity (Figure 93), and normoxic conditions (Figure 94) compared with all sites sampled from index months and strata.

DISCUSSION

Juvenile indices contribute to the assessment and management of important recreational species in Chesapeake Bay and the mid-Atlantic Bight. For example, the VIMS Trawl Survey was recognized by the Mid-Atlantic Fisheries Management Council (MAFMC) as an important source of the Summer Flounder recruitment index; the VIMS index was instrumental in shaping protective harvest regulations in Virginia for this species. Other indices utilized by management agencies include those for American Eel, Atlantic Croaker, Atlantic Menhaden, Black Sea Bass, Red Drum, Spot, and Weakfish. Although a bottom trawl is not the preferred gear with which to sample American Eel, indices from the VIMS Trawl Survey played an important role in the 2023 ASMFC American Eel Benchmark Stock Assessment (ASMFC, 2023a). In addition to management needs, the VIMS Trawl Survey also fulfills data and specimen requests from a variety of agencies, institutes, and individuals for research and educational purposes (e.g., Woodland et al., 2012; Sobocinski et al., 2013; Nys et al., 2015; Nepal and Fabrizio, 2020; Marcek et al., 2024; Nepal et al., 2024). Between July 2023 and June 2024, we responded to 86 requests for information and specimens (Appendix Table 1).

Efforts to improve recruitment indices continue and include evaluation of the size ranges and months (the index period) used in index calculations. For example, the distributional assumptions of the catch of YOY Weakfish and Atlantic Croaker were addressed in a VIMS Master's thesis (Woodward, 2009); the results showed that the nonzero catch data for Weakfish are best described by a gamma distribution and those for Atlantic Croaker appear to follow a lognormal

distribution. Such findings indicate that indices of abundance calculated for these species could benefit from further refinements. In addition, the use of different index months for Weakfish and Atlantic Croaker may improve YOY indices by ensuring fewer age-1+ fish are included in YOY index calculations. Additional work may address potential effects of depth on the distribution and catch of these species (Woodward, 2009; Tuckey and Fabrizio, 2013).

The VIMS Trawl Survey provides more than relative abundance indices used to tune stock assessments and aid in management activities. The data can also be used to investigate factors that influence species abundance that operate on time periods beyond annual recruitment cycles. For example, using fishery-independent survey data from 1968 to 2004 for estuarine-dependent species, Wood and Austin (2009) found that recruitment of anadromous species was negatively correlated with recruitment of species that spawn on the continental shelf. Furthermore, recruitment patterns favored one group over the other for periods greater than one decade and shifts between recruitment regimes occurred within a short period of time (2-3 years; Wood and Austin, 2009). Understanding that long-term recruitment cycles may dominate for decades is an important development that could inform management options.

Information from the VIMS Trawl Survey also provides a basis for monitoring species interactions. For example, annual catch rates of Channel Catfish and White Catfish have declined since 1991, while abundance of the introduced Blue Catfish has increased dramatically (Schloesser et al., 2011; Fabrizio et al., 2018). Because diets and distributions of these species overlap, the observed trends may be due to competition and thus, species interactions warrant further study. Furthermore, the shift in diet of older Blue Catfish to include other fishes may affect ecosystem function (Schloesser et al., 2011; Schmitt et al., 2017).

Changes in catches of important recreational species may be associated with degradation of estuarine nursery habitats, overfishing, poor recruitment, or a combination of these factors (Murphy et al., 1997). Although it is not possible to determine the cause of recruitment variability from trawl survey data alone, some general observations are possible. Spot recruitment indices for the Bay region have declined greatly over the past 50 years, but year-class strength of this oceanic spawner appears to be controlled by environmental factors occurring outside the Bay (Homer and Mihursky, 1991; Bodolus, 1994). Three of the highest Spot recruitment indices from the past 20 years were observed within the past ten years, indicating favorable conditions for this species in recent years. Atlantic Croaker recruitment indices show significant interannual variability with fluctuations possibly related to environmental conditions that vary annually. Norcross (1983) found that cold winters increased mortality in overwintering juvenile Atlantic Croaker and during some years may “push” the spawning population further south, thereby impeding access of early-stage juveniles to nursery areas in the Bay. The warm winter of 2011-2012 may have contributed to greater winter survival and the high index of abundance for this species observed in 2012, whereas the cold winter

of 2014 - 2015 may have contributed to the low index observed in 2015 (2014 year-class) for Atlantic Croaker. Weakfish recruitment indices remained stable from 1988 to 2013. However, the index for Weakfish for the past seven years has been below average with five of the lowest index values ever observed occurring during this time period. The lowest index ever observed for this species occurred in 2021, and the 2023 index was below-average continuing a concerning pattern. There has also been a decline in adult abundance which may be attributed to an increase in total mortality (ASMFC, 2012). Striped Bass display great recruitment variability and one or two strong year classes may dominate the population at any one time (Richards and Rago, 1999). Finally, recruitment of Summer Flounder to the Bay nurseries in 2020 was the lowest ever observed and there was only a slight increase in 2021 and 2022. Summer flounder recruitment was average in 2023 indicating a potential slow recovery. Observed declines in the abundance of the adult stock of Summer Flounder may have been due to overfishing or year-class failure (Terceiro, 2006); spawning stock biomass has been steadily decreasing in recent years and while it is not overfished, overfishing is occurring (ASMFC, 2023b). The spawner-recruit relationship in Summer Flounder is highly uncertain, suggesting that environmental conditions during the pre-recruit stage play a role in structuring year-class strength. A meta-analysis of the stock-recruitment relationship for 225 marine stocks suggests that for most stocks, environmental effects may exert a stronger influence on recruitment than spawning stock biomass (Szuwalski et al., 2014). Furthermore, environmental conditions in Bay nurseries may affect juvenile abundance through indirect effects on growth. For example, Nys et al. (2015) found that high water temperature during the summer and a high density of juvenile Summer Flounder negatively affected the observed size of juvenile Summer Flounder in Chesapeake Bay nurseries. Smaller juvenile Summer Flounder may experience higher mortality rates due to size-based predation.

The VIMS Trawl Survey program supplies critical data for management of fishery resources that use the Bay as a spawning or nursery ground. Because the Bay serves as a nursery area for many coastal migratory fish, annual recruitment data are critical for multi-state management efforts along the Atlantic Coast. Furthermore, the VIMS Trawl Survey serves as a foundation to conduct research on basic biological characteristics of Bay and tributary fishes, as well as a platform from which emerging issues may be addressed.

LITERATURE CITED

- Able, K. W. and M. P. Fahay. 1998. The first year in the life of estuarine fishes in the middle Atlantic Bight. Rutgers University Press, New Jersey.
- ASMFC. 2023a. American Eel Benchmark Stock Assessment and Peer Review Report. Atlantic States Marine Fisheries Commission.
- ASMFC. 2023b. Summer Flounder 2023 Management Track Assessment Report. Atlantic States Marine Fisheries Commission.
- ASMFC. 2012. Review of the Atlantic States Marine Fisheries Commission Fishery Management Plan for Weakfish. Weakfish Plan Review Team. Atlantic States Marine Fisheries Commission.
- Bodolus, D. A. 1994. Mechanisms of larval Spot transport and recruitment to the Chesapeake Bay. Ph. D. Dissertation. College of William and Mary, Williamsburg, VA.
- Chao, L. N. and J. A. Musick. 1977. Life history, feeding habits, and functional morphology of juvenile sciaenid fishes in the York River estuary, Virginia. Fishery Bulletin 75:657-702.
- Chesapeake Executive Council. 1988. Chesapeake Bay Program Stock Assessment Plan. Agreement Commitment Report. Annapolis, MD.
- Chittenden, M. E., Jr. 1991. Evaluation of spatial/temporal sources of variation in nekton catch and the efficacy of stratified sampling in the Chesapeake Bay. Final report to Chesapeake Bay Stock Assessment Committee & NOAA/NMFS. Virginia Institute of Marine Science, Gloucester Point, VA.
- Cochran, W. G. 1977. Sampling techniques. John Wiley & Sons. New York.
- Colvocoresses, J. A. and P. J. Geer. 1991. Estimation of relative juvenile abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R1. July 1990 to June 1991. Virginia Institute of Marine Science, Gloucester Point, VA.
- Colvocoresses, J. A., P. J. Geer and C. F. Bonzek. 1992. Estimation of relative juvenile abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104-2. July 1991 to June 1992. Virginia Institute of Marine Science, Gloucester Point, VA.
- Connelly, W. J. 2001. Growth patterns of three species of catfish (*Ictaluridae*) from three Virginia tributaries of the Chesapeake Bay. Master's Thesis. College of William and Mary, Williamsburg, VA.

- Fabrizio, M. C., T. D. Tuckey, R. J. Latour, G. C. White, and A. J. Norris. 2018. Tidal habitats support large numbers of invasive Blue Catfish in a Chesapeake Bay subestuary. *Estuaries & Coasts* 41: 827-840.
- Fabrizio, M. C., V. Nepal, and T. D. Tuckey. 2021a. Invasive Blue Catfish in the Chesapeake Bay region: a case study of competing management objectives. *North American Journal of Fisheries Management* <http://dx.doi.org/10.1002/nafm.10552>
- Fabrizio, M. C., T. D. Tuckey, A. J. Bever, and M. L. MacWilliams. 2021b. The extent of seasonally suitable habitats may limit forage fish production in a temperate estuary. *Frontiers in Marine Science* Volume 8: Article 706666.
- Fabrizio, M. C., T. D. Tuckey, J. R. Buchanan, and R. A. Fisher. 2021c. Predation impacts of invasive blue catfish on blue crabs in estuarine environments. Final Report submitted to Virginia Marine Resources Commission
<https://scholarworks.wm.edu/cgi/viewcontent.cgi?article=3519&context=reports>
- Fenske, K. H., M. J. Wilberg, D. H. Secor, and M. C. Fabrizio. 2011. An age- and sex-structured assessment model for American Eels (*Anguilla rostrata*) in the Potomac River, Maryland. *Canadian Journal of Fisheries and Aquatic Sciences* 68: 1024-1037.
- Geer, P. J. and H. M. Austin. 1994. Estimation of relative abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R4. July 1993 to June 1994. Virginia Institute of Marine Science, Gloucester Point, VA.
- Geer, P. J. and H. M. Austin. 1996. Estimation of relative abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R6. July 1995 to June 1996. Virginia Institute of Marine Science, Gloucester Point, VA.
- Geer, P. J. and H. M. Austin. 1997. Estimation of relative abundance of recreationally important finfish in the Virginia portion of Chesapeake Bay. Annual report to VMRC/USFWS Sportfish Restoration Project F104R7. July 1996 to June 1997. Virginia Institute of Marine Science, Gloucester Point, VA.
- Geer, P. J., C. F. Bonzek, and H. M. Austin. 1994. Juvenile finfish and blue crab stock assessment program bottom trawl survey annual data summary report series. Volume 1993. Virginia Institute of Marine Science Special Scientific Report No. 124. Virginia Institute of Marine Science, Gloucester Point, VA.
- Goodyear, C. P. 1985. Relationship between reported commercial landings and abundance of young Striped Bass in Chesapeake Bay, Maryland. *Transactions of the American Fisheries Society* 114: 92-96.

- Haro, A., W. Richkus, K. Whalen, W. D. Busch, S. Lary, T. Brush, and D. Dixon. 2000. Population decline of the American Eel: Implications for research and management. *Fisheries* 25(9): 7-16.
- Hedgepeth, M. Y. 1983. Age, growth and reproduction of American Eels, *Anguilla rostrata* (Lesueur), from the Chesapeake Bay area. Master's Thesis. College of William and Mary, Williamsburg, VA.
- Homer, M. L. and J. A. Mihursky. 1991. Spot. Pages 11.1-11.19 in S.L. Funderburk, J.A. Mihursky, S. J. Jordan, and D. Reiley, eds., *Habitat requirements for Chesapeake Bay Living Resources*, 2nd Edition. Living Resources Subcommittee, Chesapeake Bay Program. Annapolis, MD.
- Jenkins, R. E. and N. M. Burkhead. 1993. *Freshwater fishes of Virginia*. American Fisheries Society, Bethesda, MD.
- Jung, S. 2002. Fish community structure and the temporal variability in recruitment and biomass production in Chesapeake Bay. Ph.D. Dissertation. University of Maryland, College Park, MD.
- Lowery, W. A. and P. J. Geer. 2000. Juvenile finfish and blue crab stock assessment program bottom trawl survey annual data summary report series. Volume 1999. Virginia Institute of Marine Science Special Scientific Report No. 124. Virginia Institute of Marine Science, Gloucester Point, VA.
- Marcek, B. J., R. Humston, M. C. Fabrizio, J. Shen, and R. W. Brill. 2024. Modeling the distribution of Atlantic croaker and spot in a dynamic seascape using metabolic scope. *Estuaries & Coasts* 47: 258-275.
- McCullagh, P. and J. A. Nelder. 1989. *Generalized linear models*, 2nd edition. Chapman & Hall, New York. (p. 124-125)
- Menzel, R. W. 1945. The catfishery of Virginia. *Transactions of the American Fisheries Society* 73: 364-372.
- Montane, M. M. and H. M. Austin. 2005. Effects of hurricanes on Atlantic Croaker (*Micropogonias undulatus*) recruitment to Chesapeake Bay. Pages 185-192 in K. Sellner, ed., *Hurricane Isabel in Perspective*. Chesapeake Research Consortium, CRC Publication 05-160, Edgewater, MD.
- Morse, W. W. 1978. Biological and fisheries data on Scup, *Stenotomus chrysops* (Linnaeus). National Marine Fisheries Service, Sandy Hook Laboratory, Technical Series Report No. 12.
- Murdy, E. O., R. S. Birdsong and J. A. Musick. 1997. *Fishes of Chesapeake Bay*. Smithsonian Institution Press.

- Musick, J. A. and L. P. Mercer. 1977. Seasonal distribution of Black Sea Bass, *Centropristis striata*, in the Mid-Atlantic Bight with comments on the ecology and fisheries of the species. Transactions of the American Fisheries Society 106:12-25.
- Nepal, V. and M. C. Fabrizio 2019. High salinity tolerance of invasive Blue Catfish suggests potential for further range expansion in the Chesapeake Bay region. PLoS ONE 14(11): e0224770
- Nepal, V. and M. C. Fabrizio. 2020. Density-dependence mediates the effects of temperature on growth of juvenile Blue Catfish in nonnative habitats. Transactions of the American Fisheries Society 149:108-120.
- Nepal, V. and M. C. Fabrizio. 2021. Reproductive Characteristics Differ in Two Invasive Populations of Blue Catfish. North American Journal of Fisheries Management published online <http://doi.org/10.1002/nafm.10611>
- Nepal, V., M. C. Fabrizio, and W. J. Connelly. 2020. Phenotypic plasticity in life-history characteristics of invasive Blue Catfish, *Ictalurus furcatus*. Fisheries Research 230: 105650.
- Nepal, V., M. C. Fabrizio, R. Lavaud, and J. van der Meer. 2024. Bioenergetic strategies contributing to the invasion success of blue catfish. Ecological Modelling 496: 110830.
- Norcross, B. L. 1983. Climate scale environmental factors affecting year-class fluctuations of Atlantic Croaker, *Micropogonias undulatus* in the Chesapeake Bay, VA. Ph.D Dissertation. College of William & Mary, Williamsburg, VA.
- Nys, L. N., M. C. Fabrizio, and T. D. Tuckey. 2015. Multi-decadal variation in size of juvenile Summer Flounder (*Paralichthys dentatus*) in Chesapeake Bay. Journal of Sea Research 103: 50-58.
- Owens, S. J. and P. J. Geer. 2003. Size and age structure of American Eels in tributaries of the Virginia portion of the Chesapeake Bay. Pages 117-124 in D. A. Dixon (Editor), Biology, Management and Protection of Catadromous Eels. American Fisheries Society Symposium Series 33, Bethesda, MD.
- Parthree, D. J., C. F. Bonzek and R. J. Latour. 2008. Chesapeake Bay Trophic Interactions Laboratory Services. Project NAO6NMF4570299. VIMS, Gloucester Point, VA.
- Rhodes, S. F. 1971. Age and growth of the Silver Perch *Bairdiella chrysoura*. Master's Thesis. College of William & Mary, Williamsburg, VA.
- Richards, R. A., and P. J. Rago. 1999. A case history of effective fishery management: Chesapeake Bay Striped Bass. North American Journal of Fisheries Management 19:356-375.

- Schloesser, R. W., and M. C. Fabrizio. 2015. Relationships among proximate components and energy density of juvenile Atlantic estuarine fishes. *Transactions of the American Fisheries Society* 145: 942-955.
- Schloesser, R. W., and M. C. Fabrizio. 2016. Temporal dynamics of condition for estuarine fishes in their nursery habitats. *Marine Ecology Progress Series* 557: 207-219.
- Schloesser, R. W., and M. C. Fabrizio. 2017. Condition indices as surrogates of energy density and lipid content in juveniles of three fish species. *Transactions of the American Fisheries Society* 146: 1058-1069.
- Schloesser, R. W., M. C. Fabrizio, R. J. Latour, G. C. Garman, B. Greenlee, M. Groves, and J. Gartland. 2011. Ecological role of Blue Catfish in Chesapeake Bay communities and implications for management. In P. Michaletz and V. Travnicek, eds., *Conservation, ecology, and management of worldwide catfish populations and habitats*. American Fisheries Society Symposium 77:369-382, Bethesda, MD.
- Schmitt, J. D., E. M. Hallerman, A. Bunch, Z. Moran, J. A. Emmel, and D. J. Orth. 2017. Predation and prey selectivity by nonnative catfish on migrating alosines in an Atlantic slope estuary. *Marine and Coastal Fisheries* 9: 108-125.
- Schmitt, J. D., B. K. Peoples, L. Castello, and D. J. Orth. 2019. Feeding ecology of generalist consumers: a case study of invasive blue catfish *Ictalurus furcatus* in Chesapeake Bay, Virginia, USA. *Environmental Biology of Fishes* 102:443-465.
- Sobocinski, K. L., R. J. Orth, M. C. Fabrizio, and R. J. Latour. 2013. Historical comparison of fish community structure in lower Chesapeake Bay seagrass habitats. *Estuaries and Coasts* 36:775-794.
- Szuwalski, C. S., K. A. Vert-Pre, A. E. Punt, T. A. Branch, and R. Hilborn. 2014. Examining common assumptions about recruitment: a meta-analysis of recruitment dynamics for worldwide marine fisheries. *Fish and Fisheries* DOI:10.1111/faf.12083.
- Terceiro, M. 2006. Summer Flounder assessment and biological reference point update for 2006. Northeast Fisheries Science Center Reference Document.
- Tuckey, T. D. and M. C. Fabrizio. 2013. Influence of survey design on fish assemblages: implications from a study in Chesapeake Bay tributaries. *Transactions of the American Fisheries Society* 142:957-973.
- Tuckey, T. D. and M. C. Fabrizio. 2016. Variability in fish tissue proximate composition is consistent with indirect effects of hypoxia in Chesapeake Bay tributaries. *Marine and Coastal Fisheries* 8:1-15.

- Tuckey, T. D. and M. C. Fabrizio. 2018. Estimating relative juvenile abundance of ecologically important finfish in the Virginia portion of Chesapeake Bay. 2018 Annual Report to the Virginia Marine Resources Commission. <https://doi.org/10.25773/mmjd-zd74>.
- Tuckey, T. D., J. Swinford, M. C. Fabrizio, H. Small, and J. Shields. 2021. Penaeid shrimp in Chesapeake Bay: population growth and black gill disease syndrome. *Marine and Coastal Fisheries* 13: 159–173. <https://doi.org/10.1002/mcf2.10143>.
- Wojcik, F. J. and W. A. Van Engel. 1988. A documentation of Virginia trawl surveys, 1955 – 1984, listing pertinent variables. Volume II – York River. College of William and Mary, VIMS, Gloucester Point, VA. 198 p. (cited in Table 2)
- Wood, R. J. 2000. Synoptic scale climatic forcing of multispecies recruitment patterns in Chesapeake Bay. Ph.D. Dissertation. College of William and Mary, Williamsburg, VA.
- Wood, R. J. and H. M. Austin. 2009. Synchronous multidecadal fish recruitment patterns in Chesapeake Bay, USA. *Canadian Journal of Fisheries and Aquatic Sciences* 66:496-508.
- Woodland, R. J., D. H. Secor, M. C. Fabrizio, and M. J. Wilberg. 2012. Comparing the nursery role of inner continental shelf and estuarine habitats for temperate marine fishes. *Estuarine, Coastal, and Shelf Science* 99: 61-73.
- Woodward, J. R. 2009. Investigating the relationships between recruitment indices and estimates of adult abundance for Striped Bass, Weakfish, and Atlantic Croaker. Master's Thesis. College of William and Mary, Williamsburg, VA.
- Wyanski, D. M. 1990. Patterns of habitat utilization in 0-age Summer Flounder (*Paralichthys dentatus*). Master's Thesis. College of William and Mary, Williamsburg, VA.

TABLES

Table 1. Spatial, temporal, and length (mm) criteria used to calculate recruitment indices. Highlighted boxes indicate strata, months and sizes used to calculate indices. Cross-hatched boxes indicate that only a portion of those strata are used in index calculations.

VIMS Trawl Survey - Area / Time / Size Values by Species																						
Species - Age	VIMS SP. CODE	Strata Used								Month												
		Bay		James		York		Rapp		Size Cut-off Values (mm) - Darkened Areas Represent Index Months												
		B	L	U	L	U	L	U	L	U	January	February	March	April	May	June	July	August	September	October	November	December
		O	O	P	O	P	O	P	O	P												
		T	T	W	P	W	P	W	P	W	P											
		O	E	E	E	E	E	E	E	E												
		M	R	R	R	R	R	R	R	R												
American Eel 1+	0060										---	---	---	>152	>152	>152	>152	>152	>152	---	---	---
Atlantic Croaker (spring)	0005										0-100	0-100	0-100	0-110	0-135	0-160	0-180	0-220	0-50	0-80	0-100	0-100
Bay Anchovy Y-O-Y	0103										0-77	0-80	0-80	0-80	0-80	0-80	0-44	0-51	0-56	0-61	0-65	0-70
Black Seabass Y-O-Y	0002										0-110	0-110	0-110	0-110	0-110	0-150	0-175	0-70	0-85	0-100	0-105	0-110
Blue Catfish Y-O-Y	0314										0-165	0-165	0-165	0-50	0-50	0-75	0-100	0-115	0-130	0-130	0-130	0-165
Blue Catfish 1+	0314										>165	>165	>165	>175	>225	>250	>250	>115	>130	>130	>130	>165
Channel Catfish Y-O-Y	0040										0-130	0-130	0-130	0-140	0-150	0-50	0-80	0-105	0-120	0-130	0-130	0-130
Channel Catfish 1+	0040										>130	>130	>130	>140	>150	>50	>80	>105	>120	>130	>130	>130
Scup	0001										90-170	90-170	90-170	90-170	35-90	40-100	50-125	60-145	75-160	85-170	90-170	90-170
Silver Perch Y-O-Y	0213										0-160	0-160	0-160	0-160	0-165	0-170	0-100	0-130	0-150	0-160	0-160	0-160
Spot Y-O-Y	0033										0-200	0-200	0-50	0-75	0-100	0-135	0-160	0-180	0-200	0-200	0-200	0-200
Striped Bass Y-O-Y	0031										0-200	0-200	0-200	0-200	0-50	0-80	0-100	0-120	0-135	0-150	0-175	0-190
Summer Flounder Y-O-Y	0003										0-290	0-290	0-60	0-100	0-140	0-170	0-200	0-225	0-250	0-275	0-290	0-290
Weakfish Y-O-Y	0007										0-200	0-200	0-200	0-225	0-240	0-90	0-120	0-150	0-180	0-200	0-200	0-200
White Catfish Y-O-Y	0039										0-110	0-110	0-110	0-110	0-120	0-50	0-65	0-80	0-90	0-100	0-110	0-110
White Catfish 1+	0039										>110	>110	>110	>110	>120	>50	>65	>80	>90	>100	>110	>110
White Perch Y-O-Y	0032										0-85	0-85	0-85	0-95	0-35	0-65	0-73	0-80	0-85	0-85	0-85	0-85
White Perch 1+	0032										>86	>86	>86	>96	>36	>66	>74	>81	>86	>85	>86	>86

Table 2. Sample collection history of the VIMS Trawl Survey, 1988 – June 2024. Each entry in the table represents the number of completed tows for the regular survey (not including Mobjack Bay); YR is year, TOT is total, STAT. TYPE is station type. Other codes are below and are based on Wojcik and Van Engel (1988), Appendices A – C.

YR	TOT	MONTH												WATER SYSTEM						Vessel			Gear		STAT. TYPE		TOW DURATION/DISTANCE			
		J	F	M	A	M	J	J	A	S	O	N	D	CL	JA	PO	RA	YK	ZZ	FH	TW	OT	070	108	F	R	5	OT	DIS	
1988	889	69	69	62	48	82	82	82	82	82	82	80	69	576	97	0	105	111	0	0	0	889	889	0	313	576	885	0	4	
1989	840	61	61	61	66	76	76	76	76	76	76	59	479	108	0	124	129	0	0	0	840	840	0	361	479	840	0	0		
1990	827	61	61	61	61	76	76	77	75	76	69	76	58	473	108	0	119	127	0	279	0	548	827	0	354	473	826	0	1	
1991	930	61	25	61	61	73	94	95	95	97	97	97	74	411	108	0	120	291	0	930	0	0	0	930	357	573	928	1	1	
1992	982	79	47	79	79	97	88	88	88	89	88	88	72	404	110	0	124	344	0	982	0	0	0	982	361	621	975	7	0	
1993	915	40	73	40	71	88	89	88	88	88	88	87	75	370	110	0	126	309	0	915	0	0	0	915	365	550	914	1	0	
1994	911	40	73	40	73	88	88	88	88	88	88	88	69	368	110	0	124	309	0	911	0	0	0	911	363	548	906	5	0	
1995	993	40	73	40	73	92	88	88	88	105	105	99	102	411	96	0	201	285	0	993	0	0	0	993	314	679	984	9	0	
1996	1176	52	91	71	106	106	107	108	108	107	108	107	105	435	228	0	258	255	0	1176	0	0	0	1176	279	897	1168	6	2	
1997	1220	68	105	66	98	110	111	111	112	111	112	111	105	425	265	0	264	266	0	1220	0	0	0	1220	302	918	1217	3	0	
1998	1262	66	105	66	105	111	111	128	59	138	124	130	119	388	265	0	256	264	89	1262	0	0	0	1262	322	940	1261	1	0	
1999	1382	79	122	80	122	120	118	119	118	122	124	131	127	402	264	0	264	265	187	1382	0	0	0	1382	363	1019	1380	2	0	
2000	1367	52	129	85	101	158	111	128	125	121	141	111	105	433	250	17	266	265	136	1367	0	0	0	1367	363	1004	1367	0	0	
2001	1122	30	30	30	75	112	144	111	112	135	136	111	96	384	230	35	230	230	13	1017	0	105	0	1122	277	845	1119	1	2	
2002	1090	66	90	66	90	96	106	96	97	95	96	96	96	288	264	0	264	264	10	1090	0	0	0	1090	300	790	1089	1	0	
2003	1191	66	96	66	96	96	111	111	111	111	111	111	105	399	264	0	264	264	0	1191	0	0	0	1191	300	891	1191	0	0	
2004	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0	
2005	1211	66	105	66	105	111	111	111	111	113	111	111	90	419	264	0	264	264	0	1211	0	0	0	1211	300	911	1211	0	0	
2006	1193	66	105	66	105	111	111	111	111	113	111	78	105	423	242	0	264	264	0	1193	0	0	0	1193	292	901	1193	0	0	
2007	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0	
2008	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0	
2009	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0	
2010	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	170	1224	0	0	0	1224	300	924	1224	0	0	
2011	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	204	1224	0	0	0	1224	300	924	1224	0	0	
2012	1205	66	105	66	105	111	111	111	111	111	111	92	105	432	264	0	264	264	148	1205	0	0	0	1205	300	905	1205	0	0	
2013	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0	
2014	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	1224	0	0	0	1224	300	924	1224	0	0	
2015	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	453	771	0	0	1224	300	924	1224	0	0	
2016	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	0	1224	0	0	0	1224	300	924	1224	0	0
2017	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	0	1224	0	0	0	1224	300	924	1224	0	0
2018	1053	0	0	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	0	1053	0	0	0	1053	250	803	1053	0	0
2019	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	0	1224	0	0	0	1224	300	924	1224	0	0
2020	918	66	105	66	0	0	61	91	91	111	111	111	105	298	200	0	200	220	0	0	918	0	0	0	918	234	684	918	0	0
2021	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	0	1224	0	0	0	1224	300	924	1224	0	0
2022	1224	66	105	66	105	111	111	111	111	111	111	111	105	432	264	0	264	264	0	0	1224	0	0	0	1224	300	924	1224	0	0
2023	1202	66	105	66	105	111	111	111	111	111	111	111	83	432	242	0	264	264	0	0	1202	0	0	0	1202	292	910	1202	0	0
2024	564	66	105	66	105	111	111							168	132	0	132	132	0	0	564	0	0	0	564	102	462	564	0	0
TOT	41,579	2,250	3,350	2,360	3,425	3,801	3,881	3,794	3,722	3,865	3,865	3,767	3,499	15,130	8,049	52	8,325	9,214	957	28,569	10,064	2,382	2,556	38,459	11,264	30,315	40,968	37	10	

System: CL - Lower Chesapeake Bay (Virginia Portion), JA - James River, PO - Potomac River, RA - Rappahannock River, YK - York River, ZZ - includes: Atlantic Ocean (AT) - 1971, 78-79, 2002; Piankatank R. (PK) - 1970-71, 98-00; Mobjack Bay (MB) - 1970-73, 98-01, 10-12; Pocomoke Sound (CP) -1973-81, 98-01; Great Wicomico R. (GW) - 1998-00. **Vessel:** FH - Fish Hawk, TW - Tidewater, OT – Other **Gear Code:** 30' Gear 070 - Lined, tickler chain, 60' bridle, 54"x24" doors, 108 - Lined, tickler chain, 60' bridle, metal china-v doors. **Station Type:** F – Fixed, R – Random. **Tow type:** OT is tow duration in minutes for those not listed. DIS is distance.

Table 3. VIMS Trawl Survey pooled catch for July 2023 to June 2024 from 1,202 tows.

Adjusted Percent of Catch Excludes Bay Anchovy and Hogchoker

Species	Number of Fish (All)	Frequency	Percent of Catch	Catch Per Trawl	Adjusted Percent of Catch	Number of Fish YOY	Average Length (mm)	Standard Error (length)	Minimum Length (mm)	Maximum Length (mm)
Bay Anchovy	394,545	971	61.08	328.24	.	358,137	53	0.07	10	109
Spot	84,439	765	13.07	70.25	41	70,466	112	0.24	10	208
Atlantic Croaker	57,088	655	8.84	47.49	27.72	45,892	102	0.37	8	240
Hogchoker	45,406	637	7.03	37.78	.	12,957	82	0.2	12	173
Weakfish	16,451	475	2.55	13.69	7.99	15,538	91	0.46	7	367
Atlantic Menhaden	12,119	347	1.88	10.08	5.88	5,787	69	0.56	20	215
Silver Perch	7,142	360	1.11	5.94	3.47	6,889	103	0.35	30	195
Blue Catfish	5,824	231	0.9	4.85	2.83	384	254	1.2	54	705
White Perch	5,321	202	0.82	4.43	2.58	1,348	137	0.8	25	295
Striped Anchovy	3,460	202	0.54	2.88	1.68	3,396	72	0.42	29	129
Blackcheek Tonguefish	3,128	388	0.48	2.6	1.52	767	121	0.61	40	179
Southern Kingfish	1,279	190	0.2	1.06	0.62	1,128	111	1.53	10	339
Spotted Hake	1,142	137	0.18	0.95	0.55	1,141	83	0.86	33	231
Summer Flounder	1,020	347	0.16	0.85	0.5	830	175	1.82	22	517
Blueback Herring	899	42	0.14	0.75	0.44	895	71	0.72	31	261
Atlantic Cutlassfish	599	132	0.09	0.5	0.29	.	224	4.3	62	700
Atlantic Silverside	522	48	0.08	0.43	0.25	522	88	0.72	68	113
Butterfish	468	100	0.07	0.39	0.23	386	105	1.08	20	177
Black Sea Bass	442	163	0.07	0.37	0.21	255	112	1.4	36	217
Atlantic Moonfish	394	55	0.06	0.33	0.19	.	72	1.37	30	153
White Catfish	382	99	0.06	0.32	0.19	61	181	3.31	52	461
Northern Searobin	363	101	0.06	0.3	0.18	358	95	1.45	26	161
Striped Bass	338	57	0.05	0.28	0.16	319	79	4.31	21	476
Harvestfish	311	84	0.05	0.26	0.15	302	59	1.8	12	162
Atlantic Herring	300	16	0.05	0.25	0.15	.	51	0.85	31	82
Star Drum	270	33	0.04	0.22	0.13	.	79	1	36	152
Alewife	252	27	0.04	0.21	0.12	252	64	1.27	39	129
Banded Drum	250	43	0.04	0.21	0.12	.	94	3.74	23	220
American Shad	219	62	0.03	0.18	0.11	219	85	1.06	58	145
Oyster Toadfish	185	63	0.03	0.15	0.09	.	131	5.55	29	351
Atlantic Thread Herring	177	48	0.03	0.15	0.09	.	104	2.47	39	178
Smallmouth Flounder	125	69	0.02	0.1	0.06	83	91	1.77	41	131
Pigfish	112	49	0.02	0.09	0.05	.	130	1.65	96	185
Striped Searobin	101	46	0.02	0.08	0.05	.	61	1.92	28	141
Inshore Lizardfish	84	52	0.01	0.07	0.04	52	190	6.36	43	316
Gizzard Shad	79	27	0.01	0.07	0.04	52	199	10.22	94	413
Atlantic Spadefish	63	36	0.01	0.05	0.03	.	93	3.4	30	151
Northern Pipefish	52	47	0.01	0.04	0.03	.	154	6.23	51	235
Threadfin Shad	47	23	0.01	0.04	0.02	.	102	2.04	75	132
Hickory Shad	43	20	0.01	0.04	0.02	.	85	4.3	50	169
Chain Pipefish	37	33	0.01	0.03	0.02	.	243	8.29	120	325
Northern Puffer	34	21	0.01	0.03	0.02	20	115	9.17	19	193
Lined Seahorse	32	28	0	0.03	0.02	.	78	5.87	19	158
American Eel	29	24	0	0.02	0.01	.	302	16.3	168	561
Windowpane	26	21	0	0.02	0.01	25	109	5.26	55	160
Black Drum	22	15	0	0.02	0.01	.	205	30.16	144	772
Spiny Dogfish	22	14	0	0.02	0.01	.	638	11.28	546	755
Seaboard Goby	20	17	0	0.02	0.01	.	34	1.85	23	50
Naked Goby	17	16	0	0.01	0.01	.	43	1.9	33	57
Longnose Gar	17	13	0	0.01	0.01	.	625	49.74	348	886
Bluefish	15	8	0	0.01	0.01	.	204	10.21	118	258
Sea Lamprey	12	8	0	0.01	0.01	.	159	3.68	134	181

Table 3. (continued)

Species	Number of Fish (All)	Percent Frequency of Catch	Catch Per Trawl	Adjusted Percent of Catch	Number of Fish YOY	Average Length (mm)	Standard Error (length)	Minimum Length (mm)	Maximum Length (mm)	
Clearnose Skate	11	10	0	0.01	0.01	.	422	8.81	377	452
Spotted Seatrout	11	8	0	0.01	0.01	.	166	14.83	107	250
Skilletfish	11	8	0	0.01	0.01	.	42	3.81	17	61
Pinfish	11	7	0	0.01	0.01	.	129	4.95	86	152
Striped Cusk-eel	11	7	0	0.01	0.01	.	140	3.94	117	156
Atlantic Stingray	10	10	0	0.01	0	.	321	41.21	164	497
Fringed Flounder	10	9	0	0.01	0	.	109	6.09	67	137
Spottail Shiner	10	4	0	0.01	0	.	77	1.97	66	86
Feather Blenny	7	7	0	0.01	0	.	57	7.92	35	80
Orange Filefish	7	7	0	0.01	0	.	129	23.82	62	238
Guaguanche	7	5	0	0.01	0	.	122	18.33	55	183
Striped Burrfish	6	6	0	0	0	.	175	11.42	143	226
Spotfin Mojarra	6	5	0	0	0	.	95	6.04	78	118
Blue Runner	6	4	0	0	0	.	113	3.84	102	129
Striped Mullet	5	1	0	0	0	.	232	14.22	182	270
Red Drum	4	4	0	0	0	.	332	22.3	282	384
Spiny Butterfly Ray	4	4	0	0	0	.	595	34.49	497	651
Scup	4	3	0	0	0	3	51	2.1	45	55
Bluntnose Stingray	4	3	0	0	0	.	330	125.62	173	705
Sandbar Shark	3	3	0	0	0	.	560	32.25	521	624
Tessellated Darter	3	2	0	0	0	.	64	1	63	66
Lookdown	3	2	0	0	0	.	100	22.81	55	129
Spanish Mackerel	2	2	0	0	0	.	173	13.5	159	186
Tautog	2	2	0	0	0	.	185	2	183	187
Northern Kingfish	2	2	0	0	0	1	157	47.5	109	204
Banded Killifish	2	2	0	0	0	.	46	1.5	44	47
Green Goby	2	2	0	0	0	.	42	5	37	47
Halfbeak	2	2	0	0	0	.	67	1	66	68
Silver Seatrout	2	2	0	0	0	.	190	4.5	185	194
Creville Jack	2	1	0	0	0	.	144	0	144	144
Speckled Worm Eel	2	1	0	0	0	.	362	18	344	380
Silver Hake	1	1	0	0	0	.	158	.	158	158
Cobia	1	1	0	0	0	.	265	.	265	265
Winter Flounder	1	1	0	0	0	.	339	.	339	339
Common Carp	1	1	0	0	0	.	600	.	600	600
Sheepshead	1	1	0	0	0	.	136	.	136	136
Florida Pompano	1	1	0	0	0	.	166	.	166	166
Bighead Searobin	1	1	0	0	0	.	27	.	27	27
Brown Bullhead	1	1	0	0	0	.	163	.	163	163
Mummichog	1	1	0	0	0	.	66	.	66	66
Rough Silverside	1	1	0	0	0	.	45	.	45	45
Smooth Dogfish	1	1	0	0	0	.	345	.	345	345
Smooth Butterfly Ray	1	1	0	0	0	.	335	.	335	335
Cownose Ray	1	1	0	0	0	.	699	.	699	699
Bullnose Ray	1	1	0	0	0	.	574	.	574	574
Conger Eel	1	1	0	0	0	.	437	.	437	437
Atlantic Sturgeon	1	1	0	0	0	.	885	.	885	885
Northern Stargazer	1	1	0	0	0	.	37	.	37	37
Rock Seabass	1	1	0	0	0	.	192	.	192	192
Green Sunfish	1	1	0	0	0	.	90	.	90	90
Gray Snapper	1	1	0	0	0	.	95	.	95	95
Gulf Kingfish	1	1	0	0	0	.	129	.	129	129
Red Snapper	1	1	0	0	0	.	70	.	70	70
Freshwater Goby	1	1	0	0	0	.	74	.	74	74

All Species Combined 645,919

Table 4. American Eel indices (RSI_{Δ} ; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. ** Indicates we were unable to calculate an index for 2020 in the Rappahannock and James rivers because of a pause in sampling due to COVID-19. Average values calculated from 1988 to 2022.

Year	Rappahannock			York			James		
	Index	Prop. pos.	N	Index	Prop. pos.	N	Index	Prop. pos.	N
1988	2.31	0.20	35	1.27	0.33	40	2.32	0.30	30
1989	11.82	0.37	43	3.50	0.27	49	6.08	0.37	38
1990	13.34	0.40	43	4.90	0.30	50	9.69	0.42	38
1991	4.89	0.38	42	0.64	0.18	49	1.82	0.26	38
1992	1.95	0.28	43	0.83	0.19	47	8.99	0.42	38
1993	1.87	0.30	43	0.33	0.16	49	5.74	0.42	38
1994	3.45	0.40	43	0.33	0.16	49	2.21	0.37	38
1995	2.83	0.37	43	0.33	0.18	49	1.74	0.37	46
1996	2.54	0.36	128	0.58	0.25	126	3.90	0.41	126
1997	2.71	0.45	132	0.47	0.19	132	1.77	0.36	132
1998	2.02	0.31	124	0.48	0.19	132	1.91	0.35	132
1999	0.71	0.23	132	0.23	0.14	133	1.16	0.31	132
2000	1.38	0.32	133	0.24	0.16	133	0.87	0.28	132
2001	0.58	0.18	133	0.16	0.14	133	0.58	0.23	134
2002	0.28	0.16	132	0.24	0.15	132	0.73	0.23	132
2003	0.61	0.20	132	0.14	0.11	132	0.57	0.23	132
2004	0.44	0.25	132	0.14	0.11	132	0.46	0.16	132
2005	0.14	0.11	132	0.09	0.05	132	0.26	0.17	132
2006	0.08	0.05	132	0.04	0.04	132	0.14	0.11	132
2007	0.20	0.11	132	0.08	0.06	132	0.11	0.10	132
2008	0.47	0.22	132	0.21	0.17	132	0.17	0.13	132
2009	0.48	0.17	132	0.14	0.12	132	0.33	0.16	132
2010	0.51	0.21	132	0.16	0.14	132	0.19	0.11	132
2011	0.85	0.30	132	0.10	0.10	132	0.34	0.17	132
2012	0.85	0.27	132	0.12	0.11	132	0.18	0.09	132
2013	0.96	0.25	132	0.16	0.12	132	0.43	0.14	132
2014	0.55	0.20	132	0.18	0.16	132	0.22	0.14	132
2015*	0.32	0.17	132	0.08	0.08	132	0.11	0.10	132
2016*	0.30	0.18	132	0.07	0.07	132	0.09	0.10	132
2017*	0.18	0.14	132	0.05	0.04	132	0.15	0.11	132
2018*	0.23	0.10	132	0.06	0.06	132	0.07	0.08	132
2019*	0.25	0.13	132	0.04	0.04	132	0.08	0.06	132
2020**	-	-	-	0.02	0.02	88	-	-	-
2021*	0.18	0.11	132	0.05	0.04	132	0.03	0.04	132
2022*	0.07	0.06	132	0.03	0.02	132	0.03	0.03	132
2023*	0.01	0.02	132	0.10	0.05	132	0.05	0.05	132
Average	1.78			0.47			1.57		

Table 5. Spring Atlantic Croaker indices (RSI_{Delta}; 1987–2023). ^ indicates the 2008 Index if a single large catch of Atlantic Croaker in the Bay is included; *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015; ** Indicates we were unable to calculate an index for the 2019 year-class because of a pause in sampling due to COVID-19. Average values calculated from 1987 to 2022.

Year Class	Year	Mean all	LCI all	UCI all
1987	1988	0.95	0.27	2.30
1988	1989	14.14	1.43	36.26
1989	1990	6.40	0.60	12.01
1990	1991	28.39	4.93	113.38
1991	1992	2.80	2.17	3.51
1992	1993	7.22	3.27	13.37
1993	1994	0.52	0.26	0.83
1994	1995	2.06	1.25	3.05
1995	1996	0.03	0.01	0.05
1996	1997	65.51	8.67	218.30
1997	1998	12.68	8.42	23.21
1998	1999	4.98	2.46	10.37
1999	2000	1.17	0.70	1.70
2000	2001	1.55	0.21	6.37
2001	2002	7.64	4.61	10.79
2002	2003	0.90	0.07	2.39
2003	2004	4.36	2.90	5.95
2004	2005	2.72	1.59	4.84
2005	2006	9.46	5.79	16.17
2006	2007	6.36	4.18	9.66
2007	2008	28.06	22.21	36.08
2008	2009	10.21	6.58	15.75
^		114.71	7.32	555.36
2009	2010	29.07	6.63	73.25
2010	2011	4.43	1.36	8.36
2011	2012	56.20	39.68	84.44
2012	2013	23.23	12.01	51.44
2013	2014	56.56	2.95	522.71
2014	2015	4.74	0.76	20.11
2015*	2016	27.41	3.58	79.57
2016*	2017	15.19	6.83	34.43
2017*	2018	0.61	0.27	1.33
2018*	2019	15.64	7.29	38.18
2019**	2020	-	-	-
2020*	2021	6.87	3.45	11.60
2021*	2022	26.70	8.70	83.02
2022*	2023	50.15	29.75	79.08
2023*	2024	93.77	50.88	178.59
Average		15.28		

Table 6. Bay Anchovy indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from 1988 to 2022.

Random Stratified Index (RSI)				
Year	Geo.			
	Mean	95% C.I.'s	C.V.	N
1988	18.25	12.17 - 27.15	6.42	346
1989	52.47	36.27 - 75.71	4.54	374
1990	6.79	4.41 - 10.22	8.89	369
1991	22.51	15.05 - 33.43	6.04	491
1992	40.14	27.17 - 59.09	5.10	448
1993	43.31	28.80 - 64.89	5.23	449
1994	14.67	9.93 - 21.46	6.54	444
1995	18.36	12.84 - 26.07	5.66	540
1996	15.31	11.20 - 20.82	5.21	607
1997	18.96	13.63 - 26.23	5.19	625
1998	30.26	20.75 - 43.93	5.27	579
1999	15.47	11.20 - 21.22	5.35	606
2000	36.58	26.69 - 49.99	4.21	619
2001	9.55	6.93 - 13.04	6.06	627
2002	5.51	3.58 - 8.24	9.36	540
2003	18.03	13.17 - 24.56	5.01	624
2004	23.06	16.71 - 31.70	4.82	624
2005	22.27	16.01 - 30.85	4.98	613
2006	19.31	14.00 - 26.50	5.03	592
2007	23.76	17.33 - 32.44	4.69	624
2008	50.29	36.21 - 69.68	4.07	624
2009	30.12	22.30 - 40.55	4.21	624
2010	84.92	61.27 - 117.54	3.61	624
2011	26.56	19.20 - 36.59	4.68	624
2012	51.53	36.84 - 71.92	4.14	611
2013	23.76	16.92 - 33.19	5.03	624
2014	5.18	3.69 - 7.14	7.57	624
2015*	8.73	6.34 - 11.90	6.20	624
2016*	11.72	8.82 - 15.48	5.09	624
2017*	27.56	20.40 - 37.12	4.30	624
2018*	17.79	13.00 - 24.20	5.01	624
2019*	10.16	7.75 - 13.73	5.76	624
2020*	19.12	14.05 - 25.91	4.84	586
2021*	13.09	9.27 - 18.34	5.99	624
2022*	7.61	5.57 - 10.29	6.28	624
2023*	9.39	6.98 - 12.52	5.63	602
Average	24.08			

Table 7. Black Sea Bass indices (RSI_{GM}; Year Class 1988–2022). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015; ** Indicates we were unable to calculate an index for 2020 because of a pause in sampling due to COVID-19. Average values calculated from year class 1988 to 2022.

Year Class	Year	Random Stratified Index (RSI)			
		Geo. Mean	95% C.I.'s	C.V.	N
1988	1989	0.84	0.59 - 1.13	11.89	138
1989	1990	2.36	1.70 - 3.17	8.93	138
1990	1991	1.12	0.78 - 1.53	11.63	128
1991	1992	1.28	0.91 - 1.72	10.76	129
1992	1993	0.22	0.13 - 0.32	18.86	129
1993	1994	1.05	0.74 - 1.42	11.46	129
1994	1995	1.06	0.74 - 1.45	11.85	129
1995	1996	0.50	0.33 - 0.69	14.47	151
1996	1997	0.36	0.22 - 0.52	17.99	152
1997	1998	0.46	0.31 - 0.63	14.63	153
1998	1999	0.57	0.35 - 0.82	16.40	135
1999	2000	0.58	0.41 - 0.77	12.22	146
2000	2001	0.74	0.50 - 1.02	13.39	153
2001	2002	1.29	0.85 - 1.84	12.89	108
2002	2003	0.64	0.41 - 0.90	15.16	138
2003	2004	0.12	0.06 - 0.18	25.11	153
2004	2005	0.06	0.02 - 0.10	34.69	153
2005	2006	0.19	0.12 - 0.26	17.66	153
2006	2007	0.44	0.30 - 0.60	14.14	153
2007	2008	0.83	0.53 - 1.18	14.68	153
2008	2009	0.41	0.27 - 0.57	14.90	153
2009	2010	0.32	0.19 - 0.47	19.23	153
2010	2011	1.11	0.83 - 1.43	9.41	153
2011	2012	0.65	0.47 - 0.85	11.61	153
2012	2013	0.19	0.11 - 0.28	19.21	153
2013	2014	0.30	0.20 - 0.42	16.21	153
2014*	2015	0.15	0.09 - 0.21	19.18	153
2015*	2016	0.21	0.14 - 0.29	16.07	153
2016*	2017	0.54	0.38 - 0.71	12.28	153
2017*	2018	0.08	0.04 - 0.12	26.02	153
2018*	2019	0.60	0.43 - 0.79	11.82	153
2019**	2020	-	-	-	-
2020*	2021	0.60	0.43 - 0.78	11.87	153
2021*	2022	0.83	0.57 - 1.13	12.78	153
2022*	2023	0.74	0.53 - 0.98	11.47	153
Average		0.63			

Table 8. Blue Catfish juvenile indices (RSI_{Δ} ; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year Class	Rappahannock			York			James		
	Index	Prop. pos.	N	Index	Prop. pos.	N	Index	Prop. pos.	N
1988	0.00	0.00	34	0.00	0.00	36	0.00	0.00	24
1989	0.00	0.00	32	0.00	0.00	36	0.00	0.00	20
1990	0.00	0.00	30	0.00	0.00	32	0.00	0.00	22
1991	0.00	0.00	34	0.00	0.00	34	0.00	0.00	24
1992	0.00	0.00	34	0.00	0.00	32	0.00	0.00	24
1993	0.00	0.00	32	0.00	0.00	36	0.00	0.00	22
1994	0.00	0.00	32	0.00	0.00	36	0.00	0.00	24
1995	0.00	0.00	36	0.00	0.00	36	0.14	0.07	28
1996	0.17	0.02	92	0.00	0.00	82	6.41	0.41	78
1997	0.00	0.00	48	0.00	0.00	45	0.00	0.00	44
1998	0.00	0.00	48	0.00	0.00	45	0.16	0.04	46
1999	0.00	0.00	48	0.00	0.00	45	0.49	0.14	43
2000	0.00	0.00	49	0.00	0.00	45	0.07	0.04	45
2001	0.00	0.00	49	0.00	0.00	46	0.04	0.02	46
2002	0.00	0.00	48	0.00	0.00	45	0.16	0.02	45
2003	10.40	0.08	48	2.97	0.18	45	18.89	0.42	45
2004	0.25	0.06	48	18.56	0.22	45	7.93	0.53	45
2005	0.00	0.00	48	0.04	0.02	45	1.56	0.20	45
2006	0.02	0.02	48	0.56	0.07	45	1.51	0.37	30
2007	0.00	0.00	48	0.00	0.00	45	0.16	0.04	45
2008	0.00	0.00	48	1.69	0.04	45	0.00	0.00	45
2009	0.00	0.00	48	0.00	0.00	45	0.27	0.09	45
2010	0.00	0.00	48	0.04	0.04	45	0.00	0.00	45
2011	6.74	0.17	48	1.05	0.13	45	2.61	0.27	45
2012	0.00	0.00	48	6.12	0.09	45	0.00	0.00	45
2013	0.11	0.08	48	1.80	0.09	45	4.98	0.22	45
2014	0.40	0.04	48	1.14	0.09	45	0.12	0.04	45
2015*	0.08	0.04	48	1.81	0.13	45	0.20	0.02	45
2016*	0.00	0.00	48	3.11	0.09	45	2.94	0.13	45
2017*	0.00	0.00	48	0.07	0.02	45	0.04	0.04	45
2018*	7.75	0.33	48	9.89	0.33	45	44.30	0.58	45
2019*	0.40	0.08	48	0.86	0.13	45	0.19	0.04	45
2020*	0.02	0.02	48	12.72	0.11	45	8.11	0.27	45
2021*	0.56	0.06	48	0.56	0.09	45	0.00	0.00	45
2022*	0.00	0.00	48	0.07	0.02	45	0.00	0.00	45
2023*	0.00	0.00	48	0.00	0.00	48	0.00	0.00	48
Average	0.77			1.80			2.89		

Table 9. Blue Catfish age 1+ indices (RSI_{Δ} ; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year	Rappahannock			York			James		
	Index	Prop. pos.	N	Index	Prop. pos.	N	Index	Prop. pos.	N
1988	0.00	0.00	34	0.00	0.00	36	0.00	0.00	24
1989	0.00	0.00	32	0.00	0.00	36	0.30	0.10	20
1990	0.00	0.00	30	0.00	0.00	32	0.00	0.00	22
1991	0.00	0.00	34	0.00	0.00	34	0.00	0.00	24
1992	0.00	0.00	34	0.00	0.00	32	0.00	0.00	24
1993	0.00	0.00	32	0.00	0.00	36	0.00	0.00	22
1994	1.70	0.25	32	0.00	0.00	36	0.00	0.00	24
1995	0.31	0.11	36	0.00	0.00	36	8.97	0.29	28
1996	36.31	0.33	92	0.00	0.00	82	4.30	0.51	78
1997	137.30	0.23	48	0.02	0.02	45	33.76	0.32	44
1998	56.81	0.25	48	0.00	0.00	45	27.01	0.35	46
1999	7.91	0.17	48	0.04	0.04	45	29.11	0.63	43
2000	19.18	0.29	49	0.00	0.00	45	17.44	0.53	45
2001	6.30	0.20	49	0.04	0.04	46	4.56	0.28	46
2002	1.08	0.08	48	0.02	0.02	45	2.91	0.22	45
2003	34.66	0.63	48	0.76	0.22	45	7.26	0.69	45
2004	17.97	0.48	48	1.73	0.29	45	47.27	0.78	45
2005	37.61	0.25	48	3.44	0.18	45	47.07	0.51	45
2006	48.95	0.33	48	1.13	0.16	45	39.19	0.73	30
2007	1.95	0.15	48	0.80	0.11	45	22.66	0.47	45
2008	3.03	0.19	48	1.05	0.11	45	11.39	0.51	45
2009	13.61	0.42	48	0.49	0.11	45	38.20	0.56	45
2010	8.40	0.29	48	1.35	0.13	45	16.45	0.53	45
2011	17.18	0.73	48	1.59	0.40	45	30.45	0.71	45
2012	17.37	0.27	48	2.07	0.16	45	28.86	0.47	45
2013	69.93	0.42	48	1.19	0.18	45	24.69	0.67	45
2014	32.72	0.44	48	2.30	0.20	45	15.86	0.58	45
2015*	124.35	0.35	48	2.98	0.33	45	58.84	0.64	45
2016*	27.51	0.27	48	5.95	0.18	45	109.37	0.60	45
2017*	12.83	0.31	48	3.63	0.18	45	54.18	0.47	45
2018*	15.29	0.71	48	11.92	0.56	45	56.37	0.87	45
2019*	35.07	0.35	48	6.56	0.20	45	317.77	0.42	45
2020*	48.11	0.50	48	3.80	0.40	45	47.74	0.78	45
2021*	12.57	0.31	48	6.18	0.16	45	60.75	0.60	45
2022*	4.28	0.19	48	1.96	0.09	45	54.42	0.44	45
2023*	9.70	0.21	48	2.46	0.16	45	16.20	0.44	45
Average	24.29			1.74			34.78		

Table 10. Channel Catfish juvenile indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year	Rappahannock				York				James			
	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.
1988	0.07	9	35.33	0.02 - 0.13	0.00	12		0	1.03	8	11.03	0.74 - 1.38
1989	0.31	9	25.18	0.14 - 0.50	0.00	12		0	0.11	6	100.00	0.00 - 0.35
1990	0.18	7	16.88	0.12 - 0.25	0.00	10		0	0.25	6	40.98	0.04 - 0.50
1991	0.05	9	100.00	0.00 - 0.15	0.00	11		0	0.00	6		0
1992	0.07	9	66.96	0.00 - 0.17	0.00	10		0	0.05	6	100.00	0.00 - 0.16
1993	0.03	9	65.63	0.00 - 0.07	0.03	12	68.31	0.00 - 0.08	0.02	6	100.00	0.00 - 0.07
1994	0.10	9	47.85	0.00 - 0.21	0.01	12	100.00	0.00 - 0.03	0.04	6	100.00	0.00 - 0.13
1995	0.01	9	100.00	0.00 - 0.03	0.00	12		0	1.09	10	12.96	0.72 - 1.52
1996	0.19	27	18.12	0.12 - 0.26	0.02	26	83.63	0.00 - 0.05	0.46	27	16.28	0.29 - 0.65
1997	0.07	30	52.70	0.00 - 0.16	0.01	30	68.29	0.00 - 0.01	0.37	30	30.09	0.13 - 0.65
1998	0.05	30	47.56	0.00 - 0.09	0.00	30		0	0.21	30	40.41	0.04 - 0.42
1999	0.03	30	71.98	0.00 - 0.08	0.00	30	66.14	0.00 - 0.01	0.36	30	31.85	0.12 - 0.66
2000	0.02	31	43.32	0.00 - 0.04	0.00	29	100.00	0.00 - 0.01	0.10	30	52.35	0.00 - 0.21
2001	0.05	31	85.40	0.00 - 0.14	0.00	30		0	0.06	30	67.50	0.00 - 0.15
2002	0.06	30	93.64	0.00 - 0.12	0.00	30	100.00	0.00 - 0.01	0.03	30	80.93	0.00 - 0.09
2003	0.04	30	82.71	0.00 - 0.12	0.00	30		0	0.13	30	55.85	0.00 - 0.29
2004	0.01	30	77.15	0.00 - 0.03	0.00	30		0	0.12	30	37.83	0.03 - 0.21
2005	0.00	30		0	0.00	30		0	0.03	30	54.57	0.00 - 0.07
2006	0.02	30	88.27	0.00 - 0.07	0.00	30		0	0.02	20	89.12	0.00 - 0.06
2007	0.00	30	71.24	0.00 - 0.01	0.00	30		0	0.05	30	76.69	0.03 - 0.14
2008	0.00	30		0	0.00	30		0	0.01	30	79.68	0.00 - 0.02
2009	0.00	30		0	0.00	30		0	0.00	30	100.00	0.00 - 0.01
2010	0.00	30		0	0.00	30		0	0.00	30		0
2011	0.00	30		0	0.00	30		0	0.04	30	93.85	0.00 - 0.13
2012	0.00	30		0	0.00	30		0	0.00	30		0
2013	0.02	30	100.00	0.00 - 0.06	0.00	30		0	0.00	30		0
2014	0.00	30		0	0.00	30		0	0.00	30		0
2015*	0.00	30		0	0.00	30		0	0.00	30		0
2016*	0.00	30		0	0.00	30		0	0.00	30		0
2017*	0.00	30		0	0.00	30		0	0.00	30		0
2018*	0.00	30		0	0.00	30		0	0.01	30	78.80	0.00 - 0.03
2019*	0.00	30		0	0.00	30		0	0.00	30		0
2020*	0.00	30		0	0.00	30		0	0.00	30		0
2021*	0.00	30		0	0.00	30		0	0.00	30		0
2022*	0.00	30		0	0.00	30		0	0.00	30		0
2023*	0.00	30		0	0.00	30		0	0.00	30		0
Average	0.04				0.00				0.13			

Table 11. Channel Catfish age 1+ indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year	Rappahannock				York				James			
	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.
1988	0.00	9		0	0.00	12		0	0.00	8		0
1989	0.01	9	100.00	0.00 - 0.04	0.00	12		0	0.05	6	100.00	0.00 - 0.16
1990	0.00	7		0	0.00	10		0	0.00	6		0
1991	0.00	9		0	0.00	11		0	0.00	6		0
1992	0.05	9	100.00	0.00 - 0.17	0.00	10		0	0.00	6		0
1993	0.00	9		0	0.00	12		0	0.00	6		0
1994	0.00	9		0	0.01	12	100.00	0.00 - 0.03	0.00	6		0
1995	0.00	9		0	0.00	12		0	0.51	10	0.00	0.51 - 0.51
1996	0.00	27		0	0.00	26		0	0.10	27	67.76	0.00 - 0.26
1997	0.00	30		0	0.00	30		0	0.00	30		0
1998	0.00	30		0	0.01	30	70.71	0.00 - 0.02	0.00	30		0
1999	0.00	30		0	0.00	30		0	0.00	30		0
2000	0.00	31	100.00	0.00 - 0.01	0.00	29		0	0.00	30	100.00	0.00 - 0.01
2001	0.00	31		0	0.00	30		0	0.00	30	100.00	0.00 - 0.01
2002	0.00	30		0	0.00	30		0	0.00	30		0
2003	0.00	30	100.00	0.00 - 0.01	0.02	30	58.88	0.00 - 0.04	0.17	30	37.27	0.04 - 0.32
2004	0.00	30	100.00	0.00 - 0.01	0.00	30	66.14	0.00 - 0.01	0.01	30	100.00	0.00 - 0.02
2005	0.00	30		0	0.00	30		0	0.00	30		0
2006	0.00	30		0	0.00	30		0	0.00	20		0
2007	0.00	30		0	0.00	30		0	0.00	30		0
2008	0.00	30		0	0.00	30		0	0.00	30		0
2009	0.00	30		0	0.00	30		0	0.00	30		0
2010	0.00	30		0	0.00	30		0	0.00	30		0
2011	0.00	30		0	0.00	30		0	0.00	30		0
2012	0.00	30		0	0.00	30		0	0.00	30		0
2013	0.00	30		0	0.00	30		0	0.04	30	100.00	0.00 - 0.12
2014	0.00	30		0	0.00	30		0	0.00	30	100.00	0
2015*	0.00	30		0	0.00	30		0	0.00	30		0
2016*	0.00	30		0	0.00	30		0	0.00	30		0
2017*	0.00	30		0	0.00	30		0	0.00	30		0
2018*	0.00	30	100.00	0.00 - 0.01	0.00	30		0	0.00	30		0
2019*	0.00	30		0	0.00	30		0	0.00	30		0
2020*	0.00	30		0	0.00	30		0	0.00	30		0
2021*	0.00	30		0	0.00	30		0	0.00	30		0
2022*	0.00	30		0	0.00	30		0	0.00	30		0
2023*	0.00	30		0	0.00	30		0	0.00	30		0
Average	0.00				0.00				0.03			

Table 12. Scup indices (RSI_{GM} ; 1988–2022). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2021.

Year Class	Random Stratified Index (RSI)			
	Geo. Mean	95% C.I.'s	C.V.	N
1988	3.06	2.05 - 4.41	10.20	112
1989	4.92	3.14 - 7.45	10.03	112
1990	1.90	1.11 - 2.99	14.99	103
1991	0.65	0.41 - 0.93	15.67	104
1992	3.36	2.16 - 5.01	10.90	104
1993	0.90	0.53 - 1.35	16.67	104
1994	0.39	0.21 - 0.59	21.36	104
1995	0.54	0.29 - 0.83	20.37	104
1996	0.21	0.09 - 0.35	28.00	104
1997	0.50	0.27 - 0.75	19.83	79
1998	0.27	0.06 - 0.52	37.91	88
1999	0.13	0.02 - 0.25	41.14	105
2000	1.34	0.88 - 1.90	12.80	111
2001	0.24	0.11 - 0.37	24.52	64
2002	0.96	0.58 - 1.42	15.89	104
2003	0.46	0.28 - 0.67	17.38	104
2004	1.11	0.71 - 1.59	13.89	104
2005	1.58	0.99 - 2.36	13.77	104
2006	2.99	2.07 - 4.19	9.47	104
2007	0.20	0.09 - 0.31	25.12	104
2008	2.97	2.07 - 4.13	9.28	104
2009	4.11	2.79 - 5.89	9.14	104
2010	0.82	0.51 - 1.20	15.70	104
2011	0.22	0.07 - 0.39	33.06	104
2012	0.74	0.46 - 1.06	15.48	104
2013	0.16	0.08 - 0.25	25.35	104
2014*	1.15	0.68 - 1.76	16.14	104
2015*	0.32	0.09 - 0.60	34.75	104
2016*	0.25	0.12 - 0.39	24.99	104
2017*	0.78	0.44 - 1.20	18.35	104
2018*	0.81	0.47 - 1.22	17.55	104
2019*	0.18	0.06 - 0.31	31.36	104
2020*	0.19	0.08 - 0.31	26.82	104
2021*	0.21	0.05 - 0.41	38.54	104
2022*	0.00	0.00 - 0.01	100.00	104
Average	1.14			

Table 13. Silver Perch indices (RSI_{GM} ; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from 1988 to 2022.

Year	Random Stratified Index (RSI)			
	Geo. Mean	95% C.I.'s	C.V.	N
1988	0.61	0.35 - 0.92	18.30	172
1989	0.53	0.33 - 0.76	16.32	189
1990	0.69	0.49 - 0.92	11.94	185
1991	0.35	0.21 - 0.51	17.33	179
1992	0.81	0.49 - 1.18	15.80	178
1993	0.45	0.29 - 0.63	16.01	180
1994	0.25	0.11 - 0.40	25.42	180
1995	0.58	0.34 - 0.87	15.65	180
1996	0.59	0.38 - 0.84	15.63	304
1997	0.71	0.50 - 0.94	12.07	316
1998	0.24	0.15 - 0.33	16.77	316
1999	0.70	0.49 - 0.94	12.42	309
2000	0.68	0.46 - 0.93	13.56	317
2001	0.70	0.47 - 0.97	13.77	327
2002	0.44	0.24 - 0.67	20.16	269
2003	0.63	0.40 - 0.90	15.49	315
2004	0.34	0.22 - 0.48	16.50	315
2005	0.76	0.52 - 1.03	12.64	315
2006	1.21	0.84 - 1.64	11.31	283
2007	0.75	0.50 - 1.03	13.53	315
2008	0.49	0.34 - 0.66	13.31	315
2009	1.00	0.72 - 1.32	10.83	315
2010	1.27	0.95 - 1.65	9.29	315
2011	0.77	0.53 - 1.04	12.41	315
2012	0.80	0.55 - 1.08	12.38	302
2013	0.59	0.41 - 0.79	13.16	315
2014	1.14	0.76 - 1.60	12.82	315
2015*	0.58	0.44 - 0.74	10.36	315
2016*	1.78	1.30 - 2.35	9.19	315
2017*	1.40	0.98 - 1.91	11.06	315
2018*	1.44	1.07 - 1.87	9.13	315
2019*	0.37	0.26 - 0.48	12.76	315
2020*	0.58	0.44 - 0.73	10.19	315
2021*	0.47	0.33 - 0.62	12.98	315
2022*	0.45	0.33 - 0.59	11.69	315
2023*	1.29	0.95 - 1.68	9.56	315
Average	0.72			

Table 14. Spot indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from 1988 to 2022.

Year	Random Stratified Index (RSI)				
	Geo.	Mean	95% C.I.'s	C.V.	N
1988		67.01	46.36 - 96.67	4.29	231
1989		31.41	24.51 - 40.18	3.44	252
1990		44.78	32.34 - 61.85	4.14	248
1991		16.83	12.28 - 21.60	4.66	238
1992		1.92	1.45 - 2.49	8.20	238
1993		9.78	7.23 - 13.13	5.68	240
1994		9.23	6.88 - 12.27	5.61	240
1995		1.56	1.15 - 2.05	9.25	248
1996		5.26	4.15 - 6.60	5.30	407
1997		11.50	9.11 - 14.45	4.20	421
1998		2.51	1.92 - 3.23	7.36	374
1999		4.72	3.63 - 6.07	6.07	402
2000		3.32	2.57 - 4.23	6.51	421
2001		3.09	2.45 - 3.85	6.06	432
2002		2.89	2.10 - 3.88	8.38	360
2003		2.85	2.25 - 3.56	6.32	420
2004		3.96	3.14 - 4.95	5.68	420
2005		12.12	9.80 - 14.94	3.78	420
2006		3.37	2.71 - 4.16	5.61	420
2007		9.17	7.38 - 11.35	4.18	420
2008		19.89	15.16 - 26.01	4.22	420
2009		6.08	4.96 - 7.40	4.39	420
2010		74.97	59.30 - 94.70	2.67	420
2011		5.29	4.22 - 6.57	5.05	420
2012		17.18	13.49 - 21.82	3.92	420
2013		12.38	10.14 - 15.08	3.54	420
2014		3.26	2.60 - 4.04	5.84	420
2015*		0.83	0.61 - 1.07	10.50	420
2016*		2.39	1.90 - 2.96	6.36	420
2017*		4.46	3.57 - 5.54	5.27	420
2018*		14.70	11.76 - 18.33	3.77	420
2019*		7.11	5.55 - 9.05	5.12	420
2020*		10.53	8.43 - 13.08	4.10	382
2021*		5.60	4.54 - 6.87	4.64	420
2022*		9.65	7.48 - 12.37	4.81	420
2023*		15.03	12.37 - 18.22	3.27	420
Average		12.62			

Table 15. Striped Bass indices (RSI_{GM} ; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year class	Random Stratified Index (RSI)				
	Geo.	Mean	95% C.I.'s	C.V.	N
1988		0.18	0.17 - 0.20	3.72	38
1989		0.02	0.00 - 0.05	100.00	36
1990		0.02	0.00 - 0.06	100.00	32
1991		0.16	0.00 - 0.48	78.57	34
1992		0.00	0		34
1993		0.21	0.07 - 0.36	30.97	36
1994		0.10	0.00 - 0.32	100.00	36
1995		0.00	0		42
1996		0.68	0.10 - 1.58	40.79	107
1997		0.26	0.04 - 0.52	41.26	120
1998		0.30	0.13 - 0.49	26.33	112
1999		0.16	0.00 - 0.43	72.11	120
2000		0.72	0.36 - 1.18	21.88	120
2001		1.04	0.54 - 1.68	19.44	121
2002		0.54	0.34 - 0.77	15.92	120
2003		0.49	0.16 - 0.90	31.15	120
2004		0.79	0.35 - 1.36	24.13	120
2005		0.47	0.18 - 0.83	28.91	120
2006		0.38	0.07 - 0.78	40.11	110
2007		0.21	0.00 - 0.48	51.04	120
2008		0.68	0.14 - 1.47	37.01	120
2009		0.47	0.06 - 1.04	42.96	120
2010		0.30	0.04 - 0.62	42.08	120
2011		1.15	0.60 - 1.88	19.15	120
2012		0.09	0.00 - 0.22	62.94	120
2013		0.47	0.14 - 0.90	33.16	120
2014		0.45	0.19 - 0.77	26.42	120
2015*		0.74	0.34 - 1.27	23.83	120
2016*		0.17	0.00 - 0.55	91.33	120
2017*		0.35	0.08 - 0.68	36.54	120
2018*		0.99	0.60 - 1.48	16.00	120
2019*		0.34	0.00 - 0.82	52.28	120
2020*		0.10	0.00 - 0.25	67.67	102
2021*		0.14	0.00 - 0.36	69.75	120
2022*		0.27	0.00 - 0.75	66.87	120
2023*		0.24	0.00 - 0.59	57.76	120
Average		0.38			

Table 16. Summer Flounder indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from 1988 to 2022.

Year	Random Stratified Index (RSI)			
	Geo. Mean	95% C.I.'s	C.V.	N
1988	0.54	0.35 - 0.75	14.99	143
1989	1.24	0.94 - 1.58	8.77	162
1990	2.54	2.06 - 3.09	5.73	162
1991	2.79	2.26 - 3.41	5.66	153
1992	0.92	0.70 - 1.17	9.25	153
1993	0.52	0.38 - 0.68	11.87	153
1994	2.54	2.01 - 3.15	6.39	153
1995	0.71	0.52 - 0.92	10.89	149
1996	0.81	0.62 - 1.02	9.32	224
1997	0.89	0.69 - 1.12	8.77	226
1998	0.73	0.55 - 0.93	9.92	226
1999	0.53	0.41 - 0.67	9.94	219
2000	0.57	0.43 - 0.73	10.81	227
2001	0.47	0.34 - 0.61	11.84	236
2002	0.77	0.54 - 1.04	12.21	179
2003	0.44	0.33 - 0.56	10.95	225
2004	1.30	1.03 - 1.60	7.50	225
2005	0.35	0.25 - 0.46	13.18	225
2006	0.80	0.60 - 1.02	10.03	203
2007	1.00	0.78 - 1.24	8.22	225
2008	1.35	1.10 - 1.63	6.68	225
2009	0.75	0.58 - 0.92	8.76	225
2010	0.55	0.41 - 0.69	10.61	225
2011	0.17	0.11 - 0.23	17.54	225
2012	2.03	1.69 - 2.40	5.29	212
2013	0.82	0.65 - 1.02	8.42	225
2014	0.62	0.49 - 0.77	8.94	225
2015*	0.23	0.15 - 0.31	16.09	225
2016*	0.42	0.30 - 0.56	13.09	225
2017*	0.95	0.75 - 1.18	8.03	225
2018*	1.10	0.87 - 1.36	7.84	225
2019*	0.39	0.28 - 0.52	13.24	225
2020*	0.08	0.04 - 0.13	24.58	225
2021*	0.22	0.16 - 0.30	14.08	225
2022*	0.43	0.28 - 0.91	16.03	225
2023*	0.81	0.60 - 1.05	10.54	225
Average	0.87			

Table 17. Weakfish indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from 1988 to 2022.

Year	Random Stratified Index (RSI)			
	Geo. Mean	95% C.I.'s	C.V.	N
1988	8.13	5.37 - 12.07	8.12	173
1989	11.74	8.18 - 16.88	6.44	189
1990	4.46	3.10 - 6.26	8.44	184
1991	3.16	2.32 - 4.21	7.92	179
1992	6.78	4.74 - 9.53	7.39	178
1993	5.81	4.06 - 8.17	7.76	180
1994	2.51	1.76 - 3.47	9.59	180
1995	5.95	4.26 - 8.18	7.20	186
1996	7.26	5.33 - 9.78	6.31	305
1997	6.81	5.26 - 8.74	5.38	316
1998	7.60	5.46 - 10.45	6.65	269
1999	6.78	5.01 - 9.06	6.28	303
2000	8.35	6.34 - 10.92	5.42	316
2001	5.09	3.74 - 6.82	6.93	327
2002	6.93	4.27 - 10.94	9.89	270
2003	9.23	6.72 - 12.54	6.04	315
2004	6.66	4.94 - 8.88	6.24	315
2005	5.69	4.26 - 7.50	6.31	315
2006	6.34	4.83 - 8.25	5.80	315
2007	5.35	3.99 - 7.08	6.51	315
2008	5.77	4.33 - 7.60	6.26	315
2009	6.18	4.75 - 7.96	5.63	315
2010	14.11	1.16 - 17.78	4.00	315
2011	5.23	3.86 - 6.99	6.78	315
2012	3.02	2.30 - 3.90	7.14	315
2013	9.41	7.30 - 12.07	4.85	315
2014	3.77	2.91 - 4.82	6.39	315
2015*	1.59	1.17 - 2.09	9.36	315
2016*	1.44	1.06 - 1.88	9.42	315
2017*	2.41	1.76 - 3.22	8.70	315
2018*	2.80	2.09 - 3.68	7.80	315
2019*	1.02	0.76 - 1.32	9.68	315
2020**	2.36	1.76 - 3.08	8.08	297
2021*	0.66	0.51 - 0.82	9.35	315
2022*	1.57	1.14 - 2.09	9.64	315
2023*	2.35	1.79 - 3.01	7.52	315
Average	5.60			

Table 18. White Catfish juvenile indices (RSI_{GM}; 1988–2023). Juvenile White Catfish indices were not adjusted as too few were captured. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year class	Rappahannock				York				James			
	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.
1988	0.00	9		0	0.03	12	100.00	0.06 - 0.10	0.08	8	100.00	0.00 - 0.27
1989	0.01	9	100.00	0.00 - 0.04	0.13	12	48.32	0.00 - 0.26	0.58	6	61.32	0.00 - 1.78
1990	0.00	7		0	0.02	10	100.00	0.00 - 0.05	0.00	6		0
1991	0.00	9		0	0.00	11		0	0.00	6		0
1992	0.00	9		0	0.00	10		0	0.00	6		0
1993	0.00	9		0	0.08	12	52.52	0.00 - 0.16	0.00	6		0
1994	0.00	9		0	0.03	12	65.63	0.00 - 0.07	0.00	6		0
1995	0.00	9		0	0.01	12	100.00	0.00 - 0.02	0.38	10	0.00	0.38 - 0.39
1996	0.02	27	100.00	0.00 - 0.06	0.03	26	34.00	0.01 - 0.05	0.18	27	46.25	0.01 - 0.37
1997	0.00	30		0	0.01	30	72.57	0.00 - 0.02	0.06	30	52.32	0.00 - 0.14
1998	0.00	30		0	0.01	30	64.14	0.00 - 0.03	0.01	30	100.00	0.00 - 0.03
1999	0.00	30		0	0.00	30		0	0.00	30		0
2000	0.00	31		0	0.01	29	65.47	0.00 - 0.01	0.00	30		0
2001	0.00	31		0	0.00	30		0	0.00	30		0
2002	0.00	30		0	0.00	30		0	0.00	30		0
2003	0.00	30	69.46	0.00 - 0.01	0.05	30	22.40	0.03 - 0.07	0.01	30	72.05	0.00 - 0.03
2004	0.00	30		0	0.00	30		0	0.10	30	37.89	0.02 - 0.18
2005	0.00	30		0	0.00	30		0	0.00	30		0
2006	0.00	30		0	0.00	30		0	0.00	20		0
2007	0.00	30		0	0.00	30		0	0.00	30		0
2008	0.00	30		0	0.00	30		0	0.00	30		0
2009	0.00	30		0	0.00	30		0	0.09	30	74.02	0.00 - 0.24
2010	0.00	30		0	0.00	30		0	0.00	30		0
2011	0.00	30		0	0.00	30		0	0.00	30		0
2012	0.00	30		0	0.00	30		0	0.00	30		0
2013	0.00	30		0	0.00	30		0	0.00	30		0
2014	0.00	30		0	0.00	30		0	0.04	30	100.00	0.00 - 0.12
2015	0.00	30		0	0.00	30		0	0.07	30	89.85	0.00 - 0.22
2016	0.00	30		0	0.00	30		0	0.01	30	78.33	0.00 - 0.03
2017	0.00	30		0	0.00	30		0	0.00	30		0
2018	0.00	30		0	0.01	30	72.16	0.00 - 0.01	0.14	30	25.53	0.07 - 0.21
2019	0.00	30		0	0.00	30		0	0.01	30	100.00	0.00 - 0.04
2020	0.00	30		0	0.00	30	100.00	0.00 - 0.01	0.09	30	30.68	0.03 - 0.15
2021	0.00	30		0	0.00	30		0	0.00	30		0
2022	0.00	30		0	0.00	30		0	0.02	30	79.33	0.00 - 0.04
2023	0.00	30		0	0.00	30		0	0.00	30		0
Average	0.00				0.01				0.05			

Table 19. White Catfish age 1+ indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year	Rappahannock				York				James			
	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.
1988	0.18	9	20.26	0.10 - 0.26	0.18	12	28.57	0.07 - 0.30	0.20	8	100.00	0.00 - 0.75
1989	0.87	9	30.00	0.28 - 1.72	0.56	12	20.74	0.30 - 0.88	0.99	6	27.08	0.37 - 1.89
1990	0.32	7	44.20	0.03 - 0.70	0.46	10	22.67	0.23 - 0.74	1.83	6	14.17	1.11 - 2.81
1991	0.22	9	26.85	0.10 - 0.36	0.15	11	42.79	0.02 - 0.29	0.53	6	29.45	0.19 - 0.97
1992	0.52	9	20.32	0.28 - 0.80	0.37	10	21.08	0.20 - 0.57	0.63	6	48.39	0.02 - 1.67
1993	0.14	9	32.97	0.05 - 0.24	0.28	12	19.89	0.16 - 0.41	0.66	6	92.83	0.00 - 3.26
1994	1.06	9	22.36	0.49 - 1.84	0.28	12	22.53	0.15 - 0.43	0.06	6	50.00	0.00 - 0.13
1995	0.19	9	22.49	0.10 - 0.28	0.12	12	28.23	0.05 - 0.20	1.31	10	12.99	0.86 - 1.87
1996	0.43	27	18.32	0.25 - 0.62	0.21	26	15.74	0.14 - 0.29	0.79	27	12.82	0.54 - 1.08
1997	0.24	30	29.41	0.09 - 0.41	0.08	30	23.61	0.04 - 0.12	0.37	30	25.96	0.16 - 0.61
1998	0.13	30	24.05	0.07 - 0.20	0.05	30	23.28	0.03 - 0.07	0.22	30	34.26	0.06 - 0.40
1999	0.09	30	22.66	0.05 - 0.14	0.12	30	19.85	0.07 - 0.18	0.48	30	26.71	0.20 - 0.82
2000	0.10	31	39.54	0.02 - 0.19	0.06	29	30.44	0.02 - 0.11	0.05	30	34.98	0.02 - 0.09
2001	0.03	31	42.82	0.00 - 0.06	0.03	30	27.95	0.01 - 0.05	0.14	30	29.65	0.06 - 0.23
2002	0.02	30	52.04	0.00 - 0.04	0.03	30	23.05	0.02 - 0.05	0.16	30	30.14	0.06 - 0.27
2003	0.14	30	57.67	0.00 - 0.31	0.08	30	23.47	0.04 - 0.11	0.12	30	30.75	0.05 - 0.21
2004	0.11	30	48.09	0.00 - 0.23	0.12	30	17.33	0.08 - 0.16	0.24	30	31.44	0.08 - 0.43
2005	0.03	30	72.33	0.00 - 0.07	0.05	30	34.59	0.01 - 0.08	0.15	30	36.31	0.04 - 0.28
2006	0.02	30	58.93	0.00 - 0.04	0.04	30	32.04	0.02 - 0.07	0.10	20	48.38	0.00 - 0.21
2007	0.01	30	54.81	0.00 - 0.03	0.03	30	29.47	0.01 - 0.05	0.11	30	41.79	0.02 - 0.21
2008	0.03	30	74.14	0.00 - 0.07	0.03	30	30.29	0.01 - 0.05	0.05	30	43.00	0.01 - 0.09
2009	0.02	30	43.73	0.00 - 0.05	0.02	30	35.53	0.01 - 0.04	0.18	30	26.92	0.08 - 0.29
2010	0.02	30	49.58	0.00 - 0.04	0.02	30	45.48	0.00 - 0.04	0.41	30	33.55	0.12 - 0.78
2011	0.06	30	63.45	0.00 - 0.13	0.04	30	30.66	0.02 - 0.06	0.24	30	27.36	0.10 - 0.39
2012	0.00	30		0	0.03	30	31.31	0.01 - 0.05	0.18	30	31.16	0.06 - 0.30
2013	0.01	30	56.41	0.00 - 0.02	0.03	30	25.57	0.01 - 0.04	0.19	30	26.69	0.09 - 0.31
2014	0.01	30	77.60	0.00 - 0.02	0.03	30	28.06	0.01 - 0.05	0.34	30	36.83	0.08 - 0.66
2015*	0.01	30	47.92	0.00 - 0.01	0.02	30	40.80	0.00 - 0.03	0.20	30	27.07	0.09 - 0.32
2016*	0.00	30		0	0.01	30	57.20	0.00 - 0.01	0.13	30	28.22	0.05 - 0.21
2017*	0.00	30		0	0.00	30		0	0.16	30	30.82	0.06 - 0.27
2018*	0.04	30	74.17	0.00 - 0.04	0.02	30	26.98	0.01 - 0.04	0.17	30	19.92	0.10 - 0.25
2019*	0.00	30	100.00	0.00 - 0.01	0.02	30	36.25	0.00 - 0.03	0.23	30	39.07	0.05 - 0.44
2020*	0.03	30	95.04	0.00 - 0.08	0.01	30	43.40	0.00 - 0.03	0.32	30	19.71	0.18 - 0.47
2021*	0.01	30	70.20	0.00 - 0.02	0.01	30	48.80	0.00 - 0.02	0.19	30	30.09	0.07 - 0.32
2022*	0.00	30	67.42	0.00 - 0.00	0.00	30	100.00	0.00 - 0.01	0.16	30	30.54	0.06 - 0.27
2023*	0.00	30	100.00	0.00 - 0.01	0.02	30	45.69	0.00 - 0.03	0.21	30	19.97	0.12 - 0.30
Average	0.15				0.10				0.35			

Table 20. White Perch juvenile indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year	Rappahannock				York				James			
	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.
1988	0.00	9		0	0.01	12	100.00	0.00 - 0.03	0.00	6		0
1989	0.03	9	100.00	0.00 - 0.09	0.12	12	25.98	0.06 - 0.19	0.00	6		0
1990	0.00	7		0	0.00	10		0	0.00	6		0
1991	0.00	8		0	0.00	12		0	0.00	6		0
1992	0.01	9	100.00	0.00 - 0.04	0.00	10		0	0.00	6		0
1993	0.00	9		0	0.00	12		0	0.00	6		0
1994	0.00	9		0	0.00	12		0	0.00	6		0
1995	0.00	9		0	0.00	12		0	0.31	12	22.94	0.16 - 0.48
1996	0.12	27	81.05	0.00 - 0.33	0.05	27	28.81	0.02 - 0.08	5.00	27	5.59	3.91 - 6.33
1997	0.00	30		0	0.00	30		0	1.14	30	32.02	0.31 - 2.47
1998	0.00	22		0	0.02	30	80.74	0.00 - 0.05	1.98	30	16.21	1.09 - 3.24
1999	0.00	30		0	0.00	30		0	0.21	30	52.77	0.00 - 0.48
2000	0.00	31		0	0.03	29	60.08	0.00 - 0.07	1.52	30	8.45	1.15 - 1.94
2001	0.02	31	100.00	0.00 - 0.06	0.01	30	100.00	0.00 - 0.03	1.94	30	15.60	1.10 - 3.11
2002	0.00	30		0	0.00	30		0	0.08	30	50.00	0.00 - 0.17
2003	1.53	30	15.04	0.92 - 2.35	0.05	30	26.57	0.02 - 0.08	3.59	30	9.67	2.42 - 5.16
2004	0.12	30	66.00	0.00 - 0.31	0.03	30	34.37	0.01 - 0.05	2.90	30	9.05	2.05 - 4.00
2005	0.07	30	59.16	0.00 - 0.16	0.01	30	100.00	0.00 - 0.04	2.08	30	10.97	1.41 - 2.95
2006	0.00	30		0	0.00	30		0	1.62	30	11.27	1.11 - 2.26
2007	0.00	30		0	0.00	30		0	0.48	30	45.36	0.04 - 1.11
2008	0.00	30	100.00	0.00 - 0.01	0.00	30		0	0.20	30	86.54	0.00 - 0.63
2009	0.02	30	100.00	0.00 - 0.06	0.00	30		0	0.73	30	49.39	0.01 - 1.97
2010	0.00	30		0	0.01	30	100.00	0.00 - 0.02	0.48	30	17.00	0.30 - 0.70
2011	0.56	30	20.33	0.30 - 0.86	0.04	30	32.87	0.01 - 0.07	2.13	30	9.73	1.50 - 2.90
2012	0.00	30		0	0.00	30		0	0.23	30	69.54	0.00 - 0.63
2013	0.00	30		0	0.02	30	47.48	0.00 - 0.04	1.54	30	19.15	0.78 - 2.63
2014	0.03	30	81.99	0.00 - 0.07	0.07	30	36.80	0.02 - 0.13	0.93	30	20.83	0.47 - 1.54
2015*	0.05	30	96.24	0.00 - 0.16	0.08	30	28.86	0.03 - 0.12	0.95	30	25.39	0.39 - 1.74
2016*	0.00	30		0	0.01	30	50.00	0.00 - 0.01	1.22	30	13.30	0.80 - 1.75
2017*	0.00	30		0	0.00	30		0	1.29	30	21.59	0.60 - 2.27
2018*	1.55	30	26.39	0.55 - 3.16	0.08	30	39.85	0.01 - 0.14	4.99	30	13.71	2.66 - 8.78
2019*	0.14	30	55.17	0.00 - 0.31	0.04	30	22.28	0.02 - 0.06	0.84	30	12.93	0.57 - 1.16
2020*	0.00	22		0	0.00	30		0	1.00	20	34.51	0.24 - 2.23
2021*	0.04	30	62.97	0.00 - 0.09	0.01	30	68.64	0.00 - 0.02	0.04	30	100.00	0.00 - 0.13
2022*	0.09	30	55.92	0.00 - 0.20	0.00	30		0	0.00	30		0
2023*	0.00	30		0	0.00	30		0	0.04	30	100.00	0.00 - 0.13
Average	0.13				0.02				1.13			

Table 21. White Perch age 1+ indices (RSI_{GM}; 1988–2023). Note: *Indices were adjusted with a calibration factor to account for gear and vessel changes that occurred in 2015. Average values calculated from year class 1988 to 2022. All indices have been recalculated to reflect the new recruitment window.

Year	Rappahannock				York				James			
	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.	RSI	N	CV	95% C.I.
1988	0.40	9	13.74	0.27 - 0.53	0.06	12	70.44	0.00 - 0.15	0.00	6		0
1989	0.83	9	40.27	0.12 - 1.97	0.19	12	12.55	0.14 - 0.24	1.44	6	28.92	0.46 - 3.09
1990	0.29	7	59.98	0.00 - 0.74	0.30	10	22.99	0.15 - 0.46	0.52	6	43.09	0.06 - 1.17
1991	0.17	8	23.80	0.09 - 0.26	0.05	12	47.83	0.00 - 0.11	0.09	6	77.06	0.00 - 0.25
1992	0.35	9	34.23	0.10 - 0.66	0.05	10	60.62	0.00 - 0.10	0.33	6	73.84	0.00 - 1.04
1993	0.10	9	44.89	0.01 - 0.20	0.06	12	33.97	0.02 - 0.10	0.00	6		0
1994	0.24	9	29.15	0.09 - 0.40	0.05	12	63.25	0.00 - 0.12	0.00	6		0
1995	0.20	9	10.98	0.15 - 0.25	0.01	12	100.00	0.00 - 0.02	2.28	12	8.73	1.66 - 3.03
1996	1.26	27	13.57	0.81 - 1.81	0.10	27	23.58	0.05 - 0.16	1.73	27	12.84	1.11 - 2.53
1997	0.39	30	32.33	0.12 - 0.71	0.02	30	47.03	0.00 - 0.04	2.20	30	14.16	1.30 - 3.44
1998	0.88	22	63.40	0.00 - 3.21	0.03	30	36.03	0.01 - 0.04	1.53	30	14.86	0.92 - 2.34
1999	1.24	30	20.88	0.60 - 2.14	0.03	30	56.64	0.00 - 0.06	3.05	30	15.05	1.66 - 5.16
2000	0.81	31	21.07	0.41 - 1.32	0.06	29	38.31	0.01 - 0.10	1.85	30	25.95	0.65 - 3.9
2001	0.26	31	36.81	0.06 - 0.49	0.02	30	36.48	0.01 - 0.04	1.99	30	17.27	1.05 - 3.37
2002	0.10	30	48.47	0.00 - 0.21	0.05	30	41.05	0.01 - 0.09	1.03	30	13.43	0.68 - 1.45
2003	2.42	30	8.46	1.77 - 3.21	0.14	30	20.71	0.08 - 0.20	3.06	30	6.89	2.35 - 3.93
2004	1.02	30	19.96	0.53 - 1.68	0.10	30	21.50	0.06 - 0.15	2.36	30	18.62	1.14 - 4.27
2005	0.55	30	14.48	0.36 - 0.76	0.03	30	48.33	0.00 - 0.07	0.96	30	29.87	0.31 - 1.94
2006	0.88	30	18.30	0.49 - 1.36	0.02	30	42.19	0.00 - 0.04	1.15	30	14.95	0.71 - 1.70
2007	0.49	30	55.70	0.00 - 1.33	0.01	30	70.71	0.00 - 0.02	0.28	30	39.34	0.05 - 0.55
2008	0.73	30	32.46	0.21 - 1.47	0.01	30	46.70	0.00 - 0.03	0.50	30	52.19	0.00 - 1.30
2009	0.83	30	32.70	0.23 - 1.71	0.01	30	41.23	0.00 - 0.02	1.08	30	38.08	0.19 - 2.64
2010	0.59	30	36.02	0.14 - 1.22	0.00	30		0	0.65	30	13.51	0.44 - 0.89
2011	0.99	30	16.98	0.58 - 1.52	0.01	30	53.94	0.00 - 0.02	1.66	30	18.44	0.86 - 2.82
2012	0.84	30	23.81	0.38 - 1.47	0.03	30	42.86	0.00 - 0.05	0.89	30	22.33	0.42 - 1.51
2013	0.95	30	29.07	0.32 - 1.88	0.02	30	51.81	0.00 - 0.05	1.34	30	15.07	0.81 - 2.02
2014	0.62	30	31.96	0.19 - 1.21	0.05	30	27.91	0.02 - 0.08	1.25	30	18.07	0.68 - 2.02
2015*	0.62	30	50.57	0.00 - 1.65	0.06	30	28.75	0.03 - 0.10	2.03	30	9.96	1.43 - 2.78
2016*	0.45	30	45.94	0.03 - 1.04	0.03	30	54.04	0.00 - 0.07	2.11	30	12.33	1.35 - 3.11
2017*	0.91	30	26.21	0.36 - 1.69	0.01	30	51.81	0.00 - 0.02	1.57	30	26.88	0.55 - 3.27
2018*	1.47	30	29.14	0.46 - 3.17	0.06	30	43.91	0.01 - 0.11	2.66	30	15.49	1.45 - 4.47
2019*	0.62	30	24.86	0.28 - 1.06	0.07	30	29.63	0.03 - 0.12	0.97	30	16.52	0.58 - 1.47
2020*	0.42	22	35.57	0.11 - 0.82	0.02	30	38.39	0.01 - 0.04	1.60	20	33.58	0.37 - 3.93
2021*	0.20	30	50.06	0.00 - 4.40	0.03	30	24.44	0.02 - 0.05	0.60	30	9.07	0.47 - 0.74
2022*	0.22	30	44.00	0.02 - 0.44	0.01	30	45.13	0.00 - 0.03	0.11	30	40.15	0.02 - 0.20
2023*	2.42	30	8.46	1.77 - 3.21	0.05	30	41.05	0.01 - 0.09	3.06	30	6.89	2.35 - 3.93
Average	0.67				0.05				1.28			

FIGURES

Figure 1. The VIMS Trawl Survey random stratified design in the Chesapeake Bay. Transect lines indicate geographic regions as designated below.

Chesapeake Bay	B1	Bottom Bay
	B2	Lower Bay
	B3	Upper Bay
James River	J1	Bottom James
	J2	Lower James
	J3	Upper James
	J4	Top James
York River	Y1	Bottom York
	Y2	Lower York
	Y3	Upper York
	Y4	Top York (lower Pamunkey River)
Rappahannock River	R1	Bottom Rappahannock
	R2	Lower Rappahannock
	R3	Upper Rappahannock
	R4	Top Rappahannock
Mobjack Bay	MB	Routine monitoring established March 2010 and ending December 2012




Figure 1 (continued)




Figure 2. Weighted mean bottom water temperature (°C) and 95% confidence interval (shaded region) measured at all sites in the Rappahannock, York, and James rivers, and the Virginia portion of Chesapeake Bay from 1988 to December 2023.




Figure 3. Weighted mean bottom salinity (psu) and 95% confidence interval (shaded region) measured at all sites in the Rappahannock, York, and James rivers, and the Virginia portion of Chesapeake Bay from 1988 to December 2023.




Figure 4. Weighted mean bottom dissolved oxygen (mg/L) and 95% confidence interval (shaded region) measured at all sites in the Rappahannock, York, and James rivers, and the Virginia portion of Chesapeake Bay from 1988 to December 2023.




Figure 5. American Eel random stratified index (RSI_{Delta}) and time series averages (dotted line) based on the RSI_{Delta}'s from the Rappahannock, York, and James rivers from 1988 to 2023. NOTE: Index values could not be calculated for the Rappahannock and James rivers in 2020 due to temporary cessation of sampling due to COVID-19.




Figure 6. Distribution of index-sized American Eel in 2023, number captured, mean length and range from index strata and months.




Figure 7. American Eel - Mean bottom water temperature and 95% C.I. (°C; black line) at sites with positive catches of American Eel during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata. Note: 95% C.I. that are outside the range of observed water temperatures represent years with low numbers of positive catches of American Eel. NOTE: mean bottom water temperature could not be calculated for 2020 due to missing stations as described in the species account.




Figure 8. American Eel - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of American Eel during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata. NOTE: mean salinity could not be calculated for 2020 due to missing stations as described in the species account.




Figure 9. American Eel - Mean dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of American Eel during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata. NOTE: mean dissolved oxygen could not be calculated for 2020 due to missing stations as described in the species account.




Figure 10. Spring juvenile Atlantic Croaker random stratified index (shaded 95% C.I.) and the time series average (dotted line) from the 1987 to the 2023 year class. NOTE: An index value could not be calculated for the 2019 year class due to temporary cessation of sampling due to COVID-19.




Figure 11. Distribution of index-sized juvenile Atlantic Croaker, number captured, and mean length and range from index strata and months.




Figure 12. Atlantic Croaker - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of Atlantic Croaker during index months and strata from 1988 to 2024. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata. No data are available for the 2019 year class due to temporary cessation of sampling due to COVID-19.




Figure 13. Atlantic Croaker - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of Atlantic Croaker during index months and strata from 1988 to 2024. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata. No data are available for the 2019 year class due to temporary cessation of sampling due to COVID-19.




Figure 14. Atlantic Croaker - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of Atlantic Croaker during index months and strata from 1988 to 2024. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata. No data are available for the 2019 year class due to temporary cessation of sampling due to COVID-19.




Figure 15. Juvenile Bay Anchovy random stratified index (RSI_{GM} , 95% C.I.) and the time series average based on the RSI_{GM} (dotted line) from 1988 to 2023.




Figure 16. Distribution of juvenile Bay Anchovy, number captured, and mean length and range from index strata and months.




Figure 17. Bay Anchovy - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of Bay Anchovy during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 18. Bay Anchovy - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of Bay Anchovy during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 19. Bay Anchovy - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of Bay Anchovy during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 20. Black Sea Bass random stratified index (RSI_{GM} , 95% C.I.) and the time series average (dotted line) from 1988 to 2022. NOTE: an index for the 2019 year class could not be calculated due to temporary cessation of sampling due to COVID-19.




Figure 21. Distribution of juvenile Black Sea Bass, number captured, and mean length and range from index strata and months.




Figure 22. Black Sea Bass - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of Black Sea Bass during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 23. Black Sea Bass - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of Black Sea Bass during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 24. Black Sea Bass - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of Black Sea Bass during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 25. Juvenile Blue Catfish random stratified index (RSI_{Delta}) and time series averages (dotted lines) based on the RSI_{Delta} 's from the Rappahannock, York, and James rivers from 1988 to 2023.

Figure 26. No juvenile Blue Catfish were captured from index strata and months.




Figure 27. Juvenile Blue Catfish - Mean bottom water temperature and 95% C.I. ($^{\circ}\text{C}$; black line) at sites with positive catches of juvenile Blue Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata. Note: 95% C.I. that are outside the range of observed water temperatures represent years with low numbers of positive catches of juvenile Blue Catfish.




Figure 28. Juvenile Blue Catfish - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of juvenile Blue Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata.




Figure 29. Juvenile Blue Catfish - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of juvenile Blue Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata.




Figure 30. Age 1+ Blue Catfish random stratified index (RSI_{Delta}) and time series averages (dotted lines) based on the RSI_{Delta}'s from the Rappahannock, York, and James rivers from 1988 to 2023.




Figure 31. Distribution of Age 1+ Blue Catfish, the number captured, and mean length and range from index strata and months.




Figure 32. Age 1+ Blue Catfish - Mean bottom water temperature and 95% C.I. (°C; black line) at sites with positive catches of Age 1+ Blue Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata.




Figure 33. Age 1+ Blue Catfish - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of Age 1+ Blue Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata.




Figure 34. Age 1+ Blue Catfish - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of Age 1+ Blue Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata.




Figure 35. Juvenile Channel Catfish random stratified indices (RSI_{GM} , 95% C.I.) and time series averages (dotted lines) based on the RSI_{GM} 's from the Rappahannock, York, and James rivers from 1988 to 2023.

Figure 36. No juvenile Channel Catfish were captured from index strata and months.




Figure 37. Juvenile Channel Catfish - Mean bottom water temperature and 95% C.I. (°C; black line) at sites with positive catches of juvenile Channel Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata. Note: 95% C.I. that are outside the range of observed water temperatures represent years with low numbers of positive catches of juvenile Channel Catfish.




Figure 38. Juvenile Channel Catfish - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of juvenile Channel Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata. Note: 95% C.I. that are outside the range of observed salinity represent years with low numbers of positive catches of juvenile Channel Catfish.




Figure 39. Juvenile Channel Catfish - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of juvenile Channel Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata.




Figure 40. Age 1+ Channel Catfish random stratified indices (RSI_{GM} , 95% C.I.) and time series averages (dotted lines) based on the RSI_{GM} 's from the Rappahannock, York, and James rivers from 1988 to 2023.

Figure 41. No Age 1+ Channel Catfish were captured from index strata and months.




Figure 42. Age 1+ Channel Catfish - Mean bottom water temperature and 95% C.I. (°C; black line) at sites with positive catches of Age 1+ Channel Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata.




Figure 43. Age 1+ Channel Catfish - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of Age 1+ Channel Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata.




Figure 44. Age 1+ Channel Catfish - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of Age 1+ Channel Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata.




Figure 45. Juvenile Scup random stratified index (RSI_{GM}, 95% C.I.) and the time series average (dotted line) from 1988 to 2022.




Figure 46. Distribution of juvenile Scup, the number captured, and mean length and range from index strata and months.




Figure 47. Scup - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of Scup during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 48. Scup - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of Scup during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 49. Scup - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of Scup during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 50. Juvenile Silver Perch random stratified index (RSI_{GM} , 95% C.I.) and the time series average (dotted line) from 1988 to 2022.




Figure 51. Distribution of juvenile Silver Perch, the number captured, mean length and range from index strata and months.




Figure 52. Juvenile Silver Perch - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of juvenile Silver Perch during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 53. Juvenile Silver Perch - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of juvenile Silver Perch during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 54. Juvenile Silver Perch - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of juvenile Silver Perch during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 55. Juvenile Spot random stratified index (RSI_{GM}, 95% C.I.), the time series average (dotted line) from 1988 to 2023.




Figure 56. Distribution of juvenile Spot, the number captured, mean length and range from index strata and months.




Figure 57. Juvenile Spot - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of juvenile Spot during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 58. Juvenile Spot - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of juvenile Spot during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 59. Juvenile Spot - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of juvenile Spot during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 60. Juvenile Striped Bass random stratified index (RSI_{GM} , 95% C.I.), the time series average (dotted line) from 1988 to 2023.




Figure 61. Distribution of juvenile Striped Bass, the number captured, mean length and range from index strata and months.




Figure 62. Juvenile Striped Bass - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of juvenile Striped Bass during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 63. Juvenile Striped Bass - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of juvenile Striped Bass during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 64. Juvenile Striped Bass - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of juvenile Striped Bass during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 65. Juvenile Summer Flounder random stratified index (RSI_{GM} , 95% C.I.) and the time series average (dotted line) from 1988 to 2023.




Figure 66. Distribution of juvenile Summer Flounder, the number captured, mean length and range from index strata and months.




Figure 67. Juvenile Summer Flounder - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of juvenile Summer Flounder during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 68. Juvenile Summer Flounder - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of juvenile Summer Flounder during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 69. Juvenile Summer Flounder - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of juvenile Summer Flounder during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 70. Juvenile Weakfish random stratified index (RSI_{GM} , 95% C.I.) and the time series average (dotted line) from 1988 to 2023.




Figure 71. Distribution of juvenile Weakfish, the number captured, and mean length and range from index strata and months.




Figure 72. Juvenile Weakfish - Mean bottom water temperature (°C; black line) and 95% C.I. at sites with positive catches of juvenile Weakfish during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean water temperature measured at all sites sampled from the same index months and strata.




Figure 73. Juvenile Weakfish - Mean bottom salinity (psu; black line) and 95% C.I. at sites with positive catches of juvenile Weakfish during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean salinity measured at all sites sampled from the same index months and strata.




Figure 74. Juvenile Weakfish - Mean bottom dissolved oxygen (mg/L; black line) and 95% C.I. at sites with positive catches of juvenile Weakfish during index months and strata from 1988 to 2023. Shaded region represents the 95% C.I. for mean dissolved oxygen measured at all sites sampled from the same index months and strata.




Figure 75. Juvenile White Catfish random stratified indices (RSI_{GM} , 95% C.I.) and times series averages (dotted line) based on RSI_{GM} 's from the Rappahannock, York, and James rivers from 1988 to 2023.

Figure 76. No juvenile White Catfish were captured during index months.




Figure 77. Juvenile White Catfish - Mean bottom water temperature and 95% C.I. (°C; black line) at sites with positive catches of juvenile White Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata. Note: 95% C.I. that are outside the range of observed water temperatures represent years with low numbers of positive catches of juvenile White Catfish.




Figure 78. Juvenile White Catfish - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of juvenile White Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata.




Figure 79. Juvenile White Catfish - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of juvenile White Catfish during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata. Note: 95% C.I. that are outside the range of observed dissolved oxygen represent years with low numbers of positive catches of juvenile White Catfish.




Figure 80. Age 1+ White Catfish random stratified indices (RSI_{GM} , 95% C.I.) and time series averages (dotted line) based on RSI_{GM} 's from the Rappahannock, York, and James rivers from 1988 to 2023.




Figure 81. Distribution of White Catfish age 1+, the number captured, and mean length and range from index strata and months.




Figure 82. Age 1+ White Catfish age - Mean bottom water temperature and 95% C.I. (°C; black line) at sites with positive catches of White Catfish age 1+ during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata. Note: 95% C.I. that are outside the range of observed water temperatures represent years with low numbers of positive catches of White Catfish age 1+.




Figure 83. Age 1+ White Catfish age - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of White Catfish age 1+ during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata.




Figure 84. Age 1+ White Catfish age - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of White Catfish age 1+ during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata.




Figure 85. Juvenile White Perch random stratified indices (RSI_{GM} , 95% C.I.) and time series averages (dotted line) based on RSI_{GM} 's from the Rappahannock, York, and James rivers from 1988 to 2023.




Figure 86. Distribution of juvenile White Perch, the number captured, and mean length and range from index strata and months.




Figure 87. Juvenile White Perch - Mean bottom water temperature and 95% C.I. ($^{\circ}\text{C}$; black line) at sites with positive catches of juvenile White Perch during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata.




Figure 88. Juvenile White Perch - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of juvenile White Perch during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled from the same index months and strata.




Figure 89. Juvenile White Perch - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of juvenile White Perch during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled from the same index months and strata.




Figure 90. Age 1+ White Perch random stratified indices (RSI_{GM} , 95% C.I.) and time series averages (dotted line) based on RSI_{GM} 's from the Rappahannock, York, and James rivers from 1988 to 2023.




Figure 91. Distribution of White Perch age 1+, the number captured, and mean length and range from index strata and months.




Figure 92. Age 1+ White Perch - Mean bottom water temperature and 95% C.I. (°C; black line) at sites with positive catches of White Perch age 1+ during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean water temperature (gray line) measured at all sites sampled from the same index months and strata.




Figure 93. Age 1+ White Perch - Mean bottom salinity and 95% C.I. (psu; black line) at sites with positive catches of White Perch age 1+ during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean salinity (gray line) measured at all sites sampled during the same index months and strata.




Figure 94. Age 1+ White Perch - Mean bottom dissolved oxygen and 95% C.I. (mg/L; black line) at sites with positive catches of White Perch age 1+ during index months and strata in the Rappahannock, York, and James rivers from 1988 to 2023. Shaded region represents the minimum, maximum, and mean dissolved oxygen (gray line) measured at all sites sampled during the same index months and strata.



Appendix Table 1. Trawl Survey advisory requests, data requests, and specimen requests (n = 86) from July 2023 to June 2024.

Juvenile trawl survey specimen or data requests from 2023 - 2024 by month.			2023					2024						
Name	Agency	Nature of request	J	A	S	O	N	D	J	F	M	A	M	J
Troy Tuckey	VIMS	White Perch juveniles												
Troy Tuckey	VIMS	American Eels												
Sarah Koshak	VIMS	Ovigerous blue crabs												
Laurie Sanderson	W&M	Examples of various species												
Kevin Rosemary	MD DNR	White perch												
Nadya Mamoozadeh	MSU	American eel finclips												
Rob Aguilar	SERC	Fish for bar code initiative												
Robert Isdell	CCRM VIMS	Hooked mussels												
Mathew Thayer	CBNERR	Examples of various species												
Nihal Guennounni	VIMS	Blue crabs												
Miguel Montalvo	VIMS	Hogchokers												
Juliette Smith	VIMS	Bay anchovy												
Juliette Smith	VIMS	Atlantic menhaden												
Juliette Smith	VIMS	Naked goby												
Juliette Smith	VIMS	Blue crabs												
Juliette Smith	VIMS	White shrimp												
Juliette Smith	VIMS	Sand shrimp												
Mathew Woodstock	Morgan State	Striped bass indices												
Mathew Woodstock	Morgan State	Blue catfish indices												
Mathew Woodstock	Morgan State	American Eel indices												
Mathew Woodstock	Morgan State	Summer flounder index												
Mathew Woodstock	Morgan State	Spot index												
Mathew Woodstock	Morgan State	White perch indices												
Mathew Woodstock	Morgan State	Bay anchovy index												
Mathew Woodstock	Morgan State	Atlantic croaker index												
Mathew Woodstock	Morgan State	Gizzard shad index												
Mathew Woodstock	Morgan State	Atlantic menhaden index												
Mathew Woodstock	Morgan State	Juvenile Blue crab index												
Mathew Woodstock	Morgan State	Adult Blue crab index												
Pat Geer	VMRC	2023 Blue crab data												
Jeff Shields	VIMS	Atlantic silversides												
Jenny Dryer	VIMS	Angel wings												
Jenny Dryer	VIMS	Mya												
Jenny Dryer	VIMS	Macoma												
Jenny Dryer	VIMS	Melampus												
Jenny Dryer	VIMS	Ercolania												
Jenny Dryer	VIMS	Marenzelleria												
Jenny Dryer	VIMS	Phoronids												
Alexa Galvan	VMRC	Weakfish data												
Alexa Galvan	VMRC	Scup index												
Alexa Galvan	VMRC	Black sea bass index												
Alexa Galvan	VMRC	Summer flounder index												
Patrick McGrath	VIMS	Alosa numbers												
Somers Smott	VMRC	Spot data												
Ethan Simpson	VMRC	Horseshoe crab data												
Somers Smott	VMRC	Spot data												



Appendix Table 2. Trawl Survey comparison study summary showing the best model, number of paired comparison tows used to estimate the calibration factor (N), and the calibration factor used to multiply each tow from the R/V *Tidewater* to convert it to R/V *Fish Hawk* equivalents by species. Due to low or no catches of White and Channel Catfish in the comparison study, we applied the best model from Blue Catfish to adjust index values for these species.

Species	N	Model	Calibration Factor
American Eel age 1+	27	Binomial	0.8731
Atlantic Croaker YOY	283	Beta-binomial	1.4770
Bay Anchovy YOY	504	Beta-binomial	0.8014
Black Sea Bass YOY	25	Binomial	0.6336
Blue Catfish YOY	78	Beta-binomial	1.0023
Blue Catfish age 1+	123	Beta-binomial	1.4289
Channel Catfish YOY	0	Blue Catfish YOY	1.0023
Channel Catfish age 1+	0	Blue Catfish age 1+	1.4289
Scup	27	Beta-binomial with random paired-tow effect	1.3471
Silver Perch YOY	118	Beta-binomial	0.7524
Spot YOY	187	Beta-binomial	1.1654
Striped Bass YOY	90	Beta-binomial	1.2009
Summer Flounder YOY	146	Beta-binomial	1.0308
Weakfish YOY	220	Beta-binomial	1.1924
White Catfish YOY	10	Blue Catfish YOY	1.0023
White Catfish age 1+	32	Binomial	0.6918
White Perch YOY	161	Beta-binomial	1.0679
White Perch age 1+	210	Beta-binomial	1.1676



Appendix Figure 1. Length frequency distributions by species from July 2021 through June 2022. (Note that actual indices are calculated using a subset of months and strata. Therefore, not all index-sized fish are included in index calculations.)





Appendix Figure 1. (continued)





Appendix Figure 1. (continued)





Appendix Figure 1 (continued).



Appendix Figure 1. (continued)



Appendix Figure 1. (continued)



Appendix Figure 1. (continued)

