1979

Irreversible nongenetic temperature adaptation of oxygen uptake in clones of the sea anemone *Haliplanella luciae* (Verrill)

William Ernest Zamer
College of William & Mary - Arts & Sciences

Follow this and additional works at: <https://scholarworks.wm.edu/etd>

Part of the [Physiology Commons](#)

Recommended Citation

Zamer, William Ernest, "Irreversible nongenetic temperature adaptation of oxygen uptake in clones of the sea anemone *Haliplanella luciae* (Verrill)" (1979). *Dissertations, Theses, and Masters Projects*. Paper 1539625056.

<https://dx.doi.org/doi:10.21220/s2-47j1-be96>

This Thesis is brought to you for free and open access by the Theses, Dissertations, & Master Projects at W&M ScholarWorks. It has been accepted for inclusion in Dissertations, Theses, and Masters Projects by an authorized administrator of W&M ScholarWorks. For more information, please contact scholarworks@wm.edu.

IRREVERSIBLE NONGENETIC TEMPERATURE
" ADAPTATION OF OXYGEN UPTAKE IN CLONES OF THE
SEA ANEMONE HALIPLANELLA LUCIAE (VERRILL)

A Thesis

Presented to

The Faculty of the Department of Biology
The College of William and Mary in Virginia

In Partial Fulfillment
Of the Requirements for the Degree of
Master of Arts

by
William E. ^{Ernest} Zamer

1979

APPROVAL SHEET

This thesis is submitted in partial fulfillment of the
requirements for the degree of

Master of Arts

William E. Zamer
Author

Approved, September 1978

Charlotte P. Mangum
Charlotte P. Mangum

Robert E. L. Black
Robert E.L. Black

Bruce S. Grant
Bruce S. Grant

TABLE OF CONTENTS

	Page
ACKNOWLEDGEMENTS	iv
LIST OF TABLES	v
LIST OF FIGURES	vi
ABSTRACT	vii
INTRODUCTION	2
MATERIALS AND METHODS	5
RESULTS	11
DISCUSSION	20
LITERATURE CITED	22

ACKNOWLEDGEMENTS

I would like to express my appreciation to Dr. Charlotte P. Mangum for her patience and continued encouragement during the course of this study. The laboratory space, materials, and guidance for the electrophoretic study kindly provided by Dr. Richard Hoffmann are greatly appreciated. I would also like to thank David Reed for his help with the statistics and use of the SPSS programs used in this study. I would like to thank Drs. Robert Black and Bruce Grant, and Nick Mauro for critically reading the manuscript.

LIST OF TABLES

Table	Page
I. Oxygen uptake data	15
II. Oxygen uptake data grouped by weight class	16
III. Surface area measurements	18

LIST OF FIGURES

Figure	Page
1. Summary of cloning and experimental design	7
2. Frequency distribution of body weights - treatment groups.	12
3. Frequency distribution of body weights - developmental groups	13

ABSTRACT

Isogenic clones of H. luciae were raised at 18° and 28°C. After several generations of growth and morphogenesis at these two developmental temperatures, differences in O₂ uptake could no longer be erased by acclimation to a common temperature. Acclimated O₂ uptake rates were 18-19% higher in animals raised at the lower temperature than in animals of the same body size raised at the higher temperature. When the body size relationship is ignored, however, the influence of developmental temperature on O₂ uptake is masked due to the larger size of the 18°C raised animals.

The irreversible change is not due to an increase in the gas exchange surface area at the primary site of O₂ uptake. In addition, no evidence was found of qualitative changes in several enzymes which could influence aerobic metabolism. The change could be due to either a shift in the relative concentrations of the components of a multiple isozyme system, or to qualitative alterations in other enzymes which were not examined.

IRREVERSIBLE NONGENETIC TEMPERATURE ADAPTATION OF
OXYGEN UPTAKE IN CLONES OF THE SEA ANEMONE
HALIPLANELLA LUCIAE (VERRILL)

INTRODUCTION

For many years physiological differences have been observed among latitudinally separated populations of the same species (Bullock, 1955; Prosser, 1955; Vernberg, 1962). These differences have been attributed to a variety of phenotypic adaptations and also to genetic variation.

One of the earliest examples of physiological variation was the difference in acutely measured bell pulsation rates of the jellyfish Aurelia aurita collected from Nova Scotia and Florida (Mayer, 1914), the direction of which agreed with Krogh's (1916) prediction of increased rates in cold adapted animals compared to warm adapted animals, when the rates are measured at a common temperature. More recent studies of oxygen uptake as well as motor activity in cnidarians confirmed the existence of distinct responses in latitudinally separated populations within a species (Sassaman and Mangum, 1970; Mangum et al., 1972; Shick, 1976).

In many examples of intraspecific variation, latitudinal differences disappear with acclimation to common conditions, indicating their phenotypic origin. In several studies of latitudinally separated populations however, the differences persist despite prolonged exposure to common conditions, which has been interpreted alternatively as 1) genetic divergence and 2) "irreversible nongenetic adaptation" (Kinne, 1962). There are numerous examples of variation which is highly likely to have a genetic basis, e.g. interspecific variation (Vernberg, 1962). The possibility of nongenetic but irreversible adaptation within species is less certain.

Evidence of this phenomenon in Drosophila subobscura was presented by Smith (1956), who induced variation in heat tolerance among groups

of flies raised at different temperatures. The differences, which did not disappear completely upon acclimation to a common temperature, were adaptive in character; flies developing at the higher temperature survived longer at an upper lethal limit. However, selection, operating during development, could conceivably explain these results. Kinne (1962) reported that fish developed from eggs which had been transferred to a different salinity three to six hours after spawning exhibit lower food conversion efficiencies relative to controls developed from eggs which remained in the spawning salinity. He does not, however, compare food conversion efficiencies of fish acclimated to a common salinity which had developed at different salinities. It is not clear, therefore, whether the observed differences in food conversion efficiencies are reversible upon acclimation to common conditions. Bradley (1978) showed that copepods raised at 20°C are more tolerant of high temperatures than those raised at 10°C. However, these results could have arisen from selection during rearing or to a thermal acclimation that requires more than two days exposure to a common temperature to disappear. Schneider (1968) concluded that differences in acclimated oxygen uptake rates among populations of the crab Rhithropanopeus harrisi can be reversed only in part by breeding the crabs at a common temperature, but the full report of these data has not appeared.

The mode of reproduction in the actinian Haliplanella luciae (Verrill) offers an interesting approach to the question of nongenetic adaptation which is irreversible within a period of prolonged acclimation to common conditions. This species reproduces largely by asexual mechanisms, longitudinal fission being the most common (Shick and Lamb, 1977). The fission process in Haliplanella luciae and the effects of temperature and other environmental factors on it have been studied in detail by several

authors (Torrey and Mery, 1904; Davis, 1919; Uchida, 1932; Miyawaki, 1952; Minasian, 1976). Longitudinal fission can be induced in the laboratory by storing the animal at 10°C or below for six to eight weeks, followed by raising the temperature above 10°C (Sassaman, pers. comm.). This treatment apparently simulates increasing water temperatures during spring. After fission the daughter animals regenerate tissues to close the body wall torn in the process. Thus isogenic clones of animals can be raised under different environmental conditions and later compared, virtually eliminating the possibility of genetic variability among the experimental animals.

In the present study, three generations of isogenic anemones were reared at each of two temperatures, and their rates of oxygen uptake compared after acclimation to common thermal conditions. A few observations were also made on the electrophoretic banding patterns of five enzymes and on the dimensions of the gas exchange surface.

MATERIALS AND METHODS

Collection, Maintenance of Animals and Experimental Design

Specimens of Haliplanella luciae (Verrill) were collected from the York River estuary at the mouth of Indian Field Creek, Virginia in January 1977 ($<10^{\circ}\text{C}$). Included in this sample were animals of Uchida's (1932) color types 1 and 3, indicating genetic heterogeneity in the population. Assuming all animals to be genetically unique, single individuals were placed in 30 ml plastic beakers filled with York River water. One half of the animals were placed at 28°C and the remainder stored at 18°C . After approximately two weeks most of the animals at both temperatures had fissioned, each original individual producing two or more progeny (clonemates). When regeneration was complete, the animals were transferred to 5°C for six to eight weeks. At the end of this period, the clonemates were separated and allowed to undergo fission and subsequent formation of new tissue at different developmental temperatures: one member of each clone was randomly chosen and placed at 18°C and its clonemate was placed at 28°C . After fission and regeneration, the second generation clonemates were separated, stored at 5°C for 6-8 weeks and then returned to their developmental temperature, where a third generation was produced. Thus there are four groups of animals according to developmental temperature and the number of generations produced at a given temperature: 1) animals taken from the field that divided and formed new tissue three times at 18°C or 2) 28°C , and 3) animals that divided first at 18°C and subsequently a second and third time at 28°C or 4) first at 28°C and subsequently a second and third time at 18°C . Because of the difference in the number of fissions at

each developmental temperature (Fig. 1) a control was necessary to permit pooling the data for all clones at a particular developmental temperature. The rates for animals which fissioned twice at a developmental temperature were compared with those for animals which had fissioned three times at that developmental temperature. The important aspect of the experimental design is, however, that each individual has at least one clonemate treated in the same way but given the alternative thermal regime (Fig. 1).

Development and growth occurred at the specified temperature in natural estuary water (15-21 ‰ with members of all groups given the same salinity regime). At least four days prior to the measurement of oxygen uptake, the salinity was brought to 18 ‰. Animals were fed an excess of freshly hatched Artemia nauplii every other day.

Oxygen Uptake Measurements

Oxygen uptake ($\dot{V}O_2$) of the third generation clonemates was measured at the developmental temperature (TD), the same as the acclimation temperature (TA). These data are designated 1) TD = 18°C = TA and 2) TD = 28°C = TA. Other, but isogenic, individuals were transferred to the alternative temperature, held there for 2-4 weeks and oxygen uptake measured. These data are designated 3) TD = 18°C, TA = 28°C and 4) TD = 28°C, TA = 18°C. All measurements were made in millipore filtered water at 18 ‰ salinity by adjusting the concentration of natural water with distilled water or commercial sea salt (Dayno Corp.). Animals were starved for 60-84 hours before an experiment, and transferred to the respirometry chambers the night before an experiment to permit attachment to the wall of the chamber. Oxygen depletion was measured with a Yellow Springs Instrument Co. Model 53 Biological Oxygen Monitor, using a high sensitivity (0.0005 in) teflon membrane

Fig. 1 Cloning of H. luciae at 18°C and 28°C. The example shows two animals (A and B) assumed to be genetically unique. Solid arrows indicate transfer from one temperature to another; dotted arrows indicate fission and regeneration. Roman numerals indicate generation.

and a 1x10mm stirring bar. The animals were permitted to adjust to stirring for one hour before sealing the chamber and the data were discarded when the tentacles were retracted before completion of the measurement. All data in the PO_2 interval 110-159 mm Hg were analyzed. For each experiment a measurement of the oxygen uptake of the electrode was made using an empty chamber, and this value was subtracted from the experimental rate.

At the end of an experiment animals were stimulated to retract their tentacles, removed from the chambers, lightly blotted and weighed to the nearest tenth of a milligram (Cahn Electrobalance, Model G, Ventron Instruments Corp.).

Electrophoresis

Animals were starved for one week and then frozen whole, either singly or in groups from the same clone and treatment group, in two volumes of buffer: 0.05 M tris (titrated to pH 8 with concentrated HCl) and 0.001 M EDTA, to which NADP (4 mg/l) was added. Freezing took place quickly in a bath of dry ice and acetone, and the samples were then stored at -70°C until homogenized. Samples were thawed, homogenized and centrifuged. When necessary the supernatants were diluted as much as 50% with distilled water to obtain 25 μl which were loaded into slots of horizontal starch gels (13% w/v; Sigma starch).

Gel and electrode buffers for glucose-6-phosphate dehydrogenase (G-6-PDH, E.C. 1.1.1.49) and isocitrate dehydrogenase (IDH, E.C. 1.1.1.42) were modified from Markert and Faulhaber (1965): 0.9 M tris, 0.5 M boric acid, 0.02 M EDTA diluted 1:20 for the gel buffer and 1:6 for the electrodes; in addition, 30 mg and 10 mg NADP were added to the gel and to the cathodal buffer tray, respectively. The gel and

electrode buffers for malate dehydrogenase (MDH, E.C. 1.1.1.37) were those described by Nichols and Ruddie (1973). The discontinuous LiOH buffer system of Selander, et al. (1969) was used for phosphoglucose isomerase (PGI, E.C. 5.3.1.9) and hexokinase (HK, E.C. 2.7.1.1.). The period of electrophoresis was 3-4 hours. After electrophoresis at 350 volts, gels were sliced and stained with slight modifications of the procedures described by Shaw and Prasad (1970) and Brewer (1970). The gels were preserved overnight in 50% ethyl alcohol, blotted and stored in plastic wrap until photographed.

Gas Exchange Surface Area

Animals from the TD=18°C=TA and the TD=28°C=TA treatment groups were anesthetized with MgCl_2 . Tentacles were counted and the external surface areas of the body wall and the tentacles were approximated from measurements made at 10-25X, using formulas for the frustum of a cone and for a cylinder, respectively.

Statistical Analyses

Mean $\dot{V}\text{O}_2$ values were computed by determining the rate during each time interval of the continuous record of the change in oxygen concentration. Thus, unless specifically stated otherwise, the N value given below represents the sum of all data obtained from the members of an experimental group, which are treated as a homogeneous population.

The comparison was complicated, however, by the different body sizes of the experimental groups (see below). Two alternative procedures were used to eliminate differences due to body size from the conclusions. First, a regression coefficient of -0.43 was used to correct rates to a common weight of 1.6 mg by covariance analysis.

This coefficient was determined by performing a regression analysis on data from the TD = 18°C = TA treatment group (N = 14 animals, $\bar{r}$ = 0.75, $P < 0.001$). This relationship is homogeneous, showing no evidence of the discontinuity observed by Shick et al. (1978). Second, the data were grouped by weight classes established from an analysis of frequency distributions, and the comparison made within the classes.

Although the experimental design permits analysis of the data as paired observations the available procedures for computing Student's t require an equal number of observations in each member of the pair, a condition which was not met. The element of genetic homogeneity was maintained in the experimental design by keeping virtually constant the percent of the total number of observations made on each clone under each combination of developmental and experimental temperatures.

RESULTS

Effect of Developmental Temperature On Body Size

At 28°C fission occurred more rapidly and more frequently than at 18°C, resulting in more numerous but smaller progeny (see below). As suggested earlier by a number of investigators (Ray, 1960; Mangum, 1963; Shick, 1972) this finding suggests that the often observed phenomenon of a larger body size in the colder regions of a species range results from latitudinal differences in developmental temperature.

The frequency distribution of weight in each experimental group (Fig. 2) indicates, in general, that body size of the animals is more widely distributed in the 18°C developmental group than in the 28°C developmental group. Almost 73% of the animals in the TD = 28°C = TA treatment group weigh 1.1 mg or less, and 91% of the animals in the TD = 28°C, TA = 18°C experimental group weigh 2.0 mg or less. In contrast, only 29% of the animals in the TD = 18°C = TA group and 17% of those in the TD = 18°C, TA = 28°C treatment group weigh 1.1 mg or less. In these groups, reared at low temperature, higher frequencies of animals in the higher weight classes are also evident (Fig. 2). A comparison of the frequency distributions by weight class of the two developmental temperature groups emphasizes the trends (Fig. 3). The mean body weight of the TD = 28°C = TA and TD = 28°C, TA = 18°C animals is 1.4 mg; the mean weight of the two TD = 18°C groups is 2.0 mg. The median weight class of the two TD = 28°C groups is 0.6-0.8 mg, and the corresponding figure for the TD = 18°C groups is 2.1-2.3 mg.

Oxygen Uptake

Three clones of animals which first fissioned at 18°C and one clone

Fig. 2 Frequency distribution of body weight in treatment groups, designated by developmental (TD) and acclimation (TA) temperature.

Fig. 3 Frequency distribution of body weight in (A) all animals used in the oxygen uptake experiment, (B) TD = 28°C developmental groups only, and (C) TD = 18°C developmental groups only.

of animals which first fissioned at 28°C produced third generation animals at both developmental temperatures. These four clones were used as the experimental material. No significant differences were found between the control groups ($P > 0.05$) indicating that the responses of $\dot{V}O_2$ in animals that have fissioned twice at a particular developmental temperature is the same as in animals that have fissioned three times at that developmental temperature.

The mean weight corrected oxygen uptake rates of the two TD = 18°C groups are significantly higher ($P < 0.01$) than those of the two TD = 28°C groups at both acclimation temperatures (Table I). At the test temperature of 18°C the rates in the TD = 18°C groups are 18% higher than those of the TD = 28°C groups, and at 28°C the rates are 19% higher. However, using the raw data for the smaller animals produced at 28°C and the larger ones produced at 18°C, the difference diminished to 4% at the test temperature of 18°C and it is actually reversed at 28°C (-8%). Thus a confounding influence of developmental temperature, its effect on body size, may obscure the direct effect on the acclimated rate of oxygen uptake.

The frequency distribution by weight of all animals used in the oxygen uptake experiments is shown in Fig. 3. When the rates are compared by weight class, significant differences between the TD = 18°C and the TD = 28°C groups are found in one of the four possible comparisons at 18°C and in two of three at 28°C (Table II). The trend of higher rates in the TD = 18°C groups is uniformly consistent, but the probability levels rise due to the smaller number of observations (N).

Q_{10} values for both developmental temperature groups (Table II)

TABLE I

A. Oxygen uptake ($\mu\text{l/gm-hr}$) in isogenic Haliplanella luciae reared for 2-3 generations at different temperatures. Mean + S.E. (N). Probability values from one-way analyses of variance comparing developmental temperature groups at each test (= acclimation) temperature.

Test temperature ($^{\circ}\text{C}$)	Body wt. (mg)	Developmental temperature ($^{\circ}\text{C}$)		P
		18	28	
18	1.9 $\pm$ 0.2	243.7 $\pm$ 12.0 (111)	235.0 $\pm$ 16.2 (72)	n.s.
28	1.3 $\pm$ 0.1	487.1 $\pm$ 24.1 (36)	524.7 $\pm$ 21.6 (54)	n.s.

B. Oxygen uptake ($\mu\text{l/gm-hr}$) corrected to a common body weight (1.6 mg) by covariance. Data analysis as above.

Test temperature ($^{\circ}\text{C}$)	Developmental temperature ($^{\circ}\text{C}$)				P
	18	Q_{10}	28	Q_{10}	
18	257.0 $\pm$ 9.4 (111)		218.7 $\pm$ 13.9 (72)		P<<0.001
		1.99		1.96	
28	510.3 $\pm$ 22.6 (36)		427.8 $\pm$ 14.5 (54)		0.0018

TABLE II

Weight corrected oxygen uptake rates ($\mu\text{l/gm-hr}$), Mean $\pm$ S.E. (N). Probability (P) values from one-way analyses of variance comparing the developmental temperature group rates at each experimental (= acclimation) temperature for each weight class.

Experimental temperature ($^{\circ}\text{C}$)	Weight class		Developmental temperature ($^{\circ}\text{C}$)		P
	No.	Range (mg)	18	28	
18	1	(0.6-0.8)	270.5 $\pm$ 46.2 (10)	194.5 $\pm$ 33.2 (9)	n.s.
	2	(0.9-1.1)	269.8 $\pm$ 38.0 (11)	146.6 $\pm$ 41.4 (7)	0.0499
	3	(1.2-1.7)	218.6 $\pm$ 22.7 (23)	159.1 $\pm$ 21.8 (16)	n.s.
	4	(1.8-2.0)	-----	288.5 $\pm$ 18.7 (24)	-----
	5	(2.1-3.7)	254.7 $\pm$ 11.2 (67)	258.8 $\pm$ 4.1 (6)	n.s.
28	1	(0.6-0.8)	-----	444.0 $\pm$ 22.8 (20)	-----
	2	(0.9-1.1)	413.3 $\pm$ 38.1 (6)	417.4 $\pm$ 27.3 (20)	n.s.
	3	(1.2-1.7)	568.9 $\pm$ 13.8 (5)	408.3 $\pm$ 37.7 (9)	0.003*
	4	(1.8-2.0)	608.9 $\pm$ 34.3 (13)	439.9 $\pm$ 24.1 (5)	0.01
	5	(2.1-3.7)	427.4 $\pm$ 31.6 (12)	-----	-----

*Bartlett's Test indicates non-homogenous variances for the two developmental groups of this weight class. A Separate Variance Estimate t-test was used to make this comparison.

are similar, and they also approximate the values reported previously over the same temperature range (Sassaman and Mangum, 1970). Thus no evidence of a change in temperature sensitivity with developmental temperature is seen. No overall pattern of temperature sensitivity is apparent in the data grouped by weight class, thus supporting previous conclusions that it is not correlated with body size (Sassaman and Mangum, 1970). Nor is there a trend in Q_{10} with developmental temperature.

Gas Exchange Surface

A possible explanation of the irreversible difference between cold and warm reared anemones is a change in gas exchange surface area. This hypothesis would be supported by an increase in tenacular surface, the primary site of oxygen uptake in epifaunal anemones (Sassaman and Mangum, 1972; Shick et al., 1978) in the TD = 18°C groups. In fact, measurements of gas exchange surface (Table III) indicate the opposite relationship, suggesting that the difference would be even larger if the gas exchange surface had remained the same.

Comparing animals in the same weight class, anemones in the TD = 28°C groups have more tentacles and a consistently larger total surface area available for gas exchange than anemones in the TD = 18°C groups. Not only are the tentacle numbers greater in the TD = 28°C groups, but the average surface area per tentacle is greater as well. Moreover, the percent of the total body surface area contributed by the tentacles always exceeds 73 in the TD = 28°C groups, compared to a maximum of 57.5% in the TD = 18°C animals (weight class 2).

Electrophoresis

The enzymes examined showed no qualitative differences in banding

TABLE III

Gas exchange surface area (cm^2/gm wet wt.) in each developmental temperature group of Haliplanella luciae. N = No. animals. Weight classes as in Table II.

Developmental temperature ($^{\circ}\text{C}$)	Weight class	N	No. tentacles	Tentacle surface area	
				cm^2/gm	% Total body surface area
18	1	2	13	5.6	33.9
	2	2	16	5.0	57.5
	3	3	20	6.2	37.9
28	1	2	18	19.2	73.5
	2	4	30	14.7	72.9
	3	3	29	18.6	78.7

pattern either among the separate clones of one treatment group or among clonemates of different treatment groups. The results for all of the enzymes tested are similar and they suggest that the population of H. luciae at Indian Field Creek exhibits no variation at these loci. Of the five enzyme systems examined, only IDH was clearly polymorphic, the banding pattern suggesting fixed homozygosity at two loci. However, no qualitative differences in banding pattern were detected, either between clones or within clones. Since these loci differ in latitudinally separated populations and within populations of H. luciae (Shick and Lamb, 1977), the uniform pattern could reflect conservative selection pressures on these loci for this particular population. Regardless, there is no evidence that the different alleles were either induced or repressed by the formation of tissues at different temperatures.

DISCUSSION

Differences in aerobic metabolism that cannot be reversed by 2-4 weeks acclimation to common conditions clearly result from reproduction, regeneration and the formation of new tissues at different temperatures, when the effects of other variables are eliminated from the data. The direction of the change is generally compensatory, resulting in higher metabolic rates in the animals produced at low temperatures. The magnitude of the change, however, is not great enough to override opposing effects of developmental temperature on body size and gas exchange surface area. Oxygen uptake rates go up at low developmental temperature but down with increasing body size and decreasing tentacular surface, and body size increases and tentacular surface area decreases at low developmental temperature. The net outcome of the concomitant and counteracting trends is little or no difference between the two groups, when body size and gas exchange surface are retained as variables.

The question remains of the origin of the metabolic differences in animals of the same size. One alternative is a direct but irreversible effect of developmental temperature on the metabolic machinery within the cell. The five enzyme systems examined were selected in part for their variability in the species (and in other anemones; Manwell and Baker, 1970), and in part for their relation to aerobic metabolism. Hexokinase, phosphoglucose isomerase and isocitrate dehydrogenase vary in different populations of H. luciae (Shick and Lamb, 1977); isocitrate dehydrogenase and malate dehydrogenase catalyze reactions in the tricarboxylic acid cycle and glucose-6-phosphate dehydrogenase regulates a branchpoint of the pentose phosphate pathway.

No qualitative changes with developmental temperature or acclimation temperature were found for any of the five enzymes tested. Quantitative data, on various enzyme systems, however, are not available. Robert (1971) has shown an increase in the specific activity of glucose-6-phosphate dehydrogenase and 6-phosphogluconate dehydrogenase of the blue crab Callinectes sapidus during cold exposure, accompanied by no qualitative electrophoretic changes. A thorough test of this possibility, which we regard as a promising hypothesis, would entail an exhaustive investigation of polymorphic enzyme systems that influence the rate of oxidative reactions, directly or indirectly.

The only obvious alternative explanation would be an irreversible effect of developmental temperature on the gas exchange system. Since fluid movements on both sides of the tentacles are generated by cilia, an irreversible difference in oxygen convection would seem to be highly unlikely. An irreversible change in gas exchange surface was detected, but its effect on oxygen uptake should be the opposite of that observed. Thus a systemic explanation seems highly unlikely.

Regardless of its basis, the present findings clearly demonstrate the reality of irreversible physiological adaptation, and they raise the possibility that the compensatory responses found earlier in geographically separate populations of the species may result from developmental as well as acclimation temperature.

LITERATURE CITED

- Bradley, B.P., 1978. Increase in range of temperature tolerance by acclimation in the copepod Eurytemora affinis. Biol. Bull., 154: 177-187.
- Brewer, G.J., 1970. An introduction to isozyme techniques. Academic Press, New York, 186 pp.
- Bullock, T.H., 1955. Compensation for temperature in the metabolism and activity of poikilotherms. Biol. Rev. Camb. Phil. Soc., 30: 31-342.
- Davis, D.W., 1919. Asexual multiplication and regeneration in Sagartia luciae Verrill. J. Exp. Zool., 28: 161-263.
- Kinne, O., 1962. Irreversible nongenetic adaptation. Comp. Biochem. Physiol., 5: 265-282.
- Krogh, A., 1916. The respiratory exchange of animals and men. Longmans Green, London. 173 pp.
- Mangum, C.P., 1963. Studies on speciation in maldanid polychaetes of the North American Atlantic Coast III. Intraspecific and interspecific divergence in oxygen consumption. Comp. Biochem. Physiol., 10: 335-349.
- Mangum, C.P., M.J. Oakes, and J.M. Shick, 1972. Rate temperature responses in scyphozoan medusae and polyps. Mar. Biol. 15: 298-303.
- Manwell, C. and C.M.A. Baker, 1970. Molecular biology and the origin of species: heterosis, protein polymorphism, and animal breeding. U. of Wa. Press, Seattle, p. 161.

- Markert, C.L. and I. Faulhaber, 1965. Lactate dehydrogenase patterns of fish. J. Exp. Zool., 159: 319-332.
- Mayer, A.G., 1914. The effects of temperature upon tropical marine animals. Pap. Tortugas Lab., 6: 1-24. Publs. Carnegie Instn., 183.
- Minasian, L.L., 1976. Characteristics of asexual reproduction in the sea anemone, Haliplanella luciae (Verrill), reared in the laboratory. Pages 289-298 in G. O. Mackie, Ed., Coelenterate Ecology and Behavior. Plenum Press, New York.
- Miyawaki, M., 1952. Temperature as a factor influencing upon the fission of the orange-striped sea anemone, Diadumene luciae. J. Fac. Sci. Hokkaido Univ., Ser. 6, 11: 77-80.
- Nichols, E.A. and F.H. Ruddle, 1973. A review of enzyme polymorphism, linkage and electrophoretic conditions for mouse and somatic cell hybrids in starch gels. J. Histochem. Cytochem., 21: 1066-1081.
- Prosser, C.L., 1955. Physiological variation in animals. Biol. Rev. Camb. Phil. Soc., 30: 229-262.
- Ray, C., 1960. The application of Bergmann's and Allen's rules to the poikilotherms. J. Morph., 106: 85-108.
- Robert, M.S., 1971. Enzymatic mechanisms during temperature acclimation of the blue crab, Callinectes sapidus. Dissertation, Georgetown U., Thesis 3910.
- Sassaman, C., and C.P. Mangum, 1970. Patterns of temperature adaptation in North American Atlantic coastal actinians. Mar. Biol., 7: 123-130.
- Sassaman, C. and C.P. Mangum, 1972. Adaptations to environmental oxygen levels in infaunal and epifaunal sea anemones. Biol. Bull., 143: 657-678.

- Schneider, D.E., 1968. Temperature adaptations in latitudinally separated populations of the crab Rhithropanopeus harrisi. Amer. Zool., 8: 772. (Abstract).
- Selander, R.K., W.G. Hunt and S.Y. Yang, 1969. Protein polymorphism and genic heterozygosity in two European subspecies of the house mouse. Evolution, 23: 379-390.
- Shaw, C.R. and R. Prasad, 1970. Starch gel electrophoresis of enzymes-- a compilation of recipes. Biochem. Genet., 4: 297-320.
- Shick, J.M., 1972. Temperature sensitivity of oxygen consumption of latitudinally separated Urosalpinx cinerea (Prosobranchia: Muricidae) populations. Mar. Biol., 13: 276-283.
- Shick, J.M., 1976. Ecological physiology and genetics of the colonizing actinian Haliplanella luciae, pages 137-146 in G. O. Mackie, Ed., Coelenterate Ecology and Behavior. Plenum Press, New York.
- Shick, J.M. and A.N. Lamb, 1977. Asexual reproduction and genetic population structure in the colonizing sea anemone Haliplanella luciae. Biol. Bull., 153: 604-617.
- Shick, J.M., W.I. Brown, E.G. Dolliver, and S.R. Kayar, 1978. Oxygen uptake in sea anemones: Effects of expansion, contraction and exposure to air, and the limitations of diffusion. Physiol. Zool., 52: 50-62.
- Smith, J.M., 1956. Acclimatization to high temperatures in inbred and outbred Drosophila subobscura. J. Genet., 54: 497-505.
- Torey, H.B. and J.R. Mery, 1904. Regeneration and non-sexual reproduction in Sagartia davisi. Zoology 1(6): 211-226. U. Calif. Publs.
- Uchida, T., 1932. Occurrence in Japan of Diadumena luciae, a remarkable actinian of rapid dispersal. J. Fac. Sci. Hokkaido Univ., Ser. 6, 2: 69-85.

Vernberg, F.J., 1962. Comparative physiology: Latitudinal effects on physiological properties of animal populations. Ann. Rev. Physiol., 24: 517-546.

VITA

William Ernest Zamer

Born in Johnstown, Pennsylvania June 27, 1952. Graduated from Westmont Hilltop Senior High School in that city June, 1970. B.S., Juniata College, 1974. M.A. candidate, College of William and Mary, 1975-1979 and graduate teaching assistant in the Department of Biology 1975-1977.

In September 1978 began work as a research assistant at the Cancer Center, Medical College of Virginia.