

**Evaluation of Striped Bass Stocks in Virginia:
Monitoring and Tagging Studies, 2010-2014**

Progress Report

Contract Number: F-77-R-24
Project Period: 1 September 2010 - 31 August 2011
Principal Investigator: John M. Hoenig

Prepared by:

Philip W. Sadler, Matthew W. Smith, Shelley E. Sullivan, John M. Hoenig, Robert E. Harris, Jr. and Lydia M. Goins

**Department of Fisheries Science
School of Marine Science
Virginia Institute of Marine Science
The College of William and Mary
Gloucester Point, VA 23062-1346**

Submitted To:

**Virginia Marine Resources Commission
P.O. Box 756
Newport News, VA 23607-0756**

26 January 2012



Preface

This report presents the results of striped bass (*Morone saxatilis*) tagging and monitoring activities in Virginia during the period 1 September 2010 through 31 August 2011. It includes an assessment of the biological characteristics of striped bass taken from the 2011 spring spawning run, estimates of annual survival and fishing mortality based on annual spring tagging, and the results of the study that documents the prevalence of mycobacterial infections of striped bass in Chesapeake Bay. The information contained in this report is required by the Atlantic States Marine Fisheries Commission and is used to implement a coordinated management plan for striped bass in Virginia, and along the eastern seaboard.

Striped bass have historically supported one of the most important recreational and commercial fisheries along the Atlantic coast. In colonial times, striped bass were abundant in most coastal rivers from New Brunswick to Georgia, but overfishing, pollution and reduction of spawning habitat have resulted in periodic crashes in stocks and an overall reduction of biomass (Merriman 1941, Pearson 1938). Striped bass populations at the northern and southern extremes of the Atlantic are apparently non-migratory (Raney 1957). Presently, important sources of striped bass in their native range are found in the Roanoke, Delaware and Hudson rivers and the major tributaries of Chesapeake Bay (Lewis 1957) with the Chesapeake Bay and Hudson River being the primary sources of the coastal migratory population (Dorazio *et al.* 1994).

Examination of meristic characteristics indicate that the coastal migratory population consists of distinct sub-populations from the Hudson River, James River, Rappahannock - York rivers, and upper Chesapeake Bay (Raney 1957). The Roanoke River striped bass may represent another distinct sub-population (Raney 1957). The relative contribution of each area to the coastal population varies. Berggren and Lieberman (1978) concluded from a morphological study that Chesapeake Bay striped bass were the major contributor (90.8%) to the Atlantic coast fisheries, and the Hudson River and Roanoke River stocks were minor contributors. However, they estimated that the exceptionally strong 1970 year class constituted 40% of their total sample. Van Winkle *et al.* (1988) estimated that the Hudson River stock constituted 40% - 50% of the striped bass caught in the Atlantic coastal fishery in 1965. Regardless of the exact proportion, management of striped bass is a multi-jurisdictional concern as spawning success in one area probably influences fishing success in many areas. Furthermore, recent evidence suggests the presence of divergent migratory behavior at intra-population levels (Secor 1999). The extent to which these levels of behavioral complexity impact management strategies in Chesapeake Bay and other stocks is unknown.

Concern about the decline in striped bass landings along the Atlantic coast since the mid-1970s prompted the development of an interstate fisheries management plan (FMP) under the auspices of the Atlantic States Marine Fisheries Management Program (ASMFC 1981). Federal legislation was enacted in 1984 (Public Law 98-613, the Atlantic Striped Bass Conservation Act) which enables Federal imposition of a moratorium for an indefinite period in those states that fail to comply with the coast-wide plan. To be in compliance with the plan, coastal states have imposed restrictions on their commercial and recreational striped bass fisheries ranging from

combinations of catch quotas, size limits, closed periods and year-round moratoriums. Due to an improvement in spawning success, as judged by increases in annual values of the Maryland juvenile index, a limited fishery was established in fall, 1990. This transitional fishery existed until 1995 when spawning stock biomass reached sufficiently healthy levels (Field 1997). ASMFC subsequently declared Chesapeake Bay stocks to have reached benchmark levels and adopted Amendment 5 to the original FMP that allowed expanded state fisheries.

To document continued compliance with Federal law, the Virginia Institute of Marine Science (VIMS) has monitored the size and age composition, sex ratio and maturity schedules of the spawning striped bass stock in the Rappahannock River since December 1981 utilizing commercial pound nets and, since 1991, variable-mesh experimental gill nets. Spawning stock assessment was expanded to include the James River in 1994, utilizing commercial fyke nets and variable-mesh experimental gill nets. An experimental fyke net was established in the James River to assess its potential as a source for tagging striped bass. The use of fyke nets was discontinued after 1997. In conjunction with the monitoring studies, tagging programs have been conducted in the James and Rappahannock rivers since 1987. These studies were established to document the migration and relative contribution of these Chesapeake Bay stocks to the coastal population and to provide a means to estimate annual survival rates (S). With the re-establishment of fall recreational fisheries in 1993, the tagging studies were expanded to include the York River and western Chesapeake Bay to provide a direct estimation of the resultant fishing mortality (F). Commencing in 2005, these estimates of F were estimated from the striped bass tagged during the spring in the Rappahannock River.

Acknowledgments

We are deeply indebted to many people for their participation and/or contributions to the striped bass tagging and spawning stock assessment program. These include: the Anadromous Fishes Program staff; the cooperating commercial fishermen Ernest George, Joe Hinson, Albert and Stanley Oliff, Paul Somers, Clark Trader, Wayne France, John Wyatt and Glenn Wyatt; and Maryland Department of Natural Resources (Md DNR) staff Harry Hornick, Beth Versak, Andrea Hoover and Alexi Sharov.

Executive Summary

New Features: This year we include updated estimates of natural and fishing mortality based on the tag release and recapture data. A two-mortality period instantaneous rates model protocol was investigated and found to be superior to the single mortality instantaneous rates model first used in the 2007 report. Finally, the estimates of the progression of mycobacterial infection of Rappahannock River striped bass were refined and new estimates of mortality are introduced.

I. Assessment of the spawning stocks of striped bass in the Rappahannock and James rivers, Virginia, spring 2011.

Catch Summaries:

1. In 2011, 215 striped bass were sampled between 11 April and 2 May from two commercial pound nets in the Rappahannock River. The samples were predominantly male (61.4%) and young (63.7%). Females dominated the age nine and older age classes (72.2%). The mean age of the male striped bass was 6.0 years. The mean age of the female striped bass was 9.5 years.
2. During the 11 April – 2 May period, the 2006 and 2007 year classes were the most abundant in the Rappahannock River pound net samples and were 83.1% male. The contribution of age six and older males was only 43.2% of the total aged catch. Age seven and older females, presumably repeat spawners, were 30.6% of the total catch but represented 79.5% of all females caught.
3. In 2011, 190 striped bass were sampled between 31 March and 2 May in two experimental anchor gill nets in the Rappahannock River. The samples were predominantly male (60.0%) and young (70.0%). Females dominated the age nine and older age classes (70.2%). The mean age of the male striped bass was 5.8 years. The mean age of the female striped bass was 10.5 years.
4. During the 31 March – 2 May period, the 2006 and 2007 year classes were the most abundant in the Rappahannock River gill net samples and were 93.5% male. The contribution of age six and older males was only 29.5% of the total catch. Age seven and older females, presumably repeat spawners, were 28.9% of the total catch but were 88.7% of the total females caught.
5. In 2011, 455 striped bass were sampled between 31 March and 2 May in two experimental anchor gill nets (mile 62) in the James River. The samples were predominantly male (78.9%) and young (85.7%). Females dominated the age nine and older age classes (76.9%). The mean age of the male striped bass was 4.4 years. The mean age of the female striped bass was 9.1 years.

6. During the 31 March – 2 May period, the 2006 and 2007 year classes were the most abundant in the James River gill net samples and were 90.7% male. The contribution of age six and older males was only 17.1% of the total catch. Age seven and older females, presumably repeat spawners, were 13.2% of the total aged catch, but represented 64.5% of all females caught.

Spawning Stock Biomass Indexes (SSBI)

7. The Spawning Stock Biomass Index (SSBI) from the Rappahannock River pound nets was 17.0 kg/day for male striped bass and 27.6 kg/day for female striped bass. The male index was the sixth lowest in the 1991-2011 time series and 78.0% lower than the 2010 index. The 2011 index was 36.8% lower than the 21-year average. The 2011 female index was 56.3% lower than the 2010 index and 19.3% below the 21-year average.
8. The SSBI for the Rappahannock River gill nets was 36.8 kg/day for male striped bass and 52.2 kg/day for female striped bass. The male index was the second lowest in the 1991-2011 time series and 53.0% below the 21-year average. The female index was the sixth highest in the 1991-2011 time series and was 34.2% above the 21 -year average.
9. The SSBI for the James River gill nets was 67.5 kg/day for male striped bass and 65.0 kg/day for female striped bass. The male index was less than one half the 2010 index, and was 39.6% below the 18-year average. The female index was the sixth highest in the 18-year time series and was nearly 13.9% above the 18-year average.

Egg Production Potential Indexes (EPPI)

10. An index of potential egg production was derived from laboratory estimates of weight- and length-specific numbers of oocytes in the ovaries of mature females. The 2011 Egg Production Potential Index (EPPI, millions of eggs/day) for the Rappahannock River pound nets was 4.85 million eggs/day. This was the fourth lowest EPPI of the 2001-2011 time series. Older (8+ years) female stripers were responsible for 66.4% of the index.
11. The 2011 EPPI for the Rappahannock River gill nets was 8.41 million eggs/day. This was the fourth highest EPPI of the 2001-2011 time series. Older (8+years) female striped bass were responsible for 70.5% of the index.
12. The 2011 EPPI for the James River gill nets was 10.36 million eggs/day. This was the second highest EPPI of the 2001-2011 time series. Older (8+ years) female striped bass were responsible for 63.8% of the index.

Estimates of Annual Survival (S) based on age-specific catch rates

13. The cumulative catch rate (all age classes, sexes combined) from the Rappahannock River pound nets (8.88 fish/day) was the third lowest in the 1991-2011 time series. There was a decrease in all year classes from the 2010 values. The cumulative catch rate of male striped bass (5.43 fish/day) was the third lowest in the time series. The cumulative catch rate of female striped bass (3.46 fish/day) was the eighth lowest in the 1991-2011 time series and was 53.3% lower than the rate in 2010.
14. Year class-specific estimates of annual survival (S) for pound net data varied widely between years. The geometric mean S of the 1983-2003 year classes varied from 0.516-0.760 (mean = 0.636). The geometric mean survival rates differed between sexes. Mean survival rates for male stripers (1985-2003 year classes) varied from 0.317-0.615 (mean = 0.502) while mean survival rates of female stripers (1983-1998 year classes) varied from 0.380-0.698 (mean = 0.582).
15. The cumulative catch rate (all age classes, sexes combined) from Rappahannock River gill nets (19.10 fish/day) was the second lowest value in the 1991-2011 time series and was less than one half the rate in 2010. Cumulative catch rate of male stripers (9.90 fish/day) was the second lowest in the time series and was less than one fourth the rate in 2010. The cumulative catch rate of female striped bass (6.20 fish/day) was near the median in the time series, and 26.5% greater than the catch rate in 2010.
16. Year class-specific estimates of annual survival for gill net data varied widely between years. The geometric mean S of the 1984-2004 year classes varied from 0.408-0.747 (mean = 0.606). The mean survival rates for male stripers (1991-2004) varied from 0.216-0.701 (mean = 0.503). The mean survival rates for female stripers (1984-1997, excluding 1991) varied from 0.496-0.857 (mean = 0.651).
17. The cumulative catch rate (all age classes, sexes combined) from James River (mile 62) gill nets (45.40 fish/day) was the sixth lowest in the time series. The catch rate was one half the value from 2010. The cumulative catch rate for male striped bass (36.10 fish/day) was the seventh lowest of the time series, and was less than half the rate in 2010. The cumulative catch rate of female striped bass (9.60 fish/day) was 21.5% higher than the rate in 2010, and was the sixth highest value in the 1994-2011 time series.
18. Year class-specific estimates of annual survival in the James River varied widely between years. The geometric mean S of the 1984-2000 year classes varied from 0.338-0.711 (mean = 0.577). The mean survival rates of male stripers (1988-2000 year classes) varied from 0.286-0.612 (mean = 0.457). The mean survival rates of female stripers (1984-1998 year classes) varied from 0.339-0.853 (mean = 0.633).

Catch rate histories of the 1987-2003 year classes

19. Plots of year class-specific catch rates vs. year in the James and Rappahannock rivers from 1991-2011 showed a consistent trend of a peak in the abundance of male striped bass around age 4 or 5, followed by a steep decline. There was also a secondary peak of (mostly) female striped bass, usually around age 10.
20. The areas under the catch curves indicate that the 1993, 1994, 1996, and 1997 year classes were the strongest, and the 1990 and 1991 year classes the weakest in the Rappahannock River from 1987-2000. In the James River, the 1996, 1997, 1998, and 2000 year classes were the strongest and 1987 and 1988 year classes the weakest.

Growth rate of striped bass derived from annuli measurements

20. The scales of 222 striped bass were digitally measured and the increments between annuli were used to determine their growth history.
21. On average, striped bass grow about 137 mm fork length in their first year. The growth rate decreases with age to about 50 mm per year by age 10.
22. Striped bass were estimated to reach the minimum legal length for the resident fishery (18 in. total length) at age 3.5 and reach the minimum length for the coastal fishery (28 in. total length) at age eight.

Age determinations using scales and otoliths

23. A total of 223 specimens from 12 size ranges were aged by reading both scales and otoliths. The mean age of the otolith-aged striped bass was 0.22 years older than from the scale-aged striped bass. The two methodologies agreed on the age of the striped bass on 42.2% of the specimens and within one year 86.5% of the time.
24. Tests of symmetry applied to the age matrix indicated that the differences (higher or lower in age) between the two ageing methodologies were non-random ($p < .005$).
25. A paired t-test of the mean of the age differences produced by the two ageing methodologies found that the mean difference was not significantly different from zero ($p = .138$).
26. A Kolmogorov-Smirnov test of the age structures produced by the two ageing methodologies also indicated an overall significant difference, indicating that the two resultant age structures did represent an equivalent population.

II. Mortality estimates of striped bass (*Morone saxatilis*) that spawn in the Rappahannock River, Virginia, spring 2010-2011.

1. A total of 416 striped bass were tagged and released from pound nets in the Rappahannock River between 11 April and 9 May, 2011. Of this total, 225 were between 457-710 mm total length and considered to be predominantly resident striped bass and 188 were considered to be predominantly migrant striped bass (>710 mm TL). The median date of resident and migrant tag releases was 18 April.
2. A total of 74 striped bass (>457 mm TL), tagged during springs 1990-2010, were recaptured between 1 January and 31 December, 2010, and were used to estimate mortality. Most recaptures (58.1%) were caught within Chesapeake Bay (41.9% in Virginia, 16.2% in Maryland). However, other recaptures came from Massachusetts (14.9%), New Jersey (10.8%), New York (5.4%), Delaware and North Carolina (4.1% each) and Rhode Island (2.7% each).
3. A total of 35 migratory striped bass (>710 mm total length), tagged during springs 1990-2010, were recaptured between 1 January and 31 December, 2010, and were used to estimate the mortality. Most recaptures came from Massachusetts (31.4%), followed by New Jersey (22.9%), Chesapeake Bay (8.6% in Maryland and 5.7% in Virginia), New York, Delaware, and North Carolina (8.6% each) and Rhode Island (5.7%),.
4. The ASFMC Striped Bass Tagging Subcommittee established a data analysis protocol that involves deriving survival estimates from a suite of Seber models using program MARK. Nine of these models were applied to the recapture matrix, each reflecting a different parameterization over time. The resultant estimates of survival were 0.74 (> 457 mm TL) and 0.73 (>711 mm TL).
5. The MARK survival estimates were used to estimate exploitation rate, fishing mortality and natural mortality using Baranov's catch equation. The estimates of exploitation were 0.04 (>457 mm TL) and 0.05 (>711 mm TL). The estimates of fishing mortality were 0.04 (>457 mm TL) and 0.06 (>711 mm TL).
6. Alternatively, a suite of input models similar to the models used in program MARK were used to estimate survival, fishing and natural mortality using an instantaneous rates model. An analytical approach that allowed two periods of natural mortality was found to fit the data better than if constant natural mortality was used. The estimates of survival were 0.54 (>457 mm TL) and 0.60 (>711 mm TL). The estimates of fishing mortality were 0.05 (>457 mm TL) and 0.09 (>711 mm TL).

III. The role of Mycobacteriosis in elevated Natural Mortality of Chesapeake Bay striped bass: disease progression and developing better models for stock assessment and management.

1. Mycobacteriosis in striped bass is a chronic disease caused by various species of bacteria in the genus *Mycobacterium*. The disease appears as grey granulomatous nodules in internal organs and externally as ulcerous skin lesions. Mycobacteriosis in captive fishes is generally thought to be fatal, but this has not been established for wild striped bass.
2. The impact of the disease is poorly understood. Fundamental questions, such as mode of transmission, duration of disease stages, effects on fish movements, feeding, reproduction and mortality rates associated with the disease are unknown.
3. A total of 3,019 striped bass were tagged, assessed for external diseases indications, photographed and released from two pound nets in the upper Rappahannock (n=502) and five pound nets in the lower Rappahannock (n=2,517) River during fall, 2010. Only 29.2% of the total tagged were without any external sign of mycobacteriosis.
4. A total of 270 striped bass were tagged, assessed for external diseases indications, photographed and released from five pound nets in the lower Rappahannock River during spring, 2011. Only 38.1% of the total tagged were without any external sign of mycobacteriosis.
5. A total of 354 striped bass tagged during fall, 2010 were recaptured prior to 20 September, 2011. Unlike the results from 2006-2010, there was no prevalence of diseased striped bass in recaptures at large less than seven days that had suggested differential movement patterns between diseased and nondiseased striped bass.
6. A total of 52 striped bass tagged during spring, 2011 were recaptured prior to 20 September, 2011. One half of these recaptures were within seven days of release, but there was no evidence of differential movement between diseased and nondiseased striped bass.
7. It must be assumed that all fish have the same tag recovery rate to estimate survival rates, however, the disease severity may affect the movement of individual striped bass. It is therefore necessary to accumulate sufficient tag returns to estimate the relative survival rates.

8. Based on the recapture and reassessment of 431 tagged striped bass originally assessed as having a light or moderate mycobacterial infection, it was calculated that it takes 386 days for 100% of these striped bass to progress from light to moderate infection and 753 days for 100% progression from moderate to heavy infection.
9. The return rate for moderate and heavy mycobacteriosis-infected striped bass was less than the return rate for non-infected striped bass. The slope of the regression line of each category of infection plotted versus the non-infected striped bass produced a line with negative slope, indicating higher instantaneous natural mortality. This implies that the annual survival rates of moderate and heavy infected striped bass are 55%. Striped bass originally assessed as lightly infected had a less significant decrease in survival from the non-infected striped bass.

Table of Contents

Preface.....	ii-iii.
Acknowledgments.....	iv.
Executive summary.....	v-xi.
List of tables.....	xiv-xxi.
List of figures.....	xxii-xxvi.

I. Assessment of the spawning stocks of striped bass in the Rappahannock and James rivers, Virginia, spring 2011..... 1-112.

Introduction.....	2.
Materials and Methods.....	2-4.
Results.....	4-17.
Catch Summaries	4-7.
Spawning Stock Biomass Indexes	7-8.
Egg Production Potential Indexes.....	8.
Estimates of Annual Survival (S) based on Catch-Per-Unit-Effort.....	9-11.
Catch Rate Histories of the 1987-2003 Year Classes.....	11-15.
Growth Rate of Striped Bass Derived from Annuli Measurements	15.
Age Determinations using Scales and Otoliths	16-17.
Discussion.....	17-22.
Literature Cited.....	23-26.
Tables.....	27-91.
Figures.....	92-112.

II. Mortality estimates of striped bass (*Morone saxatilis*) that spawn in the Rappahannock River, Virginia, spring 2010-2011. 113-150.

Introduction.....	114-117.
Materials and Methods	117-120.
Capture and Tagging Protocol	117-118.
Analysis Protocol.....	118-120.
Results.....	121-123.
Spring 2011 Tag Release Summary.....	121.
Mortality Estimates 2010-2011	121-123.
Model Evaluations	123-124.
Discussion.....	124-126.
Literature Cited.....	127-129.
Tables.....	130-150.

III. The role of Mycobacteriosis in elevated Natural Mortality of Chesapeake Bay striped bass: disease progression and developing better models For stock assessment and management.	151-216.
Introduction.....	152-154.
Materials and Methods.....	154-159.
Capture and Tagging Protocol.	154.
Mycobacteriosis Assessment	155.
Analytical Approach	155-159.
Results.....	159-166.
Tag Release Summary.	159-160.
Tag Recapture Summary.	160-162.
Disease Progression in Rappahannock River Striped Bass, 2005-2010.	162-165.
Estimation of survival rates and relative survival rates.....	165-166.
Discussion.....	166-167.
Literature Cited.....	168-170.
Tables.....	171-191.
Figures.	192-215.

Appendix A. Daily river flows for the Rappahannock River, 30 March – 3 May, 1985-2011.	216-230.
Figures.....	217-230.

List of Tables

I. Assessment of the spawning stocks of striped bass in the Rappahannock and James rivers, Virginia, spring 2011.

1.	Numbers of striped bass in three age categories (year classes 2007-2009, 2003-2006 and 1992-2002) from pound nets in the Rappahannock River, by sampling date, spring, 2011.	27.
2.	Net-specific summary of catch rates and ages of striped bass (n= 215) in pound nets on the Rappahannock River, spring, 2011. Values in bold are the grand means for each column.	28.
3.	Length frequencies (TL in mm) of striped bass sampled from pound nets in the Rappahannock River, spring, 2011.	29.
4.	Mean fork length (mm), weight (g), standard deviation (SD) and CPUE (fish per day; weight per day), of striped bass from pound nets in the Rappahannock River, 11 April – 2 May, 2011.	30.
5.	Summary of the season mean (30 March – 3 May) catch rates and ages, by sex, from pound nets in the Rappahannock River, 1993-2011.	31.
6.	Numbers of striped bass in three age categories (year classes 2007-2009, 2003-2006 and 1992-2002) from gill nets in the Rappahannock River by sampling date, spring, 2011.	32.
7.	Summary of catch rates and mean ages of striped bass (n=190) from two gill nets in the Rappahannock River, spring, 2011. Values in bold are grand means for each column.	33.
8.	Length frequencies (TL in mm) of striped bass sampled from experimental gill nets in the Rappahannock River, spring, 2011.	34.
9.	Mean fork length (mm), weight (g), standard deviations (SD) and CPUE (number per day; weight per day) of striped bass from gill nets in the Rappahannock River, 31 March – 2 May, 2011.	35.
10.	Summary of the season mean (30 March – 3 May) catch rates and ages, by sex, from experimental gill nets in the Rappahannock River, 1993-2011.	36.

11.	Numbers of striped bass in three age categories (year classes 2007-2009, 2003-2006 and 1992-2002) from gill nets in the James River by sampling date, spring, 2011.	37.
12.	Summary of catch rates and mean ages of striped bass (n=455) from two gill nets in the James River, spring, 2011. Values in bold are grand means for each column.	38.
13.	Length frequencies (TL in mm) of striped bass sampled from experimental gill nets in the James River, spring, 2011.	39.
14.	Mean fork length (mm), weight (g), standard deviations (SD) and CPUE (number per day; weight per day) of striped bass from gill nets in the James River, 31 March – 2 May, 2011.	40.
15.	Summary of the season mean (30 March – 3 May) catch rates and ages, by sex, from experimental gill nets in the James River, 1995-2011.	41.
16.	Values of the spawning stock biomass index (SSBI) for male and female striped bass, by gear, in the Rappahannock River, 30 March – 3 May, 1991-2011.	42.
17.	Values of the spawning stock biomass index (SSBI) calculated from gill net catches of male and female striped bass in the James River, 30 March – 3 May, 1994-2011.	43.
18.	Predicted values of fecundity (in millions of eggs) of female striped bass with increasing fork length (mm), James and Rappahannock rivers combined.	44.
19.	Total, age-specific, estimated total egg potential (E, in millions of eggs/day) from mature (ages 4 and older) female striped bass, by river and gear type, 31 March - 2 May, 2011. The Egg Production Potential Indexes (millions of eggs/day) are in bold.	45.
20a,b.	Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	46-47.
21a,b.	Catch rates (fish/day) of year classes of male striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	48-49.
22a,b.	Catch rates (fish/day) of year classes of female striped bass sampled from pound nets in the Rappahannock River, 30 March - 3 May, 1991-2011.	50-51.

23a,b. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from pound nets in the Rappahannock River, 30 March - 3 May, 1991-2011.	52-53.
24a,b. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	54-55.
25a,b. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	56-57.
26a,b. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	58-59.
27a,b. Catch rates (fish/day) of year classes of male striped bass sampled from gill nets in the Rappahannock River, 30 March - 3 May, 1991-2011.	60-61.
28a,b. Catch rates (fish/day) of year classes of female striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	62-63.
29a,b. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	64-65.
30a,b. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	66-67.
31a,b. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.	68-69.
32a,b. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from gill nets in the James River, 30 March – 3 May, 1994-2011.	70-71.
33a,b. Catch rates (fish/day) of year classes of male striped bass sampled from gill nets in the James River, 30 March – 3 May, 1994-2011.	72-73.
34a,b. Catch rates (fish/day) of year classes of female striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011.	74-75.

35a,b. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from gill nets in the James River, 30 March – 3 May, 1994-2011.	76-77.
36a,b. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from gill nets in the James River, 30 March – 3 May, 1994-2011.	78-79.
37a,b. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from gill nets in the James River, 30 March – 3 May, 1994-2011.	80-81.
38a,b. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass sampled from pound nets in the Rappahannock River, 1991-2011.	82-83.
39a,b. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass sampled from gill nets in the Rappahannock River, 1991-2011.	84-85.
40a,b. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass sampled from gill nets in the James River, 1994-2011.	86-87.
41a,b. Back-calculated length-at-age (FL, in mm) for striped bass sampled from the James and Rappahannock rivers during spring, 2011.	88-89.
42. Data matrix comparing scale (SA) and otolith ages for chi-square test of symmetry. Values are the number of the respective readings of each combination of ages.	90.
43. Relative contributions of striped bass age classes as determined by ageing specimens (n=223) by reading both their scales and otoliths.	91.

II. Mortality estimates of striped bass (*Morone saxatilis*) that spawn in the Rappahannock River, Virginia, spring 2010-2011.

1. Summary data of striped bass tagged and released from pound nets in the Rappahannock River, spring, 2011 130.
2. Location of striped bass (> 457 mm TL) recaptured in 2010, that were originally tagged and released in the Rappahannock River during springs 1990-2010. 131.
3. Location of striped bass (> 710 mm TL) recaptured in 2010, that were originally tagged and released in the Rappahannock River during springs 1990-2010. 132.
4. Input recapture matrix for program MARK from striped bass (>457 mm TL) that were tagged and released in the Rappahannock River, springs 1990-2010. 133.
5. Input recapture matrix for program MARK from striped bass (>710 mm TL) that were tagged and released in the Rappahannock River, springs 1990-2010. 134.
6. Performance statistics (>457 mm TL), based on quasi-likelihood Akaike Information Criteria (QAIC), used to assess Seber (1970) models utilized in the ASMFC analysis protocol. 135.
7. Seber (1970) model estimates of unadjusted survival (\hat{S}) rates and adjusted rates of survival (\hat{S}_{adj}) and fishing mortality (\hat{F}) of striped bass (>457 mm TL) derived from the proportion of recaptures released alive (P_i) in the Rappahannock River, 1990-2010. 136.
8. Performance statistics (>710 mm TL), based on quasi-likelihood Akaike Information Criteria (QAIC), used to assess Seber (1970) models utilized in the ASMFC analysis protocol. 137.
9. Seber (1970) model estimates of unadjusted survival (\hat{S}) rates and adjusted rates of survival (\hat{S}_{adj}) and fishing mortality (\hat{F}) of striped bass (>710 mm TL) derived from the proportion of recaptures released alive (P_i) in the Rappahannock River, 1990-2010. 138.
10. Estimates of total mortality (Z), annual mortality (A), exploitation (U), fishing mortality (F) and natural mortality (M) from striped bass (>457 mm TL) tagged and released in the Rappahannock River, springs 1990-2010. 139.

11. Estimates of total mortality (Z), annual mortality (A), exploitation (U), fishing mortality (F) and natural mortality (M) from striped bass (>710 mm TL) tagged and released in the Rappahannock River, springs 1990-2010.	140.
12a. Input recapture matrix for IRCR analysis: from striped bass (>457 mm TL) tagged and released in the Rappahannock River, springs 1990-2010. Harvested recaptures only.	141.
12b. Input recapture matrix for IRCR analysis: from striped bass (>457 mm TL) tagged and released in the Rappahannock River, springs 1990-2010. Recaptures released with streamers cut off only.	142.
13a. Input recapture matrix for IRCR analysis: from striped bass (>710 mm TL) tagged and released in the Rappahannock River, springs 1990-2010. Harvested recaptures only.	143.
13b. Input recapture matrix for IRCR analysis: from striped bass (>710 mm TL) tagged and released in the Rappahannock River, springs 1990-2010. Recaptures released with streamers cut off only.	144.
14. Model Akaike weighting results (striped bass \geq 457 mm TL) for the 1M (constant) IRCR analysis.	145.
15. Model Akaike weighting results (striped bass \geq 457 mm TL) for the 1M (constant) and 2M IRCR analyses.	146.
16. Parameter estimates of survival (S), natural mortality (M) fishing mortality (F) and its standard error (SE) for striped bass \geq 457 mm TL from the IRCR analyses (1M and 2M), 1990-2010.	147.
17. Model Akaike weighting results (striped bass \geq 711 mm TL) for the 1M (constant) IRCR analysis.	148.
18. Model Akaike weighting results (striped bass \geq 711 mm TL) for the 1M (constant) and 2M IRCR analyses.	149.
19. Parameter estimates of survival (S), natural mortality (M) fishing mortality (F) and its standard error (SE) for striped bass \geq 711 mm TL from the IRCR analyses (1M and 2M), 1990-2010.	150.

III. The role of Mycobacteriosis in elevated Natural Mortality of Chesapeake Bay striped bass: disease progression and developing better models For stock assessment and management

1. Parameter estimates and standard errors (SE) from fitting two models to the Virginia striped bass spring tagging data (age 2 and greater).....171.
2. Tag release totals and mycobacteria infection index, by date, of striped bass in the upper and lower Rappahannock River sites, fall, 2010.172.
3. Tag release totals and mycobacteria infection index, by date, of striped bass in the upper and lower Rappahannock River sites, spring, 2011.....173.
4. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2010.....174.
5. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2010.175.
6. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2011.176.
7. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2011.177.
8. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2006 and recaptured from fall 2010 through summer 2011.178.
9. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2006 and recaptured from fall 2010 through summer 2011.179.
10. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2007 and recaptured from fall 2010 through summer 2011.....180.

11.	Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2007 and recaptured from fall 2010 through summer 2011.	181.
12.	Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2008 and recaptured from fall 2010 through summer 2011.	182.
13.	Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2008 and recaptured from fall 2010 through summer 2011.	183.
14.	Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2009 and recaptured from fall 2010 through summer 2011.	184.
15.	Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2009 and recaptured from fall 2010 through summer 2011.	185.
16.	Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2009 and recaptured from fall 2010 through summer 2011.	186.
17.	Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2009 and recaptured from fall 2010 through summer 2011.	187.
18.	Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2010 and recaptured from fall 2010 through summer 2011.	188.
19.	Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2010 and recaptured from fall 2010 through summer 2011.	189.
20.	Spatial necropsy summary of growth and days at liberty, by mycobacterial infection index and release area, of striped bass tagged in the upper and lower Rappahannock River sites through summer 2011.	190.
21.	Estimates of mortality associated with mycobacterial disease and estimated relative survival rates.	191.

List of Figures

I. Assessment of spawning stocks of striped bass in the Rappahannock and James rivers, Virginia, spring 2011.

1. Locations of the commercial pound nets and experimental gill nets sampled in spring spawning stock assessments of striped bass in the Rappahannock River, 1991-2011.92.
2. Locations of the experimental anchor gill nets sampled in spring spawning stock assessments of striped bass in the James River, springs 2003-2011.....93.
3. Daily and historic mean river flows (cf/s) for the Rappahannock River during the 30 March - 30 April spawning stock assessment period, spring 2011.....94.
4. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1987 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1991-2011.95.
5. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1988 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1991-201196.
6. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1989 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1991-2011.97.
7. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1990 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1991-2011.98.
8. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1991 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1992-2011.99.
9. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1992 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1993-2011.100.
10. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1993 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1994-2011.101.

11.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1994 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1995-2011.	102.
12.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1995 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1996-2011.	103.
13.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1996 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1997-2011.	104.
14.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1997 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1998-2011.	105.
15.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1998 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 1999-2011.	106.
16.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1999 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 2000-2011.	107.
17.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2000 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 2001-2011.	108.
18.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2001 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, spring, 2001-2011.	109.
19.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2002 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 2002-2011.....	110.
20.	Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2003 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 2003-2011.....	111.
21.	Magnitude of the age differences (otolith = 223) by reading both their scales and otoliths, spring 2011.....	112.

III. The role of Mycobacteriosis in elevated Natural Mortality of Chesapeake Bay striped bass: disease progression and developing better models For stock assessment and management

1. Gross clinical signs of mycobacteriosis in Chesapeake Bay striped bass. A) severe ulcerative dermatitis. B) multi-focal pale gray nodules within the spleen.192.
2. A spectrum of gross skin lesions attributable to mycobacteriosis in striped bass, *Morone saxatilis*. a) mild scale damage and scale loss (arrows). b) pigmented fous showing pin-point erosion through an overlying scale (arrow). c) early ulceration exhibiting focal loss of scales, mild pin-point multifocal pigmentation and underlying exposed dermis. d) large advanced shallow roughly textured ulceration exhibiting hyper-pigmentation and hemorrhage. e) late stage healing lesion exhibiting hyper-pigmentation, reformation of scales and re-epithelialization and closure of the ulcer. f) Ziel Neelsen stain of a histologic section of a skin lesion exhibiting granulomatous inflammation and acid-fast rod-shaped mycobacteria. g) histologic section showing normal healthy skin composed of epidermis, scales, dermis and underlying skeletal muscle. h) histologic section through a skin ulcer showing loss of epidermis and scales and extensive granuloma formation (G). 193.
3. a) Size distribution (fork length in mm), by infection index, of striped bass tag releases from the upper Rappahannock River, fall 2010. b) Relative proportion of each infection index, by fork length, of the tag releases.194.
4. a) Size distribution (fork length in mm), by infection index, of striped bass tag releases from the lower Rappahannock River, fall 2010. b) Relative proportion of each infection index, by fork length, of the tag releases.195.
5. a) Size distribution (fork length in mm), by infection index, of striped bass tag releases from the lower Rappahannock River, spring 2011. b) Relative proportion of each infection index, by fork length, of the tag releases.196.
6. Progression in the mycobacteriosis skin severity index, with age, of the 2002 year class of striped bass in the lower Rappahannock River, falls 2005-2010.197.
7. Progression in the mycobacteriosis skin severity index, with age, of the 2003 year class of striped bass in the lower Rappahannock River, falls 2006-2010.198.
8. Progression in the mycobacteriosis skin severity index, with age, of the 2004 year class of striped bass in the lower Rappahannock River, falls 2007-2010.199.
9. Progression in the mycobacteriosis skin severity index, with age, of the 2005 year class of striped bass in the lower Rappahannock River, falls 2008-2010.200.

10.	Progression of mycobacteriosis from lightly diseased at time of release to moderately diseased versus time-at-large for striped bass tagged and released in the Rappahannock River, fall 2005 to present (combined). Numbers next to the data points indicate number of recaptures.	201.
11.	Progression of mycobacteriosis from moderately diseased at time of release to severely diseases versus time-at-large for striped bass tagged and released in the Rappahannock River, fall 2005 to present (combined). Numbers next to the data points indicate number of recaptures.	202.
12.	Progression of pigmented foci (PF) of uninfected striped bass based on reassessment of recaptured striped bass originally tagged and released in the Rappahannock River, falls 2005-2010.	203.
13.	Mean recaptured fork length of fish tagged at the upper and lower locations of the Rappahannock River, broken down by site and release disease condition.	204-205.
14.	Boxplot of days at liberty by disease progression based on site and release disease assessment.	206-207.
15.	Progression of mycobacteriosis from diseased at time of release to the next level of severity versus time-at-large for striped bass tagged and released in the upper and lower Rappahannock River, fall 2005 to present (combined).	208-211.
16.	Logarithm of the ratio of returns of fish tagged in disease condition x and disease condition 0 (fish in condition 0 are “clean”, showing no signs of the disease) as a function of time at liberty. Numbers next to the data points are the number of tag returns. The slope of the weighted regression estimates the difference in instantaneous total mortality rates, $Z - Z_0$, which is equivalent to the difference in instantaneous natural mortality rates (because the F component of Z is assumed to be the same for both groups of fish). A) Condition 3 versus condition 0 B) Condition 2 versus condition 0. C) Condition 1 versus condition 0. D) Conditions 2 and 3 combined versus condition 0.	212-215.

Appendix A. Daily river flows for the Rappahannock River, 30 March – 3 May, 1985-2010.

1. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, spring 2010.217.
2. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2008-2009.218.
3. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2006-2007.219.
4. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2004-2005.220.
5. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2002-2003.221.
6. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2000-2001.222.
7. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1998-1999.223.
8. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1996-1997.224.
9. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1994-1995.225.
10. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1992-1993.226.

11.	Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1990-1991.	227.
12.	Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1988-1989.	228.
13.	Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1986-1987.	229.
14.	Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, spring 1985.	230.

I. Assessment of the spawning stocks of striped bass in the Rappahannock and James rivers, Virginia, spring, 2011.

Striped Bass Assessment and Monitoring Program
Department of Fisheries Science
School of Marine Science
Virginia Institute of Marine Science
The College of William and Mary
Gloucester Point, VA. 23062-1346

Introduction

Every year, striped bass migrate along the US east coast from offshore and coastal waters and then enter brackish or fresh water to spawn. Historically, the principal spawning areas in the northeastern US have been the Hudson, Delaware and Chesapeake estuarine systems (Hardy 1998). The importance of the Chesapeake Bay spawning grounds to these stocks has long been recognized (Merriman 1941, Raney 1952). In the Virginia tributaries of Chesapeake Bay, peak spawning activity is usually observed in April and is associated with rapidly rising water temperatures in the range of 13-19° C (Grant and Olney 1991). Spawning is often completed by mid-May, but may continue until June (Chapoton and Sykes 1961). Spawning grounds have been associated with rock-strewn coastal rivers characterized by rapids and strong currents on the Roanoke and the Susquehanna rivers (Pearson 1938). In Virginia, spawning occurs over the first 40 km of the tidal freshwater portions of the James, Rappahannock, Pamunkey and Mattaponi rivers (Grant and Olney 1991; Olney et al. 1991; McGovern and Olney 1996).

The Atlantic States Marine Fisheries Commission (ASMFC) declared that the Chesapeake Bay spawning stocks were fully recovered in 1995 after a period of very low stock abundance in the 1980's. This statement of recovered status was based on estimated levels of spawning stock biomass that were found in 1995 to be equal or greater than the average levels of the 1960-72 period (Rugulo et al. 1994). Thus, continued assessment of spawning stock abundance is an important component of ASMFC mandated monitoring programs. To this end, the Virginia Institute of Marine Science (VIMS) began development of spawning indexes that depict annual changes in catch rates of striped bass on the spawning grounds of the James and the Rappahannock rivers. These rivers represent the major contributors to the Chesapeake Bay stocks that originate from Virginia waters.

Materials and Methods

Samples of striped bass for biological characterization of the spring spawning stocks were obtained from the Rappahannock River from between 31 March – 3 May, 2011. This year, adverse weather conditions prevented setting of the pound nets at the start of the season. Therefore, samples from these pound nets were delayed until 11 April, 2011. In addition, one of the three pound nets normally sampled (net at mile 45) was not set this year. Due to the delay, measurements and sex of the striped bass from the net designated for the monitoring sample were recorded and the stripers greater than 18 inches then tagged and released. All undersize stripers and any striped bass of indeterminate sex were brought back to the lab. Samples (the entire catch of striped bass from each gear) were taken twice-weekly (Monday and Thursday) from among two commercial pound nets (river miles 46 and 47) in the Rappahannock River. Pound nets are fixed commercial gears that have been the historically predominant gear type used in the river and are presumed to be non size-selective in their catches of striped bass. The established protocol (Sadler *et al.* 1999) was to alternate the choice of the net sampled but weather constraints often dictated whether that net could be sampled. In addition, data from pound nets sampled in 1991 and 1992 were included to expand the time series. These samples were consistent in every respect to the 1993-2001 samples with the following exceptions in

1991: two samples (3 and 17 April) came from a pound net at river mile 25 and samples were obtained weekly vs. twice weekly.

In addition to the pound nets, samples were also obtained twice-weekly from variable-mesh experimental anchored gill nets (two at river mile 48 on the Rappahannock River and two at river mile 62 on the James River, Figures 1 and 2). The variable-mesh gill nets deployed on both rivers were constructed of ten panels, each measuring 30 feet (9.14 m) in length, and 10 feet (3.05 m) in depth. The ten stretched-mesh sizes (in inches) were 3.0, 3.75, 4.5, 5.25, 6.0, 6.5, 7.0, 8.0, 9.0, and 10.0. These mesh sizes correspond to those used for spawning stock assessment by the Maryland Department of Natural Resources. The order of the panels was determined by a randomized stratification scheme. The mesh sizes were divided into two groups, the five smallest and the five largest mesh sizes. One of the two groups was randomly chosen as the first group, and one mesh size from that group was randomly chosen as the first panel in the net. The second panel was randomly chosen from the second group, the third from the first group, and so forth, until the order was complete. The order of the panels in the first net was (in inches) 8.0, 5.25, 9.0, 3.75, 7.0, 4.5, 6.5, 6.0, 10.0, and 3.0, and in the second net the order was (in inches) 8.0, 3.0, 10.0, 5.25, 9.0, 6.0, 6.5, 3.75, 7.0, and 4.5. In 2004, a manufacturing error resulted in two nets of the first configuration being utilized.

Striped bass collected from the monitoring sites were measured and weighed on a Limnoterra FMB IV electronic fish measuring board interfaced with a Mettler PM 30000-K electronic balance. The board records lengths (FL and TL) to the nearest mm, receives weight (g) input from the balance, and allows manual input of sex and gonad maturity into a data file for subsequent analysis. Scales were collected from between the spinous and soft dorsal fins above the lateral line for subsequent aging, using the method established by Merriman (1941), except that impressions made in acetate sheets replaced the glass slide and acetone. Otoliths were extracted from a stratified subsample of the striped bass, processed for aging, and compared to their scale-derived ages. The weights of the striped bass tagged and released rather than brought to the lab were estimated using sex-specific regressions of weight vs. length.

The otolith subsample was the first 10 striped bass of each sex sampled from each of the following size ranges (fork length, in mm): <165, 166-309, 310-419, 420-495, 496-574, 575-659, 660-724, 725-779, 780-829, 830-879 and 880-900. All striped bass greater than 900 mm fork length were sampled. These size ranges roughly correspond to age classes based on previous (scale-aged) data.

The otoliths were cleansed of external tissue material by successive rinses in water immediately after extraction. The otoliths were prepared for ageing by placing the left sagitta on melted crystal bond and sectioned to a one millimeter thickness on a Buehler isomet saw. The sections were then polished on a Metaserv 2000 grinder. The polished section was immersed in a drop of mineral oil and viewed through an Olympus BX60 compound microscope at 4-20X. Each otolith was aged at least twice at different times by each of two readers using the methods described by Wischniowski and Bobko (1998).

All readable scales from the otolith-scale comparison were aged using the microcomputer program DISBCAL of Frie (1982), in conjunction with a sonic digitizer-microcomputer complex (Loesch et al. 1985). Growth increments were measured from the focus to the posterior edge of each annulus. In order to be consistent with ageing techniques of other agencies, all striped bass were considered to be one year older on 1 January of each year. Scale ages were used exclusively, except when a comparison with its companion otolith age was made.

The spawning stock biomass index (SSBI) for striped bass was defined (Sadler et al. 1999) as the 30 March - 3 May mean CPUE (kg/net day) of mature males (age 3 years and older), females (age 4 years and older) and the combined sample (males and females of the specified ages). An alternative index, based on the fecundity potential of the female striped bass sampled, was investigated and the results compared with the index based on mean female biomass.

To determine fecundity, the geometric mean of the egg counts of the gonad subsamples for each ripe female striped bass collected in 2001-2003 was calculated. A non-linear regression was fitted to data of total oocytes versus fork length. The resultant equation was then applied to the fork lengths of all mature (4+ years old) females from the pound net and gill net samples and the Egg Production Potential Index (EPPI) was defined as the mean number of eggs potentially produced per day of fishing effort by the mature female (age 4+) striped bass sampled from 30 March - 3 May.

Estimates of survival (S, the fraction surviving after becoming fully recruited to the stock) were calculated by dividing the catch rate (number/day) of a year class in year a+1 by the catch rate (number/day) of the same year class in year a. If the survival estimate between successive years was >1, the estimate was derived by interpolating to the following year. The geometric mean of S was used to estimate survival over periods exceeding one year (Ricker 1975). Separate estimates of survival were made for male and female striped bass, as well as the sexes combined.

Analysis of the differences in the ages estimated by reading the scales and otoliths from the same specimen were made using tests of symmetry (Evans and Hoenig 1998, Hoenig et al. 1995). Differences in the resultant mean ages from the two methods were tested using both two-tailed paired and unpaired t-tests (Zar 1999). The age class distributions resulting from the two ageing methods were compared using the non-parametric Kolmogorov-Smirnov two-sample test (Sokal and Rohlf 1981).

Results

Catch Summaries

Rappahannock River:

Pound nets: Striped bass (n= 215) were sampled between 11 April - 2 May, 2011 from the pound nets in the Rappahannock River. The number of striped bass sampled was 79.5% lower than the sample in 2010 (n= 1,048) and 66.2% lower than the 19-year average (n=635.5). Total

catches varied from 4-68 striped bass, with the peak catch on 11 April (Table 1). Surface water temperature increased slowly from 9.7 EC on 11 April to 16.9 EC on 29 April, and then increased to 21 EC on 2 May. River flows were below average at the start of the season, but unlike recent years there was a surge in river flows in mid-April which resulted in above average flows for the remainder of the sampling season (Fig 3). Salinities were 0.0-0.1 p.p.t. throughout the sampling season. Catches of female striped bass peaked on 11 and 14 April and were dominated by the pre-2003 year classes. Males made up 61.4% of the total catch, which was well below the 19-year average (76.6%). The 2003-2006 year classes (five to eight years old) comprised 44.7% of the total catch. In contrast, in 2010 the 2002-2005 year classes comprised 49.3% of the total catch. Males dominated the 2007-2009 year classes (97.4%) and the 2003-2006 year classes (74.0%), but females dominated the 1992-2002 year classes (72.2%).

Biomass catch rate (g/day) of male striped bass peaked on 11 April. The biomass catch rate of female striped bass peaked sharply on 14 April (Table 2). The numeric catch rate of males exceeded that of females except for 18-25 April. Unlike 2008, but consistent with most previous years, the biomass catch rates for female striped bass exceeded that for males overall (1.62:1), peaking on 25 April (4.0:1). The mean ages of male striped bass varied from 4.3-10.0 years by sampling date, with the oldest mean age occurring on 21 April. The mean ages of females varied from 8.3-10.4 years by sampling date, which was a much smaller range than in 2010 (7.3-14.0 years).

There was a broad peak in abundance of striped bass (mostly male) between 480-640 mm total lengths in the pound net samples (Table 3). This size range accounted for 34.0% of the total sampled. There was a secondary peak in abundance of predominantly female striped bass between 820-930 mm total lengths. Consistent with previous years, the striped bass from 630-710 mm total length accounted for only 10.2% of the total sample. The total contribution of striped bass greater than 710 mm total length (the minimum total length for the coastal fishery) was 51.2% (vs. 25.6% in 2010).

During the 11 April – 2 May period, the 2006 (15.3%) and 2007 (12.1%) year classes were the most abundant (Table 4). These year classes were 83.1% male. The contribution of males age six and older (the pre-2006 year classes) was 32.2% of the total aged catch. These year classes were most vulnerable to commercial and recreational exploitation within Chesapeake Bay. The contribution of females age seven and older, presumably repeat spawners, was only 30.8% of the total aged catch, but was also 79.5% of the total females captured. The catch rate (fish/day) of male striped bass was 5.5, which is 65.6% below the 19-year average (Table 5). The catch rate of female striped bass (3.5 fish/day) was below the 19-year average, and was the lowest since 2008. The biomass catch rates (kg/day) of both sexes of striped bass were well below the average of the 19-year time series. The mean age of the male striped bass was the highest in the 19 time series. The mean age of the female striped bass was lower than 2010, but above the mean value in the time series.

Experimental gill nets: Striped bass (n= 190) were also sampled between 31 March and 2 May, 2011 from two multi-mesh experimental gill nets in the Rappahannock River. The total catch was lower than the total catch in 2010 (n=486) and was 70.9% below the 19-year average. Total

catches peaked on 14 and 18 April (Table 6). Total catches of male striped bass were at the highest on 14 and 18 April. Total catches of female striped bass peaked on 18 April. Males made up 68.9% of the total catch. Males dominated the 2007-2009 year classes (97.9%) and the 2003-2006 year classes (78.8%), but the 1992-2002 year classes were 70.2% female.

Biomass catch rate (g/day) of male striped bass was highest on 14 April (Table 7). In contrast to 2010, where the catch rate (fish/day) of males exceeded that of females on every sampling date, the catch rate of female striped bass was greater from 21 April through 2 May. The mean ages of male striped bass varied from 5.0-7.5 years by sampling date, with the oldest males being most abundant on 11 April. The biomass catch rate of female striped bass (g/day) peaked sharply on 14 April. The mean ages of females varied from 7.0-15.7 years by sampling date, with the oldest females (age nine and older) being most abundant from 31 March through 7 April.

Unlike 2009, but consistent with previous years, there was a broad peak in the distribution of length frequencies of striped bass in the gill net samples between 480-630 mm TL (Table 8). There was a secondary peak of striped bass 740-820 mm TL. This secondary peak was consistent with the results of most previous years, but the secondary peak was less apparent from 2006-2009. Unlike 2008 and 2009, but consistent with previous years, the total contribution of striped bass greater than 840 mm total length from the gill nets (22.1%) was lower than from the pound nets (29.8%). The total contribution of striped bass greater than 710 mm total length was 47.9% in the gill nets compared to only 19.1% in 2010.

During the 31 March – 2 May period, the 2007 (17.9%) and 2006 (14.7%) year classes were most abundant (Table 9). These year classes were 93.5% male. The contribution of males age six and older (the pre-2006 year classes) was 29.5% of the total aged catch. These year classes were most vulnerable to commercial and recreational exploitation within Chesapeake Bay. The contribution of females age seven and older, presumably repeat spawners, was 28.9% of the total aged catch but was 88.7% of the total females captured. The catch rate of male striped bass (12.8 fish/day) was the lowest in the 19-year time series and was 70.7% below the average (Table 10). However, the catch rate of female striped bass (6.2 fish/day) was near the median in the time series and was 6.9% above the 19-year average. The biomass catch rates (g/day) for male striped bass was also the lowest in the time series and was 51.2% below the 19-year average. The biomass catch rate for female striped bass was the sixth highest in the time series and was 39.2% above the 19-year average.

James River:

Experimental gill nets: Striped bass (n= 455) were sampled between 31 March and 2 May, 2011, from two multi-mesh experimental gill nets at mile 62 in the James River. Total catches peaked on 11 and 14 April. Male, mostly young striped bass were primarily responsible for the peak catch (Table 11). Catches of female striped bass also peaked on 14 April. Males dominated the 2007-2009 year classes (97.2%) and the 2003-2006 year classes (81.1%), but the 1992-2002 year classes were predominantly female (76.9%).

Biomass catch rates (g/day) of male striped bass peaked on 11 and 14 April, but were high on all but three occasions (Table 12). The catch rates of female striped bass peaked strongly on 14 April then again on 25 April. The biomass catch rate of males exceeded that of females prior to 14 April, then the biomass catch rate for females dominated and the seasonal average was equal. The mean ages of male striped bass varied from only 4.3-7.6 years by sampling date. The mean ages of females varied from 5.9-13.7 years by sampling date.

There was a peak of striped bass 410- 620 mm total length in the gill net length frequencies (Table 13). This size range accounted for 69.9% of the total striped bass sampled. In contrast to the pound net samples from the Rappahannock River, the striped bass greater than 840 mm total length accounted for 11.9% of the total sampled. The total contribution of striped bass greater than 710 mm total length was 18.9%.

During the 31 March – 2 May period, the 2006 (25.9%) and 2007 (23.5%) year classes were the most abundant in the gill nets (Table 14). These year classes were 90.7% male. The contribution of males age six and older (the pre-2006 year classes) was only 17.4% of the total aged catch. These year classes were most vulnerable to commercial and recreational exploitation within Chesapeake Bay. The contribution of females age seven and older, presumably repeat spawners, was only 13.4% of the total aged catch, but represented 64.5% of the total females captured.

The catch rate of male striped bass (35.9 fish/day) was less than one half than the catch rate for 2010, and was 45.5% below the 17-year average (Table 15). However, the catch rate of female striped bass (9.6 fish/day) was higher than for 2010 and was 5.5% above the 17-year average. The biomass catch rate (g/day) of male striped bass was also less than one half the biomass catch rate for 2010, and was 42.8% below the 17-year average. However, the biomass catch rate of female striped bass was higher than in 2010, and was 19.7% above the 17-year average and was the highest since 2006. The mean age of male striped bass has varied from only 4.3-4.9 years by sampling year, while the mean age of female striped bass varied from 6.3-9.8 years.

Spawning Stock Biomass Indexes

Rappahannock River:

Pound nets: The Spawning Stock Biomass Index (SSBI) for spring 2011 was 17.0 kg/day for male striped bass and 27.6 kg/day for female striped bass. The index for male striped bass was the lowest since 2002 and the sixth lowest in the 21-year time series. It was 72.0% lower than the index value for 2010, and 36.8% below the 21-year average (Table 16). The magnitude of the index for male striped bass was largely determined by the 2003-2007 year classes (63.7%). The index for female striped bass was 56.3% lower than the 2010 index. It was near the median in the time series, but 19.3% below the 21-year average (Table 16). The magnitude of the index for the females was largely determined by the 1999-2002 year classes (67.4%).

Experimental gill nets: The Spawning Stock Biomass Index for spring 2011 was 36.8 kg/day for male striped bass and 52.2 kg/day for female striped bass. The index for male striped bass

was the second lowest of the time series, about one third of the 2009 index, and 53.0% below the 21-year average (Table 16). The 2002- 2004 year classes contributed 54.9% of the biomass in the male index. In contrast to the pound net index, the index for female striped bass was 6.7% above the 2010 index and was 34.2% above the 21-year average. The 1996-1998 year classes contributed 37.9% of the biomass in the female index.

James River:

Experimental gill nets: The Spawning Stock Biomass Index for spring 2011 was 67.5 kg/day for male striped bass and 65.0 kg/day for female striped bass. The male index was lowest since 1999, less than one half the 2010 index, and 39.6% below the 18-year average (Table 17). The 2005-2007 year classes contributed 63.8% of the biomass in the male index. However, the female index was the sixth highest in the time series, was 15.2% higher than the 2010 index, and was 13.9% above the 18-year average. The 1996-2001 year classes each accounted for 70.0% of the biomass in the female index.

Egg Production Potential Indexes

The number of gonads sampled, especially of the larger females, was insufficient to produce separate length-egg production estimates for each river. The pooled data (2001-2003) produce a fork length-oocyte count relationship as follows:

$$N_o = 0.000857 \times FL^{3.1373}$$

where N_o is the total number of oocytes and FL is the fork length (>400) in millimeters. Using this relationship, the predicted egg production was 125,000 oocytes for a 400-mm female and 3,719,000 oocytes for a 1180-mm female striped bass (Table 18). The 2011 Egg Production Potential Indexes (EPPI, Table 19) for the Rappahannock River were 4.85 (pound nets) and 8.41 (gill nets). The 2011 EPPI for the James River was 10.36. The indexes for both the Rappahannock and James rivers were heavily dependent on the egg production potential of the 1996-2001 year class females (66.4% in the pound nets, 70.5% in the Rappahannock gill nets and 63.8% in the James River gill nets). Previous values for the EPPI for 2001-2010 from the Rappahannock River were 3.992, 1.764, 9.829, 10.55, 6.30, 4.01, 13.792, 8.66, 6.87 and 9.87 (pound nets) and 4.039, 6.070, 3.724, 8.432, 3.06, 6.27, 9.915, 6.58, 9.04 and 7.02 (gill nets). Previous values for the EPPI for 2001-2010 from the James River were 5.286, 6.709, 6.037, 4.922, 3.24, 15.1, 8.396, 8.86, 9.52 and 8.50 respectively (Sadler et al 2001, 2002, 2003, 2004, 2005, 2006, 2007, 2008, 2009 and 2010). Thus, the EPPI values for the pound nets in the Rappahannock River signaled a decline in the status of the spawning stock from the 2010 values, while the EPPI value for the gill nets in the Rappahannock and James rivers were above their 2010 values. Modest changes in the methodology (utilizing fully mature ovaries solely rather than ovaries in various states of maturation) in the 2001-2011 indexes preclude direct comparison with the 1999 and 2000 indexes.

Estimates of Annual Survival (S) based on Catch-Per-Unit-Effort

Rappahannock River:

Pound nets: Numeric catch rates (fish/day) of individual year classes from the 1991-2011 samples are presented in Tables 20-22. The cumulative annual catch rate of all year classes for 2011 was the third lowest in the time series and was 74.5% lower than the cumulative catch rate for 2010 (Tables 20a,b). The decrease was the result of lower catch rates for almost all year classes. The catch rate of males was dominated by four through eight year olds (2003-2007 year classes, Tables 21a,b). These five age classes contributed 73.7% of the total male catch. Using the maximum catch rate of the resident males as an indicator, the 1995-1997 year classes were strongest and the 1990 and 1991 year classes were the weakest. No pre-2000 year class males were captured. The cumulative catch rate of female stripers was the eighth lowest of the time series, and was 53.3% lower than the catch rate in 2010 (Tables 22a,b). The 1999-2003 year classes accounted for 63.9% of the total female catch.

The range of overall ages was unchanged from 1991-2011, consisting mainly of 2-10 year old males and 4-16 year old females, but sex-specific changes in the age-structure have occurred. The age at which abundance peaked for males has decreased from age five (1992-1994) to age four (1997-2002, 2006-2010). The catch rate of four and five year olds were near equal in 2003 and 2004 and again in 2011, but the peak was age three in 2005. There has been an even more significant change in the age composition of the female spawning stock. From 1991-1996, the cumulative proportion of females age eight and older ranged from 0.134-0.468 (mean = 0.294) as their cumulative catch rate ranged from 0.75-2.1 fish/day (mean = 1.32). From 1997-2001 the range in the cumulative proportion of females age eight and older increased to 0.770-0.872 (mean = 0.825) as cumulative catch rates ranged from 1.4-4.5 fish/day (mean = 2.84). In 2002, the cumulative proportion of female striped bass age eight and older decreased to 0.508, then increased to 0.787-0.929 from 2003-2007. However, the cumulative catch rate dropped to 0.678 in 2008 and 0.593 in 2009, but rebounded to 0.780 in 2010, and .757 in 2011.

Estimates of annual survival (S) for the individual year classes and their overall geometric means are presented in tables 23-25. While annual survival estimates varied widely among years, due to strong or weak overall catches, the geometric mean survival rates (1991-2011) of the 1983-2003 year classes (sexes combined) varied from 0.516-0.760 (Tables 23a,b) with an overall mean survival rate of 0.636. These year classes have survival estimates across a minimum of four years. There were widely divergent estimates of annual survival of male and female striped bass. The geometric mean survival rate (1991-2011) of the 1985-2003 year classes of males varied from 0.317-0.615 (Tables 24a,b) with an overall mean survival rate of 0.502. These year classes have been the major target of the fall recreational and commercial fisheries that reopened in 1993. The geometric mean survival rate (1991-2011) of the 1983-1998 year classes of females varied from 0.380-0.698 (Tables 25a,b) with an overall mean survival rate of 0.582.

Experimental gill nets: Numeric catch rates (fish/day) of individual year classes from 1991-2011 are presented in Tables 26-28. The cumulative annual catch rate (all age classes, sexes combined) for 2011 from the gill nets was the second lowest in the time series, and was less than

one half the rate for 2010 and the lowest since 1992 (Tables 26a,b). The high cumulative catch rate in 2007 was driven by the catch rates of the 2003 and 2004 year classes (3 and 4 years of age) of striped bass. These age classes were still significant contributors in 2011, but the decrease in the cumulative catch rate was driven by low catches of 2004-2007 year class striped bass. The age of peak abundance had changed from age five (1992-1996, 2002 and 2010) to age four (1997, 1998, 2000, 2001, 2003, 2007, 2008 and 2011) and age three (1999, 2004 and 2006). The cumulative catch rate of male striped bass was also the second lowest in the time series and one fourth the catch rate in 2010 (Tables 27 a,b).

Using the maximum catch rate of the resident males as an indicator, the 1993, 1994 and 1997 year classes were the strongest and the 1990, 1991 and 2000 year classes the weakest. The catch rates of male striped bass declined rapidly after ages five or six. These age classes are the primary target of the recreational and commercial fisheries.

However, the cumulative catch rate of female striped bass was 26.5% greater than the cumulative catch rate for 2010 and near the median of the time series (Tables 28a,b). In 2004, the increased catch rates for 8-14 year-old females gave evidence of secondary peak of abundance across several year classes. This was not evident from the catches in 2005-2010. This bimodal distribution of abundance with age had been noted for the pound net catches, but has generally not been evident in the gill net catches.

The overall age structure from 1991-2011 consisted of 2-12 year old males (Tables 27a,b) and 2-14 year old females (Tables 28a,b). However, in 2011 there were small catches of the 1992-1995 year classes. The proportion of males age six and older (0.27) was less than 2010 (0.43), but consistent with the 2002-2008 values after being 0.03-0.06 from 1997-2001. The proportion of female striped bass age eight and older (0.84) was higher than 2010 (0.78) but consistent with the values since 2004. The proportion of females age eight and older increased from 0.148 to 0.652 from 1991 to 1996, declined from 0.652 to 0.315 from 1996 to 2002 (except 0.707 in 2001), then rebounded to 0.594 in 2003 and 0.786-0.839 from 2004-2011.

Estimates of annual survival (S) for the individual year classes and their overall geometric means are presented in Tables 29-31. While annual survival estimates varied widely among years, due to strong or weak overall catches, the geometric mean survival rate (1991-2011) of the 1984-2004 year classes (sexes combined) varied from 0.408-0.747 (Tables 29a,b) with an overall mean survival of 0.606. There were widely divergent estimates of annual survival of male and female striped bass. The mean survival rate (1991-2011) of the 1991-2004 year classes of males varied from 0.276-0.701 (Tables 30a,b) with an overall mean survival of 0.503. These year classes have been the major target of the fall recreational and commercial fisheries that reopened in 1993. The mean survival rate (1991-2011) of the 1984-1997 (excluding 1991) year classes of females varied from 0.496-0.857 (Tables 31a,b) with an overall mean survival rate of 0.651. The overall survival estimate of male striped bass was lower than that calculated from the pound nets. The estimate of female survival rates, although slightly greater than the pound net estimate, was based on fewer year classes than the estimate from the pound nets due to the relative rareness of the oldest females in the samples.

James River:

Experimental gill nets: Numeric catch rates (fish/day) of individual years classes from 1984-2011 are presented in Tables 32-34. The cumulative annual catch rate (all age classes, sexes combined) for 2011 was half the value from 2010 and was the sixth lowest in the time series (Tables 32a,b). The cumulative catch rate was driven by high catch rates for the three to five year old (2008-2006 year classes), mostly male striped bass.

The overall age structure of the samples has remained stable throughout the time series, starting at age two or three, and ranging up to 11-14 years, although two very old (18 and 24 years) stripers were captured in 2011 (Tables 32a,b). The age structure of male striped bass has expanded from three to six years in 1994, to up to 13 years in 2011 (Tables 33a,b). The age structure of female striped bass has been stable from 1994-2011, consisting of three to 15 year old females (Tables 34a,b). The cumulative proportion of males age six and older was 0.219, and has varied from 0.091-0.219 in 2000-2010 after peaking at 0.201-0.299 from 1996-1998. The cumulative proportion of females age eight and older, which had decreased from 0.531-0.266 from 1997-1999, rebounded to 0.426 in 2001, increased to 0.864 in 2008 and has since declined to 0.576 in 2011.

The cumulative catch rate of male striped bass mirrored the trends of the combined data with the 2011 catch rate being the seventh lowest, and less than half the cumulative catch rate for 2010 (Tables 33a,b). Using the maximum catch rate of the resident males as an indicator, the 1995-1997, 2000 and 2006 year classes were strongest and the 1992 and 1993 year classes the weakest. Male catch rates declined after ages five or six, but not as rapidly as on the Rappahannock River. However, the 2011 cumulative catch rate of female striped bass was 21.5% higher than the catch rate in 2010, and was the sixth highest in the 18-year time series (Tables 34a,b). There was no secondary peak in catch rates of females 1988-1994 year classes similar to that noted in the Rappahannock River pound net data.

Estimates of annual survival (S) for the individual year classes and their overall geometric means are presented in Tables 35-37. While annual survival estimates varied widely among years, due to strong or weak overall catches, the geometric mean survival rate (1994-2011) of the 1984 -2003 year classes (sexes combined) varied from 0.338-0.710 (Table 35), with an overall mean survival rate of 0.578. There were widely divergent estimates of annual survival of male and female striped bass. The geometric mean survival rate (1994-2011) of the 1988-2003 year classes of males varied from 0.286-0.612 (Table 36) with an overall mean survival rate of 0.463. These year classes have been the major target of the fall recreational and commercial fisheries that reopened in 1993. The geometric mean survival rate (1994-2011) of the 1984-2001 year classes (except 1999) of females varied from 0.339-0.915 (Table 37) with an overall mean survival rate of 0.666.

Catch Rate Histories of the 1987-2003 Year Classes

The catch rate histories of the 1987-2003 year classes from each sampling gear (sampling on the James River commenced in 1993) are depicted in Figures 4-19. Consistent among the year classes are a peak of male striped bass at age four or five followed by a rapid decline in the catch

rate and a secondary peak of mostly female striped bass around age 10. This secondary peak is best defined from the pound net data. The gill nets appear to be less efficient at catching larger, therefore older, striped bass. In both gears the catch rates of male striped bass was an order of magnitude greater than the catch rates of female striped bass.

Numeric catch rates for male striped bass decreased rapidly subsequent to their peak of abundance at age four or five in both gears. These fish are the primary target for the commercial and recreational fisheries within Chesapeake Bay. Catch rates of female striped bass also show a steep decline after their initial peak in abundance, presumably due to their migratory behavior, but, at least in the Rappahannock River, also exhibited a secondary peak in the catch rates of 9-11 year old females that persisted across several year classes. This secondary peak was due to the relative lack of intermediate sized (590-710 mm TL) striped bass in the samples. This pattern was not evident in the catches from 1991-1996 but has been persistent thereafter.

1987 Year class: The catch history of the 1987 year class commences at age four from the Rappahannock River and age seven from the James River. Peak abundance of male striped bass occurred at age four and the peak abundance of female striped bass occurred at age six in the Rappahannock River (Figure 4). Abundances of both sexes declined rapidly with age, although there was a distinctive secondary peak in the abundance of female striped bass captured from the pound nets. One female 1987 year class striped bass was captured in 2011 (the first since 1999).

1988 Year class: The catch history of the 1988 year class commences at age three from the Rappahannock River and age six from the James River. Age three was the apparent age of full recruitment to both sampling gears. Peak abundance of male striped bass occurred at age four (Figure 5). However, peak abundance of female striped bass was age nine from the gill nets and age 10 in the pound nets. Abundances decreased rapidly with age, although the pound net samples again had a secondary peak of female striped bass at age nine. The 1988 year class was above the mean CCA in both the pound net and gill net samples in the Rappahannock River (Tables 38 and 39). No 1988 year class striped bass were captured in 2011.

1989 Year class: The catch history of the 1989 year class, fully recruited to the gears in the Rappahannock River, commenced at age five in the James River samples. Peak abundance of male striped bass occurred at age four (pound nets) and five (gill nets in both rivers, Figure 6). Peak abundance of female striped bass occurred at age five in the Rappahannock River (both gears) and age six in the James River. There was a secondary peak in abundance of female striped bass at age nine in the pound net samples. The CCA from both gears in the Rappahannock River was below the mean (Tables 38 and 39). No 1989 year class striped bass were captured in 2011.

1990 Year class: The catch history of the 1990 year class commenced at age four in the James River. Peak abundance of male striped bass occurred at age four (gill nets) and five (pound nets) in the Rappahannock River and age four in the James River (Figure 7). The peak abundance of female striped bass occurred at age five in the gill net samples from both rivers, but was age eight in the pound net samples. The CCA was the second lowest of the time series from both gears in the Rappahannock River (Tables 38 and 39). The CCA for the James River, though

lacking values for ages two and three, was also below the mean (Table 40). No 1990 year class striped bass were captured in 2011.

1991 Year class: The catch history of the 1991 year class commenced at age three in the James River and was fully recruited to the sampling gear. Peak abundance of male striped bass occurred at age four in the James River and at age five in the Rappahannock River (both gears, Figure 8). Peak abundance of female striped bass occurred at age eight in the James River and at age 10 in the Rappahannock River. It is interesting to note that age five and six female striped bass were not caught in the same relative abundance as in the 1987-1990 year classes. The CCA was the lowest of the year classes compared to the Rappahannock River in both sampling gears (Tables 38, 39) and well below the mean in the James River (Table 40). No 1991 year class striped bass were captured in 2011.

1992 Year class: Peak abundance of male striped bass occurred at age three in the pound nets in the Rappahannock River and in the gill nets in the James River, but occurred at age five in the gill nets in the Rappahannock River (Figure 9). Peak abundance of female striped bass occurred at age seven in the James River but occurred at age nine (gill nets) and age eleven (pound nets) in the Rappahannock River. Again, there were relatively few ages five and six female striped bass captured in the Rappahannock River. Thus, what had been a secondary peak of abundance for the 1987-1989 years classes has been the primary peak in the 1990-1992 year classes. The CCA was higher than for the 1990 and 1991 year classes, but was still below the mean in the Rappahannock River (Tables 38, 39), and was the lowest value for the James River (Table 40). One female 1992 year class striped bass was captured in the Rappahannock River in 2011.

1993 Year class: Peak abundance of male striped bass occurred at age four in the Rappahannock (both gears) and the James rivers (Figure 10). Peak abundance of female striped bass occurred at age six on the James River, but not until ages nine (gill nets) and age ten (pound nets) in the Rappahannock River. Again, there were relatively few ages five and six female striped bass captured in the Rappahannock River. The CCA was the highest of all the year classes from the gill net samples, but was only near the mean from the pound net samples in the Rappahannock River (Tables 38, 39). The CCA for the James River was well below the mean (Table 40). Five female 1993 year class striped bass, three in the Rappahannock River and one in the James River were captured in 2011.

1994 Year class: Peak abundance of male striped bass occurred at age four in the Rappahannock River (both gears) and at age six in the James River (Figure 11). Peak abundance of female striped bass occurred at age five on the James River, but not until age ten in the Rappahannock River (both gears). Again, there were relatively few ages five and six female striped bass captured in the Rappahannock River. The CCA was slightly below the mean from the pound net samples but well above the mean from the gill net samples in the Rappahannock River (Tables 38, 39). The CCA for the James River was higher than for the 1991-1993 year classes but was still below the mean (Table 40). No 1994 year class striped bass were captured in 2011.

1995 Year class: Peak abundance of male striped bass occurred at age three (gill nets) and four (pound nets) in the Rappahannock River and occurred at age five in the James River (Figure 12). Peak abundance of female striped bass occurred at age four in the James River but not until age nine in the Rappahannock River (both gears). Again, there were relatively few ages five and six female striped bass captured in the Rappahannock River. The CCA was above the mean in the Rappahannock River pound nets (Table 38), but below the mean in the gill nets (Table 39). The CCA was below the mean in the James River (Table 40). The 1993-1995 year classes were characterized as having a primary peak of young, male striped bass and a secondary peak of older, female striped bass. Two female 1995 year class striped bass, one from each river, were captured in 2011.

1996 Year class: Peak abundance of male striped bass occurred at age three (gill nets) and four (pound nets) in the Rappahannock River and occurred at age four in the James River (Figure 13). Peak abundance of female striped bass occurred at age ten in the James River and at age 11 in the Rappahannock River (both gears). Again, there were relatively few ages five and six female striped bass captured in the Rappahannock River. The CCA was the highest amongst the year classes from the pound samples in the Rappahannock River (Table 38) and well above the mean in the gill net samples (Table 39). The CCA for the James River was the highest of any of the year classes (Table 40). Thirteen female 1996 year class striped bass (11 in the Rappahannock and two in the James) were captured in 2011.

1997 Year class: Peak abundance of male striped bass occurred at age three (pound nets) and age four (gill nets) in the Rappahannock River and occurred at age four in the James River (Figure 14). Age ten females showed an increase in abundance in the Rappahannock River (both gears) and the James River gill nets. The CCA was the second highest in the Rappahannock River pound nets (Table 38) and James River gill nets (Table 40), and the third highest in the Rappahannock River gill nets (Table 39). Sixteen female 1997 year class striped bass (12 in the Rappahannock and four in the James) were captured in 2011.

1998 Year class: Peak abundance of male striped bass occurred at age five (gill nets) and age six (pound nets) in the Rappahannock River and occurred at age four in the James River (Figure 15). Age nine females showed an increase in abundance versus their abundance in 2006 (at age eight) in both rivers. The CCA was the lowest since the 1992 year class in the Rappahannock River pound nets (Table 38) but only slightly below average in the gill nets (Table 39). The CCA was above average in the James River (Table 40). Fourteen (12 female and two male) 1998 year class striped bass (13 in the Rappahannock River and one in the James River) were captured in 2011.

1999 Year class: Peak abundance of male striped bass occurred at age four in the Rappahannock River gill nets and at age five in the pound nets and James River gill nets (Figure 16). The CCA at age eight was less than for the 1998 year class and was the lowest since the 1992 year class in the pound nets (Table 38) and the 1991 year class in the Rappahannock River gill nets (Table 39). The CAA for the James River was the lowest since the 1995 year class (Table 40). Nineteen (16 female and three male) 1999 year class striped bass (13 in the Rappahannock River and six in the James River) were captured in 2011.

2000 Year class: Peak abundance of male striped bass occurred at age four for all indexes for both the Rappahannock and the James rivers (Figure 17). The peak abundance of female striped bass was age five in the pound nets and age six from the gill nets in both rivers. For the third successive year class, the CCA at age eight was the lowest since the 1992 year class in the pound nets (Table 38). The CCA for the gill nets was higher than for the 1998 year class but still well below the mean (Table 39). The CCA for the James River was higher than both the 1998 and 1999 year classes and was above the overall mean (Table 40). Twenty-eight (22 female and six male) 2000 year class striped bass (13 in the Rappahannock River and 15 in the James River) were captured in 2011.

2001 Year class: Peak abundance of male striped bass occurred at age three in the Rappahannock gill nets, age four in the pound nets and age five in the James River gill nets (Figure 18). Peak abundance of female striped bass occurred at age five in the pound nets and James River gill nets, but at six in the Rappahannock River gill nets. The CCA at age eight was the highest since the 1997 year class for all three indexes (Tables 38-40). Fifty (36 female and 14 male) 2001 year class striped bass (25 in the Rappahannock River and 25 in the James River) were captured in 2011.

2002 Year class: Peak abundance of male striped bass occurred at age four and at age five for the female striped bass in the pound nets and the gill nets in both rivers (Figure 19). The CCA at age eight was near the overall average in the pound nets and James River gill nets, but was the third lowest in the Rappahannock gill nets (Tables 38-40). Fifty-two (28 male and 24 female) 2002 year class striped bass (29 in the Rappahannock River and 21 in the James River) were captured in 2011.

2003 Year class: Peak abundance of male striped bass occurred at age four in the gill nets in both rivers and at age five in the pound nets (Fig 20). Peak abundance of female striped bass occurred at age five in the James River, but at age nine from both gears in the Rappahannock River. The CCA at age eight well above the mean and the highest since 1997 from in both the pound nets and gill nets in the Rappahannock River, but was below the mean and the lowest since 1994 in the James River (Tables 38-40). There were sixty 2003 year class (39 male and 21 female) striped bass (39 in the Rappahannock River and 21 in the James River) were captured in 2011.

Growth Rate of Striped Bass Derived from Annuli Measurements

The scales of 222 striped bass were digitally measured and the increments between annuli were used to determine their growth history. The back-calculated length-at-age of striped bass was 137mm at age one (Table 41a). The rate of growth was about 100 mm in their second year and decreased gradually with age to about 80 mm in their fifth year and to about 50 mm in their 10th year (Tables 41a,b). Interestingly, the growth rates of the most recent year classes were the highest, although the growth rate of the oldest year classes were based on very few specimens. Based on these growth estimates, an 18 inch (457 mm) total length striped bass would be 3.5 years of age during the fall recreational fishery in Chesapeake Bay. These striped bass reach the 28 inch (711 mm) total length minimum for the coastal fishery at age eight.

Age Determinations using Scales and Otoliths

Tests of symmetry: A total of 223 striped bass from 12 size ranges were aged by reading both their scales and otoliths. Scale and otolith ages from the same specimen were in agreement 42.2% (94/ 223) of the time and within one year 86.5% (193/223) of the time. Differences between the two age determination methods were first analyzed utilizing tests of symmetry. A chi-square test was performed to test the hypothesis that an $m \times m$ contingency table (Table 42) consisting of two classifications of a sample into categories is symmetric about the main diagonal. The test statistic is

$$X^2 = \sum_{i=1}^{m-1} \sum_{j=i+1}^m \frac{(n_{ij} - n_{ji})^2}{n_{ij} + n_{ji}}$$

where n_{ij} = the observed frequency in the i th row and j th column and n_{ji} = the observed frequency in the j th row and i th column (Hoenig et al., 1995).

A test of symmetry that is significant indicates that there is a systematic difference between the aging methods. The number of degrees of freedom is equal to the number of non-zero age pair comparisons (here = 122). We tested the hypothesis that the observed age differences were symmetrically distributed about the main table diagonal (Table 42). The hypothesis was rejected ($X^2 = 51.74$, $p < .005$), indicating non-random differences between the two ageing methodologies. The two ageing methods were also found to be non-random in 2004, 2005 and 2007-2009, but not in 2006.

Differences between the scale and otolith age from the same specimen ranged from zero to four years (Figure 21). The otolith-derived age exceeded the scale age 27.4% of the total examined (47.3% of the non-zero differences). When the differences in ages were greater than one year, the otolith age was even more likely to be the older age (76.7%). Another test of symmetry that compared the negative and positive differences of the same magnitude (i.e. -4 and 4, -3 and 3, etc., Evans and Hoenig, 1998) accepted the hypothesis that these differences were random ($X^2 = 16.67$, $df = 5$, $p = 0.054$). This test has far fewer degrees of freedom than did the previous test of symmetry.

T-tests: Next, t-tests of the resultant means of the two ageing methods were performed. A two-tailed t-test was made to test the null hypothesis that the mean ages determined by the two methods were not different from zero. The mean age of the sample ($n=223$) determined by reading the otoliths was greater than the mean age determined by reading the scales (by 0.12 years, Table 43). The test results were:

$$\bar{Age}_{scottish} = 8.62$$

$$\bar{Age}_{scale} = 8.50$$

$$S_{scottish} = 4.07$$

$$S_{scale} = 3.61$$

$$t = 0.330$$

$$df = 444$$

$$p = .742$$

Therefore the null hypothesis was not rejected.

A paired t-test was also performed on the ages determined for each specimen by the two methodologies. The null hypothesis tested was that the mean of the difference resultant from the two methods was not different from zero. The paired t-test results were not significant ($t = 1.49$, $df = 222$, $p = .138$) and the null hypothesis was not rejected.

Kolmogorov-Smirnov test: To determine whether the distribution of age classes that resulted from the two ageing methodologies were representative of the same population, a Kolmogorov-Smirnov test was performed on the relative proportion that each assigned age class contributed to the total sample (Table 43). This compares the maximum difference in the relative proportions that an age class contributes to the test statistic ($K_{.05}$):

$$D_{max} = 0.897$$

$$K_{.05} = 1.3581$$

$$D_{.05} = \sqrt{\frac{(228) + (223)}{228^2}} = 0.1286$$

The maximum difference did not exceed the test statistic, so the null hypothesis, that the age structures derived by the two ageing methods represent the same population, was accepted. This result is consistent with the 2008-2010 results, but differs from the test results for the 2007 age comparisons.

Discussion

Striped bass stocks had recovered sufficiently by 1993 to allow the re-establishment of limited commercial and recreational fisheries in Virginia. The monitoring efforts summarized in this report were intended to document changes in the abundance and age composition of spawning stocks in the James and Rappahannock rivers during the period of managed harvest by these fisheries.

The main advantage of pound nets is that the gear provides large catches (often in excess of 100 fish per day) that are presumably not sex or size-biased. However, each pound net has a different fishing characteristic (due to differences in depth, bottom, fetch, nearness to shoals or channels, etc.), and our sampling methods (in use since 1993) may have introduced additional variability. The down-river net (mile 44) was set in a shallow, flat-bottomed portion of the river with a leader that extended farther into the bay. The upriver net (mile 47) was set in a constricted portion of the river that abutted the channel, and had a leader that extended almost to the shoreline. Ideally, each net was scheduled to be sampled weekly, but uncontrollable factors (especially tide, weather, and market conditions) affected this schedule. Since spring 2002 the down-river net has not been set and was replaced by a net across the river at mile 45. This net had been utilized since 1997 as a source for tagging striped bass, but had been excluded from the spawning stock assessment in order to keep the sampling methodology as consistent as possible with the 1991-1996 data. Weekly sampling occurred each Monday and Thursday, a schedule that translated to fishing efforts of 96 hrs (Thursday through Monday) or 72 hrs (Monday through Thursday). In 2011, persistent bad weather delayed efforts by our fishermen to establish their first net (usually done in mid-March) until 11 April (one net) and precluded setting the third net at mile 45. Hence we tagged and released all striped bass greater than 457 mm and used a sex and size-based regression to estimate biomass for our pound net index.

In past years, duration of the pound net set was as low as 24 hrs, and as large as 196 hrs, if the fisherman was unable to fish the scheduled net on the scheduled sampling date. Although these events were uncommon, we were unable to assess whether varying effort influenced estimates of catch rate. The 1997 and 1998 data include a pound net at mile 46 that had an orientation and catch characteristics similar to the net at mile 47. This net was also sampled on one date (7 April) in 2003. In 2005 this net was substituted entirely for the net at mile 47 due to extensive damage to the net at mile 47 in a maritime accident. The 1991 data included samples taken from a pound net at river mile 25 and were weekly vs. twice-weekly samples, but with similar total effort. While this net is far enough within the Rappahannock to preclude significant contamination from stocks from other rivers, it does not meet the criteria established in 1993, restricting sampling to gears located within the designated spawning grounds (above river mile 37). The catches from these other nets were similar in sex and age composition to the nets presently used and their exclusion would adversely affect our ability to assess the status of the spawning stocks in those years.

Variable-mesh gill nets were set by commercial fishermen and fished by scientists after 24 hours on designated sampling days. As a result, there were fewer instances of sampling inconsistencies, although in 2004, a manufacturing error resulted in two nets of the number one configuration being fished on both rivers. The two nets were set approximately 300 meters apart and along the same depth contours on both rivers. Although the down-river net did not always contain the greater catches, removal by one net may have affected the catch rates of its companion.

The gill nets captured proportionally more males than did the pound nets. Anecdotal information from commercial fishermen suggests that spawning males are attracted to conspecifics that have become gilled in the net meshes. Thrashing of gilled fish may emulate

spawning behavior (termed Arock fights@ by local fishermen) and enhance catches of males. The pound net catches contained a greater relative proportion of older female striped bass than did the catches from the gill nets. This trend has been persistent over several years. Thus, given the presence of large females in the spawning run, it is clear that the gill nets do not adequately sample large (900+ mm FL) striped bass.

The biological characterization of the spawning stock of striped bass in the Rappahannock River changed dramatically from 1991-2011. There was a steady decrease in the relative abundance of five to seven year-old striped bass from 1991-2001, but these ages were proportionally more abundant in 2002-2011. The males in these age classes had been the target of the recreational and commercial fisheries, but with the increase in the availability of larger striped bass in recent years, the younger striped bass may be under less fishing pressure. Current regulations protect females from harvest during their annual migration by higher minimum lengths in the coastal fishery (711 mm TL vs. 458 mm TL within Chesapeake Bay) and the closure of the fishery in the bay during the April spawning run. The result has been a general increase in the abundance of older females throughout the period. Total catches in 2011 were much lower than in 2010, especially for resident male striped bass.

Of note again in the 2010 samples was the relative abundance of 1996 year class (15 year old) male and female stripers. This year class has been above-average in abundance since recruiting to the gears at age three, which indicates that it is a very strong year class. Also, the 1993 year class, abundant in 2005-2007, but absent from the samples again in 2008 and 2009, were captured again in 2010 and 2011.

The 2011 value of the Spawning Stock Biomass Index (SSBI) for the Rappahannock River pound nets was approximately 67% lower than the SSBI for 2010 approximately 25% below the mean. The SSBI for male striped bass captured in the pound nets was the lowest since 2002 and more than 33% below the mean of the 1991-2011 time series. The SSBI for female striped bass was less than half the 2010 value and 20% below the mean of the time series. While the biomass indexes are dominated by the older age classes, there was a notable increase in four to seven year old females compared to recent years.

The 2011 SSBI for the gill nets was 42% lower than the value for 2010 and approximately 25% below the mean of the 1991-2011 time series. The male SSBI was approximately one third the value for 2010 and was the second lowest overall. However, the female SSBI was 7% higher than the value for 2010 and well over the mean for the 1991-2011 time series.

The 1991-2011 values of the SSBI in the Rappahannock River were often inconsistent between pound nets and gill nets. In the pound nets, male biomass peaked in 1993 due to strong 1988 and 1989 year classes, and again in 1999 and 2000 due to strong 1996 and 1997 year classes. The value in 2011 was driven by increased catches of 2000-2003 year classes, compared to the 2003-2006 year classes that dominated the index in 2010. The female biomass from pound nets showed no reliance upon any age groups, although the exceptionally strong 1996-1998 year classes continue to contribute highly. The male biomass from the gill nets is driven by the

number of Asuper catches@, when the net is literally filled by males, seeking to spawn, that occur differentially among the years (most notably in 1994, 1997 and 2004). Due to the highly selective nature of the gill nets (significantly fewer large females), the female SSBI from these nets is less reliable. The low biomass values from both gears of both sexes in 1992 and 1996 are probably an underestimate of spawning stock strength since water temperatures were below normal in those years. Local fishermen believe that low temperatures alter the catchability of striped bass. It is also possible that the spawning migration continued past the end of sampling in those years.

The 2011 value of the SSBI in the James River was about 33% lower than in 2010 and over 20% below the mean of the 1994-2011 time series. The male index was driven by large catches of the 2005-2007 year classes while the female index had higher catch rates of the 1996-2001 year classes. Because of the changes in location and in the methodology utilized by the new fisherman starting in 2000, the values are not directly comparable with those of previous years. The below normal river flow conditions noted for the Rappahannock River, apply to the James River as well. The relative scarcity of larger, predominantly female, striped bass from the gill nets in the James River (compared to pound net catches) implies a similar limitation in fishing power as shown in the Rappahannock River but comparative data are not available since there are no commercial pound nets on the James River.

The Egg Production Potential Index (EPPI) is an attempt to better define the reproductive potential of the spawning stocks, especially as they become more heavily dependent on fewer, but larger, female striped bass. For example, in the 2001 Rappahannock River pound net data the contribution of 8+ year old females was 75.2% of the total number of mature females (the basis of our index prior to 1998), 94.1% of the mature female biomass (the basis of the current index), and 94.3% of the calculated egg potential. The catches in 2002 were less reliant on older fish than in the preceding years so that the contribution of 8+ year old females was 46% of the total number of mature females, but still 69.1% of the female biomass and 68.4% of the potential egg production. In 2011, the contribution of 8+ year old females was 75.9% of the total number, 90.9% of the biomass, and 92.1% of the calculated egg potential. It should be noted that our fecundity estimates for individual striped bass are well below those reported by Setzler et al. (1980). Our methodology differs from the previous studies, but the relative contribution in potential egg production of the older females may be underestimated at present.

In our analysis of pound net catch rates, we observed a distinctive bimodal distribution of the striped bass. These striped bass appeared in greatest abundance at age five or six (especially males), at lower abundance at age six to eight (both sexes), and then higher abundance at ages nine to 12 (especially females). Also, prior to 1995, the peak catch rates of male and female striped bass (ages four and five) were similar. The catches of these age classes are now almost exclusively male. Thus, the 1991-1996 year classes actually showed greater abundance at ages nine to 12 years than at any other age. Age estimation of larger striped bass by scales is problematic because re-absorption or erosion of outer margins of scales may cause under-estimation of age. Under-ageing errors might tend to lump catches of old fish (>12 years) into younger categories (nine to 12 years). However, ignoring age, we also observed a bimodal size distribution, one group from 470-590 mm fork length, presumably young, and the second group

of 850-1200 mm fork length, presumably older. This trend became increasingly apparent in the 1997-2003 data and its significance has not been determined. In 2004-2011, the second group was expanded to 750-1200 mm as the strong 1996-1998 year classes were caught in abundance.

The time series of the catch rates by age class and by year class indicate that the age of peak abundance in the rivers has changed from five or six years in 1992-1994 to three to four years in 2000-2002. Changes in the annual catch rates by year class in the Rappahannock River indicated that strong year classes occurred in 1988, 1989, 1996 and 1997, and weak year classes occurred in 1990 and 1991. The relative abundance of ten-year old, 1992 year class, striped bass of both sexes in both 2001 and 2002, indicate that the 1992 year class was also strong. Likewise, the data for the James River indicated that strong year classes occurred in 1989, 1993, 1994 and 1996, and weak year classes occurred in 1990 and 1991.

The time series allows estimates of the instantaneous rates of survival of the year classes using catch curves, especially for the 1983-2003 year classes that were captured for four or five years subsequent to their peak in abundance at age four or five. The survival estimates of female striped bass of the 1983-1998 year classes in the Rappahannock River were 0.58 in pound nets and 0.65 in gill nets. The survival estimates of 1985-2003 year class male striped bass were 0.50 in pound nets and 0.50 in gill nets. The higher survival estimates for the females may be the result of their differential maturation rates. These differences cause lower peaks in abundance (usually at age five) as only fractions of each year class mature and are depicted in their lower peak abundance values. The large differences between the sexes also reflect a management strategy that targets males. Similarly, survival estimates for these year classes in the James River were approximately 0.46 for male striped bass and approximately 0.67 for female striped bass.

The catch histories of the 1987-2003 year classes in the Rappahannock River show two distinct patterns. The 1987-1990 year classes had initial peaks of abundance of both sexes at ages four or five and a secondary peak in the abundance of female striped bass after age eight. Subsequent year classes did not have the initial peak in abundance of female striped bass, but only what was the secondary peak of eight to 12 year-olds. Since catches of larger, thus older, striped bass was less consistent in the gill net catches, this pattern was less apparent in that data set. Using the area under the catch curve as an indicator of year class strength, the 1993 and 1996 year classes were the strongest and the 1990 and 1991 year classes were the weakest.

Back-calculation of the growth based on measurements between scale annuli indicated that striped bass grow about 137 mm (fork length) in their first year. Growth averaged 100 mm in their second and third years and decreased gradually to about 50 mm by age 10. Thus, striped bass reach the 18 in. (457 mm) minimum total length for the Chesapeake Bay resident fishery at 3.5 years of age (the 2007 year class in fall 2010) and the 28 in. (711 mm) minimum total length for the coastal fishery at age eight.

The ages of striped bass determined by reading both their scales and otoliths were found to differ by as much as five years. Overall, the age difference determined for the largest, and oldest, specimens was 0-5 years (14-20 years by reading the scale vs. 14-25 years by reading the otolith). The maximum age determined by reading scales has generally remained constant at 16

years since 1991 (although two 17 and one 20 year-old was aged in 2011); while there has been an annual progression in the maximum age determined by reading otoliths. Agreement between the two ageing methodologies was 42.2% and was higher than the results from 2010 (37.3%). When there was disagreement between methodologies, the scale age was 1.1 times more likely to have been aged older than the respective otolith-derived age. This result is the opposite from the results from every other year that age comparisons have been made (2003-2011). However, when the age difference was two years or greater, the otolith age was 3.2 times to be the older age. The differences were found not to be statistically non-random and different from zero. This was in contrast with the results with recent years (2004, 2005, 2007, 2008-2010). However, the relative contributions of the age classes and their overall mean age were not statistically different between the two methodologies. Previous ageing method comparison studies (Secor, et al. 1995, Welch, et al. 1993) concluded that otolith-based and scale-based ages of striped bass became increasingly divergent, with otolith ages being older, especially after 900 mm in size or 10-12 years in age. We plan to continue these comparisons in future years.

Literature Cited

- Berggren, T.J. and J.T. Lieberman. 1978. Relative contribution of Hudson, Chesapeake and Roanoke striped bass, *Morone saxatilis*, stocks to the Atlantic coast fishery. U. S. Fish. Bull. 76(2): 335-345.
- Barbieri, S.K. and L.R. Barbieri. 1993. A new method of oocyte separation and preservation for fish reproduction studies. U. S. Fish. Bull. 91: 165-170.
- Chapoton, R.B. and J.E. Sykes. 1961. Atlantic coast migration of large striped bass as evidenced by fisheries and tagging. Trans. Amer. Fish. Soc. 90(1):13-20.
- Dorazio, R.M., K.A. Hattala, C.B. McCollough and J.E. Skjveland. 1994. Tag recovery estimates of migration of striped bass from spawning areas of the Chesapeake Bay. Trans. Amer. Fish. Soc. 123: 950-963.
- Evans, G.T. and J.M. Hoenig. 1998. Testing and viewing symmetry in contingency tables, with application to readers of fish ages. Biometrics 54: 620-629.
- Field, J.D. 1997. Atlantic striped bass management: where did we go right? Fisheries 22(7): 6-8.
- Frie, R.V. 1982. Measurement of fish scales and back-calculation of body lengths using a digitizing pad and microcomputer. Fisheries 7(5): 5-8.
- Grant, G.C. and J.E. Olney. 1991. Distribution of striped bass *Morone saxatilis* (Walbaum) eggs and larvae in major Virginia rivers. U. S. Fish. Bull. 89:187-193.
- Hardy, J.D. Jr. 1978. Development of fishes of the mid-Atlantic bight. Vol. III, Aphrederidae through Rachycentridae. U. S. Fish Wildl. Serv. FWS/OBS-78/12.
- Hoenig, J.M., M.J. Morgan and C.A. Brown. 1995. Analysing differences between two age determination methods by tests of symmetry. Can. J. Fish. Aquat. Sci. 52: 354-368.
- Lewis, R.M. 1957. Comparative study of populations of the striped bass. U. S. Fish and Wildlife Service Spec. Rep. Fisheries 204:1-54.
- Loesch, J.G., W.H. Kriete, Jr., and S.M. Atran. 1985. Sonic digitizers "go fishing": fish scales reveal age by sound. Sea Tech., February 1985: 3-31.
- Mansueti, R.J. 1961. Age, growth, and movements of the striped bass, *Roccus saxatilis*, taken in size selective fishing gear in Maryland. Ches. Sci. 2: 9-36.

- McGovern, J.C. and J.E. Olney. 1996. Factors affecting survival of early life stages and subsequent recruitment of striped bass on the Pamunkey River, Virginia. *Can. J. Fish. Aquat. Sci.* 53: 1713-1726.
- Merriman, D. 1937. Notes on the life history of the striped bass (*Roccus lineatus*). *Copeia* 1:15-36.
- Merriman, D. 1941. Studies on the striped bass (*Roccus saxatilis*) of the Atlantic Coast. *Fish. Bull. U.S. Fish Wildl. Serv.* 50(35):1-77.
- Olney, J.E., J.D. Field, and J.C. McGovern. 1991. Striped bass egg mortality, production and female biomass in Virginia rivers, 1980-1989. *Trans. Amer. Fish. Soc.* 120: 354-367.
- Pearson, J.C. 1938. The life history of the striped bass, or rockfish, *Roccus saxatilis* (Walbaum). *U. S. Fish. Bull.* 49: 825-851.
- Raney, E.C. 1957. Subpopulations of the striped bass *Roccus saxatilis* (Walbaum), in tributaries in Chesapeake Bay. *U. S. Fish Wildl. Serv., Spec. Sci. Fish.* 208: 85-107.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. *Fish. Res. Bd. Can. Bull.* 191: 382 p.
- Rugolo, L.J., P.W. Jones, R.K. Schaefer, K.S. Knotts, H.T. Hornick and J.L. Markham. 1994. Estimation of Chesapeake Bay-wide exploitation rate and population abundance for the 1993 striped bass stock. Manuscript, Maryland Department of Natural Resources, Annapolis, Md.
- Sadler, P.W., R.E. Harris, J. Romine, and J.E. Olney. 1998. Evaluation of striped bass stocks in Virginia: monitoring studies, 1993-1998. Completion Report, Virginia Institute of Marine Science. 99 p.
- Sadler, P.W., R.J. Latour, R.E. Harris, and J.E. Olney. 2001. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 1999-2003. Annual Report, Virginia Institute of Marine Science: 93 p.
- Sadler, P.W., R.J. Latour, R.E. Harris, K.L. Maki and J.E. Olney. 2002. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 1999-2003. Annual Report, Virginia Institute of Marine Science: 102 p.
- Sadler, P.W., R.J. Latour, R.E. Harris, J.K. J.K. Ellis and J.E. Olney. 2003. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 1999-2003. Annual Report, Virginia Institute of Marine Science: 131 p.

- Sadler, P.W., J.M. Hoenig, R.E. Harris and B.G. Holloman. 2004. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 1999-2004. Annual Report, Virginia Institute of Marine Science: 167 p.
- Sadler, P.W., J.M. Hoenig, R.E. Harris and B.G. Holloman. 2005. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 2005-2009. Annual Report, Virginia Institute of Marine Science: 199 p.
- Sadler, P.W., J.M. Hoenig, R.E. Harris, and B.G. Holloman. 2006. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 2005-2009. Annual Report, Virginia Institute of Marine Science: 150 p.
- Sadler, P.W., J.M. Hoenig, R.E. Harris, R.J. Wilk and L.M. Goins. 2007. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 2005-2009. Annual report, Virginia Institute of Marine Science: 170pp.
- Sadler, P.W., J.M. Hoenig, R.E. Harris, M. W. Smith, R.J. Wilk and L.M. Goins. 2008. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 2005-2009. Annual report, Virginia Institute of Marine Science: 192pp
- Sadler, P.W., Smith, M.W., Hoenig, J.M., Harris, R.E., Goins, L.M. and R.J. Wilk. 2009. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 2005-2009. Annual report, Virginia Institute of Marine Science: 215pp.
- Sadler, P.W., Smith, M.W., Sullivan, S.E., Hoenig, J.M., Harris, R.E., L.M. Goins, .2010. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 2010-2014. Annual report, Virginia Institute of Marine Science: 220pp.
- Secor, D.H. 1999. Specifying divergent migrations in the concept of stock: the contingent hypothesis. Fisheries research 43: 13-34.
- Secor, D.H., T.M. Trice and H.T. Hornick. 1995. Validation of otolith-based ageing and a comparison of oolith and scale-based ageing in mark-recaptured Chesapeake Bay striped bass, *Morone saxatilis*. Fish. Bull. 93:186-190.
- Setzler, E.M., W.R. Boyton, K.V. Wood, H.H. Zion, L. Lubbers, N.K. Montford, P. Frere, L. Tucker and J.A. Mihursky. 1980. Synopsis of biological data on striped bass, *Morone saxatilis* (Walbaum). NOAA Tech. Rept. NMFS 433.
- Shepherd, G. and H. Lazar (eds). 1998. Source document to Amendment 5 to the Interstate Fishery Management Plan for striped bass. ASMFC Rep. No. 34.
- Sokal, R.R. and F.J. Rohlf. 1981. Biometry. W. H. Freeman Co. 859 p.

- Van Winkle, W., K.D. Kumar, and D.S. Vaughan. 1988. Relative contributions of Hudson River and Chesapeake Bay striped bass stocks to the Atlantic Coast population. *Amer. Fish. Soc. Mono.* 4: 255-266.
- Welch, T.J., M.J. Van Den Avyle, R.K. Betsill and E.M. Driebe. 1993. Precision and relative accuracy of striped bass age estimates from otoliths, scales, anal fin rays and spines. *N. Amer. J. Fish. Mgmt.* 13:616-620.
- Wischniowski, W. and S. Bobko. 1998. Age and growth laboratory manual. Final report Old Dominion Univ. Center for Quantitive Fisheries Ecology.
- Zar, J.H. 1999. *Biostatistical Analysis*, Fourth Edition. Prentis Hall Press. 663 pp.

Table 1. Numbers of striped bass in three age categories (year classes 2007-2009, 2003-2006 and 1992-2002) from pound nets in the Rappahannock River, by sampling date, spring, 2011. M = males, F = females.

Date	n	Year Class							
		No age		2007-2009		2003-2006		1992-2002	
		M	F	M	F	M	F	M	F
11 April	68	0	0	15	0	22	9	11	11
14 April	62	0	0	10	0	21	3	6	22
18 April	23	0	0	2	0	8	4	1	8
21 April	9	0	0	0	0	0	2	2	5
25 April	11	0	0	1	1	2	3	1	3
29 April	38	1	0	7	0	17	4	1	8
2 May	4	0	0	3	0	1	0	0	0
Total	215	1	0	38	1	71	25	22	57

Table 2. Net-specific summary of catch rates and mean ages of striped bass (n=215) in pound nets on the Rappahannock River, spring, 2011. Values in bold are the grand means for each column. M = male, F=female.

Date	Net ID	n	CPUE (fish/day)		CPUE (g/day)		Mean age	
			M	F	M	F	M	F
11 April	S473	68	16.0	6.7	53,134.1	51,360.5	6.3	9.4
14 April	S473	62	12.3	8.3	43,219.1	77,110.0	6.3	10.4
18 April	S462	23	2.8	3.0	9,463.3	24,737.0	6.4	9.8
21 April	S473	9	0.7	2.3	4,490.7	14,642.0	10.0	8.3
25 April	S462	11	1.0	1.8	2,985.5	11,887.2	6.3	8.6
29 April	S473	38	6.5	3.0	12,819.6	21,424.4	5.1	9.3
2 May	S462	4	1.3	0.0	1,719.4	0.0	4.3	
Totals	S453	0						
	S462	38	1.7	1.7	4,995.7	13,317.9	5.9	9.3
	S473	177	8.7	4.9	27,216.2	39,618.1	6.1	9.6
Season		215	5.5	3.4	17,031.8	27,563.8	6.0	9.5

Table 3. Length frequencies (TL in mm) of striped bass sampled from the pound nets in the Rappahannock River, spring, 2011.

TL	n	TL	n	TL	n	TL	n	TL	n	TL	n
250-	0	410-	3	570-	5	730-	4	890-	5	1050-	0
260-	0	420-	3	580-	2	740-	1	900-	4	1060-	1
270-	0	430-	1	590-	8	750-	2	910-	7	1070-	0
280-	0	440-	0	600-	4	760-	3	920-	3	1080-	0
290-	1	450-	1	610-	3	770-	3	930-	4	1090-	0
300-	1	460-	3	620-	3	780-	4	940-	2	1100-	0
310-	0	470-	2	630-	3	790-	4	950-	0	1110-	1
320-	1	480-	6	640-	4	800-	3	960-	3	1120-	1
330-	0	490-	4	650-	1	810-	3	970-	2	1130-	1
340-	0	500-	3	660-	1	820-	6	980-	1	1140-	0
350-	0	510-	4	670-	5	830-	7	990-	3	1150-	1
360-	0	520-	6	680-	2	840-	5	1000-	2	1160-	0
370-	0	530-	5	690-	1	850-	4	1010-	1	1170-	0
380-	0	540-	3	700-	3	860-	6	1020-	2	1180-	0
390-	1	550-	4	710-	2	870-	1	1030-	0	1190-	0
400-	2	560-	6	720-	4	880-	3	1040-	1	1200-	0

Table 4. Mean fork length (mm), weight (g), standard deviation (SD) and CPUE (fish per day; weight per day) of striped bass from pound nets in the Rappahannock River, 11 April – 2 May, 2011.

Year	Sex	n	Fork Length		Weight		CPUE	
			Mean	SD	Mean	SD	F/day	W/day
2009	male	2	280.0	5.7	270.8	63.6	0.1	22.6
2008	male	11	382.5	32.1	768.5	194.2	0.5	352.2
2007	male	25	457.5	24.4	1,299.4	209.6	1.0	1,353.6
	female	1	485.0		1,853.8		0.0	77.2
2006	male	24	520.4	26.2	1,898.2	282.8	1.0	1,898.2
	female	9	540.2	29.5	2,472.9	406.0	0.4	927.3
2005	male	16	588.1	34.7	2,716.4	468.7	0.7	1,810.9
	female	7	583.3	25.7	3,011.7	309.4	0.3	878.4
2004	male	16	671.3	46.6	4,030.6	750.3	0.7	2,687.1
	female	3	680.3	78.4	4,862.4	1,473.3	0.1	607.8
2003	male	15	718.9	39.8	4,857.1	782.6	0.6	3,035.7
	female	6	772.0	10.9	6,704.3	225.1	0.3	1,676.1
2002	male	12	765.7	48.9	5,876.4	957.9	0.5	2,938.2
	female	11	796.4	20.7	7,196.1	558.3	0.5	3,298.2
2001	male	6	778.2	69.3	6,278.2	1,266.6	0.3	1,569.5
	female	19	854.3	23.1	8,738.5	746.4	0.8	6,918.0
2000	male	4	835.8	30.6	7,462.1	776.4	0.2	1,243.7
	female	11	885.1	29.0	9,732.3	836.8	0.5	4,460.6
1999	female	6	941.2	11.5	11,514.1	302.5	0.3	2,878.5
1998	female	1	900.0		10,410.7		0.0	433.8
1997	female	4	966.0	18.9	12,471.5	529.1	0.2	2,078.6
1996	female	2	1,024.0	19.9	14,880.7	1,232.1	0.1	1,240.6
1995	female	1	1,058.0		16,075.6		0.0	669.8
1993	female	2	1,082.5	3.5	17,033.1	538.2	0.1	1,419.4
Not	male	1	600.0		2,885.0		0.0	120.2
Aged	female	0						

Table 5. Summary of the seasonal mean catch rates and ages, by sex, from the pound nets in the Rappahannock River, 30 March – 3 May, 1993-2011. M = male, F = female.

Year	n	CPUE (fish/day)		CPUE (g/day)		Mean age	
		M	F	M	F	M	F
2011	215	5.5	3.5	17,031.8	27,563.8	6.0	9.5
2010	1,048	27.5	7.4	60,615.4	63,169.0	5.2	10.1
2009	620	16.2	5.7	38,323.9	44,775.3	5.1	8.5
2008	642	16.1	2.3	23,868.6	14,975.4	4.2	8.6
2007	1,104	21.4	13.2	47,614.4	87,666.9	5.0	10.5
2006	776	18.6	3.6	25,798.2	24,752.5	4.0	9.0
2005	617	12.7	4.9	26,463.2	38,962.0	4.5	9.7
2004	951	23.5	8.3	58,561.9	65,437.0	5.3	9.4
2003	470	9.4	6.2	22,767.3	53,437.0	5.2	9.5
2002	170	3.5	1.8	7,057.2	11,422.9	4.6	7.8
2001	577	15.2	3.4	24,193.2	26,298.6	4.3	9.1
2000	1,508	37.4	1.9	42,233.1	14,704.5	3.7	8.8
1999	836	27.7	2.1	31,370.7	16,821.7	3.7	9.9
1998	401	10.3	4.0	15,598.6	32,930.6	4.0	9.5
1997	406	14.4	5.9	22,400.0	49,700.0	4.0	9.2
1996	430	10.1	2.2	14,300.0	9,400.0	3.9	7.9
1995	363	11.2	3.3	13,500.0	20,000.0	3.3	7.2
1994	375	8.4	5.4	17,400.0	30,900.0	4.5	7.2
1993	565	14.4	7.3	31,400.0	37,500.0	4.6	6.9
Mean	635.5	16.0	4.9	28,447.2	35,285.1	4.5	8.9

Table 6. Numbers of striped bass in three age categories (year classes 2007-2009, 2003-2006 and 1992-2002) from gill nets in the Rappahannock River, by sampling date, spring, 2011. M = male, F = female.

Date	n	Year Class							
		No age		2007-2009		2003-2006		1992-2002	
		M	F	M	F	M	F	M	F
31 March	7	0	0	1	0	3	0	0	3
4 April	11	0	0	5	0	3	0	1	2
7 April	35	0	0	9	0	15	1	3	7
11 April	23	0	0	4	1	8	3	3	4
14 April	39	0	0	13	0	13	0	5	8
18 April	49	0	0	13	0	19	8	4	5
21 April	7	0	0	0	0	2	0	0	5
25 April	9	0	0	1	0	2	3	0	3
28 April	5	0	0	0	0	2	0	0	3
2 May	5	0	0	1	0	0	3	1	0
Total	190	0	0	47	1	67	18	17	40

Table 7. Summary of catch rates and mean ages of striped bass (n=190) from the two gill nets in the Rappahannock River, spring 2011. Values in bold are grand means for each column.

Date	n	CPUE (fish/day)		CPUE (g/day)		Mean age	
		M	F	M	F	M	F
31 March	7	4	3	12,917.8	35,348.6	6.0	13.3
4 April	11	9	2	26,094.3	32,506.3	5.3	14.0
7 April	35	27	8	81,898.2	87,579.0	5.9	12.8
11 April	23	14	9	51,338.5	68,311.3	6.6	8.9
14 April	39	31	8	91,073.3	99,697.6	5.7	12.9
18 April	49	36	13	85,012.9	84,253.0	5.4	8.5
21 April	7	0	7	0.0	42,359.8		8.1
25 April	9	3	6	7,165.7	53,886.5	5.7	10.2
28 April	5	2	3	2,893.3	37,824.0	5.0	15.7
2 May	5	2	3	5,002.6	11,670.4	7.5	7.0
Season	190	12.8	6.2	36,339.7	55,343.7	5.8	10.5

Table 8. Length frequencies (TL in mm) of striped bass sampled from the experimental gill nets in the Rappahannock River, spring, 2011.

TL	n	TL	n	TL	n	TL	n	TL	n	TL	n
300-	0	460-	4	620-	1	780-	8	940-	0	1100-	1
310-	1	470-	1	630-	4	790-	2	950-	0	1110-	2
320-	0	480-	6	640-	2	800-	4	960-	4	1120-	0
330-	1	490-	4	650-	2	810-	4	970-	0	1130-	0
340-	2	500-	7	660-	0	820-	5	980-	5	1140-	0
350-	0	510-	6	670-	2	830-	2	990-	1	1150-	0
360-	0	520-	6	680-	1	840-	4	1000-	2	1160-	0
370-	2	530-	3	690-	2	850-	0	1010-	3	1170-	0
380-	0	540-	5	700-	1	860-	3	1020-	1	1180-	0
390-	1	550-	6	710-	0	870-	0	1030-	0	1190-	0
400-	0	560-	7	720-	1	880-	3	1040-	1	1200-	0
410-	3	570-	3	730-	2	890-	0	1050-	0	1210-	1
420-	2	580-	2	740-	4	900-	3	1060-	1	1220-	0
430-	2	590-	2	750-	4	910-	3	1070-	1	1230-	0
440-	2	600-	4	760-	9	920-	1	1080-	0	1240-	0
450-	1	610-	2	770-	3	930-	1	1090-	1	1250-	0

Table 9. Mean fork length (mm), weight (g), standard deviation (SD) and CPUE (number per day; weight per day) of striped bass from gill nets in the Rappahannock River, 31 March – 2 May, 2011.

Year Class	Sex	n	Fork Length		Weight		CPUE	
			Mean	SD	Mean	SD	F/day	W/day
2009	male	1	341.0		492.5		0.1	49.3
2008	male	13	366.8	42.9	694.5	220.3	1.3	902.9
2007	male	33	457.4	24.7	1,309.4	229.8	3.3	4,321.0
	female	1	485.0		1,447.2		0.1	144.7
2006	male	25	517.0	23.1	1,923.9	300.8	2.5	4,809.8
	female	3	534.0	6.1	2,231.7	135.3	0.3	669.5
2005	male	10	566.5	35.7	2,712.0	471.2	1.0	2,712.0
	female	3	579.3	8.5	2,926.5	291.4	0.3	878.0
2004	male	13	669.2	57.1	4,265.9	983.6	1.3	5,545.7
	female	3	647.7	4.6	3,890.1	57.7	0.3	1,167.0
2003	male	17	709.3	47.8	4,953.6	893.0	1.7	8,916.5
	female	11	753.6	25.2	6,306.5	600.1	1.1	6,937.2
2002	male	12	742.0	30.1	5,620.8	528.2	1.2	6,745.0
	female	7	807.3	25.2	7,265.6	819.2	0.7	5,085.9
2001	male	2	768.5	27.5	6,621.7	74.5	0.2	1,324.3
	female	8	840.1	16.7	8,267.6	1,122.5	0.8	6,614.1
2000	male	1	637.0		3,775.8		0.1	377.6
	female	4	889.3	24.1	9,780.0	1,068.0	0.4	3,191.2
1999	female	3	909.0	26.2	10,521.3	467.3	0.3	3,156.4
1998	male	1	900.0		11,312.7		0.1	1,131.3
	female	6	946.7	17.2	12,467.2	624.4	0.6	7,480.3
1997	female	3	982.7	30.1	13,244.0	1,617.8	0.3	3,973.2
1996	female	6	978.5	53.1	13,874.0	3,039.5	0.6	8,324.4
1995	female	1	1,023.0		15,921.5		0.1	1,592.2
1994	female	0					0.0	0.0
1993	female	2	1,094.0	76.4	19,567.2	4,878.7	0.2	3,913.4
1992	female	1	1,104.0		14,954.1		0.1	1,495.4

Table 10. Summary of the season mean (30 March – 3 May) catch rates and mean ages, by sex, from the experimental gill nets in the Rappahannock River, 1993-2011. M = males, F = female.

Year	n	CPUE (fish/day)		CPUE (g/day)		Mean age	
		M	F	M	F	M	F
2011	190	12.8	6.2	36,835.4	53,127.1	5.8	10.5
2010	486	43.7	4.9	105,816.3	48,901.0	5.3	10.9
2009	231	15.9	7.2	47,440.7	58,936.7	5.6	9.7
2008	263	21.5	4.8	52,654.9	42,860.9	5.3	10.4
2007	743	75.2	7.3	134,524.0	68,017.7	4.5	11.1
2006	335	27.9	5.6	52,966.9	39,531.5	4.7	8.8
2005	322	29.7	2.7	55,674.5	19,857.3	4.8	9.2
2004	827	79.3	7.8	170,528.8	58,098.9	4.8	8.7
2003	525	52.0	3.3	98,466.7	20,716.8	4.5	8.0
2002	323	24.5	7.8	53,606.9	40,727.5	4.8	7.0
2001	622	58.1	4.1	86,827.2	31,011.3	4.3	8.3
2000	493	47.8	3.1	64,955.7	18,196.0	3.8	7.5
1999	671	64.8	2.3	55,997.3	13,331.3	3.3	7.2
1998	603	57.1	2.9	65,500.0	12,200.0	3.9	7.3
1997	824	80.6	1.8	103,600.0	14,100.0	4.0	7.8
1996	498	45.2	4.6	54,300.0	26,600.0	3.6	6.6
1995	226	15.6	7.0	45,600.0	47,700.0	4.7	7.0
1994	516	41.5	10.1	82,700.0	54,900.0	4.7	6.9
1993	527	36.6	16.0	66,900.0	56,500.0	4.9	6.3
Mean	485.5	43.7	5.8	75,520.8	38,174.4	4.6	8.4

Table 11. Numbers of striped bass in three age categories (year classes 2007-2009, 2003-2006 and 1992-2002) from gill nets in the James River, by sampling date, spring, 2011. M = male, F = female.

Date	n	Year Class							
		No age		2007-2009		2003-2006		1992-2002	
		M	F	M	F	M	F	M	F
31 March	11	0	0	3	0	7	0	0	1
4 April	12	0	0	6	0	3	0	0	3
7 April	39	1	0	16	0	18	0	2	2
11 April	69	2	1	24	1	33	4	1	3
14 April	95	3	0	32	0	33	6	4	17
18 April	56	0	0	24	0	27	3	1	1
21 April	23	0	0	2	0	5	5	5	6
25 April	29	0	0	3	1	3	10	1	11
28 April	62	0	1	26	0	23	7	1	4
2 May	59	0	1	39	3	11	3	0	2
Total	455	6	3	175	5	163	38	15	50

Table 12. Summary of catch rates and mean ages of striped bass (n=455) from the gill nets in the James River, spring 2011. Values in bold are grand means for each column. M = male, F = female.

Date	n	CPUE (fish/day)		CPUE (g/day)		Mean age	
		M	F	M	F	M	F
31 March	11	10	1	19,703.6	11,687.7	4.9	13.0
4 April	12	9	3	15,800.4	37,338.3	4.3	13.7
7 April	39	37	2	68,108.2	32,784.4	4.6	16.0
11 April	69	60	9	115,530.8	60,146.2	4.7	7.9
14 April	95	72	23	143,755.1	215,983.8	4.9	11.2
18 April	56	52	4	101,193.9	27,970.9	4.7	8.3
21 April	23	12	11	48,176.6	82,600.3	7.6	9.4
25 April	29	7	22	15,766.9	135,185.2	5.3	8.6
28 April	62	50	12	86,891.7	60,961.2	4.7	7.1
2 May	59	50	9	61,558.6	27,966.5	4.0	5.9
Season	455	35.9	9.6	69,224.4	69,262.9	4.9	9.3

Table 13. Length frequencies (TL in mm) of striped bass sampled from the experimental gill nets in the James River, spring 2011.

TL	n	TL	n	TL	n	TL	n	TL	n	TL	n
290-	3	450-	12	610-	10	770-	2	930-	1	1090-	0
300-	3	460-	13	620-	9	780-	2	940-	2	1100-	0
310-	5	470-	14	630-	3	790-	0	950-	0	1110-	1
320-	1	480-	10	640-	3	800-	2	960-	2	1120-	0
330-	0	490-	14	650-	6	810-	2	970-	2	1130-	0
340-	2	500-	20	660-	2	820-	1	980-	1	1140-	1
350-	1	510-	13	670-	2	830-	1	990-	4	1150-	0
360-	1	520-	22	680-	2	840-	4	1000-	6	1160-	1
370-	1	530-	18	690-	1	850-	4	1010-	1	1170-	0
380-	2	540-	14	700-	2	860-	3	1020-	4	1180-	0
390-	5	550-	30	710-	2	870-	2	1030-	1	1190-	0
400-	5	560-	12	720-	4	880-	2	1040-	0	1200-	0
410-	14	570-	16	730-	1	890-	7	1050-	2	1210-	0
420-	13	580-	13	740-	8	900-	2	1060-	1	1220-	0
430-	11	590-	12	750-	3	910-	4	1070-	0	1230-	0
440-	14	600-	14	760-	0	920-	0	1080-	0	1240-	0

Table 14. Mean fork length (mm), weight (g), standard deviation (SD) and CPUE (number per day; weight per day) of striped bass from gill nets in the James River, 31 March – 2 May, 2011.

Year Class	Sex	n	Fork Length		Weight		CPUE	
			Mean	SD	Mean	SD	F/day	W/day
2009	male	4	279.5	6.5	308.8	12.3	0.4	123.5
2008	male	67	378.2	39.5	768.6	204.8	6.7	5,149.6
	female	1	318.0		433.8		0.1	43.4
2007	male	103	452.1	27.0	1,283.0	262.9	10.3	13,214.9
	female	4	467.8	20.3	1,544.5	202.9	0.4	617.8
2006	male	101	512.9	26.4	1,903.0	288.4	10.1	19,220.3
	female	17	545.8	28.5	2,557.7	428.1	1.7	4,348.1
2005	male	41	568.0	33.9	2,587.6	443.1	4.1	10,609.2
	female	11	588.7	27.4	3,242.3	350.9	1.1	3,566.3
2004	male	15	623.1	67.4	3,505.5	1,025.7	1.5	5,258.3
	female	6	661.0	36.6	4,083.8	864.0	0.6	2,450.3
2003	male	7	677.6	58.3	4,345.2	951.0	0.7	3,041.6
	female	4	723.0	32.8	5,622.8	309.3	0.4	2,249.1
2002	male	4	761.5	36.8	5,900.1	1,053.0	0.4	2,360.0
	female	6	798.0	21.6	7,425.1	725.2	0.6	4,455.1
2001	male	6	755.0	64.9	5,826.4	1,257.0	0.6	3,495.8
	female	9	837.2	19.4	8,112.2	1,056.1	0.9	7,301.1
2000	male	1	839.0		8,104.7		0.1	810.5
	female	7	868.3	25.9	9,291.0	1,228.8	0.7	6,503.7
1999	male	3	843.7	42.9	7,962.0	1,818.4	0.3	2,388.6
	female	7	919.3	47.3	10,472.7	2,114.9	0.7	7,330.9
1998	male	1	801.0		6,491.9		0.1	649.2
	female	5	956.6	27.1	12,245.6	745.9	0.5	6,122.8
1997	female	9	953.2	19.4	12,651.2	1,743.1	0.9	11,386.1
1996	female	5	963.8	74.5	13,642.2	2,685.9	0.5	6,821.1
1993	female	1	1,104.0		18,265.9		0.1	1,826.6
1987	female	1	1,077.0	9	20,219.6		0.1	2,022.0
Not	male	5	530.4	101.5	2,653.3	1,841.1	0.5	1,326.7
Aged	female	3	779.3	162.8	7,393.9	3,733.6	0.3	2,218.2

Table 15. Summary of season mean (30 March – 3 May) catch rates and ages, by sex, from experimental gill nets in the James River, 1995-2011.

Year	mile	n	CPUE (fish/day)		CPUE (g/day)		Mean age	
			M	F	M	F	M	F
2011	62	455	35.9	9.6	69,224.4	69,262.9	4.9	9.3
2010	62	890	81.2	7.8	145,647.2	56,766.9	4.5	8.9
2009	62	742	65.7	8.5	124,088.4	65,020.8	4.4	9.1
2008	62	442*	37.4	6.8	69,274.6	56,798.5	4.3	8.6
2007	62	426	40.2	7.0	69,725.9	55,447.5	4.5	9.8
2006	62	1,284	116.4	12.0	213,141.3	99,613.1	4.5	9.6
2005	62	820	79.0	3.0	147,962.7	21,585.9	4.6	8.5
2004	62	1,447	127.0	4.5	207,183.6	31,237.6	4.4	8.6
2003	62	639	132.4	8.7	234,255.6	55,043.2	4.5	7.6
2002	62	824	81.4	10.1	173,663.8	47,591.2	4.7	6.4
2001	62	1,050	98.1	6.9	181,512.7	41,347.7	4.4	7.2
2000	62	1,437	139.6	4.1	241,966.4	20,396.6	4.3	6.7
1999	55	482	25.3	22.9	45,886.4	103,362.7	4.3	6.3
1998	55	199	14.9	7.2	33,000.0	46,500.0	4.7	7.5
1997	55	160	11.1	6.7	23,900.0	44,600.0	4.9	7.8
1996	55	183	10.9	7.4	23,800.0	43,500.0	4.8	7.4
1995	55	419	24.0	22.6	52,400.0	125,300.0	4.4	6.7
Mean		700.0	65.9	9.1	120,978.4	57,845.6	4.5	8.0

* 1 sex undetermined

Table 16. Values of the spawning stock biomass index (SSBI) for male and female striped bass, by gear, in the Rappahannock River, 30 March – 3 May, 1991 – 2011.

Year	Pound nets					Gill nets				
	N		SSBI (kg/day)			N		SSBI (kg/day)		
	M	F	M	F	M+F	M	F	M	F	M+F
2011	130.0	83.0	17.0	27.6	44.6	127.0	62.0	36.8	52.2	89.0
2010	825.0	219.0	60.6	63.1	123.7	437.0	49.0	105.8	48.9	154.7
2009	437.0	180.0	38.3	44.7	83.0	159.0	72.0	47.4	58.9	106.3
2008	558.0	77.0	24.2	15.1	39.3	215.0	48.0	52.7	42.9	95.6
2007	747.0	355.0	47.6	87.6	135.2	666.0	66.0	134.1	68.0	202.1
2006	647.0	122.0	25.8	24.7	50.5	275.0	56.0	49.2	39.6	88.8
2005	438.0	177.0	26.4	39.0	65.4	291.0	27.0	55.6	19.9	75.4
2004	703.0	247.0	58.5	65.4	123.9	714.0	74.0	171.9	52.0	223.9
2003	283.0	187.0	22.8	53.6	76.4	467.0	31.0	97.3	20.7	118.0
2002	113.0	57.0	7.1	11.4	18.5	240.0	78.0	53.4	40.7	94.1
2001	470.0	105.0	24.2	27.6	51.8	572.0	41.0	88.6	30.9	119.5
2000	1,436.0	71.0	42.7	14.6	57.3	452.0	27.0	65.3	16.5	81.8
1999	738.0	61.0	30.5	19.8	50.3	532.0	21.0	51.4	13.2	64.6
1998	273.0	113.0	14.8	36.4	51.2	485.0	27.0	81.5	18.5	100.0
1997	277.0	115.0	22.2	49.6	71.7	801.0	18.0	177.8	19.1	197.0
1996	334.0	73.0	14.1	9.3	23.4	433.0	46.0	63.7	30.2	93.9
1995	207.0	76.0	12.4	19.8	32.2	162.0	69.0	43.9	56.7	100.6
1994	195.0	141.0	17.1	30.9	48.0	391.0	100.0	101.6	64.7	166.3
1993	357.0	188.0	31.2	37.5	68.7	361.0	160.0	85.6	74.1	159.6
1992	51.0	100.0	5.4	19.4	24.8	61.0	74.0	15.0	32.2	47.2
1991	153.0	70.0	21.3	21.5	42.8	406.0	47.0	65.0	17.8	83.8
Mean	446.3	134.1	26.9	34.2	61.1	392.7	56.8	78.3	38.9	117.2

Table 17. Values of the spawning stock biomass index (SSBI) calculated from gill net catches of male and female striped bass in the James River, 30 March – 3 May, 1994-2011. The 1994 catch data consisted of one gill net (GN#1) and were adjusted by the proportion of the biomass that gill net #2 captured in 1995-1998 (1.8 x GN#1 for males; 1.9 x GN#1 for females).

Year	River	n		SSBI (kg/day)		
	Mile	Male	Female	Male	Female	M+F
2011	62	354	95	67.52	64.97	132.49
2010	62	817	77	145.65	56.41	202.06
2009	62	657	84	124.10	65.00	189.10
2008	62	374	67	69.27	60.25	129.52
2007	62	361	63	69.70	55.40	125.10
2006	62	1,159	120	213.14	99.49	312.63
2005	62	781	30	147.66	21.59	169.25
2004	62	1,393	50	207.04	31.24	238.28
2003	62	590	43	145.74	35.20	180.94
2002	62	728	92	173.51	47.59	221.10
2001	62	978	68	181.40	41.31	222.71
2000	62	1,381	40	241.41	21.18	262.59
1999	55	251	211	45.81	101.98	147.79
1998	55	134	65	32.97	46.48	79.45
1997	55	100	60	23.89	44.59	68.48
1996	55	108	74	23.70	43.35	67.05
1995	55	210	202	52.10	125.15	177.25
1994	55	119	64	46.27	65.74	112.01
Mean		583.1	83.6	111.72	57.05	168.77

Table 18. Predicted values of fecundity (in millions of eggs) of female striped bass with increasing fork length (mm), James and Rappahannock rivers combined.

FL	Fecundity	FL	Fecundity	FL	Fecundity	FL	Fecundity
400	0.125	600	0.446	800	1.099	1000	2.212
420	0.146	620	0.494	820	1.187	1020	2.354
440	0.168	640	0.546	840	1.280	1040	2.502
460	0.194	660	0.601	860	1.378	1060	2.656
480	0.221	680	0.660	880	1.482	1080	2.817
500	0.251	700	0.723	900	1.590	1100	2.984
520	0.284	720	0.789	920	1.703	1120	3.157
540	0.320	740	0.860	940	1.822	1140	3.337
560	0.359	760	0.935	960	1.947	1160	3.525
580	0.401	780	1.015	980	2.077	1180	3.719

Table 19. Total, age-specific, estimated total egg potential (E, in millions of eggs/day) from mature (ages 4 and older) female striped bass, by river and gear type, 30 March – 3 May, 2011. The Egg Production Potential Indexes (millions of eggs/day) are in bold.

Age	Rappahannock River						James River		
	Pound Nets			Gill Nets			Gill Nets		
	n	E	%	n	E	%	N	E	%
4	1	0.011	0.23	1	0.018	0.21	4	0.082	0.79
5	9	0.139	2.87	3	0.093	1.11	17	0.568	5.48
6	7	0.137	2.82	3	0.120	1.43	11	0.465	4.49
7	3	0.097	2.00	3	0.170	2.02	6	0.365	3.52
8	6	0.281	5.79	11	1.005	11.95	4	0.322	3.11
9	11	0.568	11.71	7	0.793	9.43	6	0.655	6.32
10	19	1.224	25.24	8	1.026	12.20	9	1.114	10.75
11	11	0.793	16.35	4	0.613	7.29	7	0.997	9.62
12	6	0.523	10.78	3	0.493	5.87	7	1.198	11.56
13	1	0.076	1.57	6	1.119	13.31	5	0.965	9.31
14	4	0.378	7.79	3	0.630	7.49	9	1.715	16.56
15	2	0.227	4.68	6	1.250	14.87	5	1.001	9.66
16	1	0.126	2.60	1	0.238	2.83	0	0.000	0.00
17	0	0.000	0.00	0	0.000	0.00	0	0.000	0.00
18	2	0.270	5.57	2	0.591	7.03	1	0.302	2.92
19	0	0.000	0.00	1	0.249	2.96	0	0.000	0.00
24	0	0.000	0.00	0	0.000	0.00	1	0.279	2.69
n/age	0	0.000	0.00	0	0.000	0.00	3	0.333	3.22
Total	83	4.850	100.00	62	8.408	100.00	95	10.361	100.00

Table 20a. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)												
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
2000													0.76
1999											0.07	0.51	3.00
1998										0.03	2.74	1.44	3.33
1997								0.79	15.61	7.49	1.38	0.37	
1996						0.19	11.54	18.13	4.29	0.25	1.83		
1995					0.60	2.15	11.50	3.34	0.10	0.68	1.40		
1994				0.04	0.51	3.90	6.33	2.79	0.11	0.58	0.41	1.70	
1993				3.04	3.97	8.10	1.48	0.11	0.50	0.87	0.28	1.43	
1992			0.12	1.44	4.80	2.86	1.25	0.04	0.50	0.50	0.87	0.19	1.13
1991		0.20	0.57	0.48	1.00	1.63	0.05	0.52	0.43	0.40	0.81	0.06	0.33
1990	0.42	0.50	1.04	1.33	2.24	1.26	0.70	0.70	0.32	0.29	0.45	0.00	0.27
1989	0.33	0.60	3.58	4.59	0.68	0.89	0.80	0.78	0.36	0.37	0.26	0.00	0.07
1988	3.58	1.60	9.54	2.22	0.60	0.37	1.50	0.89	0.39	0.05	0.10	0.00	0.00
1987	8.00	2.75	3.65	1.15	0.68	0.37	1.00	0.89	0.43	0.05	0.00	0.03	0.03
1986	2.67	1.15	0.65	0.59	0.40	0.09	1.00	0.22	0.04	0.00	0.00	0.00	0.00
1985	1.67	0.30	0.42	0.52	0.08	0.00	0.35	0.15	0.11	0.00	0.00	0.00	0.00
1984	0.50	0.40	0.58	0.33	0.28	0.00	0.35	0.07	0.04	0.00	0.00	0.00	0.00
1983	0.25	0.20	0.46	0.33	0.08	0.03	0.20	0.00	0.00	0.00	0.00	0.00	0.00
>1983	0.75	0.45	0.73	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.58	0.30	0.38	0.56	0.60	0.32	0.50	0.44	0.54	0.32	0.00	0.00	0.00
Total	18.75	8.45	21.72	13.87	14.52	12.30	20.30	14.85	29.89	39.70	18.63	5.23	15.65

Table 20b. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2008							0.23	0.46
2007						0.07	2.63	1.08
2006					0.17	1.89	6.50	1.38
2005				0.03	4.40	5.07	10.43	0.96
2004				2.52	7.20	6.93	4.23	0.79
2003			7.89	8.55	3.26	2.15	1.53	0.88
2002		1.83	6.40	6.17	0.51	1.22	1.03	0.96
2001	3.47	5.43	3.17	1.14	0.60	1.22	1.27	1.04
2000	5.57	2.77	0.14	1.12	0.57	1.19	1.77	0.63
1999	5.90	0.71	0.51	1.51	0.29	1.19	1.10	0.25
1998	3.50	0.77	0.91	1.89	0.43	0.67	0.70	0.04
1997	2.23	1.69	0.86	2.68	0.43	0.37	0.53	0.17
1996	4.16	1.69	1.17	3.80	0.46	0.70	1.13	0.08
1995	2.33	0.94	0.23	0.71	0.00	0.00	0.13	0.04
1994	1.67	0.69	0.20	0.71	0.00	0.19	0.07	0.00
1993	1.00	0.57	0.20	0.46	0.00	0.00	0.07	0.08
1992	1.10	0.29	0.11	0.20	0.00	0.03	0.07	0.00
1991	0.17	0.09	0.00	0.00	0.00	0.00	0.00	0.00
1990	0.07	0.03	0.00	0.03	0.00	0.00	0.00	0.00
1989	0.07	0.03	0.00	0.00	0.00	0.00	0.00	0.00
1988	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1987	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.40	0.49	0.26	0.00	0.00	0.07	1.47	0.04
Total	31.64	18.05	22.05	31.52	18.35	22.96	34.89	8.88

Table 21a. Catch rates (fish/day) of year classes of male striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)												
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
2000													0.76
1999											0.07	0.44	2.93
1998										0.03	2.74	1.38	3.07
1997									0.79	15.61	7.42	1.25	0.30
1996							0.19	11.54	18.11	4.03	0.16	1.50	
1995						0.55	2.15	11.46	3.21	0.10	0.03	0.56	
1994					0.04	0.51	3.80	6.19	2.68	0.08	0.39	0.03	0.23
1993					2.88	3.83	7.50	1.37	0.07	0.26	0.16	0.00	0.07
1992			0.12	1.22	4.68	2.66	1.15	0.00	0.36	0.11	0.19	0.00	0.00
1991		0.15	0.54	0.48	0.92	1.34	0.05	0.30	0.21	0.05	0.13	0.00	0.00
1990	0.17	0.35	0.96	1.30	2.00	0.94	0.35	0.11	0.00	0.03	0.00	0.00	0.00
1989	0.17	0.40	3.46	3.52	0.08	0.43	0.55	0.04	0.04	0.03	0.00	0.00	0.00
1988	3.25	0.90	7.54	1.11	0.12	0.03	0.20	0.00	0.00	0.00	0.00	0.00	0.00
1987	6.08	0.65	1.23	0.22	0.00	0.09	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1986	2.58	0.30	0.15	0.11	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1985	0.50	0.05	0.04	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1984	0.08	0.15	0.08	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<1984	0.00	0.00	0.00	0.04	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.25	0.10	0.27	0.41	0.44	0.23	0.25	0.33	0.54	0.32	0.00	0.00	0.00
Total	13.08	3.05	14.39	8.45	11.20	10.06	14.40	10.68	27.69	37.84	15.23	3.54	9.42

Table 21b. Catch rates (fish/day) of year classes of male striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
	2004	2005	2006	2007	2008	2009	2010	2011
2008							0.13	0.46
2007						0.07	2.53	1.04
2006					0.11	1.78	6.30	1.00
2005				0.03	4.34	4.48	9.63	0.67
2004				2.49	7.03	5.48	4.03	0.67
2003			7.77	8.46	3.00	1.70	1.37	0.63
2002		1.83	6.29	5.83	0.46	1.00	0.70	0.50
2001	3.47	5.40	2.91	0.97	0.49	0.81	0.67	0.25
2000	5.47	2.49	0.09	1.03	0.37	0.48	0.27	0.17
1999	5.67	0.66	0.20	1.00	0.14	0.19	0.23	0.00
1998	3.37	0.51	0.57	0.89	0.03	0.07	0.13	0.00
1997	1.93	1.00	0.29	0.37	0.06	0.04	0.00	0.00
1996	2.23	0.43	0.03	0.29	0.03	0.70	0.10	0.00
1995	0.53	0.09	0.00	0.00	0.00	0.00	0.00	0.00
1994	0.20	0.09	0.06	0.00	0.00	0.00	0.00	0.00
1993	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1992	0.07	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1991	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1990	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.40	0.46	0.29	0.00	0.00	0.07	1.40	0.04
Total	23.44	12.96	18.50	21.36	16.09	16.87	27.50	5.43

Table 22a. Catch rates (fish/day) of year classes of female striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)												
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
1999												0.06	0.07
1998												0.06	0.27
1997											0.07	0.13	0.07
1996										0.03	0.26	0.00	0.37
1995							0.05	0.00	0.04	0.13	0.00	0.63	0.80
1994							0.10	0.15	0.11	0.03	0.19	0.38	1.47
1993					0.16	0.14	0.60	0.11	0.04	0.24	0.71	0.25	1.37
1992				0.22	0.12	0.20	0.10	0.04	0.14	0.40	0.68	0.19	1.13
1991		0.05	0.04	0.00	0.08	0.29	0.00	0.22	0.21	0.34	0.68	0.06	0.33
1990	0.25	0.15	0.08	0.04	0.24	0.31	0.35	0.59	0.32	0.26	0.45	0.00	0.26
1989	0.17	0.20	0.12	1.07	0.60	0.46	0.25	0.74	0.32	0.34	0.26	0.00	0.07
1988	0.33	0.70	2.00	1.11	0.48	0.34	1.30	0.89	0.39	0.05	0.10	0.00	0.00
1987	1.92	2.10	2.42	0.93	0.68	0.29	1.00	0.89	0.43	0.05	0.00	0.03	0.03
1986	1.08	0.85	0.50	0.48	0.36	0.09	1.00	0.22	0.04	0.00	0.00	0.00	0.00
1985	1.17	0.25	0.39	0.48	0.08	0.00	0.35	0.15	0.11	0.00	0.00	0.00	0.00
1984	0.42	0.25	0.50	0.33	0.28	0.00	0.35	0.07	0.04	0.00	0.00	0.00	0.00
1983	0.25	0.20	0.46	0.33	0.08	0.03	0.20	0.00	0.00	0.00	0.00	0.00	0.00
>1983	0.58	0.45	0.73	0.26	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.25	0.20	0.12	0.15	0.16	0.09	0.25	0.11	0.00	0.00	0.00	0.00	0.00
Total	6.42	5.40	7.36	5.40	3.32	2.24	5.90	4.18	2.19	1.87	3.40	1.79	6.24

Table 22b. Catch rates (fish/day) of year classes of female striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2008							0.10	0.00
2007							0.10	0.04
2006					0.06	0.11	0.20	0.38
2005				0.00	0.06	0.59	0.80	0.29
2004				0.03	0.17	1.44	0.20	0.13
2003			0.11	0.09	0.26	0.44	0.17	0.25
2002			0.11	0.34	0.06	0.22	0.33	0.46
2001		0.03	0.26	0.17	0.11	0.41	0.60	0.79
2000	0.10	0.29	0.06	0.09	0.20	0.70	1.50	0.46
1999	0.23	0.06	0.31	0.51	0.14	1.00	0.87	0.25
1998	0.17	0.26	0.34	1.00	0.40	0.59	0.57	0.04
1997	0.30	0.69	0.57	2.31	0.37	0.33	0.53	0.17
1996	1.93	1.26	1.14	3.51	0.43	0.70	1.03	0.08
1995	1.80	0.86	0.23	0.71	0.00	0.00	0.13	0.04
1994	1.47	0.60	0.14	0.71	0.00	0.19	0.07	0.00
1993	0.90	0.54	0.20	0.46	0.00	0.00	0.07	0.08
1992	1.03	0.29	0.11	0.20	0.00	0.04	0.07	0.00
1991	0.17	0.09	0.00	0.00	0.00	0.00	0.00	0.00
1990	0.07	0.03	0.00	0.03	0.00	0.00	0.00	0.00
1989	0.07	0.03	0.00	0.00	0.00	0.00	0.00	0.00
1988	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1987	0.00	0.03	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.00	0.03	0.00	0.00	0.00	0.00	0.07	0.00
Total	8.24	5.09	3.58	10.16	2.26	6.67	7.40	3.46

Table 23a. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year Class	Survival (S)										
	91-92	92-93	93-94	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02
2001											
2000											
1999											
1998											
1997										0.480	0.843
1996										0.237	0.990
1995									0.290	0.914	0.914
1994								0.441	0.884	0.884	0.884
1993							0.183	0.993	0.993	0.993	0.993
1992					0.596	0.437	0.983	0.983	0.983	0.983	0.983
1991						0.869	0.869	0.869	0.869	0.869	0.869
1990					0.563	0.745	0.745	0.863	0.863	0.863	0.863
1989				0.440	0.440	0.899	0.975	0.689	0.689	0.703	0.703
1988			0.233	0.877	0.877	0.877	0.593	0.438	0.506	0.506	0.000
1987	0.456	0.456	0.315	0.954	0.954	0.954	0.890	0.483	0.116	0.903	0.903
1986	0.431	0.972	0.972	0.972	0.972	0.972	0.220	0.182	0.000	-----	-----
1985	0.678	0.678	0.678	0.876	0.876	0.876	0.429	0.733	0.000	-----	-----
1984			0.881	0.881	0.881	0.881	0.200	0.571	0.000	-----	-----
1983			0.717	0.846	0.846	0.846	0.000	-----	-----	-----	-----

Table 23b. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year Class	Survival (S)									
	02-03	03-04	04-05	05-06	06-07	07-08	08-09	09-10	10-11	Mean
2004							0.963	0.610	0.187	0.479
2003						0.381	0.660	0.712	0.575	0.566
2002					0.964	0.445	0.445	0.844	0.932	0.684
2001				0.584	0.796	0.796	0.796	0.796	0.819	0.760
2000			0.497	0.914	0.914	0.914	0.914	0.914	0.356	0.732
1999			0.635	0.635	0.635	0.888	0.888	0.924	0.227	0.637
1998			0.814	0.814	0.814	0.718	0.718	0.718	0.057	0.528
1997	0.843	0.843	0.843	0.843	0.843	0.583	0.583	0.583	0.321	0.663
1996	0.990	0.990	0.970	0.970	0.970	0.668	0.668	0.668	0.071	0.611
1995	0.914	0.914	0.403	0.869	0.869	0.568	0.568	0.568	0.308	0.624
1994	0.884	0.982	0.752	0.752	0.752	0.517	0.517	0.368	0.000	0.638
1993	0.993	0.699	0.570	0.898	0.898	0.646	0.646	0.646	0.646	0.719
1992	0.983	0.973	0.264	0.830	0.830	0.705	0.705	0.705	0.000	0.718
1991	0.638	0.515	0.529	0.000	-----	-----	-----	-----	-----	0.663
1990	0.775	0.259	0.754	0.754	0.754	0.000	-----	-----	-----	0.654
1989	0.646	0.646	0.429	0.000	-----	-----	-----	-----	-----	0.584
1988	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.516
1987	0.903	0.903	0.903	0.000	-----	-----	-----	-----	-----	0.637
1986	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.621
1985	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.621
1984	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.571
1983	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.610

Table 24a. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year Class	Survival (S)										
	91-92	92-93	93-94	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02
2001											
2000											
1999											
1998											
1997										0.475	0.638
1996										0.223	0.821
1995									0.280	0.559	0.559
1994								0.433	0.381	0.381	0.768
1993							0.183	0.436	0.436	0.615	0.855
1992					0.568	0.432	0.560	0.560	0.726	0.726	0.717
1991						0.473	0.473	0.700	0.787	0.787	0.000
1990					0.470	0.372	0.315	0.522	0.522	0.000	-----
1989				0.539	0.539	0.539	0.270	0.270	0.750	0.000	-----
1988			0.147	0.565	0.505	0.565	0.000	-----	-----	-----	-----
1987	0.450	0.450	0.179	0.640	0.640	0.000	-----	-----	-----	-----	-----
1986	0.116	0.500	0.733	0.364	0.000	-----	-----	-----	-----	-----	-----
1985	0.100	0.894	0.894	0.000	-----	-----	-----	-----	-----	-----	-----
1984		0.533	0.000	-----	-----	-----	-----	-----	-----	-----	-----
1983			0.717	-----	-----	-----	-----	-----	-----	-----	-----

Table 24b. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year Class	Survival (S)									
	02-03	03-04	04-05	05-06	06-07	07-08	08-09	09-10	10-11	Mean
2004							0.780	0.735	0.166	0.457
2003						0.355	0.567	0.806	0.460	0.523
2002					0.927	0.414	0.414	0.700	0.714	0.603
2001				0.539	0.333	0.914	0.914	0.827	0.373	0.599
2000			0.455	0.643	0.643	0.683	0.683	0.563	0.630	0.609
1999			0.561	0.561	0.561	0.613	0.613	0.613	0.000	0.486
1998			0.642	0.642	0.642	0.527	0.527	0.527	0.000	0.483
1997	0.638	0.638	0.518	0.608	0.608	0.162	0.667	0.000	-----	0.477
1996	0.821	0.821	0.793	0.793	0.793	0.793	0.793	0.143	0.000	0.615
1995	0.559	0.946	0.170	0.000	-----	-----	-----	-----	-----	0.409
1994	0.768	0.870	0.450	0.667	0.000	-----	-----	-----	-----	0.500
1993	0.855	0.855	0.000	-----	-----	-----	-----	-----	-----	0.496
1992	0.717	0.717	0.000	-----	-----	-----	-----	-----	-----	0.554
1991	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.508
1990	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.353
1989	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.395
1988	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.335
1987	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.372
1986	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.317
1985	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.409
1984	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.238

Table 25a. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year Class	Survival (S)										
	91-92	92-93	93-94	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02
2001											
2000											
1999											
1998											
1997											
1996											
1995											
1994											
1993											
1992											
1991											0.697
1990								0.914	0.914	0.914	0.760
1989			0.912	0.912	0.912	0.912	0.679	0.679	0.764	0.646	
1988			0.898	0.898	0.898	0.898	0.685	0.438	0.506	0.506	0.000
1987	0.987	0.987	0.802	0.802	0.802	0.802	0.890	0.483	0.116	0.902	0.902
1986	0.743	0.743	0.987	0.987	0.987	0.987	0.220	0.182	0.000	-----	-----
1985			0.743	0.900	0.900	0.900	0.429	0.733	0.000	-----	-----
1984			0.915	0.915	0.915	0.915	0.200	0.571	0.000	-----	-----
1983			0.717	0.846	0.846	0.846	0.000	-----	-----	-----	-----

Table 25b. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from pound nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year	Survival (S)									
	02-03	03-04	04-05	05-06	06-07	07-08	08-09	09-10	10-11	Mean
2002							-----	-----	-----	
2001							-----	-----	-----	
2000							-----	-----	-----	
1999							-----	0.870	0.287	0.500
1998						0.768	0.768	0.966	0.070	0.447
1997						0.612	0.612	0.612	0.321	0.521
1996						0.665	0.665	0.665	0.071	0.380
1995			0.478	0.909	0.909	0.568	0.568	0.568	0.308	0.581
1994		0.834	0.834	0.834	0.834	0.517	0.517	0.368	0.000	0.563
1993		0.657	0.600	0.906	0.906	0.646	0.646	0.646	0.646	0.698
1992		0.912	0.282	0.830	0.830	0.705	0.705	0.705	0.000	0.589
1991	0.697	0.515	0.529	0.000	-----	-----	-----	-----	-----	0.461
1990	0.760	0.269	0.754	0.754	0.754	0.000	-----	-----	-----	0.649
1989	0.646	0.646	0.429	0.000	-----	-----	-----	-----	-----	0.655
1988	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.607
1987	0.902	0.902	0.902	0.000	-----	-----	-----	-----	-----	0.675
1986	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.646
1985	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.648
1984	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.587
1983	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.610

Table 26a. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)																		
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003						
2001													2.70						
2000												0.50	8.80						
1999											0.90	1.10	16.00						
1998									1.47	9.50	8.80	12.60							
1997									11.70	18.11	27.00	10.20	4.60						
1996									0.11	35.80	21.26	17.70	4.60	4.20					
1995									0.83	11.67	10.60	5.79	2.10	3.50	1.60				
1994									1.90	29.50	32.78	3.20	1.79	1.50	1.20	1.30			
1993									4.50	20.00	83.00	7.00	0.80	2.00	1.00	1.00	0.50		
1992									2.78	7.00	11.40	14.33	0.78	1.20	0.63	1.10	0.30	0.00	
1991									0.50	2.56	1.88	5.70	2.83	1.33	0.50	0.32	0.90	0.30	0.00
1990	0.12	0.56	1.50	8.22	7.75	3.50	2.17	0.33	0.10	0.21	0.10	0.10	0.00	0.10	0.00	0.00	0.00	0.10	
1989	1.41	0.78	8.60	27.56	4.50	2.50	0.67	0.33	0.20	0.11	0.10	0.10	0.00	0.00	0.00	0.00	0.00	0.00	
1988	9.53	1.89	25.40	8.22	2.88	1.50	1.17	0.33	0.20	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
1987	23.65	5.89	10.40	2.11	1.75	1.60	0.50	0.11	0.10	0.00	0.10	0.00	0.10	0.00	0.00	0.00	0.00	0.00	
1986	11.18	3.33	1.60	0.44	1.38	0.30	0.00	0.22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
1985	4.12	1.22	0.40	1.67	0.75	0.20	0.00	0.00	0.00	0.20	0.00	0.20	0.00	0.00	0.00	0.00	0.00	0.00	
1984	1.64	0.78	0.40	0.67	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
>1984	0.82	0.55	1.90	0.78	0.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	
N/A	0.82	0.00	1.10	2.33	1.00	1.20	2.50	2.00	2.50	0.11	0.20	0.80	0.10						
Total	53.29	15.00	51.80	57.34	33.77	49.80	137.50	57.00	67.10	51.91	62.40	32.30	52.50						

Table 26b. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2009								0.10
2008							0.50	1.30
2007							4.90	3.40
2006						1.60	6.90	2.80
2005				1.22	2.40	3.40	13.30	1.30
2004			0.40	20.67	6.00	3.70	10.50	1.60
2003		0.40	9.20	31.11	6.40	3.80	4.30	2.90
2002	4.10	4.00	8.20	7.89	2.30	1.80	2.00	1.90
2001	21.78	11.80	4.90	6.11	2.20	3.10	1.20	1.00
2000	16.22	6.60	2.80	4.00	1.40	1.60	0.70	0.50
1999	10.74	2.40	1.10	2.55	0.90	1.10	0.80	0.30
1998	10.00	1.90	1.90	2.55	1.60	1.40	0.40	0.70
1997	10.32	1.40	1.60	2.00	1.40	0.20	0.40	0.30
1996	7.58	1.30	1.80	2.33	1.10	0.80	1.00	0.60
1995	2.74	0.20	0.40	0.22	0.40	0.20	0.10	0.10
1994	1.68	0.30	0.80	0.56	0.00	0.20	0.10	0.00
1993	0.64	0.10	0.20	0.67	0.00	0.20	0.20	0.20
1992	0.42	0.10	0.00	0.56	0.00	0.00	0.20	0.10
1991	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1990	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1989	0.00	0.00	0.00	0.11	0.00	0.00	0.00	0.00
1988	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1987	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.84	0.40	0.20	0.00	0.20	0.00	1.10	0.00
Total	87.06	30.90	33.50	82.55	26.30	23.10	48.80	19.10

Table 27a. Catch rates (fish/day) of year classes of male striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)												
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
2001													2.70
2000												0.50	8.80
1999											0.90	1.10	15.90
1998										1.47	9.40	8.70	12.10
1997									11.60	18.11	27.00	8.80	4.30
1996								0.11	35.70	20.95	17.00	3.30	3.80
1995							0.83	11.67	10.60	5.68	1.90	1.40	1.20
1994					1.90	29.50	32.56	2.60	1.26	1.30	0.20	0.40	
1993				4.50	20.00	82.67	6.44	0.60	1.37	0.40	0.20	0.00	
1992				2.78	6.88	11.30	14.00	0.56	0.90	0.11	0.00	0.00	0.00
1991			0.50	2.56	1.75	5.60	2.50	0.67	0.30	0.00	0.00	0.00	0.00
1990	0.12	0.44	1.50	8.22	7.00	3.20	1.83	0.22	0.00	0.00	0.00	0.00	0.00
1989	1.29	0.78	8.30	25.33	2.63	1.40	0.50	0.00	0.00	0.00	0.00	0.00	0.00
1988	9.41	1.33	20.30	4.89	1.13	0.50	0.17	0.00	0.10	0.00	0.00	0.00	0.00
1987	22.82	2.78	4.20	0.33	0.13	0.10	0.00	0.00	0.10	0.00	0.00	0.00	0.00
1986	10.23	1.22	0.90	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1985	2.35	0.11	0.00	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1984	0.71	0.11	0.10	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
<1984	0.00	0.00	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.82	0.00	0.80	1.56	0.88	1.20	2.50	1.78	2.30	0.11	0.20	0.80	0.10
Total	47.75	6.77	36.70	46.22	24.90	45.20	134.50	54.00	64.80	49.06	58.10	25.00	49.30

Table 27b. Catch rates (fish/day) of year classes of male striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2009								0.10
2008							0.50	1.30
2007							4.90	3.30
2006						1.60	6.80	2.50
2005				1.22	2.40	3.20	13.20	1.00
2004			0.40	20.67	6.00	3.40	9.90	1.30
2003		0.40	9.20	31.00	6.20	3.10	4.10	1.80
2002	4.10	4.00	7.90	7.11	2.20	1.60	1.80	1.20
2001	21.78	11.80	4.60	5.78	2.20	2.10	0.30	0.20
2000	16.00	6.50	2.30	4.00	1.20	0.50	0.10	0.10
1999	10.52	2.40	1.00	2.11	0.40	0.30	0.50	0.00
1998	9.68	1.70	0.80	2.11	0.40	0.10	0.00	0.10
1997	9.68	1.30	0.70	0.89	0.30	0.00	0.00	0.00
1996	5.68	0.70	0.60	0.33	0.10	0.00	0.00	0.00
1995	0.64	0.10	0.10	0.00	0.00	0.00	0.00	0.00
1994	0.32	0.10	0.10	0.00	0.00	0.00	0.00	0.00
1993	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1992	0.00	0.00	0.00	0.00	0.00	0.00	0.10	0.00
1991	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.84	0.40	0.20	0.00	0.10	0.00	1.00	0.00
Total	79.24	29.40	27.90	75.22	21.50	15.90	38.80	9.90

Table 28a. Catch rates (fish/day) of year classes of female striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)												
	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
2000													
1999													0.10
1998											0.10	0.10	0.50
1997									0.10	0.00	0.00	1.40	0.30
1996									0.10	0.32	0.70	1.60	0.40
1995									0.00	0.11	0.20	2.10	0.40
1994								0.22	0.60	0.53	0.20	1.00	0.90
1993							0.33	0.56	0.20	0.63	0.60	0.80	0.50
1992					0.25	0.10	0.33	0.22	0.30	0.53	1.10	0.30	0.00
1991					0.13	0.10	0.33	0.67	0.20	0.32	0.90	0.30	0.00
1990		0.11	0.00	0.00	0.75	0.30	0.33	0.11	0.10	0.21	0.10	0.00	0.10
1989	0.12	0.00	0.30	2.22	1.88	1.10	0.17	0.33	0.20	0.11	0.10	0.00	0.00
1988	0.12	0.56	5.10	3.33	1.75	1.00	1.00	0.33	0.10	0.11	0.00	0.00	0.00
1987	0.82	3.11	6.20	1.78	1.63	1.50	0.50	0.11	0.00	0.00	0.10	0.00	0.00
1986	0.94	2.11	1.70	0.33	1.38	0.30	0.00	0.22	0.00	0.00	0.00	0.00	0.00
1985	1.76	1.11	0.40	1.33	0.75	0.20	0.00	0.00	0.20	0.00	0.00	0.00	0.00
1984	0.94	0.67	0.30	0.56	0.25	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1983	0.35	0.11	1.30	0.56	0.13	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
>1983	0.47	0.44	0.50	0.22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.00	0.00	0.30	0.78	0.13	0.00	0.00	0.22	0.20	0.00	0.00	0.00	0.00
Total	5.52	8.22	16.10	11.11	9.03	4.60	3.00	3.00	2.30	2.87	4.10	8.40	3.20

Table 28b. Catch rates (fish/day) of year classes of female striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2007								0.10
2006							0.10	0.30
2005						0.20	0.10	0.30
2004						0.30	0.60	0.30
2003				0.11	0.20	0.70	0.20	1.10
2002			0.30	0.78	0.10	0.20	0.20	0.70
2001			0.30	0.33	0.00	1.00	0.90	0.80
2000	0.22	0.10	0.50	0.00	0.20	1.10	0.10	0.40
1999	0.22	0.00	0.10	0.44	0.50	0.80	0.30	0.30
1998	0.32	0.20	1.10	0.44	1.20	1.30	0.40	0.60
1997	0.64	0.10	0.90	1.11	1.10	0.20	0.40	0.30
1996	1.90	0.60	1.20	2.00	1.00	0.80	1.00	0.60
1995	2.10	0.10	0.30	0.22	0.40	0.20	0.10	0.10
1994	1.36	0.20	0.70	0.56	0.00	0.20	0.10	0.00
1993	0.64	0.10	0.20	0.67	0.00	0.20	0.20	0.20
1992	0.42	0.10	0.00	0.56	0.00	0.00	0.10	0.10
1991	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1990	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1989	0.00	0.00	0.00	0.11	0.00	0.00	0.00	0.00
1988	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.00	0.00	0.00	0.00	0.10	0.00	0.10	0.00
Total	7.82	1.50	5.60	7.33	4.80	7.20	4.90	6.20

Table 29a. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year	Survival (S)										
	91-92	92-93	93-94	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02
2001											
2000											
1999											
1998											
1997											0.726
1996									0.546	0.833	0.754
1995								0.908	0.559	0.777	0.777
1994								0.098	0.535	0.984	0.984
1993							0.084	0.535	0.957	0.707	0.707
1992							0.289	0.289	0.878	0.957	0.894
1991						0.496	0.470	0.878	0.798	0.878	0.333
1990				0.943	0.452	0.620	0.152	0.798	0.928	0.781	0.781
1989				0.163	0.556	0.268	0.495	0.606	0.550	0.928	0.928
1988			0.324	0.350	0.521	0.780	0.282	0.606	0.969	0.000	-----
1987	0.663	0.663	0.203	0.829	0.914	0.313	0.220	0.969	-----	0.969	0.000
1986	0.298	0.480	0.929	0.929	0.217	0.856	0.856	0.000	0.802	-----	-----
1985	0.740	0.740	0.740	0.449	0.802	0.802	0.802	0.802	-----	0.802	0.000
1984	0.456	0.927	0.927	0.373	0.000	-----	-----	-----	-----	-----	-----
1983			0.431	0.232	0.000	-----	-----	-----	-----	-----	-----

Table 29b. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year	Survival (S)									
Class	02-03	03-04	04-05	05-06	06-07	07-08	08-09	09-10	10-11	Mean
2006									0.405	0.405
2005									0.100	0.100
2004						0.798	0.798	0.798	0.152	0.527
2003						0.206	0.820	0.820	0.674	0.553
2002					0.962	0.292	0.933	0.933	0.950	0.747
2001			0.542	0.720	0.720	0.712	0.712	0.387	0.833	0.644
2000			0.407	0.407	0.407	0.632	0.632	0.438	0.714	0.505
1999		0.671	0.619	0.619	0.619	0.657	0.657	0.727	0.375	0.608
1998		0.794	0.634	0.634	0.634	0.627	0.875	0.707	0.707	0.697
1997	0.726	0.726	0.579	0.579	0.579	0.700	0.535	0.535	0.750	0.638
1996	0.754	0.754	0.675	0.675	0.675	0.472	0.953	0.953	0.600	0.725
1995	0.885	0.885	0.618	0.618	0.618	0.618	0.500	0.707	0.707	0.695
1994	0.984	0.984	0.690	0.690	0.700	0.598	0.598	0.500	0.000	0.609
1993	0.923	0.923	0.923	0.923	0.923	0.739	0.739	0.739	0.739	0.678
1992	0.894	0.894	0.894	0.894	0.894	0.710	0.710	0.710	0.500	0.697
1991	0.000	-----	-----	-----	-----	-----	-----	-----	-----	0.527
1990	0.781	0.000	-----	-----	-----	-----	-----	-----	-----	0.579
1989	0.928	0.928	0.928	0.928	0.928	0.000	-----	-----	-----	0.644
1988	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.408
1987	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.569
1986	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.529
1985	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.659
1984	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.493
1983	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.208

Table 30a. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year Class	Survival (S)										
	91-92	92-93	93-94	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02
2001											
2000											
1999											
1998											
1997											
1996									0.567	0.833	0.811
1995								0.908	0.536	0.777	0.335
1994								0.080	0.707	0.984	0.707
1993							0.078	0.461	0.461	0.707	0.292
1992							0.254	0.254	0.122	0.957	0.991
1991						0.446	0.268	0.448	0.000	0.878	-----
1990				0.852	0.457	0.572	0.120	0.000	-----	0.781	-----
1989				0.104	0.532	0.357	0.000	-----	-----	0.928	-----
1988			0.241	0.231	0.442	0.340	0.767	0.767	0.000	0.000	-----
1987	0.429	0.429	0.079	0.394	0.769	0.000	-----	-----	-----	0.969	-----
1986	0.119	0.738	0.122	0.000	-----	-----	-----	-----	-----	-----	-----
1985	0.520	0.520	0.520	0.000	-----	-----	-----	-----	-----	0.802	-----
1984	0.537	0.537	0.537	0.000	-----	-----	-----	-----	-----	-----	-----
1983	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Table 30b. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year	Survival (S)									
	02-03	03-04	04-05	05-06	06-07	07-08	08-09	09-10	10-11	Mean
2007									0.673	0.673
2006									0.368	0.368
2005									0.076	0.076
2004	-----	-----	-----	-----	-----	0.782	0.782	0.782	0.131	0.500
2003	-----	-----	-----	-----	-----	0.300	0.813	0.813	0.439	0.543
2002	-----	-----	-----	-----	0.900	0.309	0.905	0.905	0.667	0.686
2001	-----	-----	0.542	0.700	0.700	0.381	0.955	0.143	0.667	0.512
2000	-----	-----	0.406	0.784	0.784	0.300	0.417	0.447	0.447	0.484
1999	-----	-----	0.228	0.938	0.938	0.619	0.619	0.619	0.000	0.529
1998	-----	0.800	0.602	0.602	0.602	0.190	0.630	0.630	0.630	0.549
1997	0.710	0.710	0.134	0.827	0.827	0.337	0.000	-----	-----	0.498
1996	0.694	0.694	0.123	0.857	0.550	0.303	0.000	-----	-----	0.501
1995	0.857	0.533	0.395	0.395	0.000	-----	-----	-----	-----	0.496
1994	0.555	0.800	0.565	0.565	0.000	-----	-----	-----	-----	0.477
1993	0.000	-----	-----	-----	-----	-----	-----	-----	-----	0.283
1992	0.991	0.991	0.991	0.991	0.991	0.991	0.991	0.991	0.00	0.701
1991	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.276
1990	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.369
1989	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.231
1988	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.373
1987	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.326
1986	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.215
1985	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.369
1984	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.380
1983	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----

Table 31a. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year Class	Survival (S)										
	91-92	92-93	93-94	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02
2001											
2000											
1999											
1998											
1997											
1996											
1995											
1994											
1993											
1992											0.894
1991											0.333
1990					0.663	0.663	0.860	0.860	0.860	0.476	0.000
1989				0.847	0.585	0.548	0.548	0.606	0.928	0.928	0.928
1988			0.653	0.526	0.756	0.756	0.330	0.577	0.577	0.000	-----
1987			0.287	0.916	0.920	0.333	0.220	0.969	0.969	0.969	0.000
1986		0.806	0.901	0.901	0.217	0.856	0.856	0.000	-----	-----	-----
1985	0.911	0.911	0.911	0.567	0.719	0.719	0.719	0.719	0.000	-----	-----
1984	0.713	0.914	0.914	0.446	0.000	-----	-----	-----	-----	-----	-----
1983			0.430	0.232	0.000	-----	-----	-----	-----	-----	-----

Table 31b. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from gill nets in the Rappahannock River, 30 March – 3 May, 1991-2011.

Year	Survival (S)									
Class	02-03	03-04	04-05	05-06	06-07	07-08	08-09	09-10	10-11	Mean
2004									0.500	0.500
2003									-----	
2002						0.973	0.973	0.973	0.973	0.973
2001						-----	-----	0.900	0.889	0.894
2000						-----	-----	0.603	0.603	0.603
1999						-----	-----	0.612	0.612	0.612
1998						-----	-----	0.679	0.679	0.679
1997	0.955	0.955	0.955	0.955	0.955	0.991	0.603	0.603	0.750	0.845
1996	-----	-----	-----	-----	-----	0.794	0.794	0.794	0.600	0.740
1995	-----	-----	0.378	0.378	0.733	0.661	0.500	0.707	0.707	0.560
1994	-----	-----	0.717	0.717	0.800	0.598	0.598	0.500	0.000	0.538
1993	0.965	0.965	0.965	0.965	0.965	0.739	0.739	0.739	0.739	0.857
1992	0.894	0.894	0.894	0.894	0.894	0.650	0.650	0.650	0.650	0.787
1991	0.000	-----	-----	-----	-----	-----	-----	-----	-----	0.155
1990	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.595
1989	0.928	0.928	0.928	0.928	0.928	0.000	-----	-----	-----	0.730
1988	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.501
1987	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.496
1986	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.605
1985	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.660
1984	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.555
1983	-----	-----	-----	-----	-----	-----	-----	-----	-----	0.207

Table 32a. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)									
	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
2001										0.86
2000									0.44	15.43
1999								0.40	3.78	31.29
1998							1.58	13.50	29.67	28.86
1997						0.20	21.58	42.40	39.33	8.00
1996						9.10	73.26	32.60	11.00	2.86
1995					1.22	10.30	38.32	8.40	2.56	1.57
1994			0.10	1.55	7.11	11.70	11.05	2.60	1.11	0.57
1993		0.67	1.70	4.44	5.22	6.10	2.10	1.60	0.89	0.86
1992		4.33	2.90	3.33	3.00	2.90	1.37	1.00	0.89	0.28
1991	2.40	9.00	4.50	2.00	1.67	2.20	0.63	1.50	0.22	0.14
1990	12.40	11.11	3.10	2.00	0.78	1.40	0.42	0.50	0.11	0.14
1989	12.00	9.78	2.60	0.89	1.11	1.20	0.11	0.00	0.00	0.14
1988	3.20	2.67	1.00	1.44	0.78	0.40	0.11	0.00	0.00	0.00
1987	0.80	2.67	1.00	1.11	0.67	1.00	0.00	0.00	0.00	0.00
1986	0.80	1.78	0.80	0.33	0.11	0.30	0.00	0.00	0.00	0.00
1985	0.80	1.22	0.30	0.22	0.11	0.10	0.00	0.00	0.00	0.00
1984	1.20	0.78	0.20	0.11	0.00	0.00	0.00	0.00	0.00	0.00
>1984	1.20	0.56	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.80	2.00	0.20	0.33	0.33	1.30	0.74	0.50	1.56	0.28
Total	35.60	46.56	18.40	17.78	22.11	48.20	151.27	105.00	91.56	91.28

Table 32b. Catch rates (fish/day) of year classes of striped bass (sexes combined) sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2009								0.40
2008							0.20	6.80
2007						0.30	5.80	10.70
2006					0.40	14.50	43.90	11.80
2005				0.11	9.80	27.90	20.70	5.20
2004			0.50	12.22	15.40	15.70	7.20	2.10
2003		0.90	27.60	12.44	6.80	7.10	2.40	1.10
2002	0.36	14.70	37.00	9.00	2.90	1.30	1.30	1.00
2001	30.54	27.50	33.70	4.66	1.80	1.10	1.00	1.50
2000	48.00	19.90	9.80	1.33	1.50	1.10	0.80	0.80
1999	28.00	7.70	3.90	1.44	0.90	1.50	0.60	1.00
1998	11.82	5.10	2.60	1.34	1.50	1.20	0.40	0.60
1997	4.08	1.60	2.90	2.00	1.30	0.80	0.80	0.90
1996	3.56	1.60	3.90	1.90	1.30	1.40	0.70	0.50
1995	1.36	0.60	1.00	0.10	0.10	0.00	0.00	0.00
1994	1.00	0.50	1.00	0.10	0.10	0.10	0.10	0.00
1993	0.28	0.30	1.10	0.40	0.20	0.20	0.20	0.10
1992	0.38	0.10	0.10	0.10	0.00	0.00	0.00	0.00
1991	0.00	0.10	0.40	0.00	0.00	0.00	0.00	0.00
1990	0.00	0.00	0.40	0.00	0.00	0.00	0.00	0.00
1989	0.00	0.00	0.00	0.10	0.00	0.00	0.00	0.00
1987	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.10
N/A	2.36	1.40	2.40	0.00	0.10	0.00	2.90	0.80
Total	131.74	82.00	128.30	47.24	44.10	74.20	89.00	45.40

Table 33a. Catch rates (fish/day) of year classes of male striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)									
	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
2001										0.86
2000									0.44	15.43
1999								0.30	3.78	31.29
1998							1.58	13.50	28.89	26.00
1997						0.20	21.47	41.90	35.56	7.57
1996						7.30	72.74	31.00	8.33	2.57
1995					1.22	8.00	37.05	7.60	2.00	1.00
1994			0.10	1.56	6.78	5.20	10.53	1.70	0.67	0.00
1993		0.67	1.70	3.89	3.78	2.50	1.68	1.10	0.11	0.14
1992		4.22	2.80	2.33	1.67	1.10	1.16	0.20	0.00	0.00
1991	2.40	7.89	3.60	1.44	1.00	0.10	0.00	0.40	0.00	0.00
1990	10.60	6.33	1.50	1.33	0.22	0.30	0.00	0.00	0.00	0.00
1989	8.00	2.33	0.70	0.44	0.00	0.00	0.00	0.00	0.00	0.00
1988	1.40	0.56	0.30	0.11	0.11	0.10	0.00	0.00	0.00	0.00
1987	0.00	0.44	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1986	0.00	0.11	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.80	1.44	0.10	0.00	0.11	0.50	0.74	0.40	1.56	0.28
Total	23.20	24.00	10.90	11.11	14.89	25.30	146.95	98.10	81.33	85.14

Table 33b. Catch rates (fish/day) of year classes of male striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2009								0.40
2008							0.10	6.70
2007						0.30	5.70	10.30
2006					0.30	14.40	43.10	10.10
2005				0.11	9.80	27.30	19.20	4.10
2004			0.50	12.22	15.40	14.30	6.70	1.50
2003		0.90	27.60	12.33	6.60	6.30	2.00	0.70
2002	0.36	14.70	36.90	8.33	2.50	1.10	1.00	0.40
2001	30.54	27.30	32.30	4.33	1.50	0.80	0.40	0.60
2000	47.82	19.60	8.70	0.89	0.70	0.60	0.20	0.10
1999	27.64	7.50	3.50	1.11	0.20	0.40	0.00	0.30
1998	10.46	4.90	2.20	0.56	0.20	0.10	0.00	0.10
1997	3.90	1.00	1.40	0.22	0.00	0.00	0.00	0.00
1996	2.28	1.20	0.60	0.10	0.10	0.10	0.00	0.00
1995	0.54	0.10	0.10	0.00	0.00	0.00	0.00	0.00
1994	1.00	0.30	0.10	0.00	0.00	0.00	0.00	0.00
1993	0.00	0.10	0.00	0.00	0.00	0.00	0.00	0.00
1992	0.10	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1991	0.00	0.00	0.10	0.00	0.00	0.00	0.00	0.00
1990	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	2.36	1.40	2.40	0.00	0.10	0.00	2.70	0.50
Total	127.00	79.00	116.40	40.20	37.40	65.70	81.10	36.10

Table 34a. Catch rates (fish/day) of year classes of female striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year Class	CPUE (fish/day)									
	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003
2000										
1999								0.10	0.00	0.00
1998								0.00	0.78	2.86
1997							0.11	0.50	3.78	0.43
1996						1.80	0.53	1.60	2.67	0.28
1995						2.30	1.26	0.80	0.56	0.57
1994					0.33	6.50	0.53	0.90	0.44	0.57
1993				0.56	1.44	3.60	0.42	0.50	0.78	0.71
1992		0.11	0.10	1.00	1.33	1.80	0.21	0.80	0.89	0.28
1991		1.11	0.90	0.56	0.67	2.10	0.63	1.10	0.22	0.14
1990	1.80	4.78	1.60	0.67	0.56	1.10	0.42	0.50	0.11	0.14
1989	4.00	7.44	1.90	0.44	1.11	1.20	0.11	0.00	0.00	0.14
1988	2.20	2.11	0.70	1.33	0.67	0.30	0.11	0.00	0.00	0.00
1987	0.80	2.22	0.90	1.11	0.67	1.00	0.00	0.00	0.00	0.00
1986	0.80	1.67	0.80	0.33	0.11	0.30	0.00	0.00	0.00	0.00
1985	0.40	1.22	0.30	0.22	0.11	0.10	0.00	0.00	0.00	0.00
1984	1.20	0.78	0.20	0.11	0.00	0.00	0.00	0.00	0.00	0.00
1983	0.80	0.33	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
1982	0.40	0.22	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
N/A	0.00	0.56	0.10	0.33	0.22	0.80	0.00	0.10	0.00	0.00
Total	12.40	22.56	7.50	6.67	7.22	22.90	4.33	6.90	10.22	6.14

Table 34b. Catch rates (fish/day) of year classes of female striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011. Maximum catch rate for each year class during the sampling period is in bold type.

Year	CPUE (fish/day)							
Class	2004	2005	2006	2007	2008	2009	2010	2011
2008							0.10	0.10
2007							0.10	0.40
2006						0.10	0.80	1.70
2005						0.60	1.50	1.10
2004						1.40	0.50	0.60
2003				0.11	0.20	0.80	0.40	0.40
2002			0.10	0.67	0.40	0.20	0.30	0.60
2001		0.20	1.40	0.33	0.30	0.30	0.60	0.90
2000	0.18	0.30	1.10	0.44	0.80	0.50	0.60	0.70
1999	0.18	0.20	0.40	0.33	0.70	1.10	0.60	0.70
1998	0.36	0.20	0.40	0.78	1.30	1.10	0.40	0.50
1997	0.18	0.60	1.50	1.78	1.30	0.80	0.80	0.90
1996	1.28	0.40	3.30	1.70	1.20	1.30	0.70	0.50
1995	0.82	0.50	0.90	0.10	0.10	0.00	0.00	0.00
1994	1.00	0.20	0.90	0.10	0.10	0.10	0.10	0.00
1993	0.28	0.20	1.10	0.40	0.20	0.20	0.20	0.10
1992	0.28	0.10	0.10	0.10	0.00	0.00	0.00	0.00
1991	0.00	0.10	0.30	0.00	0.00	0.00	0.00	0.00
1990	0.00	0.00	0.40	0.00	0.00	0.00	0.00	0.00
1989	0.00	0.00	0.10	0.10	0.00	0.00	0.00	0.00
N/A	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.40
Total	4.56	3.00	12.00	6.94	6.60	8.50	7.90	9.60

Table 35a. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from gill nets in the James River, 30 March – 3 May, 1994-2011.

Year	Survival (S)										
	Class	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02	02-03	03-04
2003											
2002											
2001											
2000											
1999											0.895
1998									0.973	0.410	
1997								0.928	0.203	0.510	
1996							0.445	0.337	0.772	0.772	
1995							0.219	0.305	0.613	0.866	
1994						0.944	0.235	0.427	0.974	0.974	
1993						0.344	0.762	0.928	0.928	0.928	
1992		0.877	0.877	0.901	0.967	0.472	0.730	0.890	0.653	0.653	
1991		0.500	0.788	0.788	0.788	0.826	0.826	0.768	0.768	0.768	
1990	0.896	0.279	0.645	0.837	0.837	0.598	0.598	0.956	0.956	0.956	
1989	0.815	0.266	0.773	0.773	0.773	0.584	0.584	0.584	0.584	0.919	
1988	0.834	0.734	0.734	0.542	0.513	0.275	0.000	-----	-----	-----	
1987		0.645	0.645	0.949	0.949	0.000	-----	-----	-----	-----	
1986		0.449	0.413	0.953	0.953	0.000	-----	-----	-----	-----	
1985		0.246	0.733	0.500	0.909	0.000	-----	-----	-----	-----	
1984	0.650	0.256	0.550	0.000	-----	-----	-----	-----	-----	-----	

Table 35b. Estimated annual and geometric mean survival (S) rates for year classes of striped bass (sexes combined) sampled from gill nets in the James River, 30 March – 3 May, 1994-2011.

Year	Survival (S)							Mean
	04-05	05-06	06-07	07-08	08-09	09-10	10-11	
2006							0.269	0.269
2005						0.742	0.251	0.432
2004						0.459	0.292	0.366
2003	-----	-----	0.451	0.755	0.755	0.338	0.458	0.525
2002	-----	-----	0.243	0.322	0.670	0.670	0.769	0.486
2001	-----	-----	0.138	0.386	0.941	0.941	0.941	0.536
2000	0.415	0.492	0.391	0.391	0.733	0.853	0.853	0.564
1999	0.275	0.506	0.727	0.727	0.727	0.816	0.816	0.650
1998	0.431	0.510	0.760	0.760	0.800	0.707	0.707	0.645
1997	0.843	0.843	0.690	0.650	0.885	0.885	0.885	0.680
1996	0.772	0.772	0.487	0.858	0.858	0.500	0.714	0.635
1995	0.857	0.857	0.316	0.316	0.000	-----	-----	0.451
1994	0.974	0.974	0.562	0.562	0.562	0.562	0.000	0.613
1993	0.928	0.928	0.364	0.794	0.794	0.794	0.500	0.710
1992	0.641	0.641	0.641	0.000	-----	-----	-----	0.667
1991	0.768	0.768	0.000	-----	-----	-----	-----	0.677
1990	0.956	0.956	0.000	-----	-----	-----	-----	0.699
1989	0.919	0.919	0.919	0.000	-----	-----	-----	0.648
1988	-----	-----	-----	-----	-----	-----	-----	0.491
1987	-----	-----	-----	-----	-----	-----	-----	0.593
1986	-----	-----	-----	-----	-----	-----	-----	0.508
1985	-----	-----	-----	-----	-----	-----	-----	0.440
1984	-----	-----	-----	-----	-----	-----	-----	0.338

Table 36a. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011.

Year	Survival (S)									
	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02	02-03	03-04
2003										
2002										
2001										
2000										
1999										0.883
1998									0.900	0.402
1997								0.849	0.213	0.515
1996							0.426	0.269	0.309	0.887
1995							0.205	0.263	0.500	0.540
1994							0.161	0.838	0.838	0.838
1993				0.972	0.661	0.672	0.655	0.357	0.357	0.845
1992		0.664	0.832	0.717	0.833	0.833	0.172	0.794	0.794	0.794
1991		0.456	0.400	0.694	0.736	0.736	0.736	0.758	0.758	0.758
1990	0.597	0.237	0.887	0.475	0.475	0.000	-----	-----	-----	-----
1989	0.291	0.300	0.629	0.000	-----	-----	-----	-----	-----	-----
1988	0.400	0.536	0.606	0.606	0.909	0.000	-----	-----	-----	-----
1987		0.227	0.000	-----	-----	-----	-----	-----	-----	-----
1986		0.000	-----	-----	-----	-----	-----	-----	-----	-----

Table 36b. Estimated annual and geometric mean survival (S) rates for year classes of male striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011.

Year Class	Survival (S)							Mean
	04-05	05-06	06-07	07-08	08-09	09-10	10-11	
2006							0.234	0.234
2005						0.703	0.214	0.388
2004	-----	-----	-----	-----	0.929	0.469	0.224	0.460
2003	-----	-----	0.447	0.535	0.955	0.317	0.350	0.479
2002	-----	-----	0.226	0.300	0.440	0.909	0.400	0.405
2001	-----	-----	0.134	0.346	0.533	0.866	0.866	0.450
2000	0.410	0.444	0.102	0.787	0.857	0.333	0.500	0.414
1999	0.271	0.467	0.317	0.600	0.600	0.866	0.866	0.559
1998	0.468	0.449	0.255	0.357	0.793	0.793	0.793	0.533
1997	0.599	0.599	0.157	0.000	-----	-----	-----	0.391
1996	0.526	0.500	0.550	0.550	0.550	0.000	-----	0.439
1995	0.430	0.430	0.000	-----	-----	-----	-----	0.326
1994	0.300	0.333	0.000	-----	-----	-----	-----	0.434
1993	0.845	0.000	-----	-----	-----	-----	-----	0.566
1992	0.000	-----	-----	-----	-----	-----	-----	0.612
1991	0.758	0.758	0.000	-----	-----	-----	-----	0.610
1990	-----	-----	-----	-----	-----	-----	-----	0.417
1989	-----	-----	-----	-----	-----	-----	-----	0.286
1988	-----	-----	-----	-----	-----	-----	-----	0.481
1987	-----	-----	-----	-----	-----	-----	-----	0.108
1986	-----	-----	-----	-----	-----	-----	-----	0.000

Table 37a. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011.

Year Class	Survival (S)									
	94-95	95-96	96-97	97-98	98-99	99-00	00-01	01-02	02-03	03-04
2002										
2001										
2000										
1999										
1998										0.854
1997									0.860	0.860
1996									-----	-----
1995						0.548	0.945	0.945	0.945	0.945
1994						0.688	0.688	0.688	0.688	0.688
1993						0.844	0.844	0.844	0.844	0.844
1992						0.791	0.791	0.791	0.561	0.561
1991						0.724	0.724	0.771	0.771	0.771
1990		0.335	0.883	0.883	0.883	0.674	0.674	0.956	0.956	0.956
1989		0.255	0.858	0.858	0.858	0.584	0.584	0.584	0.584	0.919
1988	0.959	0.794	0.794	0.504	0.448	0.367	0.000	-----	-----	-----
1987		0.707	0.707	0.949	0.949	0.000	-----	-----	-----	-----
1986		0.479	0.413	0.953	0.953	0.000	-----	-----	-----	-----
1985		0.246	0.733	0.500	0.909	0.000	-----	-----	-----	-----
1984	0.650	0.258	0.550	0.000	-----	-----	-----	-----	-----	-----
1983	0.413	0.000	-----	-----	-----	-----	-----	-----	-----	-----
1982	0.550	0.000	-----	-----	-----	-----	-----	-----	-----	-----

Table 37b. Estimated annual and geometric mean survival (S) rates for year classes of female striped bass sampled from gill nets (mile 62) in the James River, 30 March – 3 May, 1994-2011.

Year Class	Survival (S)							Mean
	04-05	05-06	06-07	07-08	08-09	09-10	10-11	
2005							0.733	0.733
2004						0.654	0.654	0.654
2003	-----	-----	-----	-----	-----	0.707	0.707	0.707
2002	-----	-----	-----	0.973	0.973	0.973	0.973	0.973
2001	-----	-----	0.915	0.915	0.915	0.915	0.915	0.915
2000	-----	-----	0.852	0.852	0.956	0.956	0.956	0.913
1999	-----	-----	-----	-----	-----	0.798	0.798	0.798
1998	0.854	0.854	0.854	0.854	0.846	0.674	0.674	0.761
1997	0.860	0.860	0.860	0.730	0.885	0.885	0.885	0.853
1996	-----	-----	0.515	0.874	0.874	0.538	0.714	0.685
1995	0.945	0.945	0.333	0.333	0.000	-----	-----	0.649
1994	0.949	0.949	0.577	0.577	0.577	0.577	0.000	0.618
1993	0.844	0.844	0.364	0.794	0.794	0.794	0.500	0.742
1992	0.709	0.709	0.709	0.000	-----	-----	-----	0.603
1991	0.771	0.771	0.000	-----	-----	-----	-----	0.638
1990	0.956	0.956	0.000	-----	-----	-----	-----	0.729
1989	0.919	0.919	0.919	0.000	-----	-----	-----	0.653
1988	-----	-----	-----	-----	-----	-----	-----	0.520
1987	-----	-----	-----	-----	-----	-----	-----	0.617
1986	-----	-----	-----	-----	-----	-----	-----	0.515
1985	-----	-----	-----	-----	-----	-----	-----	0.440
1984	-----	-----	-----	-----	-----	-----	-----	0.339
1983	-----	-----	-----	-----	-----	-----	-----	0.189
1982	-----	-----	-----	-----	-----	-----	-----	0.245

Table 38a. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass from pound nets in the Rappahannock River, 1991-2011.

age	year class										
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
2		0.2	0.3	0.3	0.7	1.5	0.3	0.3	0.1	0.4	0.0
3	3.6	0.8	1.3	0.8	5.5	5.5	4.2	2.5	11.6	16.0	2.7
4	5.2	4.4	2.6	1.8	8.4	13.6	10.5	14.0	29.8	23.5	4.2
5	14.7	8.9	4.9	3.4	9.6	15.1	13.3	17.3	34.1	24.9	7.5
6	16.9	9.6	6.1	3.5	9.7	15.2	13.4	17.4	34.3	25.3	11.0
7	17.5	10.5	6.8	4.0	10.2	15.7	14.0	18.1	36.1	27.5	11.8
8	17.9	11.3	7.5	4.4	10.7	16.6	14.4	19.5	40.3	29.2	12.7
9	19.4	12.1	7.8	4.8	11.5	16.8	16.1	21.8	42.0	30.1	14.6
10	20.3	12.5	8.1	5.7	11.7	18.3	17.8	22.7	43.2	32.8	15.0
11	20.7	12.8	8.6	5.9	12.9	19.3	18.4	22.9	47.0	33.2	15.7
12	20.7	13.1	8.6	7.0	14.0	19.8	18.6	23.6	47.5	33.5	16.4
13	20.8	13.1	8.9	8.1	14.3	20.0	19.3	23.6	48.2	34.0	16.4
14	20.8	13.2	8.9	8.4	14.4	20.5	19.3	23.6	49.3	34.2	
15	20.8	13.2	9.0	8.4	14.6	20.5	19.5	23.7	49.4		
16	20.8	13.3	9.0	8.4	14.6	20.5	19.6	23.7			
17	20.8	13.3	9.0	8.4	14.6	20.6	19.6				
18	20.8	13.3	9.0	8.4	14.7	20.7					
19	20.8	13.3	9.0	8.4	14.7						
20	20.8	13.3	9.0	8.4							
area	20.8	13.3	9.0	8.4	14.7	20.7	19.6	23.7	49.4	34.2	16.4

Table 38b. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass from pound nets in the Rappahannock River, 1991-2011.

age	year class											mean
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	
2	0.0	0.0	0.0	0.0	0.0	0.1	0.0	0.1	0.1	0.2	0.1	0.2
3	0.6	0.8	3.5	1.8	7.9	2.6	4.4	2.0	2.7	0.7		4.1
4	3.6	6.3	8.9	8.2	16.5	9.8	9.5	8.5	3.8			9.7
5	9.5	9.1	12.1	14.3	19.8	16.7	19.9	9.9				13.7
6	10.2	9.2	13.3	14.8	21.9	20.9	20.9					14.7
7	10.7	10.3	13.9	16.0	23.5	21.7						15.7
8	12.2	10.9	15.1	17.0	24.4							16.8
9	12.5	12.1	16.4	18.0								17.9
10	13.7	13.9	17.5									19.0
11	14.8	14.6										19.8
12	15.1											20.3
13												20.7
14												20.8
15												20.9
16												20.9
17												20.9
18												20.9
19												20.9
20												20.9
area	15.1	14.6	17.5	18.0	24.4	21.7	20.9	9.9	3.8	0.7	0.1	20.9

Table 39a. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass from gill nets in the Rappahannock River, 1991-2011.

age	year class										
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
2		0.7	0.3	0.3	1.4	2.3	1.0	0.4	0.1	5.9	0.7
3	9.5	1.5	1.8	2.8	8.4	22.3	30.5	12.1	35.9	24.0	10.2
4	11.4	10.1	10.0	4.7	19.8	105.3	63.2	22.7	57.1	51.0	19.0
5	36.8	37.7	17.8	10.4	34.1	112.3	66.4	28.5	74.8	61.2	31.6
6	45.0	42.2	21.3	13.2	34.9	113.1	68.2	30.6	79.4	65.8	41.6
7	47.9	44.7	23.4	14.6	36.1	115.1	69.7	34.1	83.6	76.1	43.5
8	49.4	45.3	23.8	15.1	36.7	116.1	70.9	35.7	91.2	77.5	45.4
9	50.6	45.7	23.9	15.4	37.8	117.1	72.2	38.4	92.5	79.1	47.9
10	50.9	45.9	24.1	16.3	38.1	117.6	73.9	38.6	94.3	81.1	49.5
11	51.1	46.0	24.2	16.6	38.1	118.2	74.2	39.0	96.6	82.5	50.9
12	51.2	46.1	24.2	16.6	38.6	118.3	75.0	39.2	97.7	82.7	51.3
13	51.2	46.1	24.3	16.6	38.7	118.5	75.6	39.6	98.5	83.1	52.0
14	51.2	46.1	24.3	16.6	38.7	119.2	75.6	39.8	99.5	83.4	
15	51.2	46.1	24.3	16.6	39.3	119.2	75.8	39.9	100.1		
16	51.2	46.1	24.3	16.6	39.3	119.4	75.9	40.0			
17	51.2	46.1	24.3	16.6	39.3	119.6	75.9				
18	51.2	46.2	24.3	16.6	39.5	119.8					
19	51.2	46.2	24.3	16.6	39.6						
20	51.2	46.2	24.3	16.6							
area	51.2	46.2	24.3	16.6	39.6	119.8	75.9	40.0	100.1	83.4	52.0

Table 39b. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass from gill nets in the Rappahannock River, 1991-2011.

age	year class											mean
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	
2	0.5	0.3	1.4	2.1	0.2	0.2	0.6	0.0	0.0	0.5	0.1	0.9
3	1.6	9.1	23.1	6.1	9.4	20.9	3.0	1.6	4.9	1.8		11.0
4	17.6	25.3	34.9	14.3	40.5	26.9	6.4	8.5	8.3			26.8
5	28.3	31.9	39.8	22.2	46.9	30.6	19.7	11.3				37.0
6	30.7	34.7	45.9	24.5	50.7	41.1	21.0					41.1
7	31.8	38.7	48.1	26.3	55.0	42.7						44.0
8	34.3	40.1	51.2	28.3	57.8							45.9
9	35.2	41.7	52.4	30.2								47.2
10	36.3	42.4	53.4									48.0
11	37.1	42.9										48.7
12	37.4											49.0
13												49.3
14												49.5
15												49.7
16												49.7
17												49.7
18												49.8
19												49.8
20												49.8
area	37.4	42.9	53.4	30.2	57.8	42.7	21.0	11.3	8.3	1.8	0.1	49.8

Table 40a. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass from gill nets in the James River, 1994-2011.

age	year class										
	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998
2					0.0	0.3	0.1	0.0	0.0	0.1	0.8
3				2.4	4.3	2.0	1.6	1.2	9.1	21.7	14.3
4			12.4	11.4	7.2	6.5	8.7	11.5	82.4	64.1	44.0
5		12.0	23.5	15.9	10.6	11.7	20.4	49.8	115.0	103.4	72.8
6	3.2	21.8	26.6	17.9	13.6	17.8	31.5	58.2	126.0	111.4	84.6
7	5.9	24.4	28.6	19.6	16.5	19.9	34.1	60.8	128.8	115.5	89.7
8	6.9	25.3	29.4	21.8	17.8	21.5	35.2	62.4	132.4	117.1	92.3
9	8.3	26.4	30.8	22.4	18.8	22.4	35.7	63.7	134.0	120.0	93.7
10	9.1	27.6	31.2	23.9	19.7	23.2	36.7	64.3	137.9	122.0	95.2
11	9.5	27.7	31.7	24.1	20.0	23.5	37.2	65.3	139.8	123.3	96.4
12	9.6	27.7	31.8	24.3	20.4	23.8	38.2	65.4	141.1	124.1	96.8
13	9.6	27.7	32.0	24.3	20.5	24.9	38.3	65.5	142.5	124.9	97.4
14	9.6	27.8	32.0	24.4	20.6	25.3	38.4	65.5	143.2	125.8	
15	9.6	27.8	32.0	24.8	20.7	25.5	38.5	65.5	143.7		
16	9.6	27.8	32.4	24.8	20.7	25.7	38.6	65.5			
17	9.6	27.9	32.4	24.8	20.7	25.9	38.6				
18	9.6	28.0	32.4	24.8	20.7	26.0					
19	9.6	28.0	32.4	24.8	20.7						
20	9.6	28.2	32.4	24.8							
area	9.6	28.0	32.4	24.8	20.7	26.0	38.6	65.5	143.7	125.8	97.4

Table 40b. Comparison of the area under the catch curve (fish/ day) of the 1988-2009 year classes of striped bass from gill nets in the James River, 1991-2011.

age	year class											mean
	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	
2	0.2	0.2	0.4	0.2	0.5	0.5	0.1	0.2	0.3	0.2	0.4	0.3
3	4.0	15.7	31.0	14.9	28.1	12.7	9.9	14.7	6.1	7.0		11.7
4	35.3	63.7	58.5	51.9	40.5	28.1	37.8	58.6	16.8			36.0
5	63.3	83.6	92.2	60.9	47.3	43.8	58.5	70.4				54.9
6	71.0	93.4	96.8	63.7	54.4	51.2	63.7					62.0
7	74.9	94.7	98.6	65.0	56.8	53.3						64.5
8	76.3	96.2	99.7	66.3	57.9							66.4
9	77.2	97.3	100.7	67.3								67.6
10	78.7	98.1	102.2									69.0
11	79.3	98.9										69.7
12	80.3											70.2
13												70.6
14												70.8
15												70.9
16												71.0
17												71.2
18												71.2
19												71.2
20												71.2
area	80.3	98.9	102.2	67.3	57.9	53.3	63.7	70.4	16.8	7.0	0.4	71.2

Table 41a. Back-calculated length-at-age (FL, in mm) for striped bass sampled from the James and Rappahannock rivers during spring, 2011.

Year	Class	n	length-at-age (FL, in mm)							
			1	2	3	4	5	6	7	8
2009		5	159.4							
2008		17	143.8	249.3						
2007		14	146.9	263.3	374.8					
2006		22	147.3	260.0	369.3	459.1				
2005		14	134.7	244.9	354.9	447.3	521.0			
2004		17	132.3	233.3	341.4	439.9	527.4	596.1		
2003		28	138.4	243.2	341.7	438.4	532.1	612.0	680.6	
2002		21	133.8	229.1	320.5	414.9	505.5	592.1	665.6	727.8
2001		18	133.4	223.6	317.3	409.4	496.3	581.6	658.6	725.4
2000		11	132.6	220.9	313.9	399.3	485.2	560.9	632.1	702.0
1999		13	127.3	218.3	304.5	390.5	473.3	549.6	618.5	686.7
1998		13	139.8	230.4	318.0	409.0	488.3	559.7	632.9	698.9
1997		13	134.9	221.5	312.8	392.8	477.0	550.8	619.7	676.9
1996		10	129.7	212.8	298.3	384.4	465.9	546.0	608.5	673.8
1995		1	118.5	201.9	285.4	340.0	410.6	474.8	545.4	625.6
1994		2	131.0	221.5	304.6	391.4	467.5	532.6	596.1	655.7
1993		3	127.8	219.5	292.6	368.0	436.6	505.3	566.5	632.0
all		222	137.4	236.7	332.1	422.0	501.9	575.7	644.0	704.7

Table 41b. Back-calculated length-at-age (FL, in mm) for striped bass sampled from the James and Rappahannock rivers during spring, 2011.

Year	Class	n	length-at-age (FL, in mm)																	
			9	10	11	12	13	14	15	16	17	18								
2009		5																		
2008		17																		
2007		14																		
2006		22																		
2005		14																		
2004		17																		
2003		28																		
2002		21																		
2001		18	782.5																	
2000		11	762.9	814.8																
1999		13	746.9	799.7	845.5															
1998		13	755.2	807.5	852.9	893.2														
1997		13	732.1	786.9	836.7	883.3	920.1													
1996		10	734.1	783.9	829.1	877.2	916.9	953.8												
1995		1	696.3	761.9	808.6	846.8	888.9	933.8	975.5											
1994		2	715.4	771.4	820.3	869.0	910.3	951.6	991.1	1025.										
1993		3	687.8	740.2	786.1	831.9	865.7	900.6	936.6	972.6	1007.									
all		222	754.1	800.5	839.8	878.8	914.3	944.9	972.8	993.7	1007.									

Table 42. Data matrix comparing scale (SA) and otolith ages for chi-square test of symmetry. Values are the number of the respective readings of each combination of ages. Values along the main diagonal (methods agree) are bolded for reference.

S	Otolith Age																								
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	25			
1	8																								
2		11	0																						
3			4	6	1																				
4				5	6	1																			
5					6	10	3	2																	
6						2	5	3	1																
7							1	2	7	10	0														
8								1	3	19	1	3	2												
9										16	0	6	1	0											
10											3	5	12	2	0	0		1							
11												0	7	3	2	0	0	1	0						
12													1	8	2	2	0	1	1	0					
13														1	0	2	1	4	1	0	0				
14																2	1	5	2	1	0	1			
15																	0	8	1	0	0	0			
16																	0	1	0	0	0				
17																		1	1	0	0	0			
18																			0	0	0	0	0		
19																				0	0	0	0		
20																					0	0	1		
21																						0	0		
25																							0		

Table 43. Relative contributions of striped bass age classes as determined by ageing specimens (n = 223) by reading both their scales and otoliths.

Age	scale		Otolith	
	n	prop	n	Prop
1	0	0.0000	0	0.0000
2	11	0.0493	15	0.0673
3	11	0.0493	11	0.0493
4	12	0.0538	13	0.0583
5	21	0.0942	14	0.0628
6	11	0.0493	11	0.0493
7	20	0.0897	15	0.0673
8	29	0.1300	49	0.2197
9	23	0.1031	6	0.0269
10	23	0.1031	29	0.1300
11	13	0.0583	17	0.0762
12	15	0.0673	4	0.0179
13	9	0.0404	6	0.0269
14	12	0.0538	2	0.0090
15	9	0.0404	20	0.0897
16	1	0.0045	7	0.0314
17	2	0.0090	2	0.0090
18	0	0.0000	0	0.0000
19	0	0.0000	1	0.0045
20	1	0.0045	0	0.0000
21	0	0.0000	0	0.0000
25	0	0.0000	1	0.0045
	<i>Age</i> = 8.50		<i>Age</i> = 8.62	

Figure 1. Locations of the commercial pound nets and experimental gill nets sampled in spring spawning stock assessments of striped bass in the Rappahannock River, springs 1991-2011.




Figure 2. Locations of the experimental anchor gill nets sampled in spring spawning stock assessments of striped bass in the James River, springs 2003-2011.




Figure 3. Daily and historic mean river flows (cf/s) for the Rappahannock River during the 30 March – 3 May spawning stock assessment period, spring 2010.




Figure 4. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1987 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1991-2011.




Figure 5. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1988 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1991-2011.




Figure 6. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1989 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1991-2011.




Figure 7. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1990 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1991-2011.




Figure 8. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1991 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1992-2011.




Figure 9. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1992 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1993-2011.




Figure 10. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1993 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1994-2011.




Figure 11. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1994 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1995-2011.




Figure 12. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1995 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1996-2011.




Figure 13. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1996 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1997-2011.




Figure 14. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1997 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1998-2011.




Figure 15. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1998 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 1999-2011.




Figure 16. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 1999 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 2000-2011.




Figure 17. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2000 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 2001-2011.




Figure 18. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2001 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 2001-2011.




Figure 19. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2002 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 2002-2011.




Figure 20. Age-specific catch-per-unit-effort (CPUE, fish/day) of the 2003 year class of striped bass from the Rappahannock (pound nets and experimental gill nets) and James (experimental gill nets) rivers, springs, 2003-2011.





Figure 21. Magnitude of the age differences (otolith = 223) by reading both their scales and otoliths, spring, 2011.



II. Mortality estimates of striped bass (*Morone saxatilis*) that spawn in the Rappahannock River, Virginia, spring, 2010-2011.

Striped Bass Assessment and Monitoring Program
Department of Fisheries Science
School of Marine Science
Virginia Institute of Marine Science
The College of William and Mary
Gloucester Point, VA. 23062-1346

Introduction

Striped bass (*Morone saxatilis*) have historically supported one of the most important recreational and commercial fisheries along the Atlantic coast. The species is one of the most important economical and social components of finfish catches in the Chesapeake Bay area. From 1965 to 1972, annual commercial landings of striped bass in Virginia fluctuated from about 554 to 1,271 metric tons (MT). Recreational harvests, although not well documented, may have reached equivalent levels (Field 1997). Beginning in 1973, a dramatic decrease in catches occurred, and during the period 1978 through 1985, annual commercial landings in Virginia averaged about 162 MT. This decline in Virginia's striped bass landings was reflected in similar catch statistics from Maine to North Carolina.

Concern about the decline in striped bass landings along the Atlantic coast since the mid-1970's prompted the development of an interstate fisheries management plan (FMP) under the auspices of the Atlantic States Marine Fisheries Commission (ASMFC) as part of their Interstate Fisheries Management Program (ASMFC 1981). Federal legislation was enacted in 1984 (Public Law 98-613, The Atlantic Striped Bass Conservation Act), which enables Federal imposition of a moratorium for an indefinite period in those states that fail to comply with the coastwide plan. To be in compliance with the plan, coastal states have imposed restrictions on their commercial and recreational striped bass fisheries ranging from combinations of catch quotas, size limits, and time-limited moratoriums to year-round moratoriums. The FMP was modified three times from 1984-1985 to further restrict fishing (Weaver *et al.* 1986). The first two amendments emphasized the need to reduce fishing mortality and to set target mortality rates. The third amendment was directed specifically at Chesapeake Bay stocks and focused on ensuring success of the 1982 and later year classes by recommending that states protect 95% of those females until they had the opportunity to spawn at least once.

Due to an improvement in spawning success, as judged by increases in annual values of the Maryland juvenile index, a fourth amendment to the FMP established a limited fishery in the fall of 1990. This transitional fishery existed until 1995 when spawning stock biomass in the Chesapeake Bay reached extremely healthy levels (Field 1997). The ASMFC subsequently declared Chesapeake stocks to have reached benchmark levels and the states adopted a fifth amendment to the original FMP in order to allow expanded state fisheries.

The Striped Bass Program of the Virginia Institute of Marine Science (VIMS) has monitored the size and age composition, sex ratio and maturity schedules of the spawning striped bass stock in the Rappahannock River since 1981. In conjunction with the monitoring studies, VIMS established a tagging program in 1988 to provide information on the migration, relative contribution to the coastal population, and annual survival of striped bass that spawn in the Rappahannock River. This program is part of an active cooperative tagging study that currently involves 15 state and federal agencies along the Atlantic coast. The U.S. Fish and Wildlife Service manages the coast-wide tagging database. Hence, commercial and recreational anglers that target striped bass are encouraged to report all recovered tags to that agency. The analysis protocol, as established by the ASMFC Striped Bass Tagging Subcommittee, involves fitting a

suite of reformulated Brownie models (Brownie et al. 1985; White and Burnham 1999) to the tag return data.

Although the initial purpose of the coast-wide tagging study was to evaluate efforts to restore Atlantic striped bass stocks (Wooley *et al.* 1990), tagging data are now being collected to monitor striped bass mortality rates in a recovered fishery.

This section is an update to material provided by Sadler et al. (2001). They did a comprehensive analysis of the Rappahannock River striped bass tagging data, gave a detailed description of the ASMFC analysis protocol and presented annual survival (S) estimates derived from tag-recovery models developed by Seber (1970) as well as estimates of instantaneous fishing mortality (F) that followed when S was partitioned into its components using auxiliary information.

Multi-year Tagging Models

Tag return data is generally represented by constructing an upper triangular matrix of tag recoveries, where each cell of the matrix contains the number of tag returns from a particular year of tagging and recovery. For example, a study with I years of tagging and J years of recovery would yield the following data matrix

$$R = \begin{bmatrix} r_{11} & r_{12} & \cdots & r_{1J} \\ - & r_{22} & \cdots & r_{2J} \\ \vdots & \vdots & \ddots & \vdots \\ - & - & - & r_{IJ} \end{bmatrix}, \quad (1)$$

where r_{ij} is the number of tags recovered in year j that were released in year i (note, $J \geq I$). Tagging periods do not necessarily have to be yearly intervals; however, data analysis is easiest if all periods are the same length and all tagging events are conducted at the beginning of each period.

Application of tagging models involves constructing an upper triangular matrix of expected values and comparing them to the observed data. Since the recovery data over time for each year's batch of tagged fish can be assumed to follow a multinomial distribution, the method of maximum likelihood can be used to obtain parameter estimates. Analytical solutions for the maximum likelihood parameter estimates are generally not available. Hence, several software packages that numerically maximize a product multinomial likelihood function have been developed for application of tagging models. They include programs SURVIV (White 1983), MARK (White and Burnham 1999), and AVOCADO (Hoenig et al. in prep.).

Seber models: White and Burnham (1999) reformulated the original Brownie et al. (1985) models in the way originally suggested by Seber (1970) to create a consistent framework for modeling mark-recapture data (Smith et al. 2000). This framework served as the foundation for program MARK, which is a comprehensive software package for the application of capture-recapture models. For time-specific parameterization of the Seber models, the matrix of expected values associated with equation (1) would be

$$E(R) = \begin{bmatrix} N_1(1-S_1)r_1 & N_1S_1(1-S_2)r_2 & \cdots & N_1S_1\cdots S_{J-1}(1-S_J)r_J \\ - & N_2(1-S_2)r_2 & \cdots & N_2S_2\cdots S_{J-1}(1-S_J)r_J \\ \vdots & \vdots & \ddots & \vdots \\ - & - & - & N_I(1-S_I)r_I \end{bmatrix} \quad (2)$$

where N_i is the number tagged in year i , S_i is the survival rate in year i and r_i is the probability a tag is recovered from a killed fish regardless of the source of mortality. For the 2006 estimates the updated version of MARK (version 4.3) replaced the version used in previous years (version 4.2).

The Seber models are simple and robust, but they do not yield direct information about exploitation (u) or instantaneous rates of fishing and natural mortality, which are often of interest to fisheries managers. Estimates of S can be converted to the instantaneous total mortality rate via the equation (Ricker 1975)

$$Z = -\log_e(S) \quad (3)$$

and, if information about the instantaneous natural mortality rate is available, estimates of the instantaneous fishing mortality can be recovered. Given estimates of the instantaneous rates, it is possible to recover estimates of u if the timing of the fishery (Type I or Type II) is known (Ricker 1975).

Instantaneous rate models: Hoenig et al. (1998a) modified the Brownie et al. (1985) models to allow for the estimation of instantaneous rates of fishing and natural mortality. This extension showed how information on fishing effort could be used as an auxiliary variable and also discussed generalizing the pattern of fishing within the year. The matrix of expected values corresponding to equation (1) for a model that assumes time-specific fishing mortality rates and a constant natural mortality rate would be

$$E(R) = \begin{bmatrix} N_1\phi\lambda u_1(F_1, M) & N_1\phi\lambda u_2(F_2, M)e^{-(F_1+M)} & \dots & N_1\phi\lambda u_J(F_J, M)e^{-\left(\sum_{k=1}^{J-1} F_k+(J-1)M\right)} \\ - & N_2\phi\lambda u_2(F_2, M) & \dots & N_2\phi\lambda u_J(F_J, M)e^{-\left(\sum_{k=2}^{J-1} F_k+(J-2)M\right)} \\ \vdots & \vdots & \ddots & \vdots \\ - & - & - & N_J\phi\lambda u_J(F_J, M) \end{bmatrix}$$

(4)

where ϕ is the probability of surviving being tagged and retaining the tag in the short-term, λ is the tag-reporting rate, and $u_k(F_k, M)$ is the exploitation rate in year k which, as mentioned above, depends on whether the fishery is Type I or Type II. For striped bass, a Type II (continuous) fishery is assumed. Note that ϕ and λ are considered constant over time.

These models are not as simple as the Seber models, but they do yield direct estimates of F and, depending on the information available, either M or $\phi\lambda$. Also, they can be parameterized to allow for non-mixing of newly and previously tagged animals (Hoenig *et al.* 1998b). If the goal of a particular tagging study is to estimate F and M , then auxiliary information on the tag reporting and tag-induced handling mortality rate is required to apply the instantaneous rates formulation. However, if M is known, perhaps from a study that related it to life history characteristics (e.g., Beverton and Holt 1959; Pauly 1980; Hoenig 1983; Roff 1984; Gunderson and Dygert 1988), then these models can be used to estimate F and $\phi\lambda$.

In either case, the auxiliary information needed (i.e., $\phi\lambda$ or M) can often be difficult to obtain in practice, and since F , M and $\phi\lambda$ are related functionally in the models, the reliability of the parameters being estimated is directly related to the accuracy of the estimated auxiliary parameter (Latour *et al.* 2001a).

Materials and Methods

Capture and Tagging Protocol

Each year from 1991 to 2011, during the months of March, April and May, VIMS scientists obtained samples of mature striped bass on the spawning grounds of the Rappahannock River. Samples were taken twice-weekly from pound nets owned and operated by a cooperating commercial fisherman. The pound net is a fixed trap that is presumed to be non-size selective in

its catch of striped bass, and has been historically used by commercial fishermen in the Rappahannock River.

All captured striped bass were removed from each pound net and placed into a floating holding pocket (1.2m x 2.4m x 1.2m deep, with 25.4mm mesh and a capacity of approximately 200 fish) anchored adjacent to the pound net. Fish were dip-netted from the holding pocket and examined for tagging. Fork length (FL) and total length (TL) measurements were taken and whenever possible the sex of each fish was determined. Striped bass not previously marked and larger than 458 mm TL were tagged with sequentially numbered internal anchor tags (Floy Tag and Manufacturing, Inc.). Each internal anchor tag was applied through a small incision in the abdominal cavity of the fish. A small sample of scales from between the dorsal fins and above the lateral line on the left side was removed and used to estimate age. Each fish was released at the site of capture immediately after receiving a tag.

Analysis Protocol

Program MARK: The ASMFC Striped Bass Tagging Subcommittee established a data analysis protocol that involves deriving survival estimates from a suite of Seber (1970) models. The protocol is used by each state and federal agency participating in the cooperative tagging study. Tag recoveries from striped bass greater than 457 mm total length are analyzed from known producer areas (including Chesapeake Bay). Tag recoveries from striped bass that were greater than 711 mm total length (TL) at the time of tagging are analyzed from all coastal states since those fish are believed to be fully recruited to the fishery and also because they constitute the coastal migratory population (Smith *et al.* 2000).

The protocol consists of six steps. First, prior to data analysis, a set of biologically reasonable candidate models is identified. Characteristics of the stock being studied (i.e., Chesapeake Bay, Hudson River, Delaware Bay, etc.) and time are used as factors in determining the parameterizations of the candidate models. These models are then fit to the tagging data (program MARK), and Akaike's Information Criterion (AIC) (Akaike 1973; Burnham and Anderson 1992), quasi-likelihood AIC (QAIC) (Akaike 1985), and goodness-of-fit (GOF) diagnostics are used to evaluate their fit (Burnham *et al.* 1995). The overall estimates of survival are calculated as a weighted average of survival from the best fitting models, where the weight is related to the model fit (i.e., the better the fit, the higher the weight) (Buckland *et al.* 1997; Burnham and Anderson 1998). For the 2006 analysis, the last regulatory period (2000-present in previous years), was redefined as two periods (2000-2002 and 2003-present) to reflect the adoption of the latest amendment to the Federal Management Plan (FMP). The candidate models for striped bass survival (S) and tag recovery (r) rates are:

S(.)r(.)	Survival and tag-recovery rates are constant over time.
S(t)r(t)	Survival and tag-recovery rates are time-specific.
S(.)r(t)	Survival rate is constant and tag-recovery rates are time-specific.
S(p)r(t)	Survival rates vary by regulatory periods (p=constant 1990-1994, 1995-1999, 2000-2002 and 2003-2010) and tag-recovery rates are time-specific.

S(p)r(p)	Survival and tag-recovery rates vary by regulatory period.
S(.)r(p)	Survival rate is constant and tag-recovery rates vary by regulatory periods.
S(t)r(p)	Survival rates are time-specific and tag-recovery varies by regulatory periods.
S(d)r(p)	Survival and tag-recovery rates vary over different regulatory periods (d= constant 1990-1994, 1995-1999, 2000-2002, 2003-2009 and 2010).
S(v)r(p)	Survival and tag-recovery rates vary over different regulatory periods (v= constant 1990-1994, 1995-1999, 2000-2002, 2003-2008, 2009 and 2010).

The striped bass tagging data contain a large number of tag-recoveries reflecting catch-and-release practices (i.e., the tag of a captured fish is clipped off for the reward and the fish released back into the population). Analysis utilizing these data leads to biased survival estimates if tag recoveries for re-released fish are treated as if the fish were killed. The fifth step applies a correction term (Smith *et al.* 2000) to offset the re-release-without-tag bias assuming a tag reporting rate of 0.43 (D. Kahn, Delaware Division of Fish and Wildlife, personal communication). The sixth step converts estimates of S_i to F_i via equation (3), assuming that $Z = F + M$ and M is 0.15 (Smith *et al.* 2000).

Dunning et al. (1987) quantified the rates of tag-induced mortality and tag retention for Hudson River striped bass. They found retention of internal anchor tags placed into the body cavity via an incision midway between the vent and the posterior tip of the pelvic fin was 98% for fish kept in outdoor holding pools for 180 days. Their holding experiment revealed that the survival rates of both tagged and control fish were not significantly different over a 24-hour period. A similar study conducted on resident striped bass within the York River, Virginia, yielded survival in the presence of tagging activity and short-term tag retention rates each in excess of 98% (Sadler et al. 2001). Based on these results, the ASMFC analysis protocol specifies making no attempts to adjust for the presence of short-term tag-induced mortality or acute tag-loss

Exploitation rate (R/M) method: Estimates of the exploitation rate (μ) are calculated by the recapture rate adjusted for the reporting rate:

$$\mu = (R_k + R_r * 0.08) / (\lambda M)$$

where R_k is the number or recaptures kept with tags, R_r is the number of fish released with tags, λ is the reporting rate (0.43) and M is the number of tagged striped bass released. The exploitation rate is then used to calculate the estimate of fishing mortality (F) by solving the following equation for F :

$$\mu = F / (F + M) * (1 - \exp(-M - F))$$

where natural mortality (M) is assumed to be 0.15. Other adjustments are made for tag-induced mortality (0.013) and hook-and-release mortality (0.08).

Catch equation method: Fishing and natural mortality can be estimated from the tagging data using the above described relationship between exploitation rate, fishing mortality and natural mortality. This can be rewritten as:

$$F = \mu / (S - 1) * \ln(S)$$

Survival (S) is estimated from the tagging data using the MARK models used with the estimate of μ to determine F.

Instantaneous rates methods: This method (defined in the multi-year tagging methods section) allows the estimate of natural mortality to be constant, or to vary by periods. In the 2009 analysis, two scenarios were assessed, based of the ASMFC tagging subcommittee recommendations: constant natural mortality and two periods of differing natural mortality. To determine when to separate the two periods all possible 2 period combinations were tried (1990, 1991-2008; 1990-1991, 1992-2008; ... 1990-2007, 2008) and the minimum qAIC value used as the determinant. The resultant periods were 1990-1997, 1998-2008 for striped bass > 457 mm TL and 1990-2002, 2003-2008 for striped bass > 711 mm TL. These periods were used in the models this year, with the terminal year being 2010. The candidate models for fishing mortality (F), release mortality (F') and natural mortality (M) are:

- F(t) F'(t)M(p) Fishing and release mortalities time-specific; natural mortality constant.
- F(p)F'(t)M(p) Fishing mortality period-specific (1990-1994, 1995-1999, 2000-2002 and 2003-2010); release mortality time-specific; natural mortality constant.
- F(.)F'(t)M(p) Fishing mortality constant; release mortality time-specific; natural mortality constant.
- F(t)F'(p)M(p) Fishing mortality time-specific; release mortality period-specific; natural mortality constant.
- F(t)F'(.)M(p) Fishing mortality time-specific; release and natural mortalities constant.
- F(p)F'(p)M(p) Fishing and release mortalities period-specific; natural mortality constant.
- F(.)F'(.)M(p) Fishing, release and natural mortalities constant.
- F(d)F'(d)M(p) Fishing and release mortalities vary over a different periods (1990-1994, 1995-1999, 2000-2002, 2003-2009 and 2010); natural mortality constant.
- F(v)F'(v)M(p) Fishing and release mortalities vary over different periods (1990-1994, 1995-1999, 2000-2002, 2003-2008 and 2009, 2010); natural mortality constant.

All analytical approaches were applied to striped bass greater than 457 mm total length (minimum legal size) and to striped bass greater than 710 mm TL (coastal migrants).

Results

Spring 2010 Tag Release summary

A total of 416 striped bass were tagged and released from the pound nets in the Rappahannock River between 11 April and 9 May, 2011 (Table 1). There were 225 resident striped bass (457-710 mm TL) tagged and released. These stripers were predominantly male (83.8%), but the female stripers were larger on average. The median date of these tag releases, to be used as the beginning of the 2010-2011 recapture interval, was 18 April. There were 188 migrant striped bass (>710 mm TL) tagged and released. These stripers were predominantly female (64.4%) and their average size was larger than for the male striped bass. The median date of these tag releases was 18 April. The tag release totals were only one quarter the release total for 2010, and they were well below the release targets of 700 resident and 300 migratory striped bass.

Mortality Estimates, 2010-2011

Tag recapture summary: A total of 74 striped bass (>457 mm TL) were recaptured between 1 January and 31 December, 2010. The largest source of recaptures (58.1%) was from Chesapeake Bay (41.9% in Virginia, 16.2% in Maryland, Table 2). Other recaptures came from Massachusetts (14.9%), New Jersey (10.8%), New York (5.4%), Delaware and North Carolina (4.1% each) and Rhode Island (2.7%). There were no recaptures reported from Maine, New Hampshire or Connecticut. The primary peak of recaptures was in May-July, with a secondary peak from October through December. However, there were recaptures in every month of the year except March.

A total of 35 migratory striped bass (>710 mm total length) were recaptured between 1 January and 31 December, 2010. The largest source (25.6%) of the recaptured tagged striped bass was from Massachusetts (31.4%, Table 3). Other recaptures came from New Jersey (22.9%), Chesapeake Bay (8.6% in Maryland and 5.7% in Virginia), New York, Delaware and North Carolina (8.6% each), and Rhode Island (5.7%). There were no recaptures reported from Maine, New Hampshire or Connecticut. The peak month for recaptures was in May, but some migrant striped bass were recaptured from every month of the year except March.

ASMFC protocol: Survival estimates were made utilizing the mark-recapture data for the Rappahannock River from 1990-2010. The suite of Seber (1970) models consisted of nine models that each reflected a different parameterization over time. Models that allowed parameters to be both time-specific and constant across time were specified. Since Atlantic striped bass have been subjected to a variety of harvest regulations since 1990, it was hypothesized that these harvest regulations would influence survival and catch rates. Hence, models that allowed parameters to be constant for the time periods coinciding with stable coast-wide harvest regulations were also specified. Models that allowed trends within periods and Virginia-specific models for the transition from a partial to an open fishery were eliminated prior

to the 2006 analyses after the ASMFC tagging subcommittee determined that they only poorly evaluated the data and carried no weight in the model averaging for multiple years.

Estimates of survival using MARK: Forty-seven striped bass (≥ 457 mm TL) tagged in spring 2010 and 28 striped bass tagged in previous springs were harvested during the 2011 recapture interval. These were added to complete the input matrix (Table 4) for annual estimates of survival using program MARK. Likewise, there were 17 striped bass (≥ 711 mm TL) tagged in spring 2010 and 17 striped bass tagged in previous springs harvested during the 2011 recapture interval and used to complete the input matrix (Table 5).

The suite of nine models were ranked and weighted by MARK according to their QAIC values. For striped bass ≥ 457 mm TL, the time-specific model received 96.7% of the weighting (Table 6). The 2010 estimate of survival was 0.731 which became 0.742 when adjusted for release bias (Table 7). The 2010 survival estimate was much higher than the 2009 estimate and continues trend on increasing estimates on survival from 2004-2010. The ranking and weighting among the nine models were much different in striped bass ≥ 711 mm TL with the Des period model (weighted 64.8%) and the Vic period model (22.4%) weighted highest (Table 8). The 2010 estimate of survival was 0.717 (0.732 after bias adjustment) which was also higher than the 2009 survival estimate (Table 9).

Catch equation estimates of mortality and exploitation rates: The MARK estimates of survival were used to estimate exploitation rate (U) as well as instantaneous (Z), annual (A), fishing (F) and natural (M) mortalities. The 2010 estimates for striped bass ≥ 457 mm TL were 0.30 (Z), 0.26 (A), 0.04 (U), 0.04 (F) and 0.26 (M, Table 10). The estimates of U and F have declined steadily since 2001 while the estimate of M has fluctuated, but remained well above the assumed value of 0.15 since 1996 (except 2003). The 2010 estimates for striped bass ≥ 711 mm TL were 0.31 (Z), 0.27 (A), 0.05 (U), 0.06 (F) and 0.25 (M, Table 11). The estimates of F and U have declined since 2003, but the M estimate, while lower than the value for the smaller striped bass, has also exceeded the 0.15 value since 2004.

Instantaneous rates model estimates of survival, fishing and natural mortality: The results of the iterative running of two natural mortality period scenarios resulted in the adoption of 1990-1997 and 1998-2010 M periods for striped bass ≥ 457 mm TL and 1990-2003 and 2004-2010 M periods for striped bass ≥ 711 mm TL.

Thirty striped bass (≥ 457 mm TL) tagged in spring 2010 were harvested and an additional 11 were released with the tag streamers cut off during the 2010-2011 recapture interval. In addition, there were 17 striped bass harvested and five striped bass released with their streamers cut off that were tagged in previous springs. These were added to their respective input matrixes (Tables 12a,b) for estimating survival and mortality parameters using the instantaneous rates model. Likewise there were 11 harvested and four released striped bass (≥ 711 mm TL) from striped bass tagged in spring 2010 and 10 harvested and two released striped bass tagged in previous springs during the 2010-2011 recapture interval used to complete their respective instantaneous rate model input matrixes (Tables 13a, b).

The Des period F, Des period F' model received the highest weighting in the 2010 constant M IRCR analysis for striped bass greater than 457 mm TL (Table 14). However, when these 1M suite of models were run in addition to the same models with two M periods, none of the 1M models received any weight (Table 15). The Des period model was the highest weighted model (0.984), with only minor contributions from the Vic period model and the time-specific F, period-based F' model (0.016 combined).

The IRCR estimates of survival for 2010 were 0.594 (1M) and 0.542 (2M, Table 16). The 2010 estimates of natural mortality were 0.473 and 0.560 respectively, while the 2010 estimates of fishing mortality were 0.046 and 0.052. Consistent with the 2008 and 2009 results, the two M IRCR analysis gave lower estimates of survival and higher natural mortality estimates than if constant natural mortality is assumed. The natural mortality estimates from both analyses are much higher than the 0.15 value assumed in the MARK analysis.

The Vic period model received the heaviest weighting (0.951) for the 1M IRCR analysis for striped bass ≥ 711 mm TL (Table 17). Again, the time-specific 1M model received virtually zero weight (0.001) when run in combination with the 2M variants (Table 18). Again, the Vic period model received the strongest weighting (0.571) followed by the Des variation period model (0.346) and the period model (0.080). The time-specific F, period-based F' model was only minimally weighted (0.001).

The 2010 IRCR estimates of survival were 0.701 (1M) and 0.598 (2M) respectively (Table 18). The 2010 estimates of natural mortality were 0.287 and 0.428 while the estimates of fishing mortality were 0.064 and 0.085 respectively. Again the estimates of natural mortality were much higher than the 0.15 assumed in the MARK analysis but much lower than the estimates for striped bass ≥ 458 mm TL.

Model Evaluations

Latour et al. (2001b) proposed a series of diagnostics that can be used in conjunction with AIC and GOF measures to assess the performance of tag-recovery models. In essence, they suggested that the fit of a model could be critically evaluated by analyzing model residuals and that patterns would be evident if particular assumptions were violated.

For the time-specific Seber (1970) model, Latour et al. (2002) proved the existence of several characteristics about the residuals. Specifically, they showed that row and column sums of the residuals matrix must total zero, and further, they showed that the residuals associated with the “never seen again” category must also always be zero unless parameter estimates fall on a boundary condition. Latour et al. (2001c) also scrutinized the residuals associated with the instantaneous rates model and found the residual matrix of this model possessed fewer constraints than the time-specific Seber model. Although the row sums category must total zero, the column sums and the associated residuals can assume any value.

ASMFC protocol: Given that management regulations applied to striped bass during the 1990s have specified a wide variety of harvest restrictions, it would be reasonable to assume that the time-specific models (e.g. $S(t)r(t)$, $S(p)r(t)$, $S(t)r(p)$, etc.) were most appropriate for data analysis. However, elements of the Rappahannock River tag-recovery matrix did not allow these models to adequately fit the data. The low total number tagged of striped bass releases, and the resultant low numbers of recaptures reported from the 1994 and 1996 cohorts (e.g. six from the 1996 cohort) relative to other years, may have resulted in the poor fit of the time-specific models. Unfortunately, numerical complications resulting from low sample size may have caused some of the more biologically reasonable models to not fit the Rappahannock River data well.

Discussion

In spring 2010, the release total for striped bass tagged in the Rappahannock was the highest since 1991 for all striped bass and the highest since 2007 and third highest overall for migratory striped bass. However the recapture rate of 0.022 (47/2050) for the >18 inches TL releases was the lowest of the 21 years in the release/recapture matrix that is the basis for our analyses (0.048-0.111, mean= 0.073 for 1990-2009). The recapture rate of 0.035 for the >28 inches TL releases was within the 1990-2009 range (0.015-0.152) but was also well below the mean (0.080). Comparisons with the other tagging programs did not reveal similar results, so the survival estimates for the Rappahannock striped bass were much higher than the estimates for previous years and in contrast with the estimates for the other tagging programs. It remains to be determined whether the results represent actual increased survival or are the result of atypical fishing pressure or migration patterns and it is possible that these estimates will be revised downward in subsequent analyses.

The program MARK survival estimates for 2010 were 0.731 for striped bass greater than 18 inches (457 mm) total length and 0.717 for striped bass greater than 28 inches (711 mm) total length (migratory). The survival estimate for striped bass greater than 18 inches was much higher than the estimate for 2009 and continued an apparent trend of increasing survival since 2004. Likewise, the 2010 survival estimate for striped bass greater than 28 inches was higher the 2009 estimate and the highest of the time series.

The resultant estimates of fishing mortality were below the 0.30 limit endorsed by the ASMFC. In 2006 the final period in the period-based models was redefined and partitioned into two periods (coined Des and Vic). Until this year, this redefinition of the final period produced estimates of fishing mortality that exceeded the threshold value of 0.30 endorsed by the ASMFC. Prior to 2004, the models that assume constant survival and/or reporting rate and the models that partition the time series into two periods (1990-1994 and 1995-2004) were found to best fit the data and contributed most heavily to the analysis (0.62 in 2003). These are the models that use the fewest parameters to produce the estimates of survival and fishing mortality. However, since 2004 the regulatory-based reporting rate models were the most heavily weighted. However, these new models haven't been fully evaluated and the results are contrary to the other analytical

methods. Furthermore, this analysis assumes a constant value of 0.15 for natural mortality and there is increasing evidence that natural mortality has increased and may greatly exceed this value which would result in an over estimation of fishing mortality.

The catch equation method uses the survival estimates from the MARK analysis, but rather than assume a value of natural mortality, it partitions mortality into both its natural and fishing components. This methodology produced 2010 estimates of fishing mortality of 0.04 (>18 inches) and 0.06 (>28 inches), well below the ASMFC threshold. It also produced estimates of natural mortality well above 0.15 in both size groups.

Recently, we have begun using instantaneous rates models to study mortality rates of resident striped bass as an alternative to the Seber-Brownie models. These models are more efficient in that they require fewer parameters, and they can be used to obtain estimates of current mortality rates. This provides greater flexibility in modeling mortality over time. Starting in 2008, the protocol was modified to allow for an increase in natural mortality in recent years (2M periods vs. constant M) and these models were found to better fit the data. The estimates of fishing mortality were 0.05 (1M and 2M) for striped bass >18 inches TL and 0.06 (1M) and 0.09 (2M) for striped bass >28 inches TL. The IRCR analyses also estimated that the natural mortality has greatly increased in the recent years.

A number of studies in recent years have indicated a development of mycobacteriosis, a bacterial disease in Chesapeake Bay striped bass beginning around 1997 (Vogelbein et al 1999). The disease is believed to have spread significantly thereafter. It has been suggested that mycobacteriosis might lead to an increase in striped bass mortality (Jiang et al 2007, Guathier et al 2008 and Hoenig et al 2009). Kahn and Crecco (2006) analyzed MD and VA spring tagging data for two groups of fish (fish \geq 18 inches TL and fish \geq 28 inches TL) using Program MARK and the catch equation. They reported high natural mortality rates similar to those estimated in the present analysis and suggested that their high estimates of natural mortality were related to mycobacteriosis. However, as mentioned above, the natural mortality could be overestimated if migration out of the Bay is not accounted for partially or completely.

A significant advantage of the catch equation method and the IRCR model is the ability to estimate natural mortality in addition to fishing mortality, either through the use of external model results (the catch equation uses survival estimates from Program MARK) or internally (IRCR model). As reported above, estimated values of natural mortality from both methods were substantially higher than the life-history-based fixed level of natural mortality traditionally used in the analyses (0.15 year^{-1}). A significant increase in natural mortality of striped bass in Chesapeake Bay may have a considerable effect on population dynamics and serious implications for management. An obvious effect of an increase in M is a faster decay of individual cohort size (increase in the catch curve slope) and overall decline of population abundance. A significant decline in population size should in turn affect fish availability and lead to a decline in CPUE and total harvest. However, the Bay landings reached record harvest values in 2006 and declined only slightly thereafter.

This lack of agreement between model results and observed fishery data suggests a need for careful evaluation of the tagging analysis assumptions (full mixing and equal probability of marked fish to be recovered) and interpretation of the results. What is currently interpreted in the model as total mortality can be more generally described as a rate of disappearance, where disappearance includes total mortality and emigration. Striped bass emigrate from Chesapeake Bay as they age and if the fish are moving to areas that are not fished or very lightly fished (for example, the EEZ) the probability of tagged fish being recovered becomes extremely low. In this case, the decline in the number of recovered tags is interpreted in the model as a decline in survival and increase in natural mortality. A simulation analysis is recommended to investigate the ability of the instantaneous rates model to differentiate natural mortality from emigration to areas with different or no fishing activity/tag returns.

Literature Cited

- Akaike, H. 1973. Information theory as an extension of the maximum likelihood principle. In Second International Symposium on Information Theory. Edited by B. N. Petrov and F. Csaki. Budapest. Akademiai Kiado.
- Akaike, H. 1985. Prediction and entropy. In A Celebration of Statistics. Edited by A.C. Atkinson and S.E. Fienberg. New York: Springer.
- Beverton, R.J.H., and S.J. Holt. 1959. A review of the lifespans and mortality rates of fish in nature, and their relation to growth and other physiological characteristics. Ciba Found. Colloq. Ageing 5:142-177.
- Brownie, C., D.R. Anderson, K.P. Burnham, and D.R. Robson. 1985. Statistical inference from band recovery data: a handbook, 2nd ed., U.S. Fish and Wildl. Serv. Resour. Publ. No. 156.
- Buckland, S.T., K.P. Burnham, and N.H. Augustin. 1997. Model selection: an integral part of inference. Biometrics 53:603-618.
- Burnham, K.P. and D.R. Anderson. 1992. Data-based selection of an appropriate biological model: The key to modern data analysis. In Wildlife 2001: Populations. Edited by D.R. McCullough and R.H. Barrett. London: Elsevier Science Publishers.
- Burnham, K.P. and D.R. Anderson. 1998. Model selection and inference: a practical information theoretical approach. Springer-Verlag, New York.
- Burnham, K.P., G.C. White, and D.R. Anderson. 1995. Model selection strategy in the analysis of capture-recapture data. Biometrics 51:888-898.
- Dunning, D.J., Q.E. Ross, J.F. Waldman, and M.T. Mattson. 1987. Tag retention by, and tagging mortality of, Hudson River striped bass. N. Am J. Fish. Manage. 7:535-538.
- Field, J.D. 1997. Atlantic striped bass management: where did we go right? Fisheries 22(7):6-8.
- Gauthier, DT, RJ Latour, DM Heisey, CF Bonzak, J Gartland, EJ Burge and WK Vogelbein. 2008. Mycobacteriosis-associated mortality in wild striped bass (*Morone saxatilis*) from Chesapeake Bay, USA. Ecol. Appl. 18: 1718-1727.
- Gunderson, D.R., and P.H. Dygert. 1988. Reproductive effort as a predictor of natural mortality rate. J. Cons. int. Explor. Mer 44:200-209.

- Hoening, J.M. 1983. Empirical use of longevity data to estimate mortality rates. Fish. Bull. 81:898-903.
- Hoening, J.M., N.J. Barrowman, W.S. Hearn, and K.H. Pollock. 1998a. Multiyear tagging studies incorporating fishing effort data. Can. J. Fish. Aquat. Sci. 55:1466-1476.
- Hoening, J.M., N.J. Barrowman, K.H. Pollock, E.N. Brooks, W.S. Hearn and T. Polacheck. 1998b. Models for Tagging Data that Allow for Incomplete Mixing of Newly Tagged Animals. Can. J. Fish. Aquat. Sci. 55:1477-1483.
- Hoening, JM, W Vogelbein, M Smith and P Sadler. 2009. The role of mycobacteriosis in elevated natural mortality of Chesapeake Bay striped bass: developing better models for stock assessment and management. Final Report. National Oceanic and Atmospheric Administration Chesapeake Bay Office. 24 pp.
- Jiang, H., K. H. Pollock, C. Brownie, J. M. Hoening, R. J. Latour, B. K. Wells, and J. E. Hightower. 2007. Tag return models allowing for harvest and catch and release: evidence of environmental and management impacts on striped bass fishing and natural mortality rates. North American Journal of Fisheries Management 27:387-396.
- Latour, R.J., K.H. Pollock, C.A. Wenner, and J.M. Hoening. 2001a. Estimates of fishing and natural mortality for red drum (*Sciaenops ocellatus*) in South Carolina waters. N. Am. J.Fish. Manage. 21: 733-744.
- Latour, R.J., J.M. Hoening, J.E. Olney, and K.H. Pollock. 2001b. Diagnostics for multi-year tagging models with application to Atlantic striped bass (*Morone saxatilis*). Can. J. Fish. Aquat. Sci. 5: 1716-1726.
- Latour, R.J., J.M. Hoening and K.H. Pollock. 2001c. Properties of the residuals from two tag-recovery models. Fish. Bull. In review.
- Latour, R.J., J.M. Hoening and K.H. Pollock. 2002. Properties of the residuals from two tag-recovery models. Fish. Bull. In press.
- Pauly, D. 1980. On the interrelationships between natural mortality, growth parameters, and mean environmental temperature in 175 fish stocks. J. Cons. Int. Explor. Mer. 39(2):175-192.
- Ricker, W.E. 1975. Computation and Interpretation of Biological Statistics of Fish Populations. Bull. Fish. Res. Board Can. No 191.

- Roff, D.A. 1984. The evolution of life history parameters in teleosts. *Can. J. Fish. Aquat. Sci.* 41:989-1000.
- Sadler, P.W., R.J. Latour, R.E. Harris, and J.E. Olney. 2001. Evaluation of striped bass stocks in Virginia: monitoring and tagging studies, 1999-2003. Annual Report, Virginia Institute of Marine Science: 93 p.
- Seber, G.A.F. 1970. Estimating time-specific survival and reporting rates for adult birds from band returns. *Biometrika*, 57: 313-318.
- Smith, D.R., K.P. Burnham, D.M. Kahn, X. He, C.J. Goshorn, K.A. Hattala, and A.W. Kahnle. 2000. Bias in survival estimates from tag-recovery models where catch-and-release is common, with an example from Atlantic striped bass (*Morone saxatilis*). *Can. J. Fish. Aquat. Sci.* 57:886-897.
- Vogelbein, W.K., D.E. Zwerner, H. Kator, M.W. Rhodes and J. Cardinal. 1999. Mycobacteriosis of striped bass from Chesapeake Bay. pages 53-58. In J.E. Olney (ed.), *Research on Recreational Fishes and Fisheries*, VIMS Spec. Sci. Rept. 139, 82 pp.
- Weaver, J.E., R.B. Fairbanks and C. M. Wooley. 1986. Interstate management of Atlantic coastal migratory striped bass. *MRFSS* 11: 71-85.
- White, G.C. 1983. Numerical estimation of survival rates from band-recovery and biotelemetry data. *J. Wildl. Manage.* 47:716-728.
- White, G.C. and K. P. Burnham. 1999. Program MARK survival estimation from populations of marked animals. *Bird Study* 46:120-138.
- Wooley, C.M., N.C. Parker, B.M. Florence and R.W. Miller. 1990. Striped bass restoration along the Atlantic Coast: a multistate and federal cooperative hatchery and tagging program.

Table 1. Summary data of striped bass tagged and released from pound nets in the Rappahannock River, spring 2011.

Date	N	457-710 mm TL						> 711 mm TL			
		Males		Females		Unknown		Males		Females	
		n	\overline{TL}	n	\overline{TL}		\overline{TL}		\overline{TL}		\overline{TL}
11 Apr	69	24	551.7	7	614.9	0		21	807.2	17	930.6
14 Apr	80	29	593.2	3	577.0	0		18	790.3	30	951.1
18 Apr	92	31	574.4	8	560.3	0		21	797.4	32	912.5
21 Apr	17	4	542.5	3	608.7	0		2	808.5	8	912.8
25 Apr	30	10	595.1	4	567.5	0		2	815.0	14	885.9
28 Apr	66	40	564.8	6	584.3	0		6	801.3	14	891.6
2 May	32	26	591.4	1	620.0	0		0		5	876.6
9 May	30	27	539.0	2	600.0	0		0		1	870.0

Table 2. Location of striped bass (≥ 457 mm TL), recaptured in 2010, that were originally tagged and released in the Rappahannock River during springs 1990-2010.

State	Month												total
	J	F	M	A	M	J	J	A	S	O	N	D	
New Hampshire	0	0	0	0	0	0	0	0	0	0	0	0	0
Massachusetts	0	0	0	0	0	3	6	2	0	0	0	0	11
Rhode Island	0	0	0	0	0	0	1	1	0	0	0	0	2
Connecticut	0	0	0	0	0	0	0	0	0	0	0	0	0
New York	0	0	0	0	0	1	1	0	0	1	1	0	4
New Jersey	0	0	0	0	4	1	0	0	0	0	3	0	8
Delaware	0	0	0	0	3	0	0	0	0	0	0	0	3
Maryland	0	0	0	2	0	4	1	3	1	1	0	0	12
Virginia	0	0	0	1	4	8	2	3	2	4	1	6	31
North Carolina	1	2	0	0	0	0	0	0	0	0	0	0	3
Total	1	2	0	3	11	17	11	9	3	6	5	6	74

Table 3. Location of striped bass (≥ 711 mm TL), recaptured in 2010, that were originally tagged and released in the Rappahannock River during springs 1990-2010.

State	Month												total
	J	F	M	A	M	J	J	A	S	O	N	D	
Massachusetts	0	0	0	0	0	3	6	2	0	0	0	0	11
Rhode Island	0	0	0	0	0	0	0	1	1	0	0	0	2
Connecticut	0	0	0	0	0	0	0	0	0	0	0	0	0
New York	0	0	0	0	0	1	1	0	0	1	0	0	3
New Jersey	0	0	0	0	4	1	0	0	0	0	3	0	8
Delaware	0	0	0	0	3	0	0	0	0	0	0	0	3
Maryland	0	0	0	2	0	0	0	0	1	0	0	0	3
Virginia	0	0	0	0	1	0	0	0	0	0	1	0	2
North Carolina	1	2	0	0	0	0	0	0	0	0	0	0	3
Total	1	2	0	2	8	5	7	3	2	1	4	0	35

Table 4. Input recapture matrix for program MARK: from striped bass (>457 mm TL) that were tagged and released in the Rappahannock River, springs 1990-2010.

Release		Recapture year																				
Year		90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10
1,464	1990	162	64	47	25	12	10	3	2	3	1	1	0	0	1	0	0	0	0	0	0	0
2,481	1991		167	81	53	29	6	5	2	2	4	1	0	0	1	0	0	0	0	0	0	0
130	1992			14	8	6	5	1	1	1	1	0	0	0	0	0	0	0	0	0	0	0
621	1993				50	37	17	8	9	0	0	1	0	0	0	0	0	0	0	0	0	0
195	1994					13	10	5	4	4	0	0	0	0	0	0	0	0	0	0	0	0
698	1995						55	30	20	5	4	2	3	0	1	0	1	0	0	0	0	0
376	1996							21	18	7	3	1	1	1	0	0	1	0	0	0	0	0
712	1997								47	26	14	3	0	1	2	1	0	0	0	0	0	0
784	1998									55	26	2	3	3	1	0	0	0	0	0	0	0
853	1999										66	23	9	5	3	0	0	0	0	0	0	1
1,765	2000											122	51	23	16	6	5	1	1	0	0	0
797	2001												61	23	16	7	2	2	2	0	0	0
315	2002													20	8	15	1	1	2	1	0	0
852	2003														58	37	9	4	5	3	2	3
1,477	2004															80	21	13	7	4	2	1
921	2005																44	26	10	2	5	4
668	2006																	49	11	6	6	3
1,961	2007																		117	50	24	4
523	2008																			30	9	2
867	2009																				43	10
2050	2010																					47

Table 5. Input recapture matrix for program MARK: from striped bass (>710 mm TL) that were tagged and released in the Rappahannock River, springs 1990-2010.

Release		Recapture year																				
Year		90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10
301	1990	26	9	15	2	4	6	1	0	2	1	1	0	0	1	0	0	0	0	0	0	0
390	1991		41	24	16	11	3	2	2	1	2	0	0	0	1	0	0	0	0	0	0	0
40	1992			4	3	2	2	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
212	1993				22	18	7	4	7	0	0	1	0	0	0	0	0	0	0	0	0	0
123	1994					9	7	5	1	2	0	0	0	0	0	0	0	0	0	0	0	0
210	1995						29	11	8	3	3	2	3	0	1	0	1	0	0	0	0	0
67	1996							1	3	1	0	0	1	0	0	0	0	0	0	0	0	0
212	1997								15	13	8	3	0	1	2	1	0	0	0	0	0	0
158	1998									24	13	2	3	2	0	0	0	0	0	0	0	0
162	1999										17	6	2	3	2	0	0	0	0	0	0	1
365	2000											28	19	14	9	4	3	0	1	0	0	0
269	2001												19	14	4	6	2	1	1	0	0	0
122	2002													10	6	7	1	0	2	1	0	0
400	2003														35	24	7	1	3	3	2	3
686	2004															39	12	13	5	4	2	1
284	2005																16	11	8	1	4	3
175	2006																	13	4	4	3	1
840	2007																		55	30	18	3
75	2008																			6	2	0
241	2009																				7	5
483	2010																					17

Table 6. Performance statistics (>457 mm TL), based on quasi-likelihood Akaike Information Criteria (QAIC), used to assess the Seber (1970) models utilized in the ASMFC analysis protocol. Model notations: S (f) and r (f) indicate that survival (S) and tag-reporting rate (r) are functions (f) of the factors within the parenthesis; constant parameters across time (.); parameters constant from 1990-1994, 1995-1999, 2000-2002, and 2003-2010 (p); parameters vary in 2010 (d), otherwise the same as p; parameters vary in 2009-2010 (v), otherwise the same as p; and parameters are time-specific (t).

Model	$QAIC_c$	$\Delta QAIC_c$	$QAIC_c$ weight	number of parameters
S(t)r(t)	11,265.68	0.00	0.96715	41
S(d)r(p)	11,272.80	7.12	0.02749	9
S(.)r(t)	11,276.96	11.28	0.00344	22
S(p)r(t)	11,278.45	12.77	0.00163	25
S(t)r(p)	11,282.41	36.73	0.00023	25
S(v)r(p)	11,285.36	19.68	0.00005	9
S(.)r(p)	11,296.65	30.97	0.00000	5
S(p)r(p)	11,298.23	32.55	0.00000	8
S(.)r(.)	11,380.67	114.99	0.00000	2

Table 7. Seber (1970) model estimates of unadjusted survival (\hat{S}) rates and adjusted rates of survival (\hat{S}_{adj}) and fishing mortality (\hat{F}) of striped bass (> 457 mm TL) derived from the proportion of recaptures released alive (P_l) in the Rappahannock River, 1990-2010.

Year	\hat{S}	SE (\hat{S})	P_l	Bias	\hat{S}_{adj}	\hat{F}	95% CI \hat{F}
1990	0.807	0.096	0.481	-0.143	0.942	-0.090	-0.24, 0.35
1991	0.286	0.057	0.524	-0.082	0.311	1.018	0.55, 1.59
1992	0.796	0.180	0.408	-0.142	0.928	-0.075	-0.28, 0.94
1993	0.603	0.143	0.456	-0.105	0.673	0.245	-0.08, 0.89
1994	0.568	0.139	0.381	-0.087	0.622	0.324	-0.02, 0.97
1995	0.679	0.149	0.262	-0.054	0.717	0.182	-0.09, 0.85
1996	0.636	0.146	0.274	-0.040	0.662	0.262	-0.04, 0.92
1997	0.565	0.118	0.330	-0.057	0.600	0.362	0.05, 0.89
1998	0.413	0.086	0.362	-0.059	0.439	0.673	0.31, 1.15
1999	0.379	0.072	0.243	-0.059	0.391	0.759	0.41, 1.20
2000	0.431	0.071	0.436	-0.074	0.466	0.614	0.32, 0.99
2001	0.465	0.106	0.367	-0.068	0.499	0.545	0.18, 1.08
2002	0.604	0.139	0.368	-0.063	0.645	0.289	-0.03, 0.91
2003	0.831	0.153	0.271	-0.049	0.874	-0.015	-0.18, 0.92
2004	0.351	0.071	0.281	-0.038	0.365	0.859	0.48, 1.33
2005	0.483	0.104	0.280	-0.031	0.499	0.546	0.20, 1.06
2006	0.516	0.105	0.358	-0.055	0.546	0.454	0.13, 0.94
2007	0.554	0.132	0.305	-0.043	0.574	0.396	0.05, 1.01
2008	0.521	0.155	0.208	-0.025	0.535	0.475	0.06, 1.24
2009	0.623	0.215	0.231	-0.027	0.641	0.295	-0.08, 1.38
2010	0.731	0.036	0.267	-0.015	0.742	0.148	-0.03, 0.50

Table 8. Performance statistics (>710 mm TL), based on quasi-likelihood Akaike Information Criteria (QAIC), used to assess the Seber (1970) models utilized in the ASMFC analysis protocol. Model notations: S (f) and r (f) indicate that survival (S) and tag-reporting rate (r) are functions (f) of the factors within the parenthesis; constant parameters across time (.); parameters constant from 1990-1994, 1995-1999, 2000-2002, and 2003-2010 (p); parameters vary in 2010 (d), otherwise the same as p; parameters vary in 2009 and 2010 (v), otherwise the same as p; and parameters are time-specific (t).

Model	QAIC_c	Δ QAIC_c	QAIC_c weight	number of parameters
S(d)r(p)	7,172.92	0.00	0.64752	9
S(v)r(p)	7,175.04	2.12	0.22440	9
S(.)r(t)	7,177.77	4.85	0.05738	22
S(p)r(p)	7,178.55	5.63	0.03884	8
S(.)r(p)	7,179.31	6.39	0.02651	5
S(p)r(t)	7,183.26	10.34	0.00369	25
S(t)r(t)	7,186.16	13.24	0.00086	41
S(t)r(p)	7,186.33	13.40	0.00080	25
S(.)r(.)	7,241.77	68.84	0.00000	2

Table 9. Seber (1970) model estimates (SBTC) of unadjusted survival (\hat{S}) rates and adjusted rates of survival (\hat{S}_{adj}) and fishing mortality (\hat{F}) of striped bass (> 710 mm TL) derived from the proportion of recaptures released alive (P_l) in the Rappahannock River, 1990-2010.

Year	\hat{S}	SE (\hat{S})	P_l	Bias	\hat{S}_{adj}	\hat{F}	95% CI
1990	0.629	0.024	0.577	-0.127	0.720	0.178	0.10, 0.26
1991	0.629	0.024	0.560	-0.131	0.724	0.173	0.10, 0.26
1992	0.629	0.025	0.535	-0.172	0.760	0.124	0.05, 0.21
1993	0.629	0.024	0.349	-0.093	0.694	0.216	0.14, 0.30
1994	0.629	0.024	0.318	-0.070	0.677	0.241	0.17, 0.33
1995	0.587	0.027	0.204	-0.079	0.638	0.300	0.21, 0.41
1996	0.587	0.027	0.125	-0.016	0.596	0.367	0.28, 0.47
1997	0.587	0.027	0.167	-0.036	0.609	0.346	0.26, 0.45
1998	0.587	0.027	0.217	-0.084	0.641	0.294	0.20, 0.40
1999	0.587	0.027	0.200	-0.058	0.623	0.323	0.23, 0.43
2000	0.671	0.036	0.349	-0.072	0.723	0.175	0.07, 0.31
2001	0.671	0.036	0.298	-0.052	0.708	0.196	0.09, 0.33
2002	0.671	0.036	0.295	-0.078	0.727	0.168	0.06, 0.30
2003	0.585	0.023	0.246	-0.059	0.622	0.324	0.24, 0.42
2004	0.585	0.023	0.321	-0.049	0.615	0.336	0.26, 0.43
2005	0.585	0.023	0.238	-0.035	0.606	0.350	0.27, 0.44
2006	0.585	0.023	0.282	-0.042	0.611	0.342	0.26, 0.43
2007	0.585	0.023	0.228	-0.036	0.607	0.349	0.27, 0.44
2008	0.585	0.023	0.163	-0.024	0.600	0.361	0.28, 0.46
2009	0.613	0.029	0.105	-0.009	0.619	0.330	0.16, 0.56
2010	0.717	0.047	0.235	-0.021	0.732	0.162	0.02, 0.39

Table 10. Estimates of total mortality (Z), annual mortality (A), exploitation (U), fishing mortality (F) and natural mortality (M) from striped bass (> 457 mm TL) tagged and released in the Rappahannock River, springs, 1990-2010.

Year	Z	A	U	F	M
1990	0.06	0.06	0.17	0.18	-0.12
1991	1.17	0.69	0.14	0.23	0.93
1992	0.07	0.07	0.32	0.34	-0.26
1993	0.40	0.33	0.23	0.28	0.11
1994	0.47	0.38	0.26	0.33	0.15
1995	0.33	0.28	0.19	0.22	0.11
1996	0.41	0.34	0.15	0.18	0.23
1997	0.51	0.40	0.20	0.25	0.26
1998	0.82	0.56	0.15	0.22	0.60
1999	0.91	0.60	0.13	0.20	0.71
2000	0.76	0.53	0.14	0.19	0.57
2001	0.69	0.50	0.18	0.25	0.44
2002	0.44	0.36	0.19	0.23	0.21
2003	0.14	0.13	0.19	0.21	-0.07
2004	1.01	0.64	0.12	0.20	0.81
2005	0.70	0.50	0.15	0.20	0.49
2006	0.60	0.45	0.16	0.21	0.39
2007	0.55	0.42	0.14	0.19	0.36
2008	0.63	0.47	0.10	0.13	0.49
2009	0.44	0.36	0.09	0.11	0.33
2010	0.30	0.26	0.04	0.04	0.26

Table 11. Estimates of total mortality (Z), annual mortality (A), exploitation (U), fishing mortality (F) and natural mortality (M) from striped bass (> 710 mm TL) tagged and released in the Rappahannock River, springs, 1990-2010.

Year	Z	A	U	F	M
1990	0.33	0.28	0.25	0.30	0.03
1991	0.32	0.28	0.36	0.43	-0.10
1992	0.27	0.24	0.37	0.42	-0.15
1993	0.37	0.31	0.37	0.44	-0.07
1994	0.39	0.32	0.25	0.31	0.08
1995	0.45	0.36	0.41	0.51	-0.06
1996	0.52	0.40	0.18	0.23	0.29
1997	0.50	0.39	0.38	0.48	0.02
1998	0.44	0.36	0.45	0.56	-0.12
1999	0.47	0.38	0.30	0.37	0.10
2000	0.32	0.28	0.28	0.33	0.00
2001	0.35	0.29	0.24	0.28	0.07
2002	0.32	0.27	0.35	0.41	-0.09
2003	0.47	0.38	0.28	0.36	0.12
2004	0.49	0.38	0.16	0.20	0.28
2005	0.50	0.39	0.24	0.30	0.20
2006	0.49	0.39	0.26	0.33	0.17
2007	0.50	0.39	0.19	0.25	0.25
2008	0.51	0.40	0.19	0.24	0.27
2009	0.48	0.38	0.08	0.10	0.38
2010	0.31	0.27	0.05	0.06	0.25

Table 12a. Input recapture matrix for IRCR analysis: from striped bass (>457 mm TL) tagged and released in the Rappahannock River, springs 1990-2010. Harvested recaptures only.

Release		Recapture year																				
Year		90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10
1,433	1990	21	20	24	10	8	9	2	0	0	1	1	0	0	1	0	0	0	0	0	0	0
2,457	1991		48	38	22	14	3	1	2	1	4	0	0	0	1	0	0	0	0	0	0	0
128	1992			7	4	1	3	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
617	1993				18	17	12	5	4	1	0	0	0	0	0	0	0	0	0	0	0	0
193	1994					6	7	4	1	2	0	0	0	0	0	0	0	0	0	0	0	0
674	1995						24	12	9	4	1	1	2	0	1	0	0	0	0	0	0	0
374	1996							3	10	3	2	1	1	1	0	0	1	0	0	0	0	0
703	1997								26	17	10	2	0	1	1	1	0	0	0	0	0	0
777	1998									28	16	1	3	1	0	0	0	0	0	0	0	0
844	1999										30	7	4	2	2	0	0	0	0	0	0	0
1,736	2000											44	23	11	7	4	5	1	1	0	0	0
784	2001												32	14	5	7	1	0	0	0	0	0
310	2002													10	4	6	1	1	1	1	0	0
839	2003														32	20	5	3	3	2	1	2
1,470	2004															45	14	8	4	3	1	1
916	2005																27	17	6	1	4	1
662	2006																	27	4	5	5	3
1,952	2007																		64	34	16	3
523	2008																			17	4	0
865	2009																				26	7
2047	2010																					30

Table 12b. Input recapture matrix for IRCR analysis: from striped bass (>457 mm TL) that were tagged and released in the Rappahannock River, springs 1990-2010. Recaptures released with streamers cut off only.

Release		Recapture year																				
Year		90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10
1,433	1990	60	22	15	8	1	1	0	1	2	0	0	0	0	0	0	0	0	0	0	0	0
2,457	1991		86	26	18	10	2	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
128	1992			4	3	3	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0
617	1993				26	16	3	1	0	1	0	0	0	0	0	0	0	0	0	0	0	0
193	1994					5	1	0	2	1	0	0	0	0	0	0	0	0	0	0	0	0
674	1995						14	7	7	1	0	0	0	0	0	0	0	0	0	0	0	0
374	1996							9	5	3	0	0	0	0	0	0	0	0	0	0	0	0
703	1997								9	6	2	0	0	0	0	0	0	0	0	0	0	0
777	1998									21	5	0	0	1	1	0	0	0	0	0	0	0
844	1999										19	7	1	2	0	0	0	0	0	0	0	0
1,736	2000											40	18	6	3	0	0	0	0	0	0	0
784	2001												17	6	3	0	0	0	1	0	0	0
310	2002													6	3	2	0	0	1	0	0	0
839	2003														11	9	2	1	1	0	0	0
1,470	2004															20	5	5	1	0	1	0
916	2005																12	8	2	1	0	1
662	2006																	16	5	1	1	0
1,952	2007																		31	7	1	0
523	2008																			7	2	2
865	2009																				14	2
2047	2010																					11

Table 13a. Input recapture matrix for IRCR analysis: from striped bass (>710 mm TL) that were tagged and released in the Rappahannock River, springs 1990-2010. Harvested recaptures only.

Release		Recapture year																				
Year		90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10
297	1990	10	1	6	1	3	5	1	0	0	1	1	0	0	1	0	0	0	0	0	0	0
386	1991		19	10	12	9	2	1	2	0	2	0	0	0	1	0	0	0	0	0	0	0
40	1992			2	1	1	1	0	0	0	1	0	0	0	0	0	0	0	0	0	0	0
209	1993				11	11	5	2	3	0	0	0	0	0	0	0	0	0	0	0	0	0
123	1994					4	4	4	1	0	0	0	0	0	0	0	0	0	0	0	0	0
205	1995						18	6	5	2	1	1	2	0	1	0	0	0	0	0	0	0
67	1996							0	3	1	0	0	1	0	0	0	0	0	0	0	0	0
210	1997								11	12	6	2	0	1	1	1	0	0	0	0	0	0
156	1998									16	9	1	3	1	0	0	0	0	0	0	0	0
159	1999										13	2	1	2	1	0	0	0	0	0	0	1
362	2000											13	11	6	5	3	3	0	1	0	0	0
268	2001												9	8	2	6	1	0	0	0	0	0
122	2002													7	3	5	1	0	1	1	0	0
392	2003														23	13	3	1	2	2	1	2
680	2004															21	8	8	3	3	1	1
281	2005																12	7	5	1	3	0
175	2006																	10	3	3	2	1
835	2007																		33	22	11	2
75	2008																			5	1	0
241	2009																				5	3
482	2010																					11

Table 13b. Input recapture matrix for IRCR analysis: from striped bass (>710 mm TL) that were tagged and released in the Rappahannock River, springs 1990-2010. Recaptures released with streamers cut off only.

Release		Recapture year																				
Year		90	91	92	93	94	95	96	97	98	99	00	01	02	03	04	05	06	07	08	09	10
297	1990	14	6	7	0	0	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
386	1991		19	10	4	2	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
40	1992			2	1	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
209	1993				10	7	1	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
123	1994					4	1	0	0	1	0	0	0	0	0	0	0	0	0	0	0	0
205	1995						5	2	2	1	0	0	0	0	0	0	0	0	0	0	0	0
67	1996							1	0	0	0	0	0	0	0	0	0	0	0	0	0	0
210	1997								2	1	1	0	0	0	0	0	0	0	0	0	0	0
156	1998									6	3	0	0	1	0	0	0	0	0	0	0	0
159	1999										2	1	0	1	0	0	0	0	0	0	0	0
362	2000											9	6	4	2	0	0	0	0	0	0	0
268	2001												7	4	2	0	0	0	1	0	0	0
122	2002													2	2	0	0	0	1	0	0	0
392	2003														8	6	2	0	0	0	0	0
680	2004															11	2	5	1	0	1	0
281	2005																3	4	1	0	0	1
175	2006																	2	1	1	1	0
835	2007																		11	5	1	0
75	2008																			0	0	0
241	2009																				1	1
482	2010																					4

Table 14. Model Akaike weighting results (striped bass ≥ 457 mm TL) for the 1M (constant) IRCR analysis. Model notations: Fishing mortality (F), release mortality (F') and natural mortality (M), annual estimates (t) and period estimates (4p- 1990-1994, 1995-1999, 2000-2002 and 2003-2010; d- 1990-1994, 1995-1999, 2000-2002, 2003-2009 and 2010; v- 1990-1994, 1995-1999, 2000-2002, 2003-2008 and 2009-2010).

model	weight
F(t), F'(t), 1M	0.000
F(4p), F'(t), 1M	0.000
F(.), F'(t), 1M	0.000
F(t), F'(4p), 1M	0.030
F(t), F'(.), 1M	0.000
F(4p), F'(4p), 1M	0.000
F(.), F'(.), 1M	0.000
F(d), F'(d), 1M	0.949
F(v), F'(v), 1M	0.021

Table 15. Model Akaike weighting results (striped bass ≥ 457 mm TL) for the 1M (constant) and 2M IRCR analyses. Model notations: Fishing mortality (F), release mortality (F') and natural mortality (M), annual estimates (t) and period estimates (4p- 1990-1994, 1995-1999, 2000-2002 and 2003-2010; d- 1990-1994, 1995-1999, 2000-2002, 2003-2009 and 2010; v- 1990-1994, 1995-1999, 2000-2002, 2003-2008 and 2009-2010).

2M (1990-1997, 1998-2010)	
model	weight
F(t), F'(t), 1M	0.000
F(4p),F'(t), 1M	0.000
F(.), F'(t), 1M	0.000
F(t), F'(4p), 1M	0.000
F(t), F'(.), 1M	0.000
F(4p), F'(4p), 1M	0.000
F(.), F'(.), 1M	0.000
F(d), F'(d), 1M	0.000
F(v), F'(v), M	0.000
F(t), F'(t), 2M	0.000
F(4p),F'(t), 2M	0.000
F(.), F'(t), 2M	0.000
F(t), F'(4p), 2M	0.007
F(t), F'(.), 2M	0.000
F(4p), F'(4p), 2M	0.000
F(.), F'(.), 2M	0.000
F(d), F'(d), 2M	0.984
F(v), F'(v), 2M	0.009

Table 16. Parameter estimates of survival (S), natural mortality (M), fishing mortality (F) and its standard error (SE) for striped bass ≥ 457 mm TL from the IRCR analyses (1M and 2M), 1990-2010.

Year	1M				2M			
	S	M	F	SE	S	M	F	SE
1990	0.567	0.473	0.087	0.011	0.616	0.395	0.082	0.008
1991	0.567	0.473	0.087	0.008	0.616	0.395	0.082	0.007
1992	0.566	0.473	0.089	0.010	0.616	0.395	0.083	0.008
1993	0.566	0.473	0.088	0.009	0.616	0.395	0.082	0.007
1994	0.566	0.473	0.090	0.013	0.616	0.395	0.083	0.008
1995	0.545	0.473	0.130	0.013	0.596	0.395	0.119	0.011
1996	0.546	0.473	0.128	0.014	0.596	0.395	0.118	0.012
1997	0.545	0.473	0.129	0.012	0.596	0.395	0.118	0.011
1998	0.545	0.473	0.129	0.012	0.505	0.560	0.118	0.011
1999	0.545	0.473	0.129	0.012	0.505	0.560	0.119	0.011
2000	0.571	0.473	0.083	0.009	0.519	0.560	0.090	0.010
2001	0.571	0.473	0.084	0.010	0.519	0.560	0.091	0.010
2002	0.571	0.473	0.083	0.009	0.519	0.560	0.091	0.010
2003	0.566	0.473	0.094	0.008	0.512	0.560	0.106	0.008
2004	0.566	0.473	0.094	0.008	0.512	0.560	0.106	0.008
2005	0.566	0.473	0.093	0.007	0.512	0.560	0.106	0.008
2006	0.566	0.473	0.093	0.007	0.512	0.560	0.106	0.008
2007	0.566	0.473	0.093	0.007	0.512	0.560	0.106	0.008
2008	0.566	0.473	0.093	0.007	0.512	0.560	0.106	0.008
2009	0.567	0.473	0.093	0.009	0.512	0.560	0.106	0.009
2010	0.594	0.473	0.046	0.009	0.542	0.560	0.052	0.010

Table 17. Model Akaike weighting results (striped bass ≥ 711 mm TL) for the 1M (constant) IRCR analysis. Model notations: Fishing mortality (F), release mortality (F') and natural mortality (M), annual estimates (t) and period estimates (4p- 1990-1994, 1995-1999, 2000-2002 and 2003-2010; d- 1990-1994, 1995-1999, 2000-2002, 2003-2009 and 2010; v- 1990-1994, 1995-1999, 2000-2002, 2003-2008 and 2009-2010).

model	weight
F(t), F'(t), 1M	0.000
F(4p), F'(t), 1M	0.000
F(.), F'(t), 1M	0.000
F(t), F'(4p), 1M	0.003
F(t), F'(.), 1M	0.000
F(4p), F'(4p), 1M	0.001
F(.), F'(.), 1M	0.000
F(d), F'(d), 1M	0.045
F(v), F'(v), 1M	0.951

Table 18. Model Akaike weighting results (striped bass ≥ 711 mm TL) for the 1M (constant) and 2M IRCR analyses. Model notations: Fishing mortality (F), release mortality (F') and natural mortality (M), annual estimates (t) and period estimates (4p- 1990-1994, 1995-1999, 2000-2002 and 2003-2010; d- 1990-1994, 1995-1999, 2000-2002, 2003-2009 and 2010; v- 1990-1994, 1995-1999, 2000-2002, 2003-2008 and 2009-2010).

2M (1990-2003, 2004-2010)	
model	weight
F(t), F'(t), 1M	0.000
F(4p), F'(t), 1M	0.000
F(.), F'(t), 1M	0.000
F(t), F'(4p), 1M	0.000
F(t), F'(.), 1M	0.000
F(4p), F'(4p), 1M	0.000
F(.), F'(.), 1M	0.000
F(d), F'(d), 1M	0.000
F(v), F'(v), 1M	0.001
F(t), F'(t), 2M	0.000
F(4p), F'(t), 2M	0.000
F(.), F'(t), 2M	0.000
F(t), F'(4p), 2M	0.001
F(t), F'(.), 2M	0.000
F(4p), F'(4p), 2M	0.080
F(.), F'(.), 2M	0.000
F(d), F'(d), 2M	0.346
F(v), F'(v), 2M	0.571

Table 19. Parameter estimates of survival (S), natural mortality (M), fishing mortality (F) and its standard error (SE) for striped bass ≥ 711 mm TL from the IRCR analyses (1M and 2M), 1990-2010.

Year	1M				2M			
	S	M	F	SE	S	M	F	SE
1990	0.646	0.287	0.141	0.016	0.671	0.249	0.141	0.015
1991	0.646	0.287	0.141	0.016	0.671	0.249	0.141	0.015
1992	0.646	0.287	0.141	0.015	0.671	0.249	0.141	0.015
1993	0.646	0.287	0.141	0.015	0.671	0.249	0.141	0.015
1994	0.646	0.287	0.141	0.016	0.671	0.249	0.141	0.016
1995	0.597	0.287	0.225	0.023	0.625	0.249	0.217	0.022
1996	0.597	0.287	0.225	0.023	0.625	0.249	0.217	0.023
1997	0.597	0.287	0.225	0.023	0.625	0.249	0.217	0.022
1998	0.597	0.287	0.225	0.023	0.625	0.249	0.217	0.022
1999	0.597	0.287	0.225	0.023	0.625	0.249	0.217	0.023
2000	0.663	0.287	0.120	0.016	0.692	0.249	0.114	0.015
2001	0.662	0.287	0.120	0.016	0.692	0.249	0.114	0.015
2002	0.662	0.287	0.121	0.016	0.692	0.249	0.114	0.015
2003	0.670	0.287	0.110	0.010	0.687	0.249	0.123	0.010
2004	0.670	0.287	0.110	0.009	0.575	0.428	0.123	0.010
2005	0.671	0.287	0.110	0.009	0.575	0.428	0.123	0.010
2006	0.671	0.287	0.110	0.009	0.575	0.428	0.123	0.010
2007	0.671	0.287	0.110	0.009	0.575	0.428	0.123	0.010
2008	0.670	0.287	0.110	0.009	0.575	0.428	0.123	0.010
2009	0.702	0.287	0.066	0.014	0.587	0.428	0.103	0.022
2010	0.702	0.287	0.064	0.011	0.598	0.428	0.085	0.020

III. The role of mycobacteriosis in elevated natural mortality of Chesapeake Bay striped bass: disease progression and developing better models for stock assessment and management.

Striped Bass Assessment and Monitoring Program
Department of Fisheries Science
School of Marine Science
Virginia Institute of Marine Science
The College of William and Mary
Gloucester Point, VA. 23062-1346

Introduction

During the late 1990s concern emerged among recreational and commercial fishermen about perceived declining conditions in striped bass (*Morone saxatilis*). Emaciation and ulcerative skin lesions were commonly reported and associated with a bacterial disease called mycobacteriosis. The disease is now epizootic throughout the Bay with more than 70% of striped bass in some tributaries affected. Several hypotheses have been presented to explain this emerging problem. These include stress associated with the loss of prey through recent declines in menhaden stocks (starvation), overcrowding, and loss of summer thermal refuges as a result of hypoxia and high water temperature. Recent tag-recapture analyses indicate that striped bass survival has declined significantly (~20%) over the last 10 to 15 years. This troubling decline is attributable to an increase in natural mortality and corresponds roughly with the Bay-wide outbreak of mycobacteriosis in striped bass. Current fishery management strategies do not account for changes in natural mortality over time, especially during infectious disease epizootics. Thus, the overall aim of the current study is to determine the contribution of mycobacteriosis to natural mortality in the striped bass, and thus the potential for adverse impacts by the disease on the stock.

Mycobacteriosis in fish is a chronic disease caused by various species of bacteria in the genus *Mycobacterium*. Mycobacterial disease occurs in a wide range of species of fish worldwide and is an important problem in aquacultural operations. The disease appears as grey granulomatous nodules in internal organs, especially the spleen and kidney (Figure 1b), and can also manifest itself as ulcerous skin lesions (Figure 1a). Fish with ulcerous dermal lesions in the wild sometimes have an extremely emaciated appearance.

Mycobacteriosis was first reported from Chesapeake Bay striped bass in 1997 (Vogelbein et al. 1999; Rhodes et al. 2002, 2003, 2004). Since then, the disease has spread throughout the Bay and the prevalence has risen to as high as 70 – 80% (Cardinal 2001; Vogelbein et al. 1999; this project, unpublished observations). Several species of *Mycobacterium* have been isolated from Chesapeake Bay striped bass, including several new species, but it is not yet clear which species are involved in disease processes. One recently named species, *M. shottsii*, has been observed in splenic tissues of infected striped bass at a prevalence of 50 to 70% greater than other *Mycobacterium* species (Rhodes et al. 2004, Gauthier et al. 2003). Indeed, there may be more than one pathogenic species.

Mycobacteria are slow-growing, aerobic bacteria common in terrestrial and aquatic habitats. Most are saprophytes, but certain species infect both endo- and poikilothermic animals. Mycobacterial infections are common in wild and captive fish stocks world-wide. Mycobacteriosis in fishes is a chronic, systemic disease that can result in degradation of body condition and ultimately in death (Colorni 1992). Clinical signs are nonspecific and may include scale loss, skin ulceration, emaciation, exophthalmia, pigmentation changes and spinal defects (Nigrelli & Vogel 1963; Bruno et al. 1998). Granulomatous inflammation, a host cellular response comprised largely of phagocytic cells of the immune system called macrophages, is a characteristic of the disease. In an attempt to sequester, kill and degrade mycobacteria, these

macrophages encapsulate bacteria, forming nodular structures called granulomas. Skin ulceration in most fishes is uncommon and usually represents the endstage of the disease process, as captive fish with skin lesions generally do not recover and die quickly. Hence, the presence of skin lesions is particularly alarming, as it may indicate that the fish are progressing from chronic, covert infection to active, lethal disease.

The impact of the disease on the population ecology of striped bass is poorly understood. Fundamental questions, such as mode of transmission, duration of disease stages, effects of disease on fish movements, feeding and reproduction, and mortality rates associated with disease, remain unanswered. Nonetheless, there are indications the disease may be having a significant impact on Chesapeake Bay striped bass populations. Jiang et al. (2007) analyzed striped bass tagging data from Maryland and found a significant increase in natural mortality rate at about the time when mycobacteriosis was first being detected in Chesapeake Bay striped bass. A similar analysis of Rappahannock River, Virginia, striped bass tagging data from this project also reveals an increase in natural mortality rate in recent years (see Table 1): natural mortality rate for fish age two and above was estimated to increase from $M = .231$ during the period 1990 – 1996 to $M = .407$ during the period 1997-2004. In addition, R. Latour and D. Gauthier used force-of-infection models to examine the epizootiology of mycobacteriosis in Chesapeake Bay striped bass from 2003-2005. The results of this analysis indicated that the probability a disease negative fish becomes disease positive depends on age; the inclusion of sex and season as covariates significantly improved model fit; and that there is evidence of disease associated mortality (Gauthier et al. 2008).

Mycobacteriosis in fishes is generally thought to be fatal, but this has not been established for wild striped bass. Three possible distinct disease outcomes in the case of striped bass are: 1) death, 2) recovery or reversion to a non-disease state, or 3) movement of infected fish to another location. Because of the uncertainty about the fate of the infected fish, the impact of the disease on striped bass populations is unknown. If mycobacteriosis in striped bass is ultimately fatal, the potential for significant impacts on the productivity and the quality of the Atlantic coastal migratory stock is high. Researchers, fisheries managers and commercial and recreational fishermen are therefore becoming gravely concerned. At a recent symposium entitled “*Management Issues of the Restored Stock of Striped Bass in the Chesapeake Bay: Diseases, Nutrition, Forage Base and Survival*”, Kahn (2004) reported that both Maryland and Virginia striped bass tag-recaptures have declined in recent years. This suggests that survival has declined significantly, from 60-70% in the early-mid 1990’s to 40-50% during the late 1990’s and early 2000’s. Kahn (2004) and Crecco (2003) both concluded that the 20% decline in striped bass survival was not caused by fishing mortality, but rather, by an increase in natural mortality. These analyses, however, are predicated on the assumption that tag reporting rate has not changed over time. No data are currently available to evaluate this assumption. Hypotheses presented at the Symposium to explain the decline in striped bass survival included the possible role of mycobacteriosis (May et al., 2004; Vogelbein et al., 2004). However, Jacobs et al. (2004) found that decline in striped bass nutritional status during the fall was independent of disease. Uphoff (2004) reported that abundance of forage-sized menhaden, a primary food source of striped bass, declined to near historic lows during the mid 1990’s. Similar studies indicated that

as the striped bass population has increased during the 1990's, predatory demand increased coincident with a decline in menhaden populations (Hartman, 2004; Garrison et al., 2004).

Striped bass are presently managed by attempting to control fishing mortality. Fishing mortality is determined in three ways, and each method uses a value for natural mortality rate based on the assumption that natural mortality does not change over time. (This is done because of the difficulty in estimating natural mortality rate). If natural mortality has increased over time, and if these increases have not been quantified, then estimates of fishing mortality will be too high (when they are obtained from a Virtual Population Analysis or from a Brownie-type tagging model). Thus, there is the real potential of restricting the fishery because the fishing mortality appears too high when the actual situation is that the natural mortality has risen. This is not just of theoretical concern – for the last several years the Atlantic States Marine Fisheries Commission's Striped Bass Technical Committee and Subcommittees have struggled with the problem that the total mortality rate appears to have gone up despite the fact that the fishing regulations have been stable. But information on whether diseases may be elevating the natural mortality rate is scarce and largely circumstantial (indirect) or anecdotal. To date, no one has quantified the effects of the disease on striped bass survival rate. Indeed, to our knowledge, quantitative estimates of infectious disease impacts on population dynamics have not been incorporated in the management plan of any marine finfish species.

Materials and Methods

Capture and Tagging Protocol

Striped bass for tagging were obtained from three pound nets in the upper Rappahannock River (river miles 45-47) and from five pound nets in the lower Rappahannock River (river miles 0-3). The pound net is a fixed trap that is presumed to be non-size selective in its catch of striped bass, and has been historically used by commercial fishermen in the Rappahannock River.

All captured striped bass were removed from each pound net and placed into a floating holding pocket (1.2m x 2.4m x 1.2m deep, with 25.4mm mesh and a capacity of approximately 200 fish) anchored adjacent to the pound net. Fish were dip-netted from the holding pocket and examined for tagging. Fork length (FL) and total length (TL) measurements were taken and whenever possible the sex of each fish was determined. Striped bass not previously marked and larger than 458 mm TL were tagged with sequentially numbered internal anchor tags (Floy Tag and Manufacturing, Inc.). Each internal anchor tag was applied through a small incision in the abdominal cavity of the fish. A small sample of scales from between the dorsal fins and above the lateral line on the left side was removed and used to estimate age. Each fish was released at the site of capture immediately after receiving a tag. These tags are identical to the tags issued by the U. S. Fish and Wildlife Service except that they are lime green in color and have REWARD and a VIMS phone number imprinted into them. The rewards offered were \$5 for recapture information and \$20 for donating the entire specimen, on ice, to VIMS personnel.

Mycobacteriosis Assessment

Each tagged striped bass is given a complete external disease assessment and is photographed with a digital Canon EOS 30D camera. Overview and close-up photos are made for each side to document the initial assessment and to provide a basis for comparison when project personnel obtain recaptured striped bass. We identify 3 discrete lesion categories:

PF: Pigmented focus: $\sim 1\text{mm}^2$ pale to dark brown focus (Fig. 2b)

U: Ulceration: Loss of multiple adjacent scales with erosion/excavation of underlying tissue. Hemorrhage present or absent. Pigmentation present or absent. (Fig. 2c,d)

- scale damage or extensive loss
- range of severity: single small ulcers to multi-focal, coalescing ulcers occupying large portions of the body.

H: Putative Healing: Hyper-pigmented, (may not be apparent in ventral lesions). Scales present, but incomplete or abnormally organized. (Fig. 2e)

Within the categories U and PF we assign a severity number from 1 to 3 (PF) or 4 (U and H) according to the number of pigmented foci or the number and/or size of lesions.

A skin pathology diagnostic allows distinction between diseased and healthy fish in the context of the tagging program. By this approach, the impacts of the disease will be evaluated through differential tag return rates. Survival rates of fish with pathognomonic skin pathology will be compared to survival rates of fish without skin pathology. This will provide better estimates of components of natural mortality (M) and provide inputs for future multi-species modeling efforts.

Analytical Approach:

Disease progression:

The duration of the stages (i.e., the time it takes to progress from one condition to the next) can be estimated from tagging data if it is assumed that transitions are asynchronous across the population. This means that at the time of tagging, a fish can be anywhere in the time interval it takes to progress from one stage to the next. The methodology is analogous to that used to estimate intermolt periods in crustaceans and insects (Willoughby and Hurley 1987, Restrepo and Hoenig 1988, Hoenig and Restrepo 1989, Millar and Hoenig 1997). In the crustacean molt models, the data consist of size at tagging, time at liberty, and size at recapture. If the size at recapture is greater than the size at tagging then the animal has molted. Thus, the data reduce to time at liberty and an indicator of whether the animal molted. In the case of striped bass with dermal mycobacteriosis, the data consist of condition class at tagging, time at liberty, and

condition class at recapture. Thus, the data reduce to time at liberty and an indicator of whether the animal has progressed to the next disease condition class.

The simplest model to handle this situation was developed by Munro (1974, 1983). The recaptures are tabulated by time period, say by month. Then, under the assumptions that:

- 1) the duration of a stage (condition class) is a constant, g
- 2) at the time of tagging the time elapsed since the animal entered the condition class is a uniform random variable over the interval 0 to g
- 3) the probability of recapture does not vary by condition class.

The proportion of animals, p_t , making the transition to a higher condition class at time t is a linear function of the time at liberty, t , up until g units of time have passed, and is 1.0 for $t > g$. That is,

$$p_t = \begin{cases} \frac{t}{g}, & 0 \leq t < g \\ 1.0, & t > g \end{cases} .$$

Thus, a plot of the proportion of recaptures in a time interval that show a transition to a higher condition class should describe a linear relationship with time up until the proportion reaches 100%; the slope of the regression line estimates $1/g$. The stage duration, g , is estimated by

$$g = 1/\text{slope}.$$

The categories for disease progression are defined as:

Clean:	no external sign of infection (condition 0)
Light:	PF1 and/or U1 on at least one side (condition 1)
Moderate:	PF2 and/or U2 on at least one side (condition 2)
Heavy:	PF3 and/or U3,4 on at least one side (condition 3)
Other:	all H, but without any PF or U (condition 4)

Relative return rates and spatial differentiation refine our knowledge of the effects of the disease on striped bass stocks. Comparison of the disease index (and accompanying photos) with the infection index of recaptures returned to VIMS provides a measure of disease progression (or remission) of these striped bass.

The Munro method is generally robust (Restrepo and Hoenig 1988) but it is inefficient because a) it requires recaptures to be binned into time intervals rather than using exact times of recapture, and b) it does not use the information from animals at liberty for a long period of time.

Hoenig and Restrepo (1989) developed a likelihood approach to estimating the stage duration but their model is based on the assumption that there is no individual variability in stage duration. This assumption can cause a serious positive bias in estimates of stage duration. Millar and Hoenig (1997) generalized the approach of Hoenig and Restrepo to allow for individual variability in stage duration.

Mortality estimates:

If mycobacteriosis has no impact on the fate of fish, and if tag return rate is not affected by the presence of lesions, then we would expect to recover equal proportions of tags from fish with and without external lesions. In contrast, if externally ulcerous fish have higher mortality, we might expect to see a lower tag return rate in this group. (We discuss the necessary assumptions below.) Thus, we may estimate the impact of the lesions in terms of the relative survival (or relative risk) or in terms of the odds ratio. The results of the tagging experiment can be displayed in a 2x2 contingency table, as follows:

		recovered	not recovered
lesions		<i>a</i>	<i>b</i>
no lesions		<i>c</i>	<i>d</i>

The relative survival (with lesions : without lesions) is computed as

$$relative\ survival = \frac{a/(a+b)}{c/(c+d)} = \frac{a(c+d)}{c(a+b)}$$

Thus, if 8% of the tags are recovered from fish with lesions while 16% are recovered from fish without external lesions, the relative survival is 0.5, i.e., fish with external lesions survive half as well as fish without. The odds ratio is computed as

$$odds\ ratio = ad/(bc)$$

(Rosner 1990). The odds of obtaining a tag return from a fish with lesions is a/b ; the odds ratio is simply the ratio of the odds for the two groups (fish with and without external lesions). Thus, odds ratio = $(a/b)/(c/d) = ad/bc$. The odds ratio can take on values between 0 and infinity. In the above example, the odds ratio would be 0.46. A value less than one indicates that fish with lesions have lower survival than fish without lesions.

It is of interest to examine whether the ratio of survival changes over time. If the ratio of survival is constant over time, then a plot of $\log(\text{ratio of recaptures})$ will be a linear function of time at liberty with slope equal to the difference in instantaneous mortality rates (i.e., $\exp(\text{slope})$ estimates the ratio of survival rates). Note, that for this analysis to be valid, it is necessary to assume that the *ratio* of tag reporting rates for the two groups remains constant over time but *not*

that the reporting rates for the two groups are equal nor that the rates are unchanging. Departures from a linear relationship indicate that the ratio of survival rates or the ratio of reporting rates is changing over time (or both are changing). This model is a logistic model; consequently, standard methods are available for fitting and examining the model (Hoenig et al. 1990, Hueter *et al.* 2006).

Here, we develop a logistic model of relative survival as a linear model because this approach is intuitive and provides a graphical means to see how the model performs. Better estimates can be obtained using the method of maximum likelihood (e.g., by fitting a generalized linear model) and these will be presented in the future.

Suppose the survival rate of “clean” fish is S_o and the survival rate of fish in disease condition x is S_x . We tag and release some fish in each category and the ratio of fish in condition x to condition 0 is R in the releases. We then obtain recaptures at time t , for $t = 1, 2, \dots$. Under the assumption of the model, the ratio among the recaptures at time t , R_t , should be

$$R_t = R \left(\frac{S_x}{S_o} \right)^t$$

Taking natural logarithms of both sides leads to the linear model

$$\log_e(R_t) = \log_e(R) + t \cdot \log_e \left(\frac{S_x}{S_o} \right)$$

where $\log_e(R)$ is the y-axis intercept and $\log_e(S_x/S_o)$ is the slope. Thus, exponentiating the estimated slope provides an estimate of the relative survival (ratio of survival rates). Also, letting the survival rate of fish in disease category x be expressed as $S_x = \exp(-Z_x)$ and $S_o = \exp(-Z_o)$, we have

$$\text{slope} = \log_e \left(\frac{\exp(-Z_x)}{\exp(-Z_o)} \right) = Z_o - Z_x$$

which is the difference in the instantaneous total mortality rates. Assuming both groups of fish experience the same fishing mortality, we have

$$\text{slope} = M_o - M_x$$

where M_o is the natural mortality rate of “clean” fish and M_x is the natural mortality rate of fish in disease condition x . That is, the slope estimates how much additional natural mortality is caused by mycobacteriosis.

In theory, the intercept of the linear regression line can estimate the initial ratio of fish in the two condition categories. However, if there is differential stress or mortality associated with the tagging process then an artificial situation can be created where the ratio changes substantially over the first few days after release and then stabilizes and is then subject to just differential mortality associated with the disease (and not the tagging process). Thus, it may be necessary to disregard the initial ratio at the time of tagging and the recaptures over the first few days of recapture.

In the work plan, it was proposed that relative survival be expressed by the odds ratio approach. It should be noted that the odds ratio approach is a special case of the logistic regression described above in which observations are obtained at just two points in time. That is, the data for intermediate time steps is not used.

In subsequent reports, because tagged fish will be released at two times (one year apart), it should also be possible to fit Brownie tagging models (Brownie et al. 1985) or instantaneous rates models (Hoenig et al. 1998a,b) to the data. These models allow one to estimate annual survival rate. Thus, one can compare the survival of fish tagged with and without external signs of mycobacteriosis. Two assumptions of the model are worth noting. First, tag reporting rate need not be 100%, need not be known, and need not be constant over time. However, previously tagged and newly tagged fish are assumed to have the same reporting rate. This assumption may be violated if, for example, disease severity increases in a tagged cohort over time. In this case previously tagged fish may look less appealing than newly tagged fish, thus affecting reporting rate differentially. Second, the Brownie models are based on the assumption that the population is homogeneous, i.e., that all animals have the same probability of survival. To the extent that survival is a function of the severity of the disease, there may be some heterogeneity within the defined categories of those with and without external signs of disease. Biases that may arise due to failures of these assumptions will be studied by sensitivity analysis. Information on disease progression from examination of recaptured fish and information on disease prevalence from periodic examination of samples from the pound net, will be used to guide the sensitivity analyses.

There are other potential problems to this analysis. If ulcerous fish exhibit different movement patterns than fish that do not have the skin disease, this could influence disease dynamics. This will be tested by gathering information on the location of recaptures and evaluating the spatial distribution of recaptures for the two groups of fish.

Results

Tag Release Summary

Fall 2010: A total of 3,019 striped bass were tagged, assessed for external disease indications, photographed and released from three pound nets in the upper Rappahannock (n = 502) and five pound nets in the lower Rappahannock (n = 2,517) River during fall, 2010 (Table 2). The striped

bass tagged upriver were mostly 490-580 mm in fork length, giving a higher range than the previous year (Figure 3). An increase of disease prevalence with size is observed in the upriver fish, with all fish above 610 mm displaying some external signs of the disease. There was slightly lower range in size at the lower river nets, peaking from 440-540 mm, similar to the previous year (Figure 4). The striped bass tagged in the lower Rappahannock River also showed a slight discernable trend in prevalence of infection with size, with all fish above 580 mm showing external signs of the disease. Combined, only 29.2% (882/3,019) of the total that were tagged were without any external sign of mycobacteriosis. This is an increase from last fall when 24.9% of the releases were clean. The lightly-infected group (43.1%) had the highest prevalence, while 8.6% were heavily infected. The striped bass tagged upriver had a higher prevalence of infected striped bass (79.1% vs. 69.2%). The prevalence of infection in the upper Rappahannock River striped bass were the second highest recorded to date (84.8% in 2009). In contrast the prevalence of infection from striped bass tagged in the lower Rappahannock River has decreased over the same time interval.

Spring 2011: A total of 270 striped bass were tagged, assessed, photographed and released from the pound nets in the lower Rappahannock River during May, 2011 (Table 3). The striped bass tagged in the lower Rappahannock River were similar in size to the 2010 spring releases also and showed a trend towards an increasing prevalence of infection with size (Figure 5). While some larger fish tagged in 2010 were without external signs of the disease, all fish above 580mm showed some external diseased appearance. Although greater than for the fall releases, only 38.1% (95/232) of the total that were tagged were without any external sign of mycobacteriosis. The lightly-infected group was 36.2% of the releases, while 10.4% were heavily infected. The prevalence of heavily infected striped bass increased slightly from 2010, which is consistent to the increasing trend observed since spring 2006 (7.8%).

Tag Recapture Summary

Current year:

Fall 2010 releases: A total of 354 striped bass tagged during fall 2010 were recaptured prior to 20 September, 2011 (Table 4). The overall recapture rate was 0.117 (0.106 from the lower Rappahannock river releases and 0.173 from the upper Rappahannock releases). The incidence of immediate (< 7 days) recapture was greater from the lower Rappahannock River releases (0.033 vs. 0.020) making the recapture rate beyond the initial 7 days much higher for the upper Rappahannock River released bass (0.153 vs. 0.072). Examination of the disease prevalence in the immediate (less than 7 days at large) recaptures shows that 28%, 22%, and 23% of the light, moderate, and heavily diseased recaptures occurred within 7 days compared to 30% of the clean recaptures. Previous recapture summaries have shown a higher prevalence of moderate and severe infections within the immediate recaptures, suggesting that disease may impact a fish's ability to move. However, with this year's immediate recaptures having greater occurrences of clean and lightly diseased fish, other factors (i.e. tagging induced mortality, decreased foraging activity) may be influencing the more heavily infected fish making them less likely to be recaptured immediately after tagging. The majority of recaptures occurred in the release area

(n= 207) and throughout the Rappahannock River (n= 52), with additional recaptures throughout the bay, Potomac River, and Atlantic Ocean (Table 5).

Spring 2011 releases A total of 52 striped bass tagged during spring 2011 were recaptured prior to 20 September 2011 (Table 6). One half of the recaptures were within seven days of release. This accounted for 41% of the clean recaptures, 75% of the lightly infects, 36% of moderate and 50% of the heavily infected. No obvious differences exist in the movements of the different disease classifications. These recaptures occurred at the release site primarily (n=49), with two additional in the Rappahannock and one each in the upper bay portions of Maryland and Virginia (Table 7).

Fall 2005-Spring 2010 releases:

Fall 2005 releases: No additional striped bass tagged during fall 2005 was recaptured between 21 September, 2010 and 20 September, 2011.

Spring 2006 releases: No additional recaptures of bass tagged and released in the spring of 2006 occurred between 21 September 2010 and 20 September 2011.

Fall 2006 releases: One striped bass tagged and released fall 2006 were recaptured between 21 September, 2010 and 20 September, 2011 (Table 8). New recaptures from the fall 2006 releases were released as lightly diseased and recaptures occurred at the release site (Table 9).

Spring 2007 releases: No additional striped bass tagged during Spring 2007 was recaptured between 21 September, 2010 and 20 September, 2011.

Fall 2007 releases: A total of 5 striped bass tagged during fall 2007 were recaptured between 21 September, 2010 and 20 September, 2011 (year 4 at large, Table 10). The majority of the recaptures came from clean and lightly diseased fish with 1 recapture coming from moderately diseased fish. Most recaptures occurred in the lower portion of the Chesapeake Bay, with one recapture each from the Rappahannock River and the upper Virginia portions of the Bay (Table 11).

Spring 2008 releases: No additional recaptures of bass tagged and released in the spring of 2008 occurred between 21 September 2010 and 20 September 2011.

Fall 2008 releases: A total of 37 striped bass tagged during fall 2008 were recaptured between 21 September 2010 and 20 September, 2011 (third year at large, Table 12). Of these, 15 each were released clean or lightly diseased condition. Numbers for more severely impacted released were smaller with 4 being released with moderate infections, 2 with heavy infection and 1 with other classifications. The majority were recaptured in the release area (n=9) or throughout the Rappahannock River (n=10), with smaller numbers being recorded from the upper (n=4) and lower (n=6) portions of the Maryland Chesapeake Bay, upper (n=5) and lower (n=1) Virginia portions of the Chesapeake Bay, and the Potomac River (n=2, Table 13).

Spring 2009 releases: A total of 4 striped bass tagged during spring 2009 were recaptured between 21 September, 2010 and 20 September, 2011 (1.5-2.5 years at large, Table 14). The majority of the recaptures came from clean fish (n=3), with 1 coming from lightly diseased fish. Equal numbers (n=1) were caught at the lower Rappahannock River, the lower bay portion of Maryland and the upper and lower portions of Virginia (Table 15).

Fall 2009 releases: A total of 41 striped bass tagged during fall 2009 were recaptured between 21 September 2010 and 20 September, 2011 (second year at large, Table 16). Of these, 14 were released clean and 17 for lightly infected. There were 6 fish released with a moderate infection classification and 4 with a heavy classification. The majority were recaptured in the release area (n=18), with smaller numbers being recorded from the upper Rappahannock River (n=6), the with upper (n=3) and lower (n=1) portions of the Maryland Chesapeake Bay, upper (n=4) and lower (n=4) Virginia portions of the Chesapeake Bay, and the Potomac River (n=5, Table 17).

Spring 2010 releases: A total of 13 striped bass tagged during spring 2010 were recaptured between 21 September 2010 and 20 September, 2011 (0.5-1.5 years at large, Table 18). Of these, 5 were in clean condition and 4 in light. Additionally there were 3 recaptures released with moderate infections and 1 with heavy infection. The majority were recaptured in the release area (n=11), with two in the lower portion of the Chesapeake Bay, Virginia (Table 19).

Disease progression in Rappahannock River Striped Bass, 2005-2010

Release assessments: The relative prevalence of outwardly uninfected (clean) striped bass had been increasing in the lower and upper Rappahannock River between 2005 and 2007. In 2008 both sites began experiencing significant reductions in the relative prevalence of clean bass. From 2007 to 2008 the relative prevalence of clean bass dropped about 10% at the upriver site and about 7% at the down river site. The relative prevalence of heavily infected striped bass at both sites increased from 2007 to 2008. This is expected given that there was an increase in the relative prevalence of lightly infected striped bass at both sites between 2006 and 2007. At the lower site, the relative prevalence of lightly diseased bass increased between 2007 and 2008 while the upper site remained stable. From 2008 to 2009, river wide prevalence of clean bass decreased an additional 6%. Lightly diseased prevalence increased at both sites in 2009, with a river wide increase of 3% , and moderately diseased prevalence increased over 5% throughout the river in 2009. Heavily infected disease prevalence decreased by 2% in 2009. The 2009-2010 samples showed an almost 5% increase in the prevalence of clean released fish, throughout the river. Lightly diseased fish decreased 2% riverwide, while moderately diseased increased (2 %) and heavily diseased remained approximately the same (>1% increase). This is likely related to the increased prevalence of lightly diseased fish in previous years. The observed reduction in disease prevalence is more pronounced at the upriver site. Based on previous trends however, the relative prevalence of lightly diseased bass predicts an increase in the relative prevalence of moderate and heavily diseased bass in the following year.

The relative prevalence of clean striped bass in the sample decreased rapidly to near zero by age five in the 2002 through 2005 year classes of striped bass from both locations in the Rappahannock River (Figures 6 through 9). The 2003 and 2004 year class have not shown as sharp a decline in the relative prevalence of the clean fish as the 2002 and 2005 year class; however, trends in relative prevalence at age of all year classes suggest several things:

- 1) Bass are infected by the disease at young ages as heavily diseased fish are being observed by age three.
- 2) Bass continue to become infected with the disease as they age within the age classes observed (there appears to be no reduction in susceptibility to infection with age).
- 3) Relative prevalence of more severe conditions is increasing with age.

Estimates of disease progression

A total of 772 tagged striped bass have been recaptured and returned to VIMS for necropsy and disease reassessment from fall 2005 to present. This represents 4.01% of the total tagged striped bass released. Estimates of disease progression rate could be obtained for fish released as either lightly or moderately diseased. No disease progression rate estimates could be obtained from fish released as clean because of uncertainty around whether the fish was truly disease free or simply not expressing outward signs of the disease. Likewise no estimates could be obtained for fish released in a heavily diseased state as there is no higher stage to progress to in the classification system.

There were recaptures originally 307 assessed as light and 124 recaptures originally assessed as moderate that were returned to VIMS and had their external disease status reassessed. The plot of the progression in the disease of the striped bass originally released in the light condition with time at large (grouped by season, Figure 10) was described by:

$$Y = .00259 (x) - .08571$$

which yields an estimate of 100% progression to the moderate condition at 386 days (SE = 40 days). Likewise the plot of the progression in the disease of striped bass originally assessed as moderate (Figure 11) was described by:

$$Y = .00133 (x) - .01338$$

Which yields an estimate of 100% progression to severe at 753 days (SE=118 days).

While it is impossible to obtain direct estimates of progression rate for fish released “clean” exploration of the data shows the trend that nearly all (> 95%) fish released clean in the fall of 2005 - 2010, and subsequently recaptured have progressed to a classifiable disease

condition within one year at large (Figure 12). While this is alarming questions still remain over whether this is a true indication of the incidence rate of the disease or an artifact created by the capturing and tagging process.

Spatial comparisons

Of the 772 tagged striped bass that have been recaptured and returned to VIMS for necropsy and disease reassessment, 165 were released in the upper Rappahannock, and 607 in the lower. Given the differences in physical attributes between these sites, there may be differences in the resident bass populations, including disease progression and severity. Release assessments (see prior section) of tagged fish in both portions of the river, combined with information on disease progression and growth obtained from necropsy, can provide further insight into the differences.

Fish released in the lower Rappahannock River tended to have larger recaptured fork lengths than fish from the upper Rappahannock (Figure 13, Table 20). Of fish that were released clean and recaptured as heavily diseased, the mean fork length at the lower Rappahannock was 528.0 mm compared with 497.7 mm at the upper river locations. Similar trends occur for other release disease conditions. For releases only, fish released at the lower site tended to be larger than the fish released at the upper Rappahannock River sites (mean = 490.8 mm vs 463.3 mm, respectively). Changes in fork length vary between the two sites, with the lower river having a greater change in fork length for animals progressing from clean (42.7 mm moderate and 37.2 mm severe), while the upper river had a greater change in fork length for fish remaining at their current condition (9.4 mm).

Additionally, days at liberty varies between the two sites. The variation between the changes in fork length could be attributed to longer days at liberty for fish tagged at the lower site, however on average days at liberty is greater at the upriver site (Figure 14). Fish released at the upriver site assessed as clean, had a mean days at liberty of 110 for clean recaptures, 253 for light, 447 for moderate, and 479 for severe. In contrast, fish for the downriver site had a mean days at liberty of 83 for clean recaptures, 213 for light, 385 for moderate, and 346 for severe. Again, trends continued for light and moderate releases. Extrapolating from both days at liberty and recaptured fork lengths based on disease progression, we can get a growth per day and again see that this is less for the upriver site than for the lower (Table 20). Also from this assessment we can see, in rare cases, that a diseased fish may display a negative disease progression upon its recapture. Though occurrences of this are seldom, the growth per day of these individuals tends to be much smaller than those of fish which have progressed, suggesting that some diseased fish may allocate energy to fighting the infection rather than to somatic growth. The cases occur in the lower portion of the Rappahannock River more frequently than the upper, as well as across a variety of starting disease conditions.

Similarly, progression from disease condition light to moderate is estimated to take 366 days for lower river releases but 490 days in the upper releases (Figures 15A-D). Likewise, site specific estimates of time to progression from moderate to severe vary with estimates of 507

days (upper river) and 1026 days (lower river). However, the differences in the progression estimates are not statistically significant and likely reflect difference in sample size. Combined with other observations of size, growth and days at liberty differences, there appears to be varied factors influencing the susceptibility of fish in resident vs. transient populations.

Estimation of survival rates and relative survival rates

Logistic model

The rate of return of tags from diseased fish is clearly lower than that for “clean” fish (showing no overt signs of disease). If the rate of return were equal for the two groups, a plot of the ratio of returns (or the log of the ratio) versus time would be a horizontal line. But, it can be seen in Figures 16A-D that the slope is negative indicating that diseased fish are not surviving as well as clean fish or that diseased fish are less catchable than clean fish. The slope of the regression lines in Figures 16A-D provide estimates of the difference in instantaneous natural mortality rates, i.e., of the additional mortality caused by mycobacteriosis. Estimates of the ratio of annual survival rates can be obtained by exponentiating the slope of the regression line. In computing the linear regression lines, the initial tagging ratio and the recaptures during the first seven days at liberty have not been used because of concerns that they represent an artificial situation associated with the stress of tagging (see methods section for an explanation).

Fish in disease conditions 3 and 2 have estimated elevations of natural mortality rate M above that of clean fish of .89 and .39, respectively (Table 21, Figures 16A and B). This implies annual survival rates for fish in disease conditions 3 and 2 that are 55% with a 95% confidence interval of (40% , 74%) and 68% with 95% confidence interval of (52% , 86%), respectively, of the survival of clean fish. Because the results for disease conditions 2 and 3 are similar, we combined the data from these two disease categories to boost sample sizes and increase precision. The result is an estimated difference in M between fish in conditions 2 and 3 and fish that are clean of .43; the estimated ratio of survival rates is 65%, 95% confidence interval equal to (49% , 83%) (Table 21, Figure 16D).

Fish in disease condition 1 appear to have an elevated mortality rate relative to clean fish but not as high a mortality rate as fish in disease conditions 2 and 3 (Figure 16C). The estimated difference in instantaneous natural mortality rates is 0.09 and the ratio of survival rates is 91%, 95% confidence interval of (77%, 109%) (Table 21).

The estimated impacts of the disease are not very precise but provide a compelling indication that the disease has population impacts. The estimates of the increase in mortality for fish in condition 3 (relative to clean fish) is highly statistically significant ($p < 0.001$). The estimate for condition 2 is very similar and the p -value (0.002) is also statistically significant. Combining conditions 2 and 3 to boost sample sizes gives a significant result ($p = .009$) The estimated slope for condition 1 fish indicates a relative survival rate that is four fifths that of clean fish and higher than that of fish in category 2-3. This is a reasonable result. However, the slope is not statistically significant ($p = 0.341$) so the possibility that condition 1 fish have the

same mortality rate as clean fish cannot be ruled out at this time. The past year of tag returns improved our ability to estimate the relative mortality rate of condition 1 fish versus clean fish and if the present trend continues statistically significant estimates will be available for all disease conditions after one additional year of tag returns.

Discussion

The results so far establish some important points. First, we continue to obtain excellent cooperation from commercial and sport fishers so that our rate of return of tags (about 11.45% of releases), and of tagged carcasses (4.01%), is encouraging. Second, if diseased fish are less able to withstand the stress of capture and tagging than lightly diseased or non-diseased fish, then we could have an artifact of tagging whereby an appreciable fraction of the diseased fish experience an abnormal mortality associated with the tagging process. However, our tag returns are of the same ratio as the tag releases, indicating that this is not a problem. In fact, we obtained slightly higher tag return rates from diseased fish than from fish without signs of disease. Third, it is possible that diseased fish may differ in their ability to swim and migrate as well as other behaviors from fish without signs of the disease. Fortunately, we are able to obtain detailed recapture locations from almost all fish, which can be used to further examine spatial differences and movement of diseased vs. clean fish. Finally, there is some preliminary indication of spatial differences impacting disease prevalence when examining the data from the upper and lower sites individually, suggesting the disease prevalence is increasing more rapidly at the upper river sites and also has a more drastic effect on growth when in comparison to the lower river.

While the prevalence of heavily-infected striped bass increased slightly, as did the relative prevalence of moderately diseased fish, the proportion of the striped bass examined as non-infected increased from 24.4% to 30.6%. While this is an increase from the previous year, it is the same percent as what was reported in 2008, thus resulting in no net change of non-diseased proportions in a two year period. We have recapture information from striped bass released as heavily-infected more than one year after their release, so the disease is not 100% fatal within this time frame. Some severely infected fish have been recaptured well over a year later while lightly and moderately infected fish have persisted with the disease for over two years on some occasions. Additionally the necropsies performed on returned carcasses include incidences of healing individual pigmented foci and ulcers. The slow progression and presence of healing fish may indicate that the progression in wild striped bass is slower than what has been observed in aquaculture. However the increased prevalence of diseased fish and a greater likelihood of progression over time do indicate that the disease is progressive. We have determined that the majority of striped bass will progress in disease severity on an annual basis and that very few resident (fall) striped bass remain outwardly uninfected by age five. Our current estimate of disease stage progression is 386 days for lightly infected fish to progress to moderately infected and 753 days for moderately infected to progress to severely infected. These estimates will be refined as more recaptures are returned to VIMS for reassessment; however the time to progression for lightly infected fish has decreased since the 2009 model.

The lower prevalence of mycobacterial infections in the larger, migrant striped bass indicates that the resident population is most at risk. Additionally, while time to progress to the next disease stage remains the same for light and moderate infections at the upper level (490 and 507 days, respectively), the time to progression for the lower is largely differed and over twice the time of the upper for the moderate infections (366 days light, 1026 days moderate). Since the resident striped bass form the basis of both the recreational and commercial fisheries in Virginia, the results of this study will be increasingly important.

This project has provided a direct measurement of disease-associated mortality by stage of the disease. Moderately and heavily infected fish appear to have slightly more than one half the survival rate of fish tagged without outward signs of disease. Fish with early signs of the disease appear to have slightly reduced survival relative to fish without signs of the disease. The estimated relative survival for lightly (early stage) infected fish is not statistically different at the $\alpha = .05$ level from the “clean” fish. As further tagging results are obtained the standard error can be expected to be reduced. It should be noted that the fish tagged without outward signs of disease are a mixture of uninfected fish and infected fish that are not yet showing signs of the disease. Thus, a comparison of the two groups underestimates the disease-associated mortality because some fish in the “clean” group may already be experiencing disease-related mortality.

Literature Cited

- Brownie, C., D.R. Anderson, K.P. Burnham, and D.R. Robson. 1985. Statistical inference from band recovery data: a handbook, 2nd ed., U.S. Fish and Wildl. Serv. Resour. Publ. No. 156.
- Bruno DW, J Griffiths, CG Mitchell, BP Wood, ZJ Fletcher, FA Drobniewski, and TS Hastings. 1998. Pathology attributed to *Mycobacteria chelonae* infection among farmed and laboratory - infected Atlantic salmon (*Salmo salar*). Dis Aquat Org. 33:101-109
- Cardinal JL. 2001. Mycobacteriosis in striped bass, *Morone saxatilis*, from Virginia waters of Chesapeake Bay. Master's Thesis. School of Marine Science, Virginia Institute of Marine Science. Pp.83.
- Colomi A. 1992. A systemic mycobacteriosis in the European sea bass *Dicentrarchus labrax* cultured in Eilat (Red Sea). Bamidgah – Isr J Aquacult 44:75-81
- Crecco, V. 2003. Methods of estimating fishing (F) and natural (M) mortality rates from total mortality (Z) and exploitation (u) rates for striped bass. Final Report. Connecticut Marine Fisheries Division. 40 pp.
- Garrison, LP, JS Link and G White. 2004. A multispecies modeling approach to evaluate interactions between Atlantic menhaden and its predators. Abstract: 60th Annual Northeast Fish and Wildlife Conference. 25-28 April, 2004. Ocean City, Maryland.
- Gauthier, DT, RJ Latour, DM Heisey, CF Bonzak, J Gartland, EJ Burge and WK Vogelbein. 2008. Mycobacteriosis-associated mortality in wild striped bass (*Morone saxatilis*) from Chesapeake Bay, USA. Ecol. Appl. 18: 1718-1727.
- Hartman, KJ 2004. Increases in coastal striped bass predatory demand and implications of declines in Atlantic menhaden populations. Abstract: 60th Annual Northeast Fish and Wildlife Conference. 25-28 April, 2004. Ocean City, Maryland.
- Hoening JM, NJ Barrowman, WS Hearn, and KH Pollock. 1998. Multiyear Tagging Studies Incorporating Fishing Effort Data. Can. J. Fish. Aquat. Sci. 55:1466-1476.
- Hoening JM, NJ Barrowman, KH Pollock, EN Brooks, WS Hearn and T Polacheck. 1998. Models for Tagging Data that Allow for Incomplete Mixing of Newly Tagged Animals. Can. J. Fish. Aquat. Sci. 55:1477-1483.
- Hoening JM, P Pepin, and WD Lawing. 1990. Estimating Relative Survival Rate for Two Groups of Larval Fishes from Field Data: Do Older Larvae Survive Better than Young? Fish. Bull. 88:485-491.

- Hoening, J.M. and V.R. Restrepo. 1989. Estimating the Intermolt Periods in Asynchronously Molting Crustacean Populations. *Biometrics* 45:71-82.
- Hueter, R.E., C.A. Manire, J. Tyminski, J.M. Hoening and D.A. Hepworth. 2006. Assessing Mortality of Released or Discarded Fish Using a Logistic Model of Relative Survival Derived from Tagging Data. *Trans. Am. Fish. Soc.* 135:500-508.
- Jacobs JM, HL Rogers, WF Van Heukelem, B Coakley, C Giesecker and M Matsche. 2004. Nutritional health of Chesapeake Bay striped bass *Morone saxatilis* in relation to disease. Abstract: 60th Annual Northeast Fish and Wildlife Conference. 25-28 April, 2004. Ocean City, Maryland.
- Jiang, H, K.H. Pollock, C. Brownie, R.J. Latour, J.M. Hoening, B.K. Wells, and J.E. Hightower. 2007. Tag Return Models Allowing for Harvest and Catch and Release: Evidence of Environmental and Management Impacts on Striped Bass Fishing and Natural Mortality Rates. *North American Journal of Fisheries Management.* 27:387-396.
- Kahn, DM. 2004. Tag-recapture data from Chesapeake Bay resident striped bass indicate that survival has declined. Abstract: 60th Annual Northeast Fish and Wildlife Conference. 25-28 April, 2004. Ocean City, Maryland.
- May, EB, V Pernell Lewis, AM Overton, J Jacobs and L Alade. 2004. Potential impacts of mycobacteriosis in striped bass on Chesapeake and Atlantic coastal stocks. Abstract: 60th Annual Northeast Fish and Wildlife Conference. 25-28 April, 2004. Ocean City, Maryland.
- Millar, R.B. and J.M. Hoening. 1997. A Generalized Model for Estimating Intermolt Periods of Asynchronously Molting Insects and Crustacea from Field or Laboratory Data. *J. Agric., Biol. & Environ. Statistics* 2(4):1-14.
- Munro, J.L. 1974. The biology, ecology and bionomics of spiny lobsters (Palinuridae), spider crabs (Majidae) and other crustacean resources. Part VI in J. L. Munro, ed. The biology, ecology, exploitation and management of Caribbean reef fishes: Scientific Report of the ODA/UWI Fisheries Ecology Research Project 1969-1973: University of the West Indies, Jamaica. Research Reports from the Zoology Department Number 3, University of the West Indies, Kingston. 57 pp.
- Nigrelli RF and H Vogel. 1963. Spontaneous tuberculosis in fish and other coldblooded vertebrates with special reference to *Mycobacterium fortuitum* from fish and human lesions. *Zoologica* 48:131-144.
- Restrepo, V.R. and J.M. Hoening. 1988. Munro's Method for Estimating Intermolt Periods of Tropical Decapods is Robust. *Bull. Mar. Sci.* 42:488-492.

- Rhodes MW, H Kator, S Kotob, P van Berkum, I Kaattari, WK Vogelbein, F Quinn, MM Floyd, WR Butler and CA Ottinger. 2003. *Mycobacterium shottsii* sp. nov., a slow growing species isolated from Chesapeake Bay striped bass (*Morone saxatilis*). Int. J. Syst. Evol. Micro. 53:1-5.
- Rhodes MW, H Kator, S Kotob, P van Berkum, I Kaattari, WK Vogelbein, F Quinn, MM Floyd, WR Butler, CA Ottinger. 2002. *Mycobacterium shottsii* sp. nov., a slow growing species isolated from Chesapeake Bay striped bass, (*Morone saxatilis*). Int. J. System. Environ. Microbiol. 53:421-424.
- Rhodes, MW, H Kator, I Kaattari, D Gauthier, WK Vogelbein, & C Ottinger. 2004. Isolation and characterization of *Mycobacterium* spp. from striped bass, *Morone saxatilis*, from the Chesapeake Bay. Dis. Aquat. Org. 61:41-51.
- Rosner, B. 1990. Fundamentals of Biostatistics, 3rd edition. PWS-Kent Publishing Company, Boston.
- Uphoff, JH Jr. 2004. Striped bass and Atlantic menhaden: Is there a predator-prey imbalance in Chesapeake Bay? Abstract: 60th Annual Northeast Fish and Wildlife Conference. 25-28 April, 2004. Ocean City, Maryland.
- Vogelbein WK, DE Zwerner, H Kator, MW Rhodes and J Cardinal. 1999. Mycobacteriosis of striped bass from Chesapeake Bay. pages 53-58. In J.E. Olney (ed.), *Research on Recreational Fishes and Fisheries*, VIMS Spec. Sci. Rept. 139, 82 pp.
- Vogelbein WK, DT Gauthier, MW Rhodes, H Kator, R Latour, C Bonzek and C Ottinger. 2004. Mycobacteriosis in striped bass (*Morone saxatilis*) from Chesapeake Bay. Abstract: 60th Annual Northeast Fish and Wildlife Conference. 25-28 April, 2004. Ocean City, Maryland.
- Willoughby, L.G. and M.A. Hurley. 1987. "Echo" moulting used to estimate moulting periodicity of mayflies (Ephemeroptera) and stoneflies (Plecoptera) in nature. Aquatic Insects 9:221-227.

Table 1. Parameter estimates and standard errors (SE) from fitting two models to the Virginia striped bass spring tagging data (age 2 and greater). In model (a), estimates are obtained for year-specific fishing mortality rates for killed fish in year xx, $F_k(xx)$, for fishing mortality associated with released fish experiencing hooking mortality, $Fr(xx)$, and for natural mortality rate in two time periods (1990-1996 and 1997-2004). In model (b), the same parameters are estimated but, in addition, the tag reporting rates for kept (λ_K) and released (λ_R) fish are estimated instead of being fixed at 0.43.

parameter	(a)		(b)	
	estimate	SE	estimate	SE
Fk(90)	0.122	0.023	0.182	0.057
Fk(91)	0.165	0.021	0.259	0.067
Fk(92)	0.236	0.032	0.360	0.091
Fk(93)	0.227	0.032	0.347	0.086
Fk(94)	0.263	0.043	0.428	0.107
Fk(95)	0.274	0.042	0.469	0.116
Fk(96)	0.195	0.035	0.416	0.111
Fk(97)	0.199	0.039	0.370	0.105
Fk(98)	0.306	0.058	0.645	0.179
Fk(99)	0.240	0.034	0.578	0.163
Fk(00)	0.114	0.023	0.196	0.065
Fk(01)	0.111	0.024	0.145	0.047
Fk(02)	0.252	0.057	0.286	0.084
Fr(90)	0.135	0.025	0.159	0.145
Fr(91)	0.153	0.020	0.184	0.164
Fr(92)	0.166	0.027	0.193	0.172
Fr(93)	0.209	0.031	0.241	0.218
Fr(94)	0.199	0.037	0.246	0.237
Fr(95)	0.073	0.020	0.097	0.095
Fr(96)	0.083	0.022	0.127	0.117
Fr(97)	0.101	0.027	0.137	0.125
Fr(98)	0.076	0.027	0.113	0.106
Fr(99)	0.103	0.022	0.165	0.153
Fr(00)	0.055	0.016	0.076	0.073
Fr(01)	0.064	0.018	0.069	0.065
Fr(02)	0.114	0.035	0.107	0.098
Fk(03)	0.427	0.140	0.362	0.129
Fr(03)	0.242	0.088	0.168	0.164
Fk(04)	0.924	0.556	0.684	0.329
Fr(04)	0.449	0.276	0.245	0.280
M90-96	0.231	0.019	0.083	0.177
M97-04	0.407	0.037	0.168	0.125
lambdaK	0.430	0.000	0.250	0.057
lambdaR	0.430	0.000	0.347	0.312

Table 2. Tag release totals and mycobacteria infection index, by date, of striped bass in the upper and lower Rappahannock River sites, fall, 2010

Date	release	n	infection index				
	Area		clean	light	moderate	heavy	other
20 September	Upper	102	24	41	23	14	0
23 September	Upper	54	15	19	11	9	0
29 September	Upper	68	9	35	12	12	0
6 October	Upper	113	19	45	31	18	0
11 October	Upper	66	21	20	14	11	0
13 October	Lower	312	73	143	69	27	0
18 October	Upper	99	17	50	20	11	1
20 October	Lower	356	60	219	54	21	2
25 October	Lower	190	54	85	26	24	1
28 October	Lower	126	27	52	28	17	2
3 November	Lower	415	145	168	63	32	7
5 November	Lower	272	103	97	40	28	4
10 November	Lower	300	102	128	52	16	2
15 November	Lower	136	46	52	27	6	5
18 November	Lower	113	38	35	24	12	4
22 November	Lower	166	71	64	21	8	2
23 November	Lower	131	58	47	17	4	5
totals	Upper	502	105	210	111	75	1
	Lower	2,517	777	1,090	421	195	34
	Both	3,019	882	1,300	532	260	35

Table 3. Tag release totals and mycobacteria infection index, by date, of striped bass in the upper and lower Rappahannock River sites, spring, 2011.

Date	release area	n	infection index				
			clean	light	moderate	heavy	other
12 May	Lower	69	36	22	4	5	2
16 May	Lower	81	31	26	15	7	2
19 May	Lower	39	14	14	4	5	2
23 May	Lower	29	9	14	4	2	0
26 May	Lower	21	7	8	2	4	0
2 June	Lower	22	5	11	3	3	0
6 June	Lower	9	1	3	3	2	0
Totals	Lower	270	103	98	35	28	6

Table 4. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2010.

Date	release area	n	infection index				
			clean	light	moderate	heavy	other
0-7 days	upper	10	1	4	3	2	0
	lower	84	27	37	11	8	1
Fall 2010 (>7 days)	upper	34	7	11	10	6	0
	lower	112	32	49	14	15	2
Winter 2010	upper	13	2	5	5	1	0
	lower	6	1	3	0	1	1
Spring 2011	upper	9	0	2	4	3	0
	lower	24	5	10	4	4	1
Summer 2011	upper	21	9	6	3	3	0
	lower	41	9	21	10	1	0
totals	upper	87	19	28	25	15	0
	lower	267	74	120	39	29	5
	both	354	93	148	64	44	5

Table 5. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2010.

recapture area	release area	n	infection index				
			clean	light	moderate	heavy	other
release area	upper	41	8	13	10	10	0
	lower	176	52	76	26	19	3
Rappahannock River	upper	39	8	13	14	4	0
	lower	13	4	5	2	2	0
upper Bay (Md)	upper	1	1	0	0	0	0
	lower	7	2	4	1	0	0
lower Bay (Md)	upper	2	1	1	0	0	0
	lower	31	7	14	7	1	2
Potomac River	upper	0	0	0	0	0	0
	lower	2	1	1	0	0	0
upper Bay (Va)	upper	1	0	1	0	0	0
	lower	27	7	14	2	4	0
lower Bay (Va)	upper	3	1	0	1	1	0
	lower	9	1	5	0	3	0
Atlantic Ocean	upper	0	0	0	0	0	0
	lower	2	0	1	1	0	0
totals	upper	87	19	28	25	15	0
	lower	267	74	120	39	29	5
	both	354	93	148	64	44	5

Table 6. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2011.

Date	release area	N	infection index				
			clean	light	moderate	heavy	other
0-7 days	lower	26	7	12	5	1	1
Spring 2011 (>7days)	lower	19	6	2	8	2	1
Summer 2011	lower	7	4	2	1	0	0
totals	Lower	52	17	16	14	3	2

Table 7. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2011.

recapture area	release area	n	infection index				
			clean	light	moderate	heavy	Other
release area	lower	49	17	15	9	6	2
Rappahannock River	lower	2	0	1	0	1	0
upper Bay (Md)	lower	1	1	0	0	0	0
lower Bay (Md)	lower	0	0	0	0	0	0
Potomac River	lower	0	0	0	0	0	0
upper Bay (Va)	lower	1	0	0	1	0	0
lower Bay (Va)	lower	0	0	0	0	0	0
totals	lower	53	18	16	10	7	2

Table 8. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2006 and recaptured from fall 2010 through summer 2011.

Date	release area	n	infection index				
			clean	light	moderate	heavy	other
Fall 2010	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
Winter 2010	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Spring 2011	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Summer 2011	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
totals	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
	both	1	0	1	0	0	0

Table 9. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2006 and recaptured from fall 2010 through summer 2011.

recapture area	release area	n	infection index				
			clean	light	moderate	heavy	other
release area	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
Rappahannock River	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
upper Bay (Md)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
lower Bay (Md)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Potomac River	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
upper Bay (Va)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
lower Bay (Va)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Atlantic Ocean	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
totals	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
	both	1	0	1	0	0	0

Table 10. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2007 and recaptured from fall 2010 through summer 2011.

Date	release area	n	infection index				
			Clean	light	moderate	heavy	other
Fall 2010	upper	0	0	0	0	0	0
	lower	2	1	1	0	0	0
Winter 2010	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
Spring 2011	upper	0	0	0	0	0	0
	lower	1	1	0	0	0	0
Summer 2011	upper	0	0	0	0	0	0
	lower	1	0	0	1	0	0
totals	upper	0	0	0	0	0	0
	lower	5	2	2	1	0	0
	both	5	2	2	1	0	0

Table 11. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2007 and recaptured from fall 2010 through summer 2011.

recapture area	release area	N	infection index				
			Clean	Light	Moderate	Heavy	other
release area	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Rappahannock River	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
upper Bay (Md)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
lower Bay (Md)	upper	0	0	0	0	0	0
	lower	3	2	1	0	0	0
Potomac River	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
upper Bay (Va)	upper	0	0	0	0	0	0
	lower	1	0	0	1	0	0
lower Bay (Va)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
totals	upper	0	0	0	0	0	0
	lower	5	2	2	1	0	0
	both	5	2	2	1	0	0

Table 12. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2008 and recaptured from fall 2010 through summer 2011.

Date	release	n	infection index				
	area		clean	light	moderate	heavy	other
Fall 2010	upper	4	1	2	0	0	1
	lower	15	7	6	2	0	0
Winter 2010	upper	3	1	2	0	0	1
	lower	2	1	0	0	1	0
Spring 2011	upper	0	0	0	0	0	0
	lower	4	1	1	1	1	0
Summer 2011	upper	2	1	1	0	0	0
	lower	7	3	3	1	0	0
totals	upper	9	3	5	0	0	1
	lower	28	12	10	4	2	0
	both	37	15	15	4	2	1

Table 13. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2008 and recaptured from fall 2010 through summer 2011.

recapture area	release area	n	infection index				
			clean	light	moderate	heavy	other
release area	upper	4	1	2	0	0	1
	lower	5	3	0	2	0	0
Rappahannock River	upper	4	2	2	0	0	0
	lower	6	2	3	0	1	0
upper Bay (Md)	upper	0	0	0	0	0	0
	lower	4	0	4	0	0	0
lower Bay (Md)	upper	0	0	0	0	0	0
	lower	6	5	1	0	0	0
Potomac River	upper	0	0	0	0	0	0
	lower	2	0	1	1	0	0
upper Bay (Va)	upper	1	0	1	0	0	0
	lower	4	1	1	1	1	0
lower Bay (Va)	upper	0	0	0	0	0	0
	lower	1	1	0	0	0	0
totals	upper	9	3	5	0	0	1
	lower	28	12	10	4	2	0
	both	37	15	15	4	2	1

Table 14. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2009 and recaptured from fall 2010 through summer 2011.

Date	release	n	infection index				
	area		clean	light	moderate	heavy	other
Fall 2010	upper	0	0	0	0	0	0
	lower	3	3	0	0	0	0
Winter 2010	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Spring 2011	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Summer 2011	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
totals	upper	0	0	0	0	0	0
	lower	4	3	1	0	0	0
	both	4	3	1	0	0	0

Table 15. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2009 and recaptured from fall 2010 through summer 2011.

recapture area	release area	n	infection index				
			clean	light	moderate	heavy	other
release area	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Rappahannock River	upper	0	0	0	0	0	0
	lower	1	1	0	0	1	0
upper Bay (Md)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
lower Bay (Md)	upper	0	0	0	0	0	0
	lower	1	1	0	0	0	0
Potomac River	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
upper Bay (Va)	upper	0	0	0	0	0	0
	lower	1	1	0	0	0	0
lower Bay (Va)	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
totals	upper	0	0	0	0	0	0
	lower	4	3	1	0	0	0
	both	4	3	1	0	0	0

Table 16. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2009 and recaptured from fall 2010 through summer 2011.

Date	release	n	infection index				
	area		clean	light	moderate	heavy	other
Fall 2010	upper	11	4	4	2	1	0
	lower	19	8	6	3	2	0
Winter 2010	upper	2	0	2	0	0	0
	lower	1	1	0	0	0	0
Spring 2011	upper	0	0	0	0	0	0
	lower	4	1	2	1	0	0
Summer 2011	upper	0	0	0	0	0	0
	lower	4	0	3	0	1	0
totals	upper	13	4	6	2	1	0
	lower	28	10	11	4	3	0
	both	41	14	17	6	4	0

Table 17. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during fall, 2009 and recaptured from fall 2010 through summer 2011.

recapture area	release area	n	infection index				
			clean	light	moderate	heavy	other
release area	upper	6	2	3	1	0	0
	lower	12	4	5	3	0	0
Rappahannock River	upper	6	1	3	1	1	0
	lower	0	0	0	0	0	0
upper Bay (Md)	upper	0	0	0	0	0	0
	lower	3	0	2	0	1	0
lower Bay (Md)	upper	0	0	0	0	0	0
	lower	1	0	1	0	0	0
Potomac River	upper	1	0	1	0	0	0
	lower	4	3	1	0	0	0
upper Bay (Va)	upper	0	0	0	0	0	0
	lower	4	0	1	1	2	0
lower Bay (Va)	upper	0	0	0	0	0	0
	lower	4	3	1	0	0	0
totals	upper	13	3	7	2	1	0
	lower	28	10	11	4	3	0
	both	41	13	18	6	4	0

Table 18. Seasonal recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2010 and recaptured from fall 2010 through summer 2011.

Date	release	n	infection index				
	area		clean	light	moderate	heavy	other
Fall 2010	upper	0	0	0	0	0	0
	lower	8	4	3	1	0	0
Winter 2010	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Spring 2011	upper	0	0	0	0	0	0
	lower	4	1	1	1	1	0
Summer 2011	upper	0	0	0	0	0	0
	lower	1	0	0	1	0	0
totals	upper	0	0	0	0	0	0
	lower	13	5	4	3	1	0
	both	13	5	4	3	1	0

Table 19. Spatial recapture summary, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites during spring, 2010 and recaptured from fall 2010 through summer 2011.

recapture area	release area	n	infection index				
			clean	light	moderate	heavy	other
release area	upper	0	0	0	0	0	0
	lower	11	4	4	2	1	0
Rappahannock River	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
upper Bay (Md)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
lower Bay (Md)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
Potomac River	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
upper Bay (Va)	upper	0	0	0	0	0	0
	lower	0	0	0	0	0	0
lower Bay (Va)	upper	0	0	0	0	0	0
	lower	2	1	0	1	0	0
totals	upper	0	0	0	0	0	0
	lower	13	5	4	3	1	0
	both	13	5	4	3	1	0

Table 20. Spatial necropsy summary of growth and days at liberty, by mycobacteria infection index and release area, of striped bass tagged and released in the upper and lower Rappahannock River sites through summer 2011.

Release site and disease	Recaptured disease	Mean recaptured Fork Length (mm)	change in Fork Length	Mean days at Liberty	Growth per Day
Upper clean	clean	460.6	9.4	110	0.13526
	Light	493.9	19.3	253	0.09922
	Moderate	510.4	39.5	447	0.08013
	Severe	497.7	33.86	479	0.07298
Lower clean	clean	482.7	3.31	83	-0.37917
	Light	509.7	17.31	213	0.42939
	Moderate	522.5	42.7	385	0.09987
	Severe	528.0	37.2	346	0.09728
Upper Light	Light	518.2	12.9	241	0.31093
	Moderate	521.3	31.49	387	0.08718
	Severe	487.6	18.08	454	0.04721
Lower Light	clean	493.6	0.33	138	0.08425
	Light	511.3	8.23	140	-0.37257
	Moderate	525.0	31.87	343	0.08641
	Severe	525.8	35.33	393	0.08892
Upper Moderate MModerate	Light	487.33	9.0	86	0.29323
	Moderate	520.46	14.07	301	0.10931
	Severe	491.6	22.53	371	0.06917
Lower Moderate Release	Light	509.30	13.09	185	-0.30264
	Moderate	512.8	12.08	166	-0.13073
	Severe	512.15	22.1	322	0.07553

Table 21. Estimates of mortality associated with mycobacterial disease and estimated relative survival rates. The slope of the regression line of log(ratio of recaptures) versus time estimates the difference in natural mortality rate (M for clean fish - M for diseased fish). The exponentiated slope estimates the ratio of finite (annual) survival rates (S for diseased fish/ S for clean fish).

Comparison	Slope	S.E.	P-value	exp (slope)
heavy vs. clean	-0.59	0.16	<0.001	0.55
moderate vs. clean	-0.39	0.13	0.002	0.68
light vs. clean	-0.09	0.08	0.341	0.91
moderate + heavy vs. clean	-0.43	0.13	0.009	0.65

Figure 1. Gross clinical signs of mycobacteriosis in Chesapeake Bay striped bass. A) Severe ulcerative dermatitis. Note shallow, rough textured hemorrhagic and hyper-pigmented (dorsal lesions) ulcers. B) Multi-focal pale gray nodules within the spleen.



Figure 2. A spectrum of gross skin lesions attributable to mycobacteriosis in the striped bass, *Morone saxatilis*. a) mild scale damage and scale loss (arrows). b) pigmented foci (arrows). Inset: higher magnification of a pigmented focus showing pin-point erosion through an overlying scale (arrow). c) early ulceration exhibiting focal loss of scales, mild pin-point multifocal pigmentation and underlying exposed dermis. d) large advanced shallow roughly textured ulceration exhibiting hyper-pigmentation and hemorrhage. e) late stage healing lesion exhibiting hyper-pigmentation, reformation of scales and re-epithelialization and closure of the ulcer. f) Ziehl Neelsen stain of a histologic section of a skin lesion exhibiting granulomatous inflammation and acid-fast rod-shaped mycobacteria (staining red). g) histologic section showing normal healthy skin composed of epidermis (Ep), scales (Sc), dermis (D) and underlying skeletal muscle (M). h) histologic section through a skin ulcer showing loss of epidermis and scales and extensive granuloma formation (G).




Figure 3. a) Size distribution (fork length in mm), by infection index, of striped bass tag releases from the upper Rappahannock River, fall 2010. b) Relative proportion of each infection index, by fork length, of the tag releases.





Figure 4. a) Size distribution (fork length in mm), by infection index, of striped bass tag releases from the lower Rappahannock River, fall 2010. b) Relative proportion of each infection index, by fork length, of the tag releases.





Figure 5. a) Size distribution (fork length in mm), by infection index, of striped bass tag releases from the lower Rappahannock River, spring 2011. b) Relative proportion of each infection index, by fork length, of the tag releases.





Figure 6. Progression in the mycobacteriosis skin severity index, with age, of the 2002 year class of striped bass in the lower Rappahannock River, falls 2005-2010.




Figure 7. Progression in the mycobacteriosis skin severity index, with age, of the 2003 year class of striped bass in the lower Rappahannock River, falls 2006-2010.




Figure 8. Progression in the mycobacteriosis skin severity index, with age, of the 2004 year class of striped bass in the lower Rappahannock River, falls 2007 - 2010.




Figure 9. Progression in the mycobacteriosis skin severity index, with age, of the 2005 year class of striped bass in the lower Rappahannock River, falls 2008 - 2010.




Figure 10. Progression of mycobacteriosis from lightly diseased at time of release to moderately diseased versus time-at-large for striped bass tagged and released in the Rappahannock River, fall 2005 to present (combined). Numbers next to the data points indicate number of recaptures.




Figure 11. Progression of mycobacteriosis from moderately diseased at time of release to severely diseased versus time-at-large for striped bass tagged and released in the Rappahannock River, fall 2005 to present (combined). Numbers next to the data points indicate number of recaptures.




Figure 12. Progression of pigmented foci (PF) of uninfected striped bass based on reassessment of recaptured striped bass originally tagged and released in the Rappahannock River, falls 2005-2010.




Figure 13. Mean recaptured fork length of fish tagged at the upper and lower locations of the Rapphannock River, broken down by site and release disease condition. A) Original disease assessment of clean. B) Original disease assessment of light. Note the presence of fish “healing” or displaying negative disease progression C) Original disease assessment of Moderate, again with some fish displaying negative disease progression.

Figure 13a.




Figure 13b.




Figure 13c.




Figure 14. Boxplot of days at liberty by disease progression based site and release disease assessment. A) Fish released as clean with no external signs of the disease. B) Fish with an original assessment of light, note the presence of fish “healing” or displaying negative disease progression. C) Fish released with a disease assessment of moderate, again with some fish displaying negative disease progression.

Figure 14a.




Figure 14b.





Figure 14c.

Figure 15. Progression of mycobacteriosis from diseased at time of release to next level of severity versus time-at-large for striped bass tagged and released in the upper and lower Rappahannock River, fall 2005 to present (combined). Numbers next to the data points indicate number of recaptures.

Figure 15a: Upper Rappahannock released as disease condition 1 (light) progressing to disease condition 2 (moderate)




Figure 15b: Lower Rappahannock released as disease condition 1 (light) progressing to disease condition 2 (moderate)




Figure 15c: Upper Rappahannock released as disease condition 2 (moderate) progressing to disease condition 3 (severe)




Figure 15d: Lower Rappahannock released as disease condition 2 (moderate) progressing to disease condition 3 (severe)




Figure 16. Logarithm of the ratio of returns of fish tagged in disease condition x and disease condition 0 (fish in condition 0 are “clean”, showing no signs of the disease) as a function of time at liberty. Numbers next to the data points are the number of tag returns. The slope of the weighted regression estimates the difference in instantaneous total mortality rates, $Z_0 - Z_x$, which is equivalent to the difference in instantaneous natural mortality rates (because the F component of Z is assumed to be the same for both groups of fish). A) Condition 3 versus condition 0. B) Condition 2 versus condition 0. C) Condition 1 versus condition 0. D) Conditions 2 and 3 combined versus condition 0.

Figure 16a.




Figure 16b.




Figure 16c.





Figure 16d.



**Appendix A. Daily flow rates of the Rappahannock River,
30 March – 3 May, 1985-2010.**

Striped Bass Assessment and Monitoring Program
Department of Fisheries Science
School of Marine Science
Virginia Institute of Marine Science
The College of William and Mary
Gloucester Point, VA. 23062-1346

Figure 1. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, spring 2010.




Figure 2. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2008-2009.





Figure 3. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2006-2007.





Figure 4. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2004-2005.





Figure 5. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2002-2003.





Figure 6. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 2000-2001.





Figure 7. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1998-1999.





Figure 8. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1996-1997.





Figure 9. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1994-1995.





Figure 10. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1992-1993.





Figure 11. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1990-1991.





Figure 12. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1988-1989





Figure 13. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, springs 1986-1987.





Figure 14. Daily and historic mean river flows (cf/s) for the Rappahannock River during the spawning stock assessment period, spring 1985.

