

**Summary of Colonial Nesting Herons Within
the Colonial National Historic Park Boundaries,
2006 Breeding Season**



**The Center for Conservation Biology
The College of William and Mary**

August 2006

Summary of Colonial Nesting Herons Within the Colonial National Historic Park Boundaries, 2006 Breeding Season

Center for Conservation Biology
College of William and Mary
Williamsburg, VA 23187-8795

Cover photo of Great Blue Heron by Josh Nemeth



This paper is funded by grants from the Colonial National Historic Park. The views expressed herein are those of the authors and do not necessarily reflect the views of the Colonial National Historic Park.

Summary of Colonial Nesting Herons Within the Colonial National Historic Park Boundaries, 2006 Breeding Season

2006

**Fletcher M. Smith
Center for Conservation Biology
College of William and Mary
Williamsburg, VA 23187-8795**

Recommended Citation:

Smith, F.M. 2006. Summary of Colonial Nesting Herons Within the Colonial National Historic Park Boundaries, 2006 Breeding Season. Technical Report Series, CCBTR-06-10. Center for Conservation Biology, College of William and Mary, Williamsburg, VA. 13 pp.

Project Funded By:

Colonial National Historic Park



The Center for Conservation Biology is an organization dedicated to discovering innovative solutions to environmental problems that are both scientifically sound and practical within today's social context. Our philosophy has been to use a general systems approach to locate critical information needs and to plot a deliberate course of action to reach what we believe are essential information endpoints.

TABLE OF CONTENTS

Heron Colony Nest Counts Overview	1
Survey Methods	1
Habitat Descriptions and Colony Summaries	1
Jamestown Colony Overview.....	1
Yorktown Colony Overview.....	3
Swanns Point Colony Overview.....	4
College Creek Colony Overview.....	5
Ringfield Colony Overview.....	6
ACKNOWLEDGMENTS	7
APPENDIX I	8
APPENDIX II	8
APPENDIX III	9

HERON COLONY NEST COUNTS OVERVIEW

Year	Yorktown	Jamestown	Swanns Point	Ringfield	College Creek
1993	375/70*	160	90	0	0
1997	355/117*	220	92	0	0
2003	196/25*	225	135	14	22
2005	126/9*	110	55	21	9
2006	46/1*	20/1*	56/1*	13	11

Table 1. Overview of colony counts in recent years.

**Second number reflects the number of Great Egret nests.*

SURVEY METHODS

Ground surveys were used to count the number of nests in each colony. Surveys were conducted in mid-June. Trees were marked with flagging to avoid double counting of nest trees. The colony boundaries were mapped using aerial imagery.

Data fields recorded at each site include numbers of occupied trees, numbers of nests per tree (See Appendix II), status of each nest tree (e.g. living, dead), species of nest tree, and activity status of each nest if it could be determined (See Appendix III). General habitat characteristics were recorded at each colony site.

Activity status for each nest was described according to the following categories:

1. Adult incubating.
2. Presence of adult in a shading or a brooding posture, or the presence of small young seen from below. Presence of fresh egg shells under the nest was also used to determine nest status.
3. Presence of large young.
4. Presence of dead young, either in the nest or on the ground underneath the nest.
5. Partial nest. Either under construction or abandoned.
6. Adult present but not incubating or brooding.
7. Unknown status. Nest present but no sign of eggs, young, or adults visible from ground.

HABITAT DESCRIPTIONS AND COLONY SUMMARIES

Jamestown Colony: The Jamestown Island heronry contains some of the largest loblolly pines of any of the sites. This site contains many trees in the 35-45 inch diameter range. The colonies are spread between 4 different ridges on the southern end of Jamestown Island. The only ridge occupied in 2006 is the largest and most central ridge (referred to as Ridge 1 in Figure 1). In contrast to the 2003 heron surveys, Ridge 1 has much more green-

brier and a thick wax myrtle shrub layer. Many of the large canopy trees on the ridge died during Hurricane Isabel in the fall of that year. The ridge due west from Ridge 1 has in the past contained heron nests (which we will refer to as Ridge 2). The Ridge 2 habitat is much denser than Ridge 1, with far more greenbrier and wax myrtle than Ridge 1. Ridges 3, 4, and 5 are smaller than Ridges 1 and 2 in both length and width and have a very dense understory layer. The Jamestown colony declined substantially during the past few breeding seasons, from 225 pairs in 2003, to 21 in 2006. This is due to a tendency of Great Blue Herons in recent years to nest in smaller, more isolated colonies (Watts, pers. com.). In 2006, the birds occupied 1 of 4 ridges occupied in 2003 and 1 of 3 ridges occupied in 2005.



Figure 1. Location of the Great Blue Heron colony on Jamestown Island. Colony boundaries are in white. Ridges 2, 3, 4, and 5 formerly had breeding herons but in 2006 had no nesting herons.

Yorktown Colony: The Yorktown heronry continues to decline over time, from a high of 375 pairs in 1993, to the 46 pairs that occupy the colony now.. All nests within the colony are in mature living pines. The colony is also bordered on three sides by mature living pines, suggesting that lack of nesting substrate is not the cause of the decline. The colony is bordered to the west by Beaverdam Creek (See Figure 2 for the colony boundaries map). The Yorktown site has an open understory.

Colony density in 2006 was down and the majority of nesting trees had fewer than 3 nests. No young heron carcasses were found below the nests, and few depredated eggs were found. This suggests that productivity was good at the Yorktown Colony in 2006.




Figure 2. Boundary of the Yorktown heronry. Colony boundaries are in white.

Swanns Point Colony: The Swanns point colony held steady between 2005 (55 pairs detected) and 2006 (56 pairs detected) but has declined substantially since the 2003 survey (135 pairs detected). The main colony is still largely centered in the mature pines of Black Duck Gut, but has shifted to the east and encompasses some of the mature bald cypress trees as well. The understory is open and very park like in the pine section of the colony, and quite swampy with an open understory in the section composed of cypress trees. A few nests were found to the northeast of the main colony and directly east of the main colony (See Figure 3 for a map of colony boundaries).



Figure 3. Location of colonies on Swanns Point. Colony boundaries are in white.

College Creek Colony: The College Creek heronry has declined from a high of 22 pairs detected in 2003 to 11 pairs detected in 2006. The colony was first discovered during the 2000 survey. The colony remains spread across the tips of two shoreline points separated by a small tidal gut (See Figure 4 for a map of colony boundaries). The site is very near the Colonial National Historic Parkway, and one nest this year is in a loblolly adjacent to the CNHP.



Figure 4. Location of College Creek heronry. Colony boundaries are in white.

Ringfield Colony: This colony was first discovered in 2003. This colony saw a decrease in the number of nesting pairs from 21 pairs in 2005 to 13 pairs in 2006. The colony is spread over the tips of two shoreline points separated by a small tidal gut (See Figure 5 for a map of colony boundaries). The colony trees are all healthy.




Figure 5. Location of Ringfield colony.

ACKNOWLEDGMENTS

We would like to thank the employees of the Colonial National Historic Park for their continued support. This project would not have been possible without the efforts of many people. Lydia Whitaker, Carlton Adams, Renee Peace, Anne Womack, Gloria Sciole, Mark Roberts, and Cheryl Pope provided important administrative support from the College of William and Mary. This study was funded through a grant from the Colonial National Historic Park and the Center for Conservation Biology at the College of William and Mary.

Appendix I. Overview of the current years survey results and of historic data. Data presented for Colonial National Historic Park properties along the lower James and York Rivers. Numbers represent nest counts, and by inference nesting pairs.

Year	Jamestown	Swanns Point	Yorktown	College Creek	Ringfield
1982	-	-	355	-	-
1983	-	-	368/5*	-	-
1984	19	-	335/8*	-	-
1985	21	-	350/6*	-	-
1986	25	-	390/11*	-	-
1987	14	16	415/6*	-	-
1988	13	13	470/15*	-	-
1993	160	90	375/70*	-	-
1997	220	92	355/117*	-	-
2003	225	135	196/25*	22	14
2005	110	55	126/10*	9	21
2006	19/1*	55/1*	45/1*	11	13

**All numbers represent the number of Great Blue Herons followed by the number of Great Egrets.*

Appendix II. The following is a table depicting the number of nests per tree within each site. Under each column is the number of nest trees containing that number of nests.

Site	1	2	3	4	5	6
Jamestown						
Great Blue Heron	16	-	1	-	-	-
Great Egret	1	-	-	-	-	-
Swanns Point						
Great Blue Heron	13	5	3	3	1	1
Great Egret	1	-	-	-	-	-
Yorktown						
Great Blue Heron	11	3	5	2	1	-
Great Egret	1	-	-	-	-	-
College Creek	5	3	-	-	-	-
Ringfield	10	-	1	-	-	-

Appendix III. The following table provides an overview of the activity status recorded at each nest. The numbers in each column are the percentages of each activity status recorded at each site.

Site	Adult Incubating	Adult Brooding	Adult Present	Large Young	Dead Young	Partial Nest	Unknown Status
Jamestown	-	-	65%	-	-	-	35%
Swanns Point	-	-	18%	27%	2%	-	53%
Yorktown	-	-	4%	-	7%	2%	87%
Ringfield	-	-	8%	23%	-	-	69%
College Creek	-	-	27%	9%	-	-	64%