

Achievements of Soviet Scientists in Investigations of the Helminthofauna of Marine Mammals of the World Ocean^{1,2}

S. L. DELAMURE and A. S. SKRIABIN³

ABSTRACT

During the last 40 yr (1940 to 1980), Soviet helminthologists have examined about 9,680 specimens of marine mammals in the Northern and Southern Hemispheres, using uniform or standard methods in most cases. As a result, 20 new species of trematodes [digeneids], 19 cestodes, 9 acanthocephalans, and 23 nematodes have been described. Also several new supraspecific taxa have been established. Systematic revisions of Campulidae, Tetrabothriidae, Pseudallidae, and Anisakidae have been undertaken.

The morphology, adaptation of helminths to their environments and to the host's mode of life, infection rates and characteristics, and the seasonal and age dynamics of the helminthofauna have been studied. Methods of distinguishing host populations using quantitative and qualitative differences as well as the geographic variability of the helminthofauna have been developed. Consequently local stocks of mammals have been revealed, as has the pathogenicity of several helminth species.

All of these data have been summarized from ecological, zoogeographical, and phylogenetic points of view. Results of these studies are now employed by zoologists interested in sea mammals.

In 1780 Fabricius published a scientific description of the cestode *Pyramicocephalus phocarum* (Fabricius, 1780), the first of the helminth species to become known to science as a parasite of marine mammals. Some time later Karl Rudolphi (Rudolphi 1802 to 1819) described another eight species (trematodes [digeneids], cestodes, and acanthocephalans) parasitizing these animals. Without becoming too immersed in the history of the development of knowledge of the parasites of marine mammals (for details see Delamure 1955), we wish to indicate that over the last 200 yr (1780-1980) some 276 species of helminths infecting marine mammals of the World Ocean have been discovered by Soviet and foreign scientists.

Today the parasites of these animals are of interest, not only to parasitologists but also to mammalogists studying marine mammals. They are attracted by the possibilities of using helminthological data for the purposes of studying the ecology, phylogeny, dynamical statistics [population dynamics?], and the population structure of the [various] species of marine mammals, including those of commercial importance.

Helminths of marine mammals are not without significance to medical and veterinary specialists, especially since it is known that there are pathogenic species common to pinnipeds, cetaceans, wild and domestic terrestrial mammals, and also to man: *Opisthorchis felinus* (see Kurochkin 1960); *Diphyllobothrium cordatum* (see Yamaguti 1959); *D. pacificus* (see Atias and Catton 1976); *Pyramicocephalus phocarum* (see Rausch et al. 1967); *Taenia solium* larvae (see Graff et al. 1980); *Trichinella spiralis* (see

Rausch et al. 1956; Fay 1960; Smirnov 1963; Kozlov 1966, 1971; Treshev and Serdukov 1968; Arvy 1970); larval Anisakidae (Kurochkin and Mamaev 1972; Shiraki 1974; and others).

The interest which has developed in studying helminths of marine mammals will be illustrated by examining the dynamics of recent publications by 10-yr periods.

Before 1940 Soviet scientists had published eight works in this one area. Between 1941 and 1950 they published 15, from 1951 to 1960—31, from 1961 to 1970—91, and from 1971 to 1980—143 articles.

According to the summary of Scheffer and Rice (1963), the World Ocean is inhabited by 31 species of pinnipeds and 80 species of cetaceans. Not all of them have been studied in helminthological respects. Better known than others are the helminthofaunas of the walrus and long-eared seals of the Northern Hemisphere, the true seals and major commercial whales of temperate and cold waters of the Northern and Southern Hemispheres, and, likewise, certain species of small whales. This is due, in no small measure, to Soviet scientists who for the first time investigated the helminthofauna of the gray whale (Treshev 1966a); Bryde's whale (Skriabin 1970a); harbor seals inhabiting the Bering Sea (Yurakhno 1970); island seals (Nikolsky and Kazikanov 1970); small seals of the Baltic Sea and Lake Ladoga (Barysheva and Bauer 1957; Delamure et al. 1980); Caspian seals (Schupakov 1936; Kurochkin 1958; and others); and the Baikal seal (Sudarikov and Ryzhikov 1951; Delamure et al. 1979a; and others).

In this connection it should be indicated that in the last 40 yr Soviet helminthologists have examined 9,680 marine mammals from various zones of the World Ocean (pinnipeds—6,222 and cetaceans—3,458). The large majority of those examinations (8,432) were carried out using the same accepted methodology (Delamure and Skriabin 1965b), by colleagues of the chairman of Zoology of the Crimean Pedagogical Institute (since 1972, the Simferopol State University), by participants in the 33 expeditions to the Black, Azov, Caspian, Baltic, Greenland, Barents, White, Eastern Siberian, Chukchi, Bering, and Okhotsk Seas;



¹Virginia Institute of Marine Science Translation Series No. 30.

²The text and Literature Cited sections of this paper were translated from the Russian by W. J. Hargis, Jr., with assistance of fellow students in the 1982 Russian scientific reading course of the School of Marine Science of the College of William and Mary. Despite the assistance of colleagues, the translator-editor must assume full responsibility for its accuracy and faithfulness of style. The abstract portion of the paper was translated by O. N. Bauer of the Zoological Institute of the Academy of Sciences of the U.S.S.R.

³The M. V. Frunze Simferopol State University, Crimea, U.S.S.R.

Lakes Ladoga and Baikal; and expanses of the Southern Hemisphere—including the waters of the Antarctic. Regarding our waters, we have not as yet been able to investigate the marine mammals of the Karskogo Sea and the Sea of Laptevsk.

Turning now to the most common results of the efforts of Soviet scientists in faunal areas, it is interesting to compare them with analogous investigations performed abroad. Calculation has shown that until 1900 only 16 species of trematodes, parasitic in pinnipeds and cetaceans, were known, but now there are 72. During the last 40 yr 27 species have been described (principally of the families Campulidae, Notocotylidae, and Microphallidae). Of that number, 20 or 74% have been by Russian scientists (Skriabin 1944, 1947; Krotov and Delamure 1952; Gubanov V. Kh.: [in:] Delamure 1955;⁴ Delamure and Kleinenberg 1958; Kurochkin 1958, 1962; Skriabin 1959, 1969a, 1970a; Petrov 1963; Treshev 1966a, 1968; Yurakhno 1967, 1968, 1969; Treshev et al. 1969; Kurochkin and Nikolsky 1972. New taxa have been established—the genera *Oschmarinella* Skriabin 1947; *Leucasiella* Krotov and Delamure 1952; *Cetitrema* Skriabin 1969a; and the family Campuloidea (A. Skriabin 1976).

Significant progress has been made in the studies of the cestodes parasitizing marine mammals. Known at present are 71 species of cestodes (versus only 20 in 1900). These discoveries include interesting tetrabothriid species (of the genera *Tetrabothrius*, *Priapocephalus*, and *Trigonocotyle*) and diphyllobothriids (of the genera *Diphyllobothrium*, *Diplogonoporus*, *Tetragonoporus*, *Hexagonoporus*, and *Polygonoporus*). Of the 32 cestode species described in the last 40 yr, some 19 (or 59%) have been by Soviet scientists (Gubanov in Delamure 1955; Delamure 1955, 1971; Belopolskaya 1960; Skriabin 1961, 1967; Delamure et al. 1964; Delamure and Skriabin 1966; Delamure and Parukhin 1968; Muraeva and Treshev 1970; Yurakhno 1973). New genera have been established (*Tetragonoporus* Skriabin, 1961; *Hexagonoporus* Gubanov in Delamure 1955; *Polygonoporus* Gubanov in Delamure 1955; and *Polygonoporus* A. Skriabin 1968); and a subfamily, *Polygonoporinae* Delamure and Skriabin 1968. Occupying a special place is the discovery by A. Skriabin of a 30 m long diphyllobothriid, *Polygonoporus giganticus* A. Skriabin, 1967, parasitizing the cachalot, with strobilae about 5 cm wide, having some members [strobilae] with 14 gonads in the sexual complex.⁵ Diplo- and polygonadal diphyllobothriids have been the subject of special investigations which attempt to clarify their systematics and phylogeny (Delamure and Skriabin 1966, 1968). There have been studies of larval forms or stages of cestodes invading cetaceans (Delamure 1955; Skriabin 1965, 1970a, 1971).

Research has added significant information about acanthocephalans. At present, 35 species parasitizing pinnipeds and cetaceans are known (versus 8 in 1900). Since 1940, 19 species have been described (of the genera *Bolbosoma* and *Corynosoma*). Of these, nine have been described by Soviet scientists (Krotov and Delamure 1952; Gubanov in Delamure 1955; Skriabin 1959; Treshev 1966a, 1970; Nikolsky 1974a).

By 1900, 23 species of nematodes were known from marine mammals; at present there are 98. In the last 40 yr, 46 species have been discovered (of the families Pseudaliidae, Folaroididae, Crassicaudidae, Anisakidae, and Capillariidae), of which 23 (50%) were described by Russian scientists (Skriabin 1933, 1942; Delamure in Skriabin 1942; Delamure 1946, 1951a; Gubanov

1951; Mozgovoy 1951; Gubanov in Delamure 1955; Kurochkin and Zablotski 1958; Skriabin 1959, 1966, 1969a; Delamure and Alekseev 1966; Delamure and Kleinenberg 1968; Zablotski 1971; Yurakhno and Skriabin 1971; Nikolski 1974a). The new genera, *Otophocaenurus* Skriabin, 1942; *Skrjbinalius* Delamure in Skriabin, 1942, *Placentonema* Gubanov, 1951; and *Delamurella* Gubanov in Delamure 1955, were established. The discovery of the 9 m long nematode (*Placentotrema gigantissima* Gubanov, 1951), which parasitizes the placenta of cachalots by H. Gubanov deserves special attention.

At present 276 species of helminths parasitizing pinnipeds and cetaceans of the World Ocean are known. In addition to the Soviet scientists contributing significantly to knowledge of the helminth fauna during the last decade, one must acknowledge investigators from other countries, especially Baylis, Baer, Davey, Dailey, Dollfus, Dougherty, Margolis, Markowski, Rausch, Stunkard, Shultz, and Yamaguti.

The study of helminths of marine mammals has not been limited to discovery of new species, to redescription of old ones, to creation and revision of higher taxa, to studies of morphology and systematics, to development of better characteristics to clarify synonymies and permit more exact definitions, or to the discovery of principal hosts. In recent times there has been a change not only toward consideration of the morphological variables of the helminths of marine animals but also toward comparison of the factors of development, host-parasite relationships, zoogeography, and variables of routes and intensity of infection. Manifestations of the variability of different, separate indicators and also the causes of that variability have important meanings for the systematics of helminths and for the study of the population structure of their hosts—pinnipeds and cetaceans.

Study of the accumulated materials permitted revision of the taxonomic characteristics of marine mammals according to the taxa of helminths such as ampulids, tetrabothriids, pseudaliids, and anasakids (Skriabin 1942; A. Skriabin 1970a; Kontrimavichus et al. 1976; Temirova and Skriabin 1978; and others).

Helminths are not second in any respect to other animals in the number of their adaptations toward their environments and the life histories of their different hosts; e.g., those of the pseudaliid infecting the respiratory organs of dolphins (Delamure 1951a, 1955), as well as those of the crassicaudid nematode which infects the genito-urinary systems of whales (Skriabin 1969a). The origin of their adaptations is closely connected with those changes which the whales underwent in the course of their evolution.

The adaptations of gigantic cestodes and nematodes, which assure survival of the species over the wide expanses of the ocean, are very interesting (Delamure and Skriabin 1965a, 1968). Adaptation of the parasite to the anatomical and physiological peculiarities of the principal hosts together with the damage it causes have been studied by Skriabin (1960) using *Placentonema gigantissima* as a model or example. The importance of such investigations to the understanding of the host-parasite system is evident. Attempts have been made to study the life cycles of helminths of marine mammals. The studies of A. V. Uspenskaya (1960), who deciphered the life cycle of the nematode *Terranova decipiens*, and T. I. Popova, A. A. Mosgovoy, and M. A. Dimitrienko (1964), who devoted attention to the same question, are relevant. A. A. Mozgovoy, T. I. Popova, and V. I. Shakhmatova (1963) studied the life cycle of *Parafilaroides gymnurus* and B. E. Sudarikov and K. M. Rhyzhikov (1951) attempted to comprehend the cycle of *Contracaecum osculatum baicalensis* under the conditions found in Lake Baikal. These investigations supplement the findings of H. M. Pronin et al. (1980). Incidentally, comparison of

⁴V. Kh. seems to translate best as "in the work" or "in the work of," which can be reduced to "in" as it has been throughout this translation.

⁵This phrase concerning the sexual complex was particularly difficult to translate. It is hoped that it is correct, but may not be.

the helminth fauna of the Baikal seal, *Pusa siberica*, with the helminth fauna of other seals disclose their similarity to the species of the genus *Phoca* inhabiting the Asiatic sector of the northern Pacific and the Arctic and in no way supports their similarity to the helminth fauna of the Caspian seal, *Pusa caspica* (Mozgovoy and Ryzhikov 1950; Delamure et al. 1978). Comparative studies of the helminth fauna of the Caspian seals over an extensive period of time confirm the thesis of V. A. Dogiel (1947) regarding the relict parasite fauna having adapted its patterns to those of the host (Delamure et al. 1964, and others).

Massive examination of pinnipeds conducted over the last few years allows us to approach, for the first time, study of the intensity and extensiveness [prevalence] of invasion or infestation, seasonality, and growth dynamics of the helminthofauna of commercial marine animals inhabiting the waters of the Soviet Union and adjacent waters. Walruses, polar bears, sea lions, greenland seals, ringed seals, common seals (harbor seals), island seals, ribbon seals, bearded seals, and hooded seals were investigated from these points of view (Delamure and Treshev 1966; Treshev and Popov 1970, 1975; Delamure and Popov 1975; Kovalenko 1975; Popov 1975; Delamure et al. 1968, 1976).

It was established that the harbor seals inhabiting the Bering Sea begin to be infected by helminths at about 1 to 1.5 m in size when they cease feeding from their mothers and proceed to live and acquire nourishment independently. By the age of 2 mo the seal is usually attacked by two to four species of helminths. Later the infections increase very quickly and by the age of 4 [yr] the seals are struck [infected] by up to 12 species of helminths. In animals older than 12 yr of age a certain decline can be detected (9 to 10 species) [down to 9 to 10 species], which can possibly be explained by the changing ecology of the host. It is characteristic that in several species, infections become more intensive and also extensive [prevalent] as the seals grow but they are especially great or large in the young and also in the very old animals (Delamure and Alekseev 1962; Delamure et al. 1978; Treshev 1969; Yurakhno 1970b).

The generalized methodology for differentiating the populations of marine mammals, based upon the study of the qualitative and quantitative differences in their helminthofauna and also the geographical variability of the helminths (Delamure et al. 1979a; Popov and Marichev 1979) by comparison of the helminthofaunas and the characteristics of infestation of the different populations of marine mammals, demonstrated localized herds of walrus, Greenland seal, ringed seal, harbor seal, bearded seal, and the striped whale of the northern Pacific. It has also unearthed or disclosed the boundaries of the summer habitations [ranges, grounds, or habitats] of a certain population of whales of the Southern Hemisphere (Skriabin 1958, 1970a, 1972, 1975, 1978; Shustov 1965; Treshev 1970, 1978; Yurakhno 1970a; Kosygin and Popov 1972; Delamure and Popov 1978, 1980; Delamure et al. 1978, 1979a). These investigations confirm the opinions of mammalogists that existing stocks of pinnipeds and cetaceans must be protected. A comparison of the character [or nature] of infection with the findings (or data) of mammalogists on the mortality of seals permitted the assumption that helminths are one of the causes of their deaths.

Combining these data with histopathological investigations confirms the fact that pinnipeds whose hearts are stricken by the nematode *Dipetalonema spirocauda* perish during the first years [first year or early years] of life (Delamure et al. 1965; Delamure and Treshev 1966; Treshev et al. 1967; and others). Intensive infections of Baikal seal by the nematode *Contraecaecum osculatum baicalensis* lead to the formation of large ulcers and

fistulas in the walls of their stomachs and cause internal bleeding. Such a picture has been established from those animals which were killed during commercial harvesting and those which perished naturally and washed up on the shores of the lake by waves (Zhaltsanova 1980). Information on the damage which can be done to the health of sea animals is found in a number of other publications (Vsevolodov 1948; Chupakhina 1969; Zablotski 1971; Temirova and Usik 1971; Gurevich and Evdokimov 1972; Nikolski and Kandinski 1974; Popov 1975; Starostin 1975; Delamure et al. 1976; Kovalaeva et al. 1978; and others). The data which have been introduced unquestionably confirm that the helminths emerge as one of the primary factors in the mortality of marine mammals, a fact that must be taken into account when one is establishing or studying their numerical or population dynamics.

Having gathered and analyzed a large amount of material we naturally arrived at the necessity of forming generalizations about it from the viewpoints of ecology, zoogeography, and phylogeny (Delamure 1952, 1955, 1956; Delamure and Skriabin 1958; Skriabin 1958). The results established the rules [regularities] of the patterns of the helminth faunas of the pinnipeds and cetaceans and likewise the patterns of distribution of helminths in the animals of the World Ocean. This showed that among the species and genera of helminths there are some which are dispersed in bipolar, amphiboreal, and amphi-Pacific [fashion]. This was accomplished for the first time on the basis of helminthological material and it was also explained in light of the history of the free-living fauna of the World Ocean (Berg 1922, 1934; Andriashv 1939; Zenkovich 1947, 1951; and others).

Comparison between the helminthofauna and the large taxa of their hosts in light of their ecology and phylogeny is of great interest. Here it has been established that, though the helminthofauna of pinnipeds shows characteristics of independence, it was developed principally from ancestors derived from the helminthofauna of terrestrial animals. In contrast with that of pinnipeds, the helminthofauna of cetaceans possesses clearly expressed traits of independence; it includes such helminth families as Pholeteridae, Brauninidae, Crassicaudidae, Pseudaliidae, and Campulidae almost completely. Investigations of the composition of the helminthofauna clearly reflect fundamental differences in ecology and phylogeny of these unique animals (Delamure 1955).

Sharp differences were demonstrated also when comparing the helminthofauna of toothed and baleen whales. Because of the small number of species (11), narrow specialization and feeding only at the surface of the sea on the not very varied [on the relatively homogeneous] plankton organisms—sometimes on mollusks and sometimes on schooling fishes, and other peculiarities of baleen whales; they have developed in themselves a helminthofauna which has no marked originality. It is totally different with the toothed whales. The numerous species of these cetaceans (69 species), which have settled widely in the World Ocean—even in brackish and freshwaters, have won for themselves the most varied ecological niches and accustomed themselves to feeding at different depths on varied kinds of food. In the process of evolutionary development of the toothed whales, they had a much greater opportunity than the baleen whales to become specific hosts for numerous species and genera and even a few families of helminths. There are essential differences between the helminthofauna of toothed and baleen whales which result from the long process of ecological selection, and it is possible to use them to compare the phylogenetic differences between the sub-orders of whales and also to compare their relationships with other animals (Delamure 1970).

Comparing the helminths of dolphins and beaked whales with the helminths of the Viverridae family [civets] in the historical [evolutionary] aspects, allows us to conclude [postulate] that these groups of hosts are close in phylogenetic respects (Delamure 1960).

Helminthological workers have discovered many valuable and interesting answers in their investigations of native mammals, which have involved their comprehensive studies of the parasites and also the characteristics of pinnipeds and cetaceans indigenous to the waters of the U.S.S.R. and also [those from] beyond their borders (Kleinenberg 1956; Tomilin 1957; Kleinenberg et al. 1964; Berzin 1971; Yablokov et al. 1972; Fedoseev 1975; Geytner et al. 1976; and others). These publications utilize and also exemplify the scientific investigations of parasites of marine mammals. It is not possible to encompass the subject without their numerous works in this area.

We have presented the work of the last 10 decades by Soviet helminthologists before many All-Union scientific conferences as well as those of several republics and now before the Soviet-American conference, conducted within the framework of the agreement, in surroundings that provide immeasurable support and assistance.

From this accounting it is evident that studies of the helminths of pinnipeds and cetaceans are of notably practical importance. It is possible to report with confidence that study of the helminths of animals, especially those of industrial importance living in the waters of the Soviet Union, compare favorably with those of our friends, and that this is possible thanks to the support and assistance provided by the Laboratory of Helminthology of the U.S.S.R. Academy of Sciences, and the Ministry of Fisheries of the U.S.S.R., the Ichthyological Commission, VNIRO and the foundation related to it, the Zoological Institute of the U.S.S.R. Academy of Science, Vsesoyuznogo Institute of Helminthology, the Limnological Institute CO [SO] Academy of Sciences of U.S.S.R., the Astrakhan-Zapovednika, Ministry of Enlightenment [or Instruction] of the U.S.S.R., and the administration of the Simferopol University and the institutions of our associates.

LITERATURE CITED⁶

- ANDRIASHEV, A. P.
1939. On the amphipacific (Japan to Oregon) distribution of the marine fauna of the northern regions of the Pacific Ocean. [In Russ.] Zool. Zh. 2: 181-194.
- ARVY, L.
1970. Trichinosis in cetaceans. Invest. Cetacea. V. 10. Berne, p. 325-330.
- BARYSHEVA, A. F., and O. N. BAUER.
1957. Parasites of fishes of Lake Ladoga. [In Russ.] Izv. Vniorkh. 42:175-226.
- BELOPOLSKAYA, M. M.
1960. The helminth fauna of the harbor seal (*Phoca vitulina largha* Pall.). [In Russ.] Vestnik. Leningr. Gos. un-ta. Ser. Viol. 15(3):113-121.
- BERG, L. S.
1934. On the amphiboreal distribution of the marine fauna of the northern hemisphere. [In Russ.] Izv. Geogr. o.-va. 66(1):69-78.
1944. Climate and life. [In Russ.] Gos. izd-vo Geograf. Literaturny, 356 p.
- BERZIN, A. A.
1971. Cachalot. [In Russ.] Izd. "Pishcheyevaya promyshlennosti M", 367 p.
- CHUPAKHINA, T. I.
1969. Helminth fauna of fur seals and harbor seals. [In Russ.] V. sb. chyetyer'toye Vses. sovyeshchaniye po izuch. morskikh mylepityayushchikh. Tyez. Dokl. M., p. 166-168.
- DELAMURE, S. L.
1946. Three species of single host (?) nematodes of dolphins of the Black and Azov Seas. [In Russ.] Helminthol. Sborn. Posvyashch. Akad. K. I. Skriabin Izd. A. N. SSSR, p. 104-114.
1950. Phylogenetic affinities of dolphins and terrestrial predators (Mustelidae) in the light of helminthological science. [In Russ.] Dokl. A. N. SSSR 73:237-239.
1951a. A new pseudaliid parasite of the lungs of the white-sided dolphin. [In Russ.] Tr. Gel'mintol. Lab. A. N. SSSR 5:93-97.
1951b. Origin and adaptation of the pseudaliid parasitic in the respiratory organs of whales. [In Russ.] Tr. Gel'mintol. Lab. A. N. SSSR 5:98-104.
1952. Zoogeographical characteristics of the helminth fauna of pinnipeds and whales. [In Russ.] Tr. Gel'mintol. Lab. A. N. SSSR 6:235-250.
1955. The helminth fauna of marine mammals of the world, their ecology and phylogeny. [In Russ.] Izd. A. N. SSSR, 517 p.
1956. On the amphiboreal and bipolar distribution of the helminths of marine mammals. [In Russ.] Dokl. A. N. SSSR 107:621-623.
1970. The helminth fauna of toothed and baleen whales. [In Russ.] Voprosy Morskoy Parazitol. Kiev. Naukova dumka, p. 18-21.
1971. *Diphyllobothrium ponticum* n. sp. a new diphyllbothriid of the bottlenose dolphin (*Tursiops truncatus* Montagu). [In Russ.] V. sb. Raboty po Gel'mintol. Posvashch. 90-lyetiyu Akad. K. I. Skriabin Izd. "Kolos", M., p. 123-125.
- DELAMURE, S. L., and E. V. ALEKSEEV.
1963. Survey of the helminth fauna of White Sea herds of Greenland seal. [In Russ.] V. sb. Nauchno-issl. Rabot Syev. PINRO, Arkangelysk, p. 73-100.
1966. *Parafilaroides arcticus* n. sp., parasite of ringed seals of the Chukchi Sea. [In Russ.] V. sb. Parazity, promezhutchnye khozyayeva i pyeryenoschiki. Kiev. Naukova dumka, p. 11-15.
- DELAMURE, S. L., and S. E. KLEINENBERG.
1958. New facts on the helminth fauna of the beluga. [In Russ.] Byullyeteny Mosk. ob-va Isp. Prirody. Otd Biol. 63(3):25-32.
- DELAMURE, S. L., Yu. V. KUROCHKIN, and A. S. SKRIABIN.
1964. On the helminth fauna of the Caspian Seal (*Phoca caspica* Gm.). [In Russ.] Tr. Astrakansk. Zapovednika. 9:105-118.
- DELAMURE, S. L., E. C. MIKHALEV, V. N. POPOV, and V. D. PASTUKHOV.
1978. Information on the helminth fauna of the Baikal seal. [In Russ.] Morskoye mlyekopityayushchiye. Tyez. Dokl. 7-go vseye Sovyeshchonia, M., p. 103-104.
- DELAMURE, S. L., and A. M. PARUKHIN.
1968. A new diphyllbothriid parasite of the southern ocean fur seal. [In Russ.] V. sb. Biologiya Morya, 14, Parazity Morskikh Fivotnykh. Kiev. Izd. A. N. USSR, p. 25-33.
- DELAMURE, S. L., and V. N. POPOV.
1978. Helminthological facts on local seal herds living in the Okhotsk Sea. [In Russ.] Tyez. Dokl. I-go sbyeza Parazitotsyenogov. Ch. 3, Kiev, Naukova dumka, p. 30-32.
1980. Helminthological facts on local populations of harbor seals of the Okhotsk Sea. [In Russ.] Dyevyataya Konf. Ukrainskogo Parazitol. ob-va. Kiev, Naukova dumka, p. 29-30.
- DELAMURE, S. L., V. N. POPOV, and A. N. TRASHCHYENKOV.
1980. On the knowledge of the helminth fauna of seals of the Baltic Sea and Lake Ladoga. [In Russ.] Naukhn. Dokl. Vysshyye Shkoly Biol. 7:41-45.
- DELAMURE, S. L., and A. S. SKRIABIN.
1958. Principal laws governing the geographical distribution of the helminths of marine mammals. [In Russ.] Izv. Krymsk. otd. Geogr. ob-va SSSR, 5:244-250.
1965a. Origin and morphological peculiarities of the polygonadic diphyllbothriids. Tyez. Dokl. ryepublikanskoy Naukhn. Konf. po problemye "Biologichyekiye osnovy rationallynogo ispolyzovaiya i okhrany rastyetely nogo i zhivotnogo mira." [In Russ.] Simfyeropoly, p. 210-213.
1965b. On the methods of making helminthological dissections of marine mammals. (Peculiarities of collecting helminthological materials from pinnipeds and cetaceans). [In Russ.] V. sb. Morskoye mlyekopityayushchiye. M. Nauka, p. 302-310.
1966. A new diphyllbothriid — *Diphyllobothrium pierocephalum* sp. nova. parasitic in the hooded seal — *Cystophora cristata*. Helminthologia T. 7. [In Russ.] Academia Scientiarum Slovaca, Bratislava, p. 65-70.
1968. Origin and systematic position of diplogonadal and polygonadal diphyllbothriids. [In Russ.] V. kh. Gyelyminty Chyelyvyecka Zhivotnykh Rastyenyi i Myery Boryby s Nimi Izd. Nauka, p. 159-166.
- DELAMURE, S. L., and V. V. TRESHEV.
1966. On the helminth fauna of the hooded seal (*Cystophora cristata* Erleben), inhabiting the seas of Greenland. [In Russ.] Zool. Zh. 45:1867-1871.
- DELAMURE, S. L., M. V. YURAKHNO, and V. N. POPOV.
1976. On the helminth fauna of Bering Sea pinnipeds in Peter-the-Great Bay. [In Russ.] Zh. Parazitol. Nauka 10(4):325-332.

⁶Not all citations mentioned in the text are included here. Also, not all of the Literature Cited are mentioned in the text.

- 1979a. On perspectives obtained from morphological investigations of the helminths of marine mammals toward knowledge of the internal structure of their hosts. [In Russ.] V. sb. Sostoyaniye i pyerspektivy razvitiya morfologii. Matyerialy k vses. sobyeshchanyu. M. Nauka, p. 154-155.
- 1979b. Composition of helminths as indicators of seal populations. [In Russ.] Zh. Priroda 5:111-112.
- DELAMURE, S. L., D. D. ZAVALEEVA, and G. A. FEDOSEEV.
1965. Materials on the fauna of parasitic worms of the ringed seal (*Pusa hispida krascheninikovi*). [In Russ.] Tyez. Dokl. Itogovoy Nauk. Konf. Professororoko - Pripodvatyatsyelskogo Sostava Krymsk, Goc. Ped. in-ta Simfyeropoly, p. 251-253.
- DOGIEL, V. A.
1947. A course in general parasitology. [In Russ.] Uchpyedgiz, 371 p.
- FABRICIUS, O. F.
1780. Fauna Groenlandica, Systematice sistens Animalia Groenlandicae Occidentalis Hactenus Indagata, etc. Hafnia and Lipsiae.
- FEDOSEEV, G. A.
1975. Structure and dynamics of pinniped populations as bases governing their numbers and availability. [In Russ.] V. sb. Morskiye mlyekopitayushchiye. Matyerialy 6-go Vses. sovyeshchaniya. Kiev, Naukova dumka 2:41-43.
- GEPTNER, B. G., K. K. CHAPSKY, B. A. ARSENEV, and B. E. SOKOLOV.
1976. Mammals of the Soviet Union. [In Russ.] T. Z. Izd. "Vysshaya shkola", 717 p.
- GRAFF, A. S., P. D. SHAUGHNESSY, R. M. McCULLY, and A. VESTER.
1980. Occurrence of *Taenia solium* in a cape fur seal (*Arctocephalus pusillus*). Onderstepoort J. Vet. Res. 47:119-120.
- GUBANOV, A. M.
1951. A gigantic nematode of the placenta of whales, *Placentonema gigantissima* n.g. et n. sp. [In Russ.] Dokl. A. N. SSSR 77:1123-1125.
- GUREVICH, V. S., and V. N. EVDOKIMOV.
1972. The importance of x-irradiation in the control of lung pneumonia of dolphins. [In Russ.] Tyez. Dokl. 5-go vses. sovyeshchaniya po izuchenyiyu morskikh mlyekopitayushchikh Ch. 2. Makachkala, p. 68-70.
- KLEINENBERG, S. E.
1956. Mammals of the Black and Azov Seas. (Opyt biologo - promyslovogo issledovaniya). [In Russ.] Izd. A. N. SSSR. 286 p.
- KLEINENBERG, S. E., A. V. YABLOKOV, V. M. BELKOVICH, and M. M. TARASEVICH.
1964. Belukha. [In Russ.] Opyt monograficheskogo isslyedoveniya vida. Izd. Nauka. M., 455 p.
- KOVALENKO, L. M.
1975. On the helminth fauna of long-eared seals inhabiting the Kurile Islands. [In Russ.] V. sb. Morskiye mlyekopitayushchiye. Ch. I. Matyerialy 6-go Vses. Sovyeshchaniya. Kiev, Naukova dumka, p. 137-140.
- KOZLOV, D. P.
1966. Trichinellids of walrus of the Soviet Arctic. [In Russ.] Matyerialy Konf. vsyes. ob-va gel'mintol. M., p. 131-133.
1971. Questions on the routes of infection of trichinellids of pinnipeds. V. sb. Voprosy biologii fiziologii i biokhimii gylmintov zhirotnykh i rastyenyi. Tr. Gel'mintol. Lab. A. N. SSSR 21:36-40.
- KOLYEVATOVA, A. I., L. A. SAFRONOVA, E. Y. SEREBRANNIKOV, and V. V. FORMIN.
1978. Uncinarial-caused fatalities in the fur seals of the Komandorsky Islands. [In Russ.] V. sb. morskiye mlyekopitayushchiye Tyez. Dokl. 7-go vsyes. Sovyeshchaniya. M., p. 160-162.
- KONTRIMAVICHUS, V. L., S. L. DELAMURE, and S. N. BOEV.
1976. Metastrongyloids living in wild animals. [In Russ.] Izd. Nauka. M., 237 p.
- KROTOV, A. I., and S. L. DELAMURE.
1952. Materials on the parasitic worm fauna of birds and mammals of the USSR. [In Russ.] Tr. Gel'mintol. Lab. A. N. SSSR 6:278-292.
- KOSYGIN, G. M., and V. N. POPOV.
1972. On the local herds of bearded seals. [In Russ.] Tyez. Dokl. 5-go vsyes. Sovyeshchaniya po izuchenyiyu morskikh mlyekopitayushchikh. Ch. I. Makhachkala, p. 38-40.
- KUROCHKIN, Yu. V.
1958. The helminth fauna of Caspian seals during their reproductive period. [In Russ.] V. sb. Raboty po gylmintologii K. 80-lyetiya Akad. K. I. Skriabin. Izd. A. N. SSSR, p. 188-194.
1960. On the Caspian source of opisthorchosis. [In Russ.] Tyez. Dokl. Matyerialy Konf. po vopr. zoogyelografii sushi. Izd. A. N. KazSSR, Alma-ata, p. 80-81.
1962. Helminth fauna of Caspian seals during autumn breeding. [In Russ.] Gel'mintol. sb. (matyerialy 315-oy STE). Tr. Astrakhansk Zapovyednika. 6:119-126.
- KUROCHKIN, Yu. V., and Yu. L. MAMAIEV.
1972. Conclusions and perspectives on knowledge of the parasites and diseases of fish of the Pacific Ocean. [In Russ.] V. sb. Parazity. M. Nauka.
- KUROCHKIN, Yu. V., and O. P. NIKOLSKY.
1972. On the trematode fauna of Antarctic seals. [In Russ.] Izv. Tikhookeanskiy NII ryb. Kh-va i Okyeaanogr. 81:266-269.
- KUROCHKIN, Yu. V., and V. I. ZABLOTSKI.
1958. On the helminth fauna of Caspian seals. [In Russ.] Tr. Astrakhansk. Zapovyednika 4:337-343.
- MOZGOVOY, A. A., T. I. POPOVA, and V. I. SHAKHMATOVA.
1963. On the biology of *Parafilaroides gymnuris* (Railliet, 1889). [In Russ.] V. Kn. Stroityelystro gylmintologicheskoy nauki i praktiki v. SSSR. T. Z., Izdat. A. N. SSSR (tsit po K. I. Skriabin s soavt., 1963).
- MOZGOVOY, A. A., and K. M. RYZHIKOV.
1950. Questions on the origin of Baikal seals in the light of helminthological science. [In Russ.] Dokl. A. N. SSSR 72:997-999.
- MURAEVA, S. I., and V. V. TRESHEV.
1970. A new cestode — *Priapocephalus eschrichtii* sp. n. (Cestoda:Tetrabothriidae) — parasite of the gray whale of the Chukotsk (Chukchi) Sea. [In Russ.] Zh. Vystnik Zoologii, Naukova dumka, I:84-86.
- NIKOLSKI, O. P.
- 1974a. The acanthocephalan fauna of pinnipeds of the Pacific Ocean section of the Antarctic Sea. [In Russ.] Izv. TINRO 88:101-106.
- 1974b. *Contracaecum mirounga* sp. n. (Nematoda: Anisakidae) — new nematode of the elephant seal of the Antarctic. [In Russ.] Izv. TINRO 88:107-109.
- NIKOLSKI, O. P., and P. A. KANDINSKI.
1974. Certain pathological modifications of the organism [organs?] of parasitized seals with their etiology. [In Russ.] V. sb. Isslyed. po biologii ryb i promysl, okyeanogr., p. 122-128.
- PETROV, A. M.
1963. A new trematode — *Zalophotrema lubimori* nov. sp. — of the livers of seal lions. [In Russ.] Tr. Vses. in-ta Gel'mintol. 10:26-29.
- POPOV, V. N.
1975. On the helminth fauna of Sakhalin populations of *Phoca vitulina largha*. [In Russ., Engl. summ.] Zool. Zh. 54:769-770.
- POPOV, V. N., and V. S. MARICHEV.
1979. Variability of the morphological characteristics of *Corynosoma strumosum* from the ringed seal from various chastyey areas. [In Russ.] V. sb. Sostoyaniye k vsyes. covyeshchaniyu razvitiya morfologii Matyerialy k vsyes. sovyeshchaniya. M. "Nauka", p. 368-369.
- POPOVA, T. I., A. A. MOZGOVOY, and M. G. DMITRIYENKO.
1964. On biological studies of ascarids of animals of the White Sea. [In Russ.] V. sb. Eksperymental'naya i ekologicheskaya gylminologiya. Tr. Gel'mintol. Lab. A. N. SSSR 14:163-169.
- PRONIN, N. M., D. ZHALTSANOVA, L. N. BRYKOVA, and A. P. GLADYSH.
1980. On the ecology of *Contracaecum osculatum baicalensis* — parasite of seals and fishes of Lake Baikal. [In Russ.] Gylmintozy chyelyovye, shivotnykh, rastyenyi i myery bory bory s nimi. Tzez Dokl. Vses. ob-va Gylmintol. M., p. 144-145.
- RAUSCH, R. L., B. B. BABERO, V. R. RAUSCH, and E. L. SCHILLER.
1956. Studies on the helminth fauna of Alaska. 27. The occurrence of larvae of *Trichinella spiralis* in Alaskan mammals. J. Parasitol. 42:259-271.
- RAUSCH, R. L., E. M. SCOTT, and V. R. RAUSCH.
1967. Helminths in Eskimos in Western Alaska, with particular reference to *Diphyllobothrium* infection and anemia. Trans. R. Soc. Trop. Med. Hyg. 61: 351-357.
- RUDOLPHI, K. A.
1819. Entozoorum synopsis cui accedunt mantissa duplex et indices locuplittissimi. Berolini, 811 p.
- SCHEFFER, V. B., and D. W. RICE.
1963. A list of the marine mammals of the world. U.S. Fish Wildl. Serv., Spec. Sci. Rep. Fish. 431, 12 p.
- SCHILLER, E. L.
1954. Studies on the helminth fauna of Alaska. 27. Notes on the intermediate stages of some helminth parasites of the sea otter. Biol. Bull. (Woods Hole) 106:107-121.
- SCHUPAKOV, I. G.
1936. Parasite fauna of the Caspian harbor seal. On questions of relict parasite faunas. [In Russ.] Uchenyye zap. Lyeningr. Un-ta. Syer. Biol. 7(3):134-143.
- SHIRAKI, T.
1974. Larval nematodes of family Anisakidae (Nematoda) in the northern Sea of Japan as a causative agent of eosinophilic phlegmone or granuloma in the human gastro-intestinal tract. Acta Med. Biol. 22:57-98.

- SHUSTOV, A. P.
1965. Certain peculiarities of the helminth fauna of ribbon seal of the Bering Sea. [In Russ.] *Izv. Tikhovkyansk. n.-i in-ta Rybn. kh-va i Okyengr.: Magadan*, 59:193.
- SKRIABIN, A. S.
1958. Zoogeographical characteristics of the helminth fauna of marine mammals of the boreo-Pacific subregion. [In Russ.] *Izv. Krymsk. Gos. Pyed. in-ta Simferopol*, p. 5-45.
1959. New species of helminths of marine mammals of the Pacific Ocean and the Far-eastern seas. [In Russ.] *Izv. Krymsk. Gos. Ped. in-ta Simferopol* 34:99-118.
1961. A new diphyllbothriid parasite of the cachalot — *Tetragonoporos calyptiocephalus* n. sp. *Helminth.* 3:311-315.
1965. Larval forms of helminths of marine mammals of far-eastern seas. [In Russ.] V. sb. *Morskiye mlyekopitayushchiye*. Nauka, M., p. 311-316.
1966. A new crassicauda (*Crassicauda delamureana* n. sp.) — parasite of the sea whale. [In Russ.] V. sb. *Krayevaya parazitologiya i prirodna ochagovosty transmissiia bolyexnyey*. Kiev. Naukova dumka, p. 100-107.
1967. A gigantic diphyllbothriid, *Polygonoporos giganticus* n.g., n. sp. — parasite of the cachalot. [In Russ.] *Zh. Parazitol. I(I):131-136*.
1969a. A new species of trematodes of the genus *Ogmogaster* (Notocotylidae) — parasite of whalebone whales. [In Russ.] *Zool. Zh.* 48:1882-1885.
1969b. A new nematode *Crassicauda costata* sp. n. — parasite of the right whale. [In Russ.] *Zh. Parazitol.* 3, 3, *Izd. Nauka*, p. 258-265.
1969c. Adaptations of the nematode genus, *Crassicauda* (Spirurida), for parasitizing the organs and tissues of cetaceans. [In Russ.] V. sb. *Problemy Parazitol.* Kiev, Nauka dumka, p. 227-229.
1970a. On the helminth fauna of Bryde's whale (*Balaenoptera edeni*) of the northwestern sector of the Indian Ocean. [In Russ.] V. sb. *Voprosy Morskoy Parazitol.* Kiev, Naukova dumka, p. 112-113.
1970b. A new trematode, *Cetitrema foliiformis* g.n., sp. n. — parasite of the liver of Bryde's whale. *Zh. Parasitol.* 4(I):34-38.
1970c. Original discoveries of the larval cestode, *Monogryma delphini* (Tetra- phylliidea: Phyllobothriidae) of Antarctic whales. [In Russ.] *Zool. Zh.* 49 (II):1728-1799.
1971. A contribution to the teratology of *Phyllobothrium delphini* larvae (Cestoda). [In Russ.] *Zool. Zh.* 50:582-584.
1972. On studies of the helminth fauna of Cachalots of the southern hemisphere. [In Russ.] *Tyez. Dokl. 5-go Vsesoyeshchaniya po Izuchenyu Morskikh Mlyekopitayushchikha*. Ch. 2, *Maknachkala*, p. 225-229.
1975. On studies of the helminth fauna of sei whales of the southern hemisphere. [In Russ.] V. sb. *Morskiye Mlyekopitayushchiye*. Ch. 2, *Materyialy 6-go Vsesoyeshchaniya*. Kiev, Naukova dumka, p. 79-81.
1978. Remarks on the systematics of the genus *Antisakis* Dujardin, 1845 (order Ascaridida). [In Russ.] V. sb. *Morskiye Mlyekopitayushchikha*. *Tyez. Dokl. 7-go Vsesoyeshchaniya*, M., p. 289-291.
- SKRIABIN, K. I.
1933. *Kutassicaulus* n.g., nuveau représentant des Nématodes de la sous-famille des *Dictyocaulinae* Skrzabin 1933 [*Kutassicaulus* n.g. new representative nematode of the sub-family Dictyocaulinae?]. *Ann. Parasitol. Hum. Comp.* 22:359-363.
1942. The ways of the phylogenetic evolution of nematodes of the family *Pseudaliidae* — parasitic of the auditory apparatus, circulatory system and respiratory organs of marine mammals. [In Russ.] *Dokl. A. N. SSSR* 37: 41-46.
1944. On the trematode genus *Odhneriella* Skriabin, 1915 of marine mammals. [In Russ.] *Dokl. A. N. SSSR* 41(7):331-332.
1947. *Oschmarinella sobolevi* n.g., n. sp. — new trematode of the liver of whales. [In Russ.] *Dokl. A. N. SSSR* 57:857-859.
- SMIRNOV, G. G.
1963. Nature and symptoms of trichinellosis of the Arctic. [In Russ.] *Zool. Zh.* 42(3):338-344.
- STAROSTIN, N. M.
1972. *Untsinariz* [studies?] of fur seals of the Bering Sea. [In Russ.] 8-y *Vsesoy. Konf. po prirodnoy ochagovosti bolyezney zhivotnykh i okhrange ikh chislennosti z, Kirov*.
- SUDARIKOV, V. E., and K. M. RYZHIKOV.
1951. On the biology of *Contracecum osculatum baicalensis*, nematode of Baikal seals. [In Russ.] *Tr. Gel'mintol. Lab. A. N. SSSR* 5:59-66.
- TEMIROVA, P. V., and V. D. USIK.
1971. Pathomorphological modification in the lungs of Azov and Black Sea seals (*Phocoena phocoena* L.) caused by the pseudaliid parasite, *Halocercus ponticus* Delamure, 1946. [In Russ.] V. sb. *Materyialy Nauchn. Konf. Vsesoy. ob-va Gel'mintol. M., A. N. SSSR* 22:256-260.
- TEMIROVA, S. I., and A. S. SKRIABIN.
1978. The suborder Tetrabothriata (Ariola, 1899) Skriabin, 1940. [In Russ.] V. kn. *Osnoby tsyestologii*. 9, M. Nauka, p. 1-117.
- TOMILIN, A. G.
1957. Animals of the SSSR and adjacent regions. Cetaceans. *Izd. A. N. SSSR*, M., 756 p.
- TRESHEV, V. V.
1966a. New species of the genus *Ogmogaster* Jägerskiöld, 1891 (*Ogmogaster delamurei* n. sp.). [In Russ.] V. sb. *Parazity promyehzhuotochnyye khozayeva i pyeryenoschiki*. Kiev, Naukova dumka, p. 22-25.
1966b. Species of *Skriabina* of whales of the Chkotskogo Sea. [In Russ.] *Ryepublikanskiy myezhvyedomstvyennyy sbornik*. *Syernya "Problemy parazitologiyii"*. Kiev, Naukova dumka, p. 112-115.
1968. A new campulid, *Orthosplanchnus albamarinus* sp. n. (Trematoda: Campulidae) — parasite of the beluga. [In Russ.] *Zool. Zh.* 47(6):937-940.
1969. On studies of the helminth fauna of White Sea herds of the Greenland seal (*Phagophoca groenlandica* Erleben, 1777). [In Russ.] V. sb. *Voprosy morskoy biologii, Tyez. 2-go Vses. cimpoziuma molodykh uchenykh*. Kiev, Naukova dumka, p. 129-130.
1970. On questions of localization of herds of the Greenland seal (*Phagophoca groenlandica* Erleben, 1777). [In Russ.] V. sb. *Voprosy morskoy parazitologii*. Kiev, Naukova dumka, p. 119-121.
1980. On studies of skriabinid parasites of marine mammals of the Arctic [In Russ.] *Dyevyataya Konf. Ukrainskogo parazitol ob-va*. Kiev, Naukova dumka, 74:96-97.
- TRESHEV, V. V., and V. N. POPOV.
1970. On the helminth fauna of the ringed seal of the northern part of the Sea of Okhotsk. [In Russ.] V. sb. *Voprosy morskoy parazitologii*. Kiev, Naukova dumka, p. 121-124.
1975. Helminths of the ringed seal of the south-eastern section of the Barents Sea. [In Russ.] V. sb. *Morskiye Mlyekopitayushchiye Ch. 2. Materyialy 6-go Vsesoyeshchaniya*. Kiev, Naukova dumka, p. 131-132.
- TRESHEV, V. V., V. A. POTELOV, and D. D. ZAVALEEVA.
1967. On knowledge of the helminth fauna of the ringed seal and the hooded seal. [In Russ.] *Tr. Polyarogo Nauchno-Isslyedov. Vi Protektnogo in-ta Morskogo Rybnogo-kh-va i Okyeanogr.* 21:194-197.
- TRESHEV, V. V., and A. M. SERDUKOV.
1968. On problems of trichinellids of the Arctic. [In Russ.] V. sb. *Biologiya morya*, 14. *Parazity morskikh zhivotnykh*. *Izd. A. N. SSSR*, p. 35-37.
- TRESHEV, V. V., M. V. YURAKHNO, and V. V. ZIMUSHKO.
1969. On the helminth fauna of the grey whale of the Bering and Chukchi Seas. [In Russ.] V. sb. *voprosy morskoy biologii, Tyez. 2-go. Vsesoyeshchaniya molodykh uchenykh*. Kiev, Naukova dumka, p. 130-131.
- USPENSKAYA, A. V.
1960. Parasitofaune des crustacés benthiques de la mer du Barents. [Parasite fauna of the benthic crustaceans of the Barents Sea]. *Ann. Parasitol. Hum. Comp.* 35:221-242.
- YABLOKOV, A. V., V. M. BELKOVICH, and V. M. BORISOV.
1972. Whales and dolphins. [In Russ.] *Izd. "Nauka"*, 472 p.
- YAMAGUTI, S.
1959. *Systema helminthum*. V. 2. The cestodes of vertebrates. Interscience, N.Y., 860 p.
- YURAKHNO, M. V.
1967. New species of the genus *orthosplanchnus* — *Orthosplanchnus pigmaeus* (Trematoda: Campulidae) — parasite of whales. [In Russ.] *Zh. Vyestnik Zool. Kiev, Naukova dumka*, 3:79-82.
1968. A new trematode, *Microphallus orientalis* sp. n. (Trematoda: Microphallida) — parasite of Pacific Ocean walrus and bearded seal. [In Russ.] *Zool. Zh.* 47(4):630-631.
1969. A new trematode, *Orthosplanchnus osculatus* sp. n. (Trematoda: Campulidae) — parasite of Pacific Ocean walrus. [In Russ.] *Zh. Vyestnik zoologii*. Kiev, Naukova dumka, 4:29-31.
1970a. New information on the helminth fauna of Bering Sea harbor seals. [In Russ.] V. sb. *Voprosy morskoy parazitol.* Kiev, Naukova dumka, p. 149-152.
1970b. Summary of certain knowledge of the helminth fauna of Bering Sea ribbon seal. [In Russ.] V. sb. *Voprosy morskoy parazitol.*, Kiev, Naukova dumka, p. 152-155.
- YURAKHNO, M. V., and A. S. SKRIABIN.
1971. The krascheninnikova parafilaroides (*Parafilaroides krascheninnikove*) sp. n. — parasite of the ringed seal (*Pusa hispida krascheninnikova* Naumov et Smirnov). [In Russ.] *Zh. Vyestnik Zool. Kiev, Naukova dumka* I:32-36.
- ZABLOTSKI, B. I.
1971. *Capallaria delamurei* sp. n. (Nematoda: Trichocephalata) of Caspian Sea seals. *Zh. Parazitol.* 5(3):251-253.

ZENKEVICH, L. A.

1947. Fauna and biological productivity of the sea. Izd. "Sovetskaya Nauka", T. 2., 588 p.

1951. Fauna and biological productivity of the sea. Izd. "Sovetskaya Nauka", T. 1., 505 p.

57
11
A44672
No. 25

NOAA Technical Report NMFS 25



Parasitology and Pathology of Marine Organisms of the World Ocean

William J. Hargis, Jr. (Editor)

March 1985

U.S. DEPARTMENT OF COMMERCE
Malcolm Baldrige, Secretary
National Oceanic and Atmospheric Administration
John V. Byrne, Administrator
National Marine Fisheries Service
William G. Gordon, Assistant Administrator for Fisheries

